



POLITECNICO DI TORINO  
Repository ISTITUZIONALE

A multiscale approach to study the stability of long waves in near-parallel flows

*Original*

A multiscale approach to study the stability of long waves in near-parallel flows / S. Scarsoglio; D. Tordella; W. O. Criminale. - (2007). ((Intervento presentato al convegno 60th Annual Meeting Division of Fluid Dynamics (APS-DFD) tenutosi a Salt Lake City, Utah nel November 18-20, 2007.

*Availability:*

This version is available at: 11583/1649117 since:

*Publisher:*

*Published*

DOI:

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

Abstract Submitted  
for the DFD07 Meeting of  
The American Physical Society

Sorting Category: 22. (T)

**A multiscale approach to study the stability of long waves in near-parallel flows** STEFANIA SCARSOGLIO, DANIELA TORDELLA, Politecnico di Torino, WILLIAM CRIMINALE, University of Washington — The linear stability of a two-dimensional non-parallel flow is considered as an initial-value problem. A spatio-temporal multiscale approach is assumed. The choice of the polar wavenumber ( $k \rightarrow 0$ ) as the small parameter (Blossey, Criminale & Fisher 2007) leads to a regular perturbation scheme. The introduction, in the perturbation Fourier decomposition, of a complex longitudinal wavenumber (Scarsoglio, Tordella & Criminale 2007) makes the problem well-posed at any order. By imposing arbitrary three-dimensional disturbances in terms of the vorticity, both the early transient as well as the asymptotic fate can be observed (Criminale & Drazin 1990). An example concerning the stability of a growing wake is presented (basic flow as  $U(x, y), V(x, y)$ , Tordella & Belan 2003). A summary of significant early time transients is shown. In the longitudinal perturbation case, asymptotic temporal results are compared with multiscale normal mode analyses (small parameter  $1/R$ ) for the intermediate and far wake (Tordella, Scarsoglio & Belan 2006; Belan & Tordella 2006).

Prefer Oral Session  
 Prefer Poster Session

Daniela Tordella  
daniela.tordella@polito.it  
Politecnico di Torino

Date submitted: 30 Jul 2007

Electronic form version 1.4