

Smooth surfaces with non smooth nullity

Original

Smooth surfaces with non smooth nullity / DI SCALA, ANTONIO JOSE'. - (2011), pp. 1-5.

Availability:

This version is available at: 11583/1896746 since: 2015-12-01T10:02:18Z

Publisher:

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



A FLAT BUT NON-SMOOTHLY RULED SURFACE


ANTONIO J. DI SCALA

As for everything else, so for a mathematical theory:
beauty can be perceived but not explained.

A. Cayley ^a

^aQuoted in J. R. Newman, *The World of Mathematics* (New York 1956).

Here we discuss an example of flat but not C^∞ -ruled surface given at page 93 of *Differential Geometry Curves-Surfaces-Manifolds* by Wolfgang Kühnel, Second Edition, AMS 2006. A similar, but not explicit example, due to E. Heintze is in 3.9.4 pag.68-39 of Klingenberg's book "A course in Differential Geometry", Springer, New York, 1978.



The surface is constructed by gluing the red cone \mathcal{C}_1 with the green cone \mathcal{C}_2 along a common generatrix (the y -axis). The blue curve \mathbf{c} , the generatrix of both cones, is in the xz plane. The vertex of \mathcal{C}_1 is the point $(0, 1, 0)$ while the vertex of \mathcal{C}_2 is the point $(0, -1, 0)$. The resulting surface is C^∞ near the origin $(0, 0, 0)$ but is not C^∞ -ruled around $(0, 0, 0)$. See below for more 3D-plots of the surface. Let $\mathbf{c} \in \mathbb{R}^3$ be a curve in the xz -plane give by

$$\mathbf{c}(x) = (x, 0, z(x))$$

where the function $z(x)$ is C^∞ , $z(0) = 0$ and $\frac{d^n z}{dx^n} \Big|_0 = 0$ for $n = 1, 2, \dots$. We assume also that $\frac{d^2 z(x)}{dx^2} > 0$ for $x \neq 0$.

Let \mathcal{C}_1 be the cone containing \mathbf{c} with vertex at the point $(0, 1, 0)$. The tangent plane $T_O \mathcal{C}_1$ at the origin O is the xy -plane. Since \mathcal{C}_1 is a C^∞ surface near O there exist a C^∞ -function $f_1(x, y)$ such that near O the cone \mathcal{C}_1 is given by the graph $(x, y, f_1(x, y))$.

Date: October 25, 2012.

I thank Fabrizio Catanese who asked to me for an example as in Proposition **A**. I would like to thank Fabio Nicola and Paolo Tilli for discussions about the issue.

A simple calculation shows that near $(0, 0)$

$$f_1(x, y) = z\left(\frac{x}{1-y}\right)(1-y).$$

So for every $n, m \in \mathbb{N}$ we have

$$(1) \quad \left. \frac{\partial^{n+m} f_1}{\partial x^n \partial y^m} \right|_{(0,y)} = 0$$

Let now \mathcal{C}_2 be the cone containing \mathbf{c} but with vertex at the point $(0, -1, 0)$. The cone \mathcal{C}_2 is also given by a graph of a function $f_2(x, y)$ near O . By the same argument f_2 also satisfy equation (1).

Then the function f defined as

$$f(x, y) = \begin{cases} f_1(x, y) & \text{if } x \leq 0 \\ f_2(x, y) & \text{if } x > 0 \end{cases}$$

is a C^∞ -function near $(0, 0)$.

Claim: *The graph $(x, y, f(x, y))$ is an example of C^∞ -surface with zero Gaussian curvature but is not a C^∞ -ruled surface near the origin O .*

Proof. We already seen that f is C^∞ . Since the graph is the union of two cones and the Gauss curvature $\kappa(x, y)$ is smooth it follows that $\kappa(x, y) \equiv 0$ since κ restricted to each cone is identically zero.

Assume that the graph of f is ruled near O i.e. the graph is the image of a map $(s, t) \rightarrow \alpha(t) + s\beta(t)$, with $\alpha, \beta \in C^\infty$, $\alpha(0) = O$ and $\alpha'(0) \wedge \beta(0) \neq 0$. So $\beta(0) = (0, b, 0)$ with $b \neq 0$. Since $\alpha'(0)$ is tangent to the xy -plane it follows that the projection of α to the xy -plane is transversal to the y -axis at $(0, 0)$. This imply the existence of C^∞ -functions $t(x), y(x)$ such that

$$\alpha(t(x)) = (x, y(x), f(x, y(x))).$$

Notice that the hypothesis $\frac{d^2 z(x)}{dx^2} > 0$ imply that the only straight lines contained in the cones $\mathcal{C}_1, \mathcal{C}_2$ are the straight lines connecting one of the vertices with the generatrix \mathbf{c} . Indeed, the shape operator of the graph in a point $(x, y, f(x, y))$ is not zero if $x \neq 0$. Hence any segment contained in the graph must be tangent to the kernel of the shape operator i.e. must be part of a straight lines connecting one of the vertices with the generatrix \mathbf{c} as claimed. More precisely, along $\alpha(t(x))$ we have the following continuous vector field $R(x)$ pointing in the direction of the rulings:

$$R(x) = \begin{cases} (x, y(x) - 1, f(x, y(x))) & \text{if } x < 0 \\ (-x, -y(x) - 1, -f(x, y(x))) & \text{if } x \geq 0 \end{cases}$$


Notice that $R(x)$ never vanish.



Then there exist a function $m(x)$ such that


$$\beta(t(x)) = m(x)R(x).$$

Since both $\beta(t(x))$ and $R(x)$ are continuous and never vanish along α it follows that $m(x)$ is continuous and never zero. The first component of $\beta(t(x))$ is $-m(x)|x|$ which is not derivable at $x = 0$ because $m(0) \neq 0$ since $\beta(0) \neq 0$. This is a contradiction since $\beta(t(x))$ is C^∞ . \square

Proposition A. *The above non-ruled C^∞ -surface has the property that each of its points is part of a segment included in the surface.*







Dipartimento di Scienze Matematiche,
Politecnico di Torino,
Corso Duca degli Abruzzi 24, 10129 Torino, Italy.
antonio.discal@polito.it
<http://calvino.polito.it/~adiscal/>