

Turbulent diffusion in different spatial dimensions.

Original

Turbulent diffusion in different spatial dimensions / Ducasse, Lauris; von Hardenberg, J.; Iovieno, Michele; Tordella, Daniela. - 8:(2010), pp. S6-32-S6-32. (Intervento presentato al convegno 8th European Fluid Mechanics Conference tenutosi a Bad Reichenhall, Germany nel September 13-16, 2010).

Availability:

This version is available at: 11583/2366565 since:

Publisher:

European Mechanics Society

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Turbulent diffusion in different spatial dimensions.

L. Ducasse*, J.von Hardenberg[†], M. Iovieno* and D. Tordella*

The diffusion properties of a mixing layer in shear-free turbulence is studied numerically in two and three dimensional flows. In both cases, the system consists of two turbulent flows with different initial kinetic energies, E_1 and E_2 , put alongside and separated by a finite gradient through a mixing layer of thickness $\Delta(t)$. The presented results focus on the differences in behavior of the system as a function of the dimensionality.

Previous studies on the problem, wether experimental ¹ or numerical ^{2,3} have shown that the velocity field was highly intermittent, which resulted in non Gaussian statistics characterized by high values of the skewness and kurtosis. Indeed, large scale eddies coming from the high energy region, penetrate intermittently into the low energy one through the mixing layer causing this latter to grow. At the beginning of the simulation, the typical thickness of the mixing layer is of the same order as the integral scale which is about 1/80 of the total domain.

Our results show that the thickness of the mixing layer evolves asymptotically in time as a power law, $\Delta(t) \propto t^\alpha$. The exponent α can thus be considered as a measure of the mixing velocity of the system. Figure 1, shows the evolution of $\Delta(t)$ as a function of time for the two dimensional (a) and the three dimensional (b) cases. The exponent α is found to be around two times higher in the first case ($\alpha \approx 0.7$ in 2D and $\alpha \approx 0.33$ in 3D), suggesting that the mixing process is faster in two dimensional turbulence.

*Dipartimento di Ingegneria Aeronautica e Spaziale, Politecnico di Torino, Torino, Italy

[†]Istituto di Scienze dell'Atmosfera e del Clima, CNR, Torino, Italy

¹Veeravalli and Warhaft, *J. Fluid Mech.* **207**, 191-229 (1989).

²Briggs, Ferziger, Koseff and Monismith, *J. Fluid Mech.* **310**, 215-241 (1996).

³Tordella and Iovieno, *J. Fluid Mech.* **549**, 429-441 (2006).




Figure 1: Mixing layer thickness as a function of time : (a) 2D case, (b) 3D case.