

Weakly uniform rank two vector bundles on multiprojective spaces

*Original*

Weakly uniform rank two vector bundles on multiprojective spaces / Ballico, E.; Malaspina, Francesco. - In: BULLETIN OF THE AUSTRALIAN MATHEMATICAL SOCIETY. - ISSN 0004-9727. - STAMPA. - 84:2(2011), pp. 255-260.  
[10.1017/S0004972711002243]

*Availability:*

This version is available at: 11583/2440866 since: 2015-12-16T16:30:06Z

*Publisher:*

Cambridge University press

*Published*

DOI:10.1017/S0004972711002243

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# Weakly uniform rank two vector bundles on multiprojective spaces

Edoardo Ballico and Francesco Malaspina

Università di Trento  
38123 Povo (TN), Italy  
e-mail: ballico@science.unitn.it

Politecnico di Torino  
Corso Duca degli Abruzzi 24, 10129 Torino, Italy  
e-mail: francesco.malaspina@polito.it

## Abstract

Here we classify the weakly uniform rank two vector bundles on multiprojective spaces. Moreover we show that every rank  $r > 2$  weakly uniform vector bundle with splitting type  $a_{1,1} = \dots = a_{r,s} = 0$  is trivial and every rank  $r > 2$  uniform vector bundle with splitting type  $a_1 > \dots > a_r$ , splits.

## 1 Introduction

We denote by  $\mathbb{P}^n$  the  $n$ -dimensional projective space over an algebraic field of characteristic zero. A rank  $r$  vector bundle  $E$  on  $\mathbb{P}^n$  is said to be uniform if there is a sequence of integers  $(a_1, \dots, a_r)$  with  $a_1 \geq \dots \geq a_r$  such that for every line  $L$  on  $\mathbb{P}^n$ ,  $E|_L \cong \bigoplus_{i=1}^r \mathcal{O}(a_i)$ . The sequence  $(a_1, \dots, a_r)$  is called the splitting type of  $E$ .

The classification of these bundles is known in many cases: rank  $E \leq n$  with  $n \geq 2$  (see [10], [9], [4]); rank  $E = n + 1$  for  $n = 2$  and  $n = 3$  (see [3], [5]); rank  $E = 5$  for  $n = 3$  (see [1]). Nevertheless there are uniform vector bundles (of rank  $2n$ ) which are not homogeneous (see [7]).

In [2] the authors gave the notion of weakly uniform bundle on  $\mathbb{P}^1 \times \mathbb{P}^1$ . For the study of rank two weakly uniform vector bundles on  $(\mathbb{P}^1)^s$ , see [11], [6] and [2].

Here we are interested on vector bundles on multiprojective spaces. Fix integers  $s \geq 2$  and  $n_i \geq 1$ . Let  $X := \mathbb{P}^{n_1} \times \dots \times \mathbb{P}^{n_s}$  be a multiprojective space. Let

$$u_i : X \rightarrow \mathbb{P}^{n_i}$$

be the projection on the  $i$ -th factor. For all  $1 < i < j$  let

$$u_{ij} : X \rightarrow \mathbb{P}^{n_i} \times \mathbb{P}^{n_j}$$

---

Mathematics Subject Classification 2010: 14F05, 14J60.

keywords: uniform vector bundles, splitting type, multiprojective spaces

denote the projection onto the product of the  $i$ -th factor and the  $j$ -th factor. Set  $\mathcal{O} := \mathcal{O}_X$ . For all integers  $b_1, \dots, b_s$  set  $\mathcal{O}(b_1, \dots, b_s) := \otimes_{i=1}^s u_i^*(\mathcal{O}_{\mathbb{P}^{n_i}}(b_i))$ . We recall that every line bundle on  $X$  is isomorphic to a unique line bundle  $\mathcal{O}(b_1, \dots, b_s)$ . Set  $X_i := \prod_{j \neq i} \mathbb{P}^{n_j}$ . Let

$$\pi_i : X \rightarrow X_i$$

be the projection. Hence  $\pi_i^{-1}(P) \cong \mathbb{P}^{n_i}$  for each  $P \in X_i$ . Let  $E$  be a rank  $r$  vector bundle on  $X$ . We say that  $E$  is *weakly uniform* with splitting type  $(a_{h,i})$ ,  $1 \leq h \leq r$ ,  $1 \leq i \leq s$ , if for all  $i \in \{1, \dots, s\}$ , every  $P \in X_i$  and every line  $D \subseteq \pi_i^{-1}(P)$  the vector bundle  $E|_D$  on  $D \cong \mathbb{P}^1$  has splitting type  $a_{1,i} \geq \dots \geq a_{r,i}$ . A weakly uniform vector bundle  $E$  on  $X$  is called *uniform* if there is a line bundles  $(a_1, \dots, a_s)$  such that the splitting types of  $E(a_1, \dots, a_s)$  with respect to all  $\pi_i$  are the same. In this case a splitting type of  $E$  is the splitting type  $c_1 \geq \dots \geq c_r$ ,  $r := \text{rank}(E)$ , of  $E(a_1, \dots, a_s)$ . Notice that the  $r$ -ple of integers  $(c_1, \dots, c_r)$  is not uniquely determined by  $E$ , but that the  $(s-1)$ -ple  $(c_1 - c_2, \dots, c_{s-1} - c_s)$  depends only from  $E$ . Indeed, a rank  $r$  weakly uniform vector bundle  $E$  of splitting type  $(a_{h,i})$ ,  $1 \leq h \leq r$ ,  $1 \leq i \leq s$ , is uniform if and only if there are  $s-1$  integers  $d_j$ ,  $2 \leq j \leq s$ , such that  $a_{h,i} = a_{h,1} + d_i$  for all  $i \in \{2, \dots, s\}$ . If  $E$  is uniform, then the  $r$ -ples  $(a_{1,1} + y, \dots, a_{r,1} + y)$ ,  $y \in \mathbb{Z}$ , are exactly the splitting types of  $E$ . If  $E$  is uniform it is usually better to consider  $E(0, a_{1,2} - a_{1,1}, \dots, a_{1,s} - a_{1,1})$  instead of  $E$ , because all the splitting types of  $E(0, a_{1,2} - a_{1,1}, \dots, a_{1,s} - a_{1,1})$  as a weakly uniform vector bundle are the same.

In this paper we prove the following result:

**Theorem 1.1.** *Let  $E$  be a rank 2 vector bundle on  $X$ .  $E$  is weakly uniform if and only if there are  $L \in \text{Pic}(X)$ , indices  $1 \leq i < j \leq s$  and a rank 2 weakly uniform vector bundle  $G$  on  $\mathbb{P}^{n_i} \times \mathbb{P}^{n_j}$  such that  $E \otimes L \cong u_{ij}^*(G)$ .  $E$  splits if either  $n_i \geq 3$  or  $n_j \geq 3$ . If  $1 \leq n_1 \leq 2$ ,  $1 \leq n_2 \leq 2$  and  $(n_1, n_2) \neq (1, 1)$ , then  $E$  splits unless there is  $h \in \{1, 2\}$  such that  $n_h = 2$  and  $E \otimes L \cong u_h^*(T\mathbb{P}^2)$  for some  $L \in \text{Pic}(X)$ .*

Moreover we discuss the case of higher rank. We show that every rank  $r > 2$  weakly uniform vector bundle with splitting type  $a_{1,1} = \dots = a_{r,s} = 0$  is trivial and every rank  $r > 2$  uniform vector bundle with splitting type  $a_1 > \dots > a_r$ , splits. Our methods did not allowed us to attack other splitting types.

## 2 Weakly uniform rank two vector bundles

In order to prove Theorem 1.1 we need a few lemmas.

We first consider the case  $s = 2$ .

**Lemma 2.1.** *Assume  $s = 2$ ,  $n_1 = 1$  and  $n_2 = 2$ . Let  $E$  be a rank 2 vector bundle on  $\mathbb{P}^1 \times \mathbb{P}^2$ .  $E$  is weakly uniform if and only if either  $E$  splits as the direct sum of 2 line bundles or there is a line bundle  $L$  on  $\mathbb{P}^1 \times \mathbb{P}^2$  such that  $E \cong L \otimes \pi_2^*(T\mathbb{P}^2)$ .*

*Proof.* Since the “if” part is obvious, it is sufficient to prove the “only if” part. Let  $(a_{h,i})$ ,  $1 \leq h \leq 2$ ,  $1 \leq i \leq s$ , be the splitting type of  $E$ . Up to a twist by a line bundle we may assume  $a_{1,1} = a_{1,2} = 0$ . By rigidity or looking at the Chern classes  $c_i(E|_{\{Q\} \times \mathbb{P}^2})$ ,  $i = 1, 2$ , it is easy to see that if one of these two cases occurs for some  $Q$ , then it occurs for all  $Q$ . First assume  $a_{2,2} = 0$ . Since the trivial line bundle on  $\mathbb{P}^1$  is spanned, the theorem of changing basis implies that  $F := \pi_{2*}(E)$  is a rank 2 vector bundle on  $\mathbb{P}^2$  and that the natural map  $\pi_2^*(F) \rightarrow E$  is an isomorphism ([8], p. 11). Since  $E$  is weakly uniform,  $F$  is uniform. The

classification of all rank 2 uniform vector bundles on  $\mathbb{P}^2$  shows that either  $F$  splits or it is isomorphic to a twist of  $T\mathbb{P}^2$  (see [4]), concluding the proof in the case  $a_{2,2} = 0$ . Similarly, if  $a_{2,1} = 0$ , there is a rank 2 vector bundle  $G$  on  $\mathbb{P}^1$  such that  $\pi_1^*(G) \cong E$ . Since every vector bundle on  $\mathbb{P}^1$  splits, we have that also  $E$  splits. Now we may assume  $a_{2,2} < 0$  and  $a_{2,1} < 0$ . Since  $a_{2,2} < 0$ , the base-change theorem gives that  $\pi_{2*}(E)$  is a line bundle, say of degree  $b_2$ , and that the natural map  $\pi_2^*\pi_{2*}(E) \rightarrow E$  has locally free cokernel ([8], p. 11). Thus in this case  $E$  fits in an exact sequence

$$0 \rightarrow \mathcal{O}(0, b_2) \rightarrow E \rightarrow \mathcal{O}(a_{2,1}, -b_2 - a_{2,2}) \rightarrow 0 \quad (1)$$

The term  $a_{2,1}$  in the last line bundle of (1) comes from  $c_1(E)$ . If (1) splits, then we are done. Since  $a_{2,1} \leq 1$ , Künneth's formula gives  $H^1(\mathbb{P}^1 \times \mathbb{P}^2, \mathcal{O}(-a_{2,1}, 2b_2 + a_{2,2})) = 0$ . Hence (1) splits.  $\square$

**Lemma 2.2.** *Assume  $s = 2$ ,  $n_1 = 1$  and  $n_2 \geq 3$ . Then every rank two weakly uniform vector bundle on  $X$  is the direct sum of two line bundles.*

*Proof.* We copy the proof of Lemma 2.1. Every rank 2 uniform vector bundle on  $\mathbb{P}^m$ ,  $m \geq 3$ , splits. Hence  $E$  splits even in the case  $a_{2,2} = 0$ .  $\square$

**Lemma 2.3.** *Assume  $s = 2$  and  $n_1 = n_2 = 2$ . Let  $E$  be a rank 2 indecomposable weakly uniform vector bundle on  $X$ . Then either  $E \cong u_1^*(T\mathbb{P}^2)(u, v)$  or  $E \cong u_2^*(T\mathbb{P}^2)(u, v)$ .*

*Proof.* Let  $(a_{h,i})$  be the splitting type of  $E$ . Up to a twist by a line bundle we may assume  $a_{1,1} = a_{1,2} = 0$ . As in the proof of Lemma 2.1 the theorem of changing basis gives that either  $E \cong u_1^*(T\mathbb{P}^2(-2))$  or  $E$  splits if  $a_{2,1} = 0$  and that  $E \cong u_2^*(T\mathbb{P}^2(-2))$  or  $E$  splits if  $a_{2,2} = 0$ . If  $a_{2,1} < 0$  and  $a_{2,2} < 0$ , then we apply  $\pi_{2*}$  and get an exact sequence (1). Here Künneth's formula gives that (1) splits, without using any information on the integer  $a_{2,2}$ .  $\square$

**Lemma 2.4.** *Assume  $s = 2$ ,  $n_1 \geq 3$  and  $n_2 = 2$ . Let  $E$  be a rank 2 weakly uniform vector bundle on  $X$ . Then either  $E$  splits or  $E \cong u_2^*(T\mathbb{P}^2)(u, v)$  for some integers  $u, v$ .*

*Proof.* Let  $(a_{hi})$  be the splitting type of  $E$ . Up to a twist by a line bundle we may assume  $a_{1,1} = a_{1,2} = 0$ . As in the proof of Lemma 2.1 the theorem of changing basis gives that  $E \cong u_1^*(T\mathbb{P}^2(-2))$  or  $E$  splits if  $a_{2,1} = 0$  and that  $E$  splits in the case  $a_{1,2} < 0$ , because (1) splits by Künneth's formula.  $\square$

**Lemma 2.5.** *Assume  $s = 2$ ,  $n_1 \geq 3$  and  $n_2 \geq 3$ . Let  $E$  be a rank 2 weakly uniform vector bundle on  $X$ . Then  $E$  splits.*

*Proof.* Let  $(a_{hi})$  be the splitting type of  $E$ . Up to a twist by a line bundle we may assume  $a_{1,1} = a_{1,2} = 0$ . If  $a_{2,2} = 0$ , then base change gives  $E \cong u_2^*(F)$  for some uniform vector bundle on  $\mathbb{P}^2$ . Thus we may assume  $a_{2,2} < 0$ . We have again the extension (1). Here again (1) splits by Künneth's formula.  $\square$

Now we are ready to prove the main theorem:

*Proof of Theorem 1.1.* First assume  $s = 2$ . Theorem 1.1 says nothing in the case  $n_1 = n_2 = 1$  for which a full classification is not known ([2] shows that moduli arises). Lemmas 2.1, 2.2, 2.3, 2.4 and 2.5 cover all cases with  $s = 2$ . Hence we may assume  $s \geq 3$  and use induction on  $s$ . If  $n_i = 1$  for all  $i$ , then we may apply [2], Theorem 4. For arbitrary  $n_i$  the proof of [2], Theorem 4, works verbatim, but for reader's sake we repeat that proof. Let  $(a_{hi})$

be the splitting type of  $E$ . Up to a twist by a line bundle we may assume  $a_{1i} = 0$  for all  $i$ . If  $a_{2i} = 0$  for some  $i$ , then the base-change theorem gives  $E \cong \pi_i^*(F)$  for some weakly uniform vector bundle  $F$  on  $X_i$ . If  $s = 3$ , then we are done. In the general case we reduce to the case  $s' := s - 1$ . Thus to complete the proof it is sufficient either to obtain a contradiction or to get that  $E$  splits under the additional condition that  $a_{2i} < 0$  for all  $i$  and  $s \geq 3$ . Applying the base-change theorem to  $\pi_{1*}$  we get that  $E$  fits in the following extension

$$0 \rightarrow \mathcal{O}(0, c_2, \dots, c_s) \rightarrow E \rightarrow \mathcal{O}(a_{1,2}, d_2, \dots, d_s) \rightarrow 0 \quad (2)$$

Since  $-a_{1,2} \geq 0$ , Künneth's formula shows that (2) splits unless  $n_i = 1$  for all  $i \geq 2$ . Using  $\pi_{2*}$  instead of  $\pi_{1*}$  we get that  $E$  splits, unless  $n_1 = 1$ .  $\square$

### 3 Higher rank weakly uniform vector bundles

Now we consider higher rank weakly uniform vector bundles.

**Proposition 3.1.** *Let  $E$  be a rank  $r$  weakly uniform vector bundle on  $X$  with splitting type  $(0, \dots, 0)$ . Then  $E$  is trivial.*

*Proof.* The case  $s = 1$  is true by [8], Theorem 3.2.1. Hence we may assume  $s \geq 2$  and use induction on  $s$ . By the inductive assumption  $E|_{\pi_1^{-1}(P)}$  is trivial for each  $P \in \mathbb{P}^{n_1}$ . By the base-change theorem  $F := \pi_{1*}(E)$  is a rank  $r$  vector bundle on  $X_1$  and the natural map  $\pi_1^*(F) \rightarrow E$  is an isomorphism. This isomorphism implies that  $F$  is uniform of splitting type  $(0, \dots, 0)$ . Hence the inductive assumption gives that  $F$  is trivial. Thus  $E$  is trivial.  $\square$

In order to study uniform vector bundles with  $a_1 > \dots > a_r$  we need the following lemmas:

**Lemma 3.2.** *Fix an integer  $r \geq 2$  and a rank  $r$  vector bundle on  $X$ . Assume the existence of an integer  $i \in \{1, \dots, s\}$  such that  $E|_{\pi_i^{-1}(P)}$  is the direct sum of line bundles for all  $P \in X_i$ . If  $n_i = 1$  assume that the splitting type of  $E|_{\pi_i^{-1}(P)}$  is the same for all  $P \in X_i$ . Let  $(a_1, \dots, a_r) = (b_1^{m_1}, \dots, b_k^{m_k})$ ,  $b_1 > \dots > b_k$ ,  $m_1 + \dots + m_k = r$ , be the splitting type of  $E|_{\pi^{-1}(P)}$  for any  $P \in X_i$ . Then there are  $k$  vector bundles  $F_1, \dots, F_k$  on  $X_i$  and  $k$  vector bundles  $E_1, \dots, E_k$  on  $X$  such that  $\text{rank}(F_i) = m_i$ ,  $E_k = E$ ,  $E_{i-1}$  is a subbundle of  $E_i$  and  $E_i/E_{i-1} \cong \pi_i^*(F_i)(-b_i)$  (with the convention  $E_0 = 0$ ).*

*Proof.* Notice that even in the case  $n_i \geq 2$  the splitting type of  $E|_{\pi^{-1}(P)}$  does not depend from the choice of  $P \in X_i$  (e.g. use Chern classes or local rigidity of direct sums of line bundles). Thus  $E|_{\pi_i^{-1}(P)} \cong \bigoplus_{j=1}^k \mathcal{O}_{\pi_i^{-1}(P)}(b_j)^{\oplus m_j}$  for all  $P \in X_i$ .

Set  $F_1 := \pi_{i*}(E(0, \dots, -b_1, \dots, 0))$ . By the base-change theorem  $F_1$  is a rank  $m_1$  vector bundle on  $X_i$  and the natural map  $\rho : \pi_i^*(F_1)(0, \dots, b_1, \dots) \rightarrow E$  is a vector bundle embedding, i.e. either  $\rho$  is an isomorphism (case  $r = m_1$ ) or  $\text{Coker}(\rho)$  is a rank  $r - m_1$  vector bundle on  $X$ . If  $m_1 = r$ , then  $k = 1$  and we are won. Now assume  $k \geq 2$ , i.e.  $m_1 < r$ . Fix any  $P \in X_i$ . By definition  $\text{Coker}(\rho)$  fits in an exact sequence of vector bundles on  $X$ :

$$0 \rightarrow \pi_i^*(F_1)(0, \dots, b_1, \dots, 0) \rightarrow E \rightarrow \text{Coker}(\rho) \rightarrow 0 \quad (3)$$

and the restriction to  $\pi_i^{-1}(P)$  of the injective map of (3) induces an embedding of vector bundles  $j_P : \mathcal{O}_{\pi_i^{-1}(P)}(b_1)^{\oplus m_1} \rightarrow \bigoplus_{j=1}^k \mathcal{O}_{\pi_i^{-1}(P)}(b_j)^{\oplus m_j}$ . Since  $b_1 > b_j$  for all  $j > 1$ , we get  $\text{Coker}(j_P) \cong \bigoplus_{j=2}^k \mathcal{O}_{\pi_i^{-1}(P)}(b_j)^{\oplus m_j}$ . We apply to  $\text{Coker}(\rho)$  the inductive assumption on  $k$ .  $\square$

**Lemma 3.3.** *Assume  $s = 2$  and  $n_1 \geq 2$ ,  $n_2 \geq 3$ . Fix an integer  $r$  such that  $3 \leq r \leq n_2$  and a rank  $r$  uniform vector bundle  $E$  with splitting type  $a_1 > \cdots > a_r$ . Then  $E$  is isomorphic to a direct sum of  $r$  line bundles.*

*Proof.* Since  $r \geq 3$ , we have  $a_r \leq a_1 - 2$ . Thus the classification of uniform vector bundles on  $\mathbb{P}^{n_2}$  with rank  $r \leq n_2$ , gives  $E|_{\pi_1^{-1}(P)} \cong \bigoplus_{i=1}^r \mathcal{O}_{\pi_1^{-1}(P)}(a_i)$  for all  $P \in \mathbb{P}^{n_1}$ . Apply Lemma 3.2 with respect to the integers  $i = 1$  and  $k = r$  and let  $F_i, E_i$ ,  $1 \leq i \leq r$ , be the vector bundles given by the lemma. Since  $E_r = E$ , it is sufficient to prove that each  $E_i$  is a direct sum of  $i$  line bundles. Since  $\text{rank}(E_i) = i$ , the latter assertion is obvious if  $i = 1$ . Fix an integer  $i$  such that  $1 \leq i < r$  and assume that  $E_i$  is isomorphic to a direct sum of  $i$  line bundles. Lemma 3.2 gives an extension

$$0 \rightarrow E_i \rightarrow E_{i+1} \rightarrow L \rightarrow 0$$

with  $L$  a line bundle on  $\mathbb{P}^{n_1} \times \mathbb{P}^{n_2}$ . Since  $n_1 \geq 2$  and  $n_2 \geq 2$ , Künneth's formula gives that any extension of two line bundles on  $\mathbb{P}^{n_1} \times \mathbb{P}^{n_2}$  splits. Thus  $E_{i+1}$  is a direct sum of  $i + 1$  line bundles.  $\square$

**Proposition 3.4.** *Fix an integer  $r \geq 3$  and a rank  $r$  uniform vector bundle on  $X$  with splitting type  $a_1 > \cdots > a_r$ . Assume  $s \geq 2$ ,  $n_2 \geq r$  and  $n_i \geq 2$  for all  $i \neq 2$ . Then  $E$  is isomorphic to a direct sum of  $r$  line bundles.*

*Proof.* The case  $s = 2$  is Lemma 3.3. Thus we may assume  $s \geq 3$  and that the proposition is true for  $\mathbb{P}^{n_1} \times \cdots \times \mathbb{P}^{n_{s-1}}$ . By the inductive assumption  $E|_{u_s^{-1}(P)} \cong \bigoplus_{i=1}^r \mathcal{O}_{u_s^{-1}(P)}(a_i, \dots, a_i)$  for all  $P \in \mathbb{P}^{n_s}$ . As in the proof of Lemma 3.2 taking instead of  $\pi_i$  the projection  $u_i : X \rightarrow \mathbb{P}^{n_i}$  we get line bundles  $L_i$ ,  $1 \leq i \leq r$  of  $\mathbb{P}^{n_s}$ , (i.e. line bundles  $u_i^*(L) \cong \mathcal{O}(0, \dots, 0, c_i, 0, \dots, 0)$  on  $X$ ) and subbundles  $E_1 \subset E_2 \subset \cdots \subset E_r = E$  such that  $E_i/E_{i-1} \cong \mathcal{O}_X(a_{i-1}, \dots, a_{i-1}, c_i)$  (with the convention  $E_0 = 0$ ). It is sufficient to prove that each  $E_i$  is isomorphic to a direct sum of  $i$  line bundles. Since this is obvious for  $i = 1$ , we may use induction on  $i$ . Fix an integer  $i \in \{2, \dots, r\}$ . Our assumption on  $X$  implies that the extension of any two line bundles splits. Hence  $E_i \cong E_{i-1} \oplus \mathcal{O}_X(a_{i-1}, \dots, a_{i-1}, c_i)$ .  $\square$

## References

- [1] E. Ballico and P. Ellia, Fibrés uniformes de rang 5 sur  $\mathbb{P}^3$ . Bull. Soc. Math. France 111 (1983), 59–87.
- [2] E. Ballico and P. E. Newstead, Uniform bundles on quadric surfaces and some related varieties. J. London Math. Soc. (2) 31 (1985), no. 2, 211–223.
- [3] G. Elencwajg, Les fibres uniformes de rang 3 sur  $\mathbf{P}_2(\mathbf{C})$  sont homogènes. Math. Ann. 231 (1978), 217–227.
- [4] G. Elencwajg, A. Hirschowitz and M. Schneider, Les fibres uniformes de rang au plus  $n$  sur  $\mathbf{P}_n(\mathbf{C})$  sont ceux qu'on croit. Vector bundles and differential equations (Proc. Conf., Nice, 1979), pp. 37–63, Progress in Math. 7, Birkhäuser, Boston, Mass., 1980.
- [5] P. Ellia, Fibrés uniformes de rang  $n + 1$  sur  $\mathbb{P}^n$ . Mem. Soc. Math. France 7 (1982).
- [6] P. E. Newstead and R. L. E. Schwarzenberger, Reducible vector bundles on a quadric surface. Proc. Cambridge Philos. Soc. 60 (1964), 421–424.

- [7] J. M. Drezet, Example de fibrés uniformes non omogénés. C. R. Acad. Sci. Paris Sér A 291 (1980), 125–128.
- [8] Ch. Okonek, M. Schneider and H. Spindler, Vector bundles on complex projective spaces. Progress in Math. 3, Birkhäuser, Boston, Mass., 1980.
- [9] E. Sato, Uniform vector bundles on a projective space. J. Math. Soc. Japan. 28 (1976), 123–132.
- [10] A. Van de Ven, On uniform vector bundles. Math. Ann. 195 (1972), 245–248.
- [11] R. L. E. Schwarzenberger, Reducible vector bundles on a quadric surface, Proc. Cambridge Philos. Soc. 58 (1962), 209–216.