

Combinatorial presentation of multidimensional persistent homology

Original

Combinatorial presentation of multidimensional persistent homology / Wojciech, Chacholski; Martina, Scolamiero; Vaccarino, Francesco. - (2014).

Availability:

This version is available at: 11583/2591369 since:

Publisher:

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

COMBINATORIAL PRESENTATION OF MULTIDIMENSIONAL PERSISTENT HOMOLOGY

W. CHACHOLSKI¹ M. SCOLAMIERO F. VACCARINO²

ABSTRACT. A multifiltration is a functor indexed by \mathbb{N}^r that maps any morphism to a monomorphism. The goal of this paper is to describe in an explicit and combinatorial way the natural \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module structure on the homology of a multifiltration of simplicial complexes. To do that we study multifiltrations of sets and vector spaces. We prove in particular that the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules that can occur as R -spans of multifiltrations of sets are the direct sums of monomial ideals.

1. INTRODUCTION

Let \mathbb{N}^r be the poset of r -tuples of natural numbers with partial order given by $(v_1, \dots, v_r) \leq (w_1, \dots, w_r)$ if and only if $v_i \leq w_i$ for all $1 \leq i \leq r$. A functor $F: \mathbb{N}^r \rightarrow \text{Spaces}$, with values in the category of simplicial complexes, is called a **multifiltration** if, for any $v \leq w$ in \mathbb{N}^r , the map $F(v \leq w): F(v) \rightarrow F(w)$ is a monomorphism. Such a multifiltration is called **compact** if $\text{colim}_{\mathbb{N}^r} F$ is a finite complex. Compact multifiltrations are the main objects we are studying in this article. By applying homology with coefficients in a ring R to F we obtain a functor $H_n(F, R): \mathbb{N}^r \rightarrow R\text{-Mod}$ with values in the category of R -modules. The category of functors indexed by \mathbb{N}^r with values in $R\text{-Mod}$ is equivalent to the category of \mathbb{N}^r -graded modules over the polynomial ring $R[x_1, \dots, x_r]$. The aim of this paper is to describe this $R[x_1, \dots, x_r]$ -module structure on $H_n(F, R)$ in a way that is suitable for calculations. One very efficient way of doing it would be to give the minimal free presentation of $H_n(F, R)$. This however we are unable to do directly. Instead we are going to describe two homomorphisms of finitely generated and free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules $\mathbf{A} \rightarrow \mathbf{B} \rightarrow \mathbf{C}$ whose composition is the zero homomorphism (this sequence is a chain complex), and $H_n(F, R)$ is isomorphic to the homology of this complex. Since the modules involved are finitely generated and free and the homomorphisms preserve grading, these homomorphisms are simply given by matrices of elements in R . In our case the coefficients of the matrices are either 1, -1 or 0 and they can be explicitly expressed in terms of the multifiltration (we give a polynomial time procedure of how to do that in Section 5). One can then use standard computer algebra packages to study algebraic invariants of the module $H_n(F, R)$, in particular one can get its minimal free presentation as well as a minimal resolution, the set of Betti numbers and the Hilbert function. These invariants can be used then to study point clouds according to the theory of multidimensional persistence (see for example [1, 2]). Our procedure reduces the

¹Partially supported by Göran Gustafsson Stiftelse and VR grant 2009-6102.

²Partially supported by the TOPDRIM project funded by the Future and Emerging Technologies program of the European Commission under Contract IST-318121.

Date: September 30, 2014.

computation of $H_n(F, R)$ to the computation of the homology of a chain complex of free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules. This is the starting point in [2] where the authors explain how to calculate this homology in polynomial time. One of our aims has been to show that such calculations can be done effectively for arbitrary compact multifiltrations and not only for the so called one critical which are studied in [2].

The theory of multidimensional persistence is interesting both from an applied and theoretical point of view. From the applied perspective it is useful to construct algorithms that characterize and distinguish multifiltrations of data sets or networks according to topological features (see [4]). From a theoretical point of view, multidimensional persistence modules are \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules built from a multifiltration. It is interesting to study how the combinatorial structure of the multifiltration is reflected in the module structure and that is what we address in this paper. We start with discussing the multifiltrations of sets in Section 3. We recall the structure of such multifiltrations, how can they be decomposed into indecomposable parts and how to classify the indecomposable pieces. We use it to give an algorithm for producing a free presentation of a multifiltration of sets. In Section 4 we then study the effect of taking the R -span functor on multifiltrations of sets. We explain why the obtained multifiltrations of R -modules are rather special and prove that they correspond to sums of monomial ideals. Since the R -span functor commutes with colimits, free presentations for multifiltrations of sets can be used to obtain free presentations of monomial ideals. These presentations are used in Section 5 to obtain the desired description of the $R[x_1, \dots, x_r]$ -module structure on $H_n(F, R)$. We conclude by pointing out, in Section 6, that for multifiltrations indexed by \mathbb{N}^2 a presentation of the module $H_n(F, R)$, as the cokernel of a homogeneous homomorphism, is an easier task. In this case, the kernel of $\mathbf{B} \rightarrow \mathbf{C}$ is free and therefore, given our previous results, it is sufficient to choose a set of free generators of this kernel to find a presentation of $H_n(F, R)$. The problem of identifying such a set of free generators in an algorithmic and combinatorial way is left as an open question.

2. NOTATION

2.1. The symbols Sets, Spaces, and R -Mod denote the categories of respectively sets, simplicial complexes, and R -modules, where we always assume that R is a commutative ring with identity. The R -linear span functor which assigns to a set S the free module $R(S) = \bigoplus_S R$ is denoted by $R: \text{Sets} \rightarrow R\text{-Mod}$.

2.2. By definition a **simplicial complex** X is a collection of subsets of a set X_0 (called the set of vertices of X) such that: for any x in X_0 , $\{x\} \in X$ and if $\sigma \in X$ and $\tau \subset \sigma$, then $\tau \in X$. An element σ in X is called a **simplex** of dimension $|\sigma| - 1$. A complex is called **finite** if X_0 is a finite set. A morphism between two simplicial complexes $f: X \rightarrow Y$ is by definition a map of sets $f: X_0 \rightarrow Y_0$ such that $f(\sigma)$ is a simplex in Y for any simplex σ in X . A morphism $f: X \rightarrow Y$ is a **monomorphism** if and only if the function $f: X_0 \rightarrow Y_0$ is injective.

Let us choose an order $<$ on the set X_0 . For $n \geq 0$, the symbol X_n denotes the set of strictly increasing sequences $x_0 < \dots < x_n$ of elements in X_0 for which the subset $\{x_0, \dots, x_n\} \subset X_0$ is a simplex in X . Such a sequence is called an **ordered simplex** of dimension n . For $0 \leq i \leq n + 1$, by forgetting the i -th element in a sequence $x_0 < \dots < x_{n+1}$ we get an element in X_n . The obtained map is denoted

by $d_i: X_{n+1} \rightarrow X_n$. By applying the R -span functor and taking the alternating sum of the induced maps we obtain:

$$RX_{n+1} \xrightarrow{\partial_{n+1} := \sum_{i=0}^{n+1} (-1)^i d_i} RX_n \xrightarrow{\partial_n := \sum_{i=0}^n (-1)^i d_i} RX_{n-1}$$

where for $n = 0$, the R -module RX_{-1} is taken to be trivial. It is a standard fact that the composition $\partial_n \partial_{n+1}$ is the trivial map and hence the image $\text{im}(\partial_{n+1})$ is a submodule of the kernel $\ker(\partial_n)$. The quotient $\ker(\partial_n)/\text{im}(\partial_{n+1})$ is called the n -th homology of X and is denoted by $H_n(X, R)$. The isomorphism type of this module does not depend on the chosen ordering on X_0 . Note that this is not a functor on the entire category of simplicial complexes. However if $f: X \rightarrow Y$ is a monomorphism, then we can choose first an ordering on Y_0 and then use it to induce an ordering on X_0 so the function $f: X_0 \rightarrow Y_0$ is order preserving. With these choices, by applying f to ordered sequences element-wise, we obtain a map of sets $f_n: X_n \rightarrow Y_n$ which commutes with the maps d_i . In this way we get an induced map of homology modules that we denote by $H_n(f, R): H_n(X, R) \rightarrow H_n(Y, R)$.

2.3. The symbol $R[x_1, \dots, x_r]$ denotes the \mathbb{N}^r -graded polynomial ring with coefficients in a ring R . The category of \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules with the degree preserving homomorphisms is denoted by $R[x_1, \dots, x_r]\text{-Mod}$ and we use bold face letters to denote such modules.

A monomial in $R[x_1, \dots, x_r]$ is a polynomial of the form $x_1^{v_1} \cdots x_r^{v_r}$. Its grade is given by $v = (v_1, \dots, v_r)$. Such a monomial is also written as x^v . An \mathbb{N}^r -graded ideal in $R[x_1, \dots, x_r]$ is called **monomial** if it is generated by monomials. An \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module isomorphic to the ideal of $R[x_1, \dots, x_r]$ generated by a single monomial x^v is called **free on one generator** v and denoted by $\langle x^v \rangle$. An \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module which is isomorphic to a direct sum of free modules on one generator is called **free**. The R -module $\text{Hom}(\langle x^v \rangle, \langle x^w \rangle)$ is either trivial if $v \not\geq w$, or is isomorphic to R if $v \geq w$. We use this to identify the R -module of homomorphisms between free modules $\text{Hom}(\oplus_{t \in T} \langle x^{v_t} \rangle, \oplus_{s \in S} \langle x^{w_s} \rangle)$ with the set of $S \times T$ matrices of elements in R whose (s, t) entry is 0 if $v_t \not\geq w_s$. Thus to describe a degree preserving homomorphism between two finitely generated and free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules we need to specify:

- A matrix M of elements in R .
- Two functions, one that assigns to every row of M an element in \mathbb{N}^r and the other that assigns to every column of M an element in \mathbb{N}^r . The values of these functions are called grades of the respective rows and columns. The grades of the columns correspond to the grades of the generators of the domain of the homomorphism and the grades of the rows correspond to the grades of the generators of the range of the homomorphism.
- The matrix M should satisfy the following property: the entry corresponding to a row with grade w and a column with grade v is zero if $v \not\geq w$.

2.4. Let I be a small category. The symbol $\text{Fun}(I, \mathcal{C})$ denotes the category of functors indexed by I with values in a category \mathcal{C} and natural transformations as morphisms. We use the symbol $\text{Nat}_{\mathcal{C}}(F, G)$ to denote the set of natural transformations between two functors $F, G: I \rightarrow \mathcal{C}$. Recall [3] that the colimit of a functor $F: I \rightarrow \mathcal{C}$ is an object $\text{colim}_I F$ in \mathcal{C} together with morphisms $p_i: F(i) \rightarrow \text{colim}_I F$, for any object i in I . These morphisms are required to satisfy the following universal property. First, for any $\alpha: i \rightarrow j$ in I , $p_j F(\alpha: i \rightarrow j) = p_i$. Second, if

$q_i: F(i) \rightarrow X$ is a sequence of morphisms in \mathcal{C} indexed by objects of I fulfilling the equality $q_j F(\alpha: i \rightarrow j) = q_i$ for any morphism α in I , then there is a unique $f: \text{colim}_I F \rightarrow X$ such that $q_i = fp_i$ for any object i in I .

If I is the empty category, $\text{colim}_I F$ is called the **initial** object and denoted by \emptyset . The initial object has the property that, for any object X in \mathcal{C} , the set of morphisms $\text{mor}_{\mathcal{C}}(\emptyset, X)$ has exactly one element. If I is a discrete category, then $\text{colim}_I F$ is called the **coproduct** and denoted either by $\coprod_{i \in I} F(i)$ or $\bigoplus_{i \in I} F(i)$. The second notation is used only in the case the coproduct is taken in an additive or abelian category, as for example in $R\text{-Mod}$.

2.5. An object X in \mathcal{C} is called **decomposable** if it is isomorphic to a sum $X_1 \coprod X_2$ where neither X_1 nor X_2 is the initial object. It is **indecomposable** if it is neither initial nor decomposable. An object X is called **uniquely decomposable** if the following two conditions hold. First, it is isomorphic to a coproduct $\coprod_{i \in I} X_i$ where X_i is indecomposable for any i . Second, if X is isomorphic to $\coprod_{i \in I} X_i$ and to $\coprod_{j \in J} Y_j$, where X_i 's and Y_j 's are indecomposable, then there is a bijection $\phi: I \rightarrow J$ such that X_i and $Y_{\phi(i)}$ are isomorphic for any i in I .

In the category of sets the initial object is the empty set, the coproduct is the disjoint union, a set is decomposable if it contains at least two elements, and is indecomposable if it contains exactly one element. For $F: I \rightarrow \text{Sets}$, its colimit is the quotient of $\coprod_{i \in I} F(i)$ by the equivalence relation generated by x_i in $F(i)$ is related to x_j in $F(j)$ if there are morphisms $\alpha: i \rightarrow k$ and $\beta: j \rightarrow k$ in I for which $F(\alpha)(x_i) = F(\beta)(x_j)$.

2.6. The symbol \mathbb{N}^r denotes the poset of r -tuples of natural numbers with partial order given by $(v_1, \dots, v_r) \leq (w_1, \dots, w_r)$ if and only if $v_i \leq w_i$ for all $1 \leq i \leq r$. The initial element $(0, \dots, 0)$ in \mathbb{N}^r is denoted simply by 0 . Recall that the partial order on \mathbb{N}^r is a lattice. This means that for any finite set of elements S in \mathbb{N}^r , there are elements $\min(S)$ and $\max(S)$ in \mathbb{N}^r (not necessarily in S) with the following properties. First, for any v in S , $\min(S) \leq v \leq \max(S)$. Second, if u and w are elements in \mathbb{N}^r for which $u \leq v \leq w$, for any v in S , then $u \leq \min(S)$ and $\max(S) \leq w$. Furthermore any non-empty subset S of \mathbb{N}^r has an element v such that if $w < v$, then w is not in S . Such elements are called minimal in S and may not be unique. A functor F indexed by the poset \mathbb{N}^r that maps any morphism to a monomorphism is called a **multifiltration**. We will denote the colimit of a functor F indexed by \mathbb{N}^r by $\text{colim } F$. A multifiltration $F: \mathbb{N}^r \rightarrow \text{Sets}/R\text{-Mod}$ is called **one critical** if for any element x in $\text{colim } F$, the set $\{v \in \mathbb{N}^r \mid x \text{ is in the image of } p_v: F(v) \rightarrow \text{colim } F\}$ has a unique minimal element which we denote by v_x and call the **critical coordinate** of x (see [2]). A functor $F: \mathbb{N}^r \rightarrow \text{Sets}/\text{Spaces}$ is called **compact** if $\text{colim } F$ is a finite set/simplicial complex.

2.7. Let v be an element in \mathbb{N}^r . The functor $\text{mor}_{\mathbb{N}^r}(v, -): \mathbb{N}^r \rightarrow \text{Sets}$ is called **free on one generator**. For example $\text{mor}_{\mathbb{N}^r}(0, -): \mathbb{N}^r \rightarrow \text{Sets}$ is the constant functor with value the one point set. Since \mathbb{N}^r is a poset, the values of a free functor on one generator are either empty, or the one point set. A functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ is called **free** if it is isomorphic to a disjoint union of free functors on one generator. Note that any free functor is a multifiltration.

Composition with the R -span functor $R: \text{Sets} \rightarrow R\text{-Mod}$, is denoted by the same symbol $R: \text{Fun}(\mathbb{N}^r, \text{Sets}) \rightarrow \text{Fun}(\mathbb{N}^r, R\text{-Mod})$ and called by the same name the R -span functor. Recall that this R -span functor is the left adjoint to the forget the

R -module structure functor. This implies that the R -span functor commutes with colimits, in particular it maps the initial object to the initial object and commutes with coproducts.

The functor $R\text{mor}_{\mathbb{N}^r}(v, -): \mathbb{N}^r \rightarrow R\text{-Mod}$ is also called **free on one generator**. A functor $F: \mathbb{N}^r \rightarrow R\text{-Mod}$ is called **free** if it is isomorphic to the R -span of a free functor with values in Sets or equivalent, if it is isomorphic to a direct sum of free functors on one generator.

2.8. Recall that the category of functors $\text{Fun}(\mathbb{N}^r, R\text{-Mod})$ is equivalent to the category of \mathbb{N}^r -graded modules $R[x_1, \dots, x_r]\text{-Mod}$. We are going to identify these categories using the following explicit equivalence which assigns to a functor $F: \mathbb{N}^r \rightarrow R\text{-Mod}$, the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module given by $\mathbf{F} := \bigoplus_{v \in \mathbb{N}^r} F(v)$, where x_i acts on component $F(v)$ via the map $F(v \leq v + e_i)$ where e_i is the i -th vector in the standard base. Via this identification, the free functor $R\text{mor}_{\mathbb{N}^r}(v, -): \mathbb{N}^r \rightarrow R\text{-Mod}$ is mapped to the free module $\langle x^v \rangle$.

3. FUNCTORS WITH VALUES IN SETS

The aim of this section is to prove several basic properties of functors of the form $F: \mathbb{N}^r \rightarrow \text{Sets}$. Many of these properties are well known. We start with:

3.1. Proposition. *A functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ is indecomposable (see 2.5) if and only if the set $\text{colim } F$ contains exactly one element.*

Proof. If the values of F are not all empty, then $\text{colim } F$ is not empty. Further more if $F = G \coprod H$, then $\text{colim } F = (\text{colim } G) \coprod (\text{colim } H)$. This shows that if $\text{colim } F$ contains exactly one point, then F is indecomposable. On the other hand we can decompose F as $\coprod_{x \in \text{colim } F} F[x]$ where, for any point x in $\text{colim } F$, $F[x]: \mathbb{N}^r \rightarrow \text{Sets}$ is the subfunctor of F whose values are given by $F[x](v) := \{y \in F(v) \mid p_v(y) = x\}$ (see 2.4). Observe that not all the values of $F[x]$ are empty. This describes F as a coproduct of indecomposable functors. Thus if F is indecomposable, then $\text{colim } F$ has to contain only one element. \square

The argument in the above proof shows more:

3.2. Corollary. *Any functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ is uniquely decomposable as $F = \coprod_{x \in \text{colim } F} F[x]$.*

In this paper we are not interested in all functors indexed by \mathbb{N}^r with values in Sets, but those that map any morphism to a monomorphism. Such functors are called multifiltrations of sets (see 2.6) and here is their characterization:

3.3. Proposition. *A functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ is a multifiltration if and only if the map $p_v: F(v) \rightarrow \text{colim } F$ is a monomorphism for any v in \mathbb{N}^r .*

Proof. Recall that $\text{colim } F$ is the quotient of $\coprod_{v \in \mathbb{N}^r} F(v)$ by the equivalence relation generated by x_v in $F(v)$ is related to x_w in $F(w)$, if there is $u \geq v$ and $u \geq w$ such that $F(v \leq u)(x_v) = F(w \leq u)(x_w)$. Note that since \mathbb{N}^r is a lattice, the described relation is already an equivalence relation. Thus two elements of $F(v)$ are mapped to the same element in $\text{colim } F$ if and only if they are mapped to the same element via $F(v \leq u)$ for some u and the proposition follows. \square

3.4. Corollary. *A functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ is an indecomposable multifiltration if and only if the set $F(v)$ has at most one element for any v in \mathbb{N}^r and there is u for which $F(u)$ is not empty.*

Proof. Assume first F is an indecomposable multifiltration. By Proposition 3.1, $\text{colim } F$ is the one point set. The multifiltration assumption implies that $F(v)$ is a subset of $\text{colim } F$ for any v (see 3.3). Consequently the set $F(v)$ can not contain more than one element. Since $\text{colim } F$ is not empty, the values of F can not be all empty either. This shows one implication.

Recall that any element in $\text{colim } F$ is of the form $p_v(x)$ for some v in \mathbb{N}^r and x in $F(v)$. Assume that $\text{colim } F$ has at least two elements, which we write as $p_v(x)$ and $p_w(y)$. The elements $F(v \leq \max\{v, w\})(x)$ and $F(w \leq \max\{v, w\})(y)$ therefore also have to be different. Consequently the set $F(\max\{v, w\})$ has more than one element. \square

Indecomposable multifiltrations of sets are therefore exactly the non empty sub-functors of the free functor $\text{mor}_{\mathbb{N}^r}(0, -)$ on one generator given by the origin 0 in \mathbb{N}^r (see 2.7).

Note that since there is a unique map from any set to the one point set, according to 3.4, if $F: \mathbb{N}^r \rightarrow \text{Sets}$ is an indecomposable multifiltration, then, for any $G: \mathbb{N}^r \rightarrow \text{Sets}$, there is at most one natural transformation $G \rightarrow F$. Thus the full subcategory of $\text{Fun}(\mathbb{N}^r, \text{Sets})$ given by the indecomposable multifiltrations is a poset. This is the inclusion poset of all the non empty sub-functors of the free functor $\text{mor}_{\mathbb{N}^r}(0, -)$. Our next goal is to describe this poset. We do that using the notion of the **support** of a functor $F: \mathbb{N}^r \rightarrow \text{Sets}$:

$$\text{supp}(F) := \{v \in \mathbb{N}^r \mid F(v) \neq \emptyset\}$$

For example $\text{supp}(\text{mor}_{\mathbb{N}^r}(v, -)) = \{w \in \mathbb{N}^r \mid v \leq w\}$. Not all subsets of \mathbb{N}^r can be a support. If v belongs to $\text{supp}(F)$, then so does any $w \geq v$. Subsets of \mathbb{N}^r that satisfy this property are called **saturated**.

3.5. Proposition. *The function $(F: \mathbb{N}^r \rightarrow \text{Sets}) \mapsto \text{supp}(F)$ is an isomorphism between the poset of indecomposable multifiltrations of sets and the inclusion poset of saturated non-empty subsets of \mathbb{N}^r .*

Proof. Observe first that if there is a natural transformation $F \rightarrow G$, then if $F(v)$ is not empty, then neither is $G(v)$. This means that $\text{supp}(F) \subset \text{supp}(G)$ which shows that the function $F \mapsto \text{supp}(F)$ is a function of posets.

To define the inverse of the support function, choose a saturated subset S in \mathbb{N}^r and an element v in \mathbb{N}^r . Set:

$$\Psi(S)(v) := \begin{cases} \{v\} & \text{if } v \in S \\ \emptyset & \text{if } v \notin S \end{cases}$$

Since S is saturated, if $\Psi(S)(v)$ is not empty, then neither is $\Psi(S)(w)$ for any $v \leq w$. We can therefore define $\Psi(S)(v \leq w): \Psi(S)(v) \rightarrow \Psi(S)(w)$ to be the unique map. This defines a functor which by Corollary 3.4 is an indecomposable multifiltration. The construction Ψ gives a map of posets between the saturated subsets in \mathbb{N}^r and indecomposable multifiltrations.

Note that $\text{supp}(\Psi(S)) = S$. Furthermore, for any $F: \mathbb{N}^r \rightarrow \text{Sets}$, there is a unique natural transformation $F \rightarrow \Psi(\text{supp}(F))$ which becomes an isomorphism if F is an indecomposable multifiltration. This shows that Ψ is the inverse of the support function. \square

Our next step is to describe the set of saturated subsets of \mathbb{N}^r . For any subset S of \mathbb{N}^r define $\text{gen}(S) := \{v \in S \mid \text{if } w < v, \text{ then } w \notin S\}$ and call it the **minimal set**

of generators of S . For example $\text{gen}(\text{supp}(\text{mor}_{\mathbb{N}^r}(v, -))) = \{v\}$. Furthermore 3.5 implies that an indecomposable multifiltration $F: \mathbb{N}^r \rightarrow \text{Sets}$ is free (necessarily on one generator) if and only if $\text{gen}(\text{supp}(F))$ consists of one element. This can be generalised to arbitrary multifiltrations:

3.6. Proposition. *A multifiltration $F: \mathbb{N}^r \rightarrow \text{Sets}$ is free if and only if it is one critical (see 2.6).*

Proof. We have a decomposition $F = \coprod_{x \in \text{colim } F} F[x]$. Note that $\text{supp}(F[x]) = \{v \in \mathbb{N}^r \mid x \text{ is in the image of } p_v: F(v) \rightarrow \text{colim } F\}$. Thus by definition, F is one critical if and only if $\text{gen}(\text{supp}(F[x]))$ are one element sets, i.e., if the functors $F[x]$ are free on one generator, for every x in $\text{colim } F$. \square

Directly from the definition of the minimal set of generators it follows that: (1) elements in $\text{gen}(S)$ are not comparable; (2) any element in S is comparable to some element in $\text{gen}(S)$. This first property implies $\text{gen}(S)$ is finite, since:

3.7. Lemma. *If S is an infinite subset in \mathbb{N}^r , then it contains an infinite chain, i.e., a sequence of the form $v_1 < v_2 < \dots$.*

Proof. We argue by induction on r . The case $r = 1$ is clear since \mathbb{N} is totally ordered. Assume $r > 1$. Consider the projection onto the last $r - 1$ components $\text{pr}: \mathbb{N}^r \rightarrow \mathbb{N}^{r-1}$. If the image $\text{pr}(S)$ is finite, then for some v in \mathbb{N}^{r-1} the intersection $S \cap \text{pr}^{-1}(v)$ is infinite so it contains an infinite chain as it can be identified with a subset of \mathbb{N} . Assume $\text{pr}(S)$ is infinite. By induction, it contains an infinite chain $v_1 < v_2 < \dots$. It follows that there is a sequence of elements in S of the form $(a_1, v_1), (a_2, v_2), \dots$. Define i_1 to be an index for which $a_{i_1} = \min\{a_1, a_2, \dots\}$ and set $x_1 := (a_{i_1}, v_{i_1})$. Define i_2 to be an index for which $a_{i_2} = \min\{a_j \mid j > i_1\}$ and set $x_2 := (a_{i_2}, v_{i_2})$. Note that $x_1 < x_2$. Continue by induction to obtain a chain $x_1 < x_2 < \dots$ in S . \square

3.8. Proposition. *The function $S \mapsto \text{gen}(S)$ is a bijection between the set of saturated subsets of \mathbb{N}^r and the set of all finite subsets of \mathbb{N}^r whose elements are not comparable.*

Proof. For a subset T in \mathbb{N}^r , define:

$$\text{sat}(T) := \{v \mid \text{there is } u \text{ in } T \text{ such that } v \geq u\}$$

We are going to prove that the function $T \mapsto \text{sat}(T)$ is the inverse to $S \mapsto \text{gen}(S)$. Since any element in S is comparable to some element in $\text{gen}(S)$, it follows that $S \subset \text{sat}(\text{gen}(S))$. In the case S is saturated, $\text{sat}(\text{gen}(S)) \subset S$ and hence these two sets are equal.

Consider an element v in $\text{gen}(\text{sat}(T))$. Since v is in $\text{sat}(T)$, $u \leq v$ for some u in T . If $u \neq v$, then by definition of $\text{gen}(\text{sat}(T))$, u could not belong to $\text{sat}(T)$, which is a contradiction. Thus $u = v$ and v belongs to T . This shows the inclusion $\text{gen}(\text{sat}(T)) \subset T$. Assume T consists of non-comparable elements. Let v be in T and $w < v$. Then w can not belong to $\text{sat}(T)$, otherwise, for some u in T , $u \leq w$ and we would have two comparable elements v and u in T . It follows that v belongs to $\text{gen}(\text{sat}(T))$. We can conclude that $T \subset \text{gen}(\text{sat}(T))$ and hence these two sets are equal. \square

3.9. Corollary. *Let R be a commutative ring with a unit. The poset of indecomposable multifiltrations of sets is isomorphic to the inclusion poset of monomial ideals in $R[x_1, \dots, x_r]$.*

Proof. Let $F: \mathbb{N}^r \rightarrow \text{Sets}$ be a functor. Define $\Psi(F)$ to be the monomial ideal in $R[x_1, \dots, x_r]$ given by:

$$\Psi(F) := \langle x^v \mid v \in \text{gen}(\text{supp}(F)) \rangle$$

If there is a natural transformation $F \rightarrow G$, then $\text{supp}(F) \subset \text{supp}(G)$. We claim that in this case there is an inclusion:

$$\Psi(F) = \langle x^v \mid v \in \text{gen}(\text{supp}(F)) \rangle \subset \langle x^v \mid v \in \text{gen}(\text{supp}(G)) \rangle = \Psi(G)$$

To see this let v be in $\text{gen}(\text{supp}(F))$. We show that there is u in $\text{gen}(\text{supp}(G))$ such that $u \leq v$. That would imply x^v is divisible by x^u proving the claim. If v belongs to $\text{gen}(\text{supp}(G))$ there is nothing to prove. Assume that this is not the case. Since v belongs to $\text{supp}(G)$, there is u in $\text{supp}(G)$ for which indeed $u \leq v$. In this way Ψ defines a functor from $\text{Fun}(\mathbb{N}^r, \text{Sets})$ to the inclusion poset of monomial ideals in $R[x_1, \dots, x_r]$. The restriction of Ψ to indecomposable multifiltrations is a function of posets.

On the other hand let I be a monomial ideal in $R[x_1, \dots, x_r]$, consider the set $S_I := \{v \in \mathbb{N}^r \mid x^v \in I\}$. This is a saturated subset of \mathbb{N}^r because if $u \leq v$ and x^u is in I then x^v must also be in I . We define $\Phi(I)$ to be the indecomposable multifiltration associated to S_I (see 3.5). If there is an inclusion of ideals $I \subseteq J$, then $S_I \subseteq S_J$ and again by 3.5 we have an inclusion $\Phi(I) \subseteq \Phi(J)$. In this way we obtain a functor Φ between the poset of monomial ideals to the poset of indecomposable multifiltrations. Given a functor $F: \mathbb{N}^r \rightarrow \text{Sets}$ there is a unique natural transformation $F \rightarrow \Phi(\Psi(F))$ and this is an isomorphism if F is an indecomposable multifiltration as both of these functors have the same support. If I is a monomial ideal in $R[x_1, \dots, x_r]$ then it is also immediate to verify that $\Psi(\Phi(I)) = I$. \square

According to Propositions 3.5 and 3.8 the function $F \mapsto \text{gen}(\text{supp}(F))$ is a bijection between the set of indecomposable multifiltrations of sets and finite non-empty subsets of \mathbb{N}^r whose elements are not comparable. We finish this section with giving a constructive formula for the inverse to this function. Let T be a subset of \mathbb{N}^r . Define $F_T: \mathbb{N}^r \rightarrow \text{Sets}$ to be a functor given by the following coequalizer in $\text{Fun}(\mathbb{N}^r, \text{Sets})$:

$$F_T := \text{colim} \left(\coprod_{v_0 \neq v_1 \in T} \text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -) \begin{array}{c} \xrightarrow{\pi_0} \\ \xrightarrow{\pi_1} \end{array} \coprod_{v \in T} \text{mor}_{\mathbb{N}^r}(v, -) \right)$$

where on the component indexed by $v_0 \neq v_1 \in T$, the map π_i , is given by the unique natural transformation $\text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -) \rightarrow \text{mor}_{\mathbb{N}^r}(v_i, -)$ induced by $v_i \leq \max\{v_0, v_1\}$.

3.10. Proposition. *If $T \subset \mathbb{N}^r$ is not empty, then the functor F_T is an indecomposable multifiltration whose support is given by $\text{sat}(T)$.*

Proof. Let u be an element in \mathbb{N}^r . The set $F_T(u)$ is a quotient of $\coprod_{v \in T} \text{mor}_{\mathbb{N}^r}(v, u)$ and hence $F_T(u) \neq \emptyset$ if and only if $\coprod_{v \in T} \text{mor}_{\mathbb{N}^r}(v, u) \neq \emptyset$, implying the equality $\text{supp}(F_T) = \text{sat}(T)$. In particular if T is non-empty, then neither is $\text{supp}(F_T)$.

Let $v_0 \leq u$ and $v_1 \leq u$ be two different elements in $\coprod_{v \in T} \text{mor}_{\mathbb{N}^r}(v, u)$. These inequalities give an element $\max\{v_0, v_1\} \leq u$ in $\coprod_{v_0 \neq v_1 \in T} \text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -)$ which is mapped via π_i to $v_i \leq u$. The elements $v_0 \leq u$ and $v_1 \leq u$ are therefore

sent, via the quotient map, to the same element in $F_T(u)$. The set $F_T(u)$ can therefore have at most one element and hence, according to 3.4, F_T is an indecomposable multifiltration. \square

3.11. Corollary. *If F is an indecomposable multifiltration, then it is isomorphic to $F_{\text{gen}(\text{supp}(F))}$.*

We can use the above construction to give a presentation of any multifiltration. Here is a procedure of how to do that. Let $F: \mathbb{N}^r \rightarrow \text{Sets}$ be a multifiltration. For any v in \mathbb{N}^r , index elements of $F(v)$ by elements of $\text{colim } F$ as follows: y in $F(v)$ has index x in $\text{colim } F$ if $p_v(y) = x$. Let $F[x]$ be the subfunctor of F whose elements have index $x \in \text{colim } F$ (see the proof of 3.1). It is an indecomposable multifiltration. Recall that $F = \coprod_{x \in \text{colim } F} F[x]$. The functor F is then isomorphic to:

$$\coprod_{x \in \text{colim } F} F_{\text{gen}(\text{supp}(F[x]))}$$

Since we are going to use this presentation, we need to introduce notation describing the involved functors.

- For any x in $\text{colim } F$, define:

$$\mathcal{G}F[x] := \coprod_{v \in \text{gen}(\text{supp}(F[x]))} \text{mor}_{\mathbb{N}^r}(v, -)$$

$$\mathcal{K}F[x] := \coprod_{v_0 \neq v_1 \in \text{gen}(\text{supp}(F[x]))} \text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -)$$

- Recall that there are natural transformations $\pi_0[x], \pi_1[x]: \mathcal{K}F[x] \rightarrow \mathcal{G}F[x]$ induced by $v_0 \leq \max\{v_0, v_1\}$ and $v_1 \leq \max\{v_0, v_1\}$.
- Since $F[x]$ is indecomposable, there is a unique natural transformation denoted by $p_{F,x}: \mathcal{G}F[x] \rightarrow F[x]$. This natural transformation has the universal property describing $F[x]$ as the colimit of the diagram:

$$\mathcal{K}F[x] \begin{array}{c} \xrightarrow{\pi_0[x]} \\ \xrightarrow{\pi_1[x]} \end{array} \mathcal{G}F[x]$$

By summing over all x in $\text{colim } F$, we obtain functors $\mathcal{G}F := \coprod_{x \in \text{colim } F} \mathcal{G}F[x]$, $\mathcal{K}F := \coprod_{x \in \text{colim } F} \mathcal{K}F[x]$ and natural transformations $\pi_0, \pi_1: \mathcal{K}F \rightarrow \mathcal{G}F$ and $p_F := \coprod_{x \in \text{colim } F} p_{F,x}: \mathcal{G}F \rightarrow F$. The natural transformation p_F has the universal property describing F as the colimit of the diagram:

$$\mathcal{K}F \begin{array}{c} \xrightarrow{\pi_0} \\ \xrightarrow{\pi_1} \end{array} \mathcal{G}F$$

Although the natural transformations $p_{F,x}$ are unique, the construction $F \mapsto \mathcal{G}F$ is not functorial. Nevertheless we attempt to define it also for a natural transformation $\alpha: F \rightarrow G$. Consider the map of sets $\text{colim } \alpha: \text{colim } F \rightarrow \text{colim } G$. Since for any v in \mathbb{N}^r , the following square commutes, we get an inclusion $\alpha(F[x]) \subseteq G[\text{colim } \alpha(x)]$

$$\begin{array}{ccc} F(v) & \xrightarrow{\alpha(v)} & G(v) \\ p_v \downarrow & & \downarrow p_v \\ \text{colim } F & \xrightarrow{\text{colim } \alpha} & \text{colim } G \end{array}$$

It follows that the set $\{w \in \text{gen}(\text{supp}(G[\text{colim } \alpha(x)])) \mid w \leq v\}$ is not empty for any v in $\text{gen}(\text{supp}(F[x]))$. We can order this set using the lexicographical order and define $w_{\alpha,x,v}$ to be the smallest element of this set. Since $w_{\alpha,x,v} \leq v$, there is a unique natural transformation $\text{mor}_{\mathbb{N}^r}(v, -) \rightarrow \text{mor}_{\mathbb{N}^r}(w_{\alpha,x,v}, -)$. Define $\bar{\alpha}: \mathcal{G}F \rightarrow \mathcal{G}G$ to be the natural transformation which on the summand $\text{mor}_{\mathbb{N}^r}(v, -)$ indexed by x in $\text{colim } F$ and v in $\text{gen}(\text{supp}(F[x]))$ is given by the composition of $\text{mor}_{\mathbb{N}^r}(v, -) \rightarrow \text{mor}_{\mathbb{N}^r}(w_{\alpha,x,v}, -)$ and the inclusion into $\mathcal{G}G$ of the summand $\text{mor}_{\mathbb{N}^r}(w_{\alpha,x,v}, -)$ indexed by $\text{colim } \alpha(x)$ in $\text{colim } G$ and $w_{\alpha,x,v}$ in $\text{gen}(\text{supp}(G[\text{colim } \alpha(x)]))$. Because of these choices we obtain a commutative diagram of natural transformations:

$$\begin{array}{ccc} \mathcal{G}F & \xrightarrow{\bar{\alpha}} & \mathcal{G}G \\ p_F \downarrow & & \downarrow p_G \\ F & \xrightarrow{\alpha} & G \end{array}$$

Explicitly:

$$\begin{array}{ccc} \text{mor}_{\mathbb{N}^r}(v, -) & \xrightarrow{\quad} & \text{mor}_{\mathbb{N}^r}(w_{\alpha,x,v}, -) \\ \downarrow \text{summand indexed by } x \text{ and } v & & \downarrow \text{summand indexed by } \text{colim } \alpha(x) \text{ and } w_{\alpha,x,v} \\ \coprod_{x \in \text{colim } F} \coprod_{v \in \text{gen}(\text{supp}(F[x]))} \text{mor}_{\mathbb{N}^r}(v, -) & \xrightarrow{\bar{\alpha}} & \coprod_{x \in \text{colim } G} \coprod_{v \in \text{gen}(\text{supp}(G[x]))} \text{mor}_{\mathbb{N}^r}(v, -) \\ p_F \downarrow & & \downarrow p_G \\ \coprod_{x \in \text{colim } F} F[x] & \xrightarrow{\alpha} & \coprod_{x \in \text{colim } G} G[x] \end{array}$$

It is important to point out that the assignment $(\alpha: F \rightarrow G) \mapsto (\bar{\alpha}: \mathcal{G}F \rightarrow \mathcal{G}G)$ is not a functor. It is not true in general that $\overline{\beta\alpha}$ equals $\bar{\beta}\bar{\alpha}$.

4. SET VALUED VS. R -MOD VALUED FUNCTORS

Let R be a commutative ring with identity. Recall that we identify the category of functors $\text{Fun}(\mathbb{N}^r, R\text{-Mod})$ with the category of \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules by assigning to $F: \mathbb{N}^r \rightarrow R\text{-Mod}$ the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module given by $\mathbf{F} = \bigoplus_{v \in \mathbb{N}^r} F(v)$ (see 2.8). Via the above identification the free functor $R\text{mor}_{\mathbb{N}^r}(0, -)$ (see 2.7) is mapped to the module $R[x_1, \dots, x_r]$. Thus sub-functors of $R\text{mor}_{\mathbb{N}^r}(0, -)$ are identified with \mathbb{N}^r -graded ideals in $R[x_1, \dots, x_r]$. Among these sub-functors there are the R -spans of indecomposable multifiltrations of sets and among the \mathbb{N}^r -graded ideals in $R[x_1, \dots, x_r]$ there are the monomial ideals. Note that for an indecomposable multifiltration of sets $F: \mathbb{N}^r \rightarrow \text{Sets}$, the \mathbb{N}^r -graded ideal $\mathbf{R}F \subset R[x_1, \dots, x_r]$ coincides with the monomial ideal $\Psi(F)$ given in the proof of Corollary 3.9. It thus follows from this corollary that the sub-functors of $R\text{mor}_{\mathbb{N}^r}(0, -)$ that are identified with monomial ideals are exactly the R -spans of indecomposable multifiltrations of sets. Since monomial ideals are indecomposable $R[x_1, \dots, x_r]$ -modules, then so are the R -spans of indecomposable multifiltrations of sets. These are the easiest indecomposable multifiltrations of R -modules. The following is a key fact about their finite sums:

4.1. Proposition. *Let $\{F_i: \mathbb{N}^r \rightarrow \text{Sets}\}_{1 \leq i \leq n}$ and $\{G_j: \mathbb{N}^r \rightarrow \text{Sets}\}_{1 \leq j \leq m}$ be two finite families of indecomposable multifiltrations of sets. If $\bigoplus_{i=1}^n RF_i: \mathbb{N}^r \rightarrow R\text{-Mod}$ and $\bigoplus_{j=1}^m RG_j: \mathbb{N}^r \rightarrow R\text{-Mod}$ are isomorphic as functors with values in $R\text{-Mod}$, then $n = m$, and there is a permutation σ of $\{1, \dots, n\}$ for which $F_i: \mathbb{N}^r \rightarrow \text{Sets}$ and $G_{\sigma(i)}: \mathbb{N}^r \rightarrow \text{Sets}$ are isomorphic for any i .*

Proof. First note that if $F, G: \mathbb{N}^r \rightarrow \text{Sets}$ are indecomposable multifiltrations, then the map $R\text{Nat}_{\text{Sets}}(F, G) \rightarrow \text{Nat}_{R\text{-Mod}}(RF, RG)$, induced by the R -span functor, is an isomorphism of R modules (this is not true if F is a multifiltration but not indecomposable). Consequently the R module $\text{Nat}_{R\text{-Mod}}(RF, RG)$ is isomorphic to R if $\text{supp}(F) \subset \text{supp}(G)$ or it is trivial if $\text{supp}(F) \not\subset \text{supp}(G)$.

We proceed by induction on n to prove the proposition. Assume $n = 1$. Since RF and $\bigoplus_{j=1}^m RG_j$ are isomorphic, then so are their colimits which as R modules are isomorphic to respectively R and $\bigoplus_{j=1}^m R$. For commutative rings the rank of a free module is a well define invariant and hence $m = 1$. The functors RF and RG_1 are therefore isomorphic and by the discussion above $\text{supp}(F)$ and $\text{supp}(G)$ are the same subsets of \mathbb{N}^r . We can then use 3.5 to get F and G are isomorphic.

Assume $n > 1$. Consider the subsets $\text{supp}(F_i) \subset \mathbb{N}^r$ for $1 \leq i \leq n$ and choose among them a maximal one T with respect to the inclusion. By permuting we can assume that:

$$\text{supp}(F_i) = T, \text{ if } 1 \leq i \leq n' \quad \text{and} \quad \text{supp}(F_i) \neq T, \text{ if } n' < i \leq n$$

Let $\phi: \bigoplus_{i=1}^n RF_i \rightarrow \bigoplus_{j=1}^m RG_j$ and $\psi: \bigoplus_{j=1}^m RG_j \rightarrow \bigoplus_{i=1}^n RF_i$ be inverse isomorphisms. Since the restriction of ϕ to F_1 is non trivial, there is j such that $T = \text{supp}(F_1) \subset \text{supp}(G_j)$. By the same argument, since the restriction of ψ to G_j is not trivial, there is l for which $\text{supp}(G_j) \subset \text{supp}(F_l)$. As we chose T to be a maximal among the supports of F_i 's, we get $l \leq n'$ and $\text{supp}(G_j) = T$. Again by permuting if necessary we can assume that:

$$T = \text{supp}(G_i), \text{ if } 1 \leq i \leq m' \quad \text{and} \quad T \not\subset \text{supp}(G_i), \text{ if } m' < i \leq m$$

This means that ϕ maps the submodule $\bigoplus_{i=1}^{n'} RF_i \subset \bigoplus_{i=1}^n RF_i$ to the submodule $\bigoplus_{j=1}^{m'} RG_j \subset \bigoplus_{j=1}^m RG_j$. Furthermore the restriction of $\phi: \bigoplus_{i=1}^{n'} RF_i \rightarrow \bigoplus_{j=1}^{m'} RG_j$ is an isomorphism whose inverse is given by the restriction of ψ . We therefore get that their colimits $\bigoplus_{i=1}^{n'} R$ and $\bigoplus_{j=1}^{m'} R$ are also isomorphic and hence $n' = m'$. Moreover, by taking the quotients, we obtain an isomorphism between $\bigoplus_{i>n'}^n RF_i$ and $\bigoplus_{j>m'}^m RG_j$. The proposition now follows from the inductive assumption. \square


The above proposition can be restated in the form:

4.2. Corollary.

- (1) *Let $\{I_i\}_{1 \leq i \leq n}$ and $\{J_j\}_{1 \leq j \leq m}$ be monomial ideals in $R[x_1, \dots, x_r]$. If the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ modules $\bigoplus_{i=1}^n I_i$ and $\bigoplus_{j=1}^m J_j$ are isomorphic, then $m = n$ and there is a permutation σ of $\{1, \dots, n\}$ for which $I_i = J_{\sigma(i)}$.*
- (2) *Let $F, G: \mathbb{N}^r \rightarrow \text{Sets}$ be compact multifiltrations (see 2.6). Then F and G are isomorphic if and only if their R -spans $RF, RG: \mathbb{N}^r \rightarrow R\text{-Mod}$ are isomorphic.*

The statement 4.2.(2) is not true if the functors F and G are not multifiltrations:


4.3. **Example.** Let $F_1, F_2: \mathbb{N} \rightarrow \text{Sets}$ be functors with the same values $F_1(0) = F_2(0) = \{a, b, c, d\}$, $F_1(1) = F_2(1) = \{e, f\}$ and $F_1(n) = F_2(n) = \{g\}$ for $n \geq 2$, however with different maps which are given by the following diagrams:



Although the functors F_1 and F_2 are not isomorphic, their R -spans RF_1 and RF_2 are.

The following example illustrates the fact that not all (indecomposable) multifiltrations of R -modules are R -spans of (indecomposable) multifiltrations of sets.


4.4. **Example.** Consider the multifiltration $F: \mathbb{N}^2 \rightarrow R\text{-Mod}$ which on the square $\{v \leq (2, 2)\} \subset \mathbb{N}^2$ is given by the following commutative diagram:



and for w in $\mathbb{N}^2 \setminus \{v \leq (2, 2)\}$, the map $F(\min(w, (2, 2)) \leq w)$ is an isomorphism. Assume further that α, β , and γ are monomorphisms and their images are pairwise different submodules of $R \oplus R$. Then this functor is not isomorphic to the R -span of any functor with values in Sets. Note further that in this case F is an indecomposable multifiltration of R -modules whose colimit is free of rank 2 (compare with 3.1).

Being one critical (see 2.6) for multifiltrations of sets is equivalent to being free (see 3.6). This is not true for multifiltrations of R -modules if $r > 2$:

4.5. **Example.** Consider the multifiltration $F: \mathbb{N}^3 \rightarrow R\text{-Mod}$ which on the cube $\{v \leq (1, 1, 1)\} \subset \mathbb{N}^3$ is given by the following commutative diagram:



and for w in $\mathbb{N}^3 \setminus \{v \leq (1, 1, 1)\}$ the map $F(\min(w, (1, 1, 1)) \leq w)$ is an isomorphism. Then this functor is one critical, it is not free, and it is not the R -span of a multifiltration of sets.

For bifiltrations ($r = 2$) we have the following positive result:

4.6. Proposition. *Assume R is a field. A bifiltration $F: \mathbb{N}^2 \rightarrow R\text{-Mod}$ is free if and only if it is one critical.*

Proof. One implication holds more generally for all r . If $F: \mathbb{N}^r \rightarrow R\text{-Mod}$ is free, it is the R -span of a free functor $G: \mathbb{N}^r \rightarrow \text{Sets}$. Thus F is isomorphic to $\bigoplus_{x \in \text{colim } G} R\text{mor}(v_x, -)$. Since the R -span functor commutes with colimits, we can identify $\text{colim } F$ with $R(\text{colim } G)$. Consider an element $y = \sum_{i=1}^n c_i x_i$ in $\text{colim } F$ where x_i belongs to $\text{colim } G$. Note that:

$$\{v \in \mathbb{N}^2 \mid y \in F(v)\} = \bigcap_{i=1}^n \{v \in \mathbb{N}^2 \mid x_i \in G(v)\}$$

It follows that this set has a unique minimal element given by $\max\{v_{x_i} \mid 1 \leq i \leq n\}$. This shows that F is one critical.

Assume now that $F: \mathbb{N}^2 \rightarrow R\text{-Mod}$ is one critical. To show that it is free it would be enough to prove that it is the R -span of a multifiltration of sets since in this case this multifiltration of sets would be also one critical and therefore free by 3.6. Define $G(0, 0)$ to be a base of $F(0, 0)$. Since $F((0, 0) \leq (1, 0)): F(0, 0) \rightarrow F(1, 0)$ is an inclusion, we can extend that base of $F(0, 0)$ to a base $G(1, 0)$ of $F(1, 0)$. We can proceed by induction on n and define in this way a sequence of sets

$$G(0, 0) \subset G(1, 0) \subset \cdots \subset G(n, 0) \subset \cdots$$

whose R -span gives the functor F restricted to $\mathbb{N} \times \{0\} \subset \mathbb{N}^2$. We continue again by induction. Assume that $k > 1$ and we have constructed a functor:

$$G: \mathbb{N} \times \{v \in \mathbb{N} \mid v < k\} \rightarrow \text{Sets}$$

whose R -span is isomorphic to the restriction of F . By the same argument as before, since $F((0, k-1) \leq (0, k)): F(0, k-1) \rightarrow F(0, k)$ is an inclusion we can extend the base $G(0, k-1)$ of $F(0, k-1)$ to a base $G(0, k)$ of $F(0, k)$. Assume $n > 0$ and that we have defined a functor:

$$G: \mathbb{N} \times \{v \in \mathbb{N} \mid v < k\} \cup \{v \in \mathbb{N} \mid v < n\} \times \{v \in \mathbb{N} \mid v \leq k\} \rightarrow \text{Sets}$$

whose R -span is isomorphic to the restriction of F . Since F is one critical the intersection of the images of $F(n-1, k)$ and $F(n, k-1)$ in $F(n, k)$ coincide with the image of $F(n-1, k-1)$. It follows that the induced map:

$$\text{colim}(F(n, k-1) \hookrightarrow F(n-1, k-1) \hookrightarrow F(n-1, k)) \rightarrow F(n, k)$$

is an inclusion. Here the assumption $r = 2$ is crucial. We can then extend the subset:

$$\text{colim}(G(n, k-1) \hookrightarrow G(n-1, k-1) \hookrightarrow G(n-1, k)) \hookrightarrow F(n, k)$$

to a base $G(n, k)$ of $F(n, k)$. In this way we get a desired functor

$$G: \mathbb{N} \times \{v \in \mathbb{N} \mid v \leq k\} \rightarrow \text{Sets}$$

whose R -span is isomorphic to F . □

We finish this section with a procedure of obtaining a free presentation of the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module \mathbf{RF} associated to the R -span of a multifiltration $F: \mathbb{N}^r \rightarrow \text{Sets}$. In the first 3 steps we recall from the end of Section 3 how to build a presentation of F .

- Decompose F into indecomposable components $\coprod_{x \in \text{colim } F} F[x]$.
- For any x , find the set $T_x := \text{gen}(\text{supp}(F[x]))$.
- Recall that F can be described as the coequalizer of two natural transformations $\pi_0, \pi_1 : \mathcal{K}F \rightarrow \mathcal{G}F$ between free functors. Explicitly F is isomorphic to the colimit of the following diagram:

$$\coprod_{x \in \text{colim } F} \left(\coprod_{v_0 \neq v_1 \in T_x} \text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -) \begin{array}{c} \xrightarrow{\pi_0[x]} \\ \xrightarrow{\pi_1[x]} \end{array} \coprod_{v \in T_x} \text{mor}_{\mathbb{N}^r}(v, -) \right)$$

where on the component indexed by $v_0 \neq v_1 \in T_x$, the map π_i , is given by the unique natural transformation $\text{mor}_{\mathbb{N}^r}(\max\{v_0, v_1\}, -) \rightarrow \text{mor}_{\mathbb{N}^r}(v_i, -)$ induced by $v_i \leq \max\{v_0, v_1\}$.

- Since the R -span functor commutes with colimits, we get that the module $\mathbf{R}F$ is isomorphic to the coequalizer of the following two maps π_0 and π_1 between free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules (see 2.3):

$$\bigoplus_{x \in \text{colim } F} \left(\bigoplus_{v_0 \neq v_1 \in T_x} \langle x^{\max\{v_0, v_1\}} \rangle \begin{array}{c} \xrightarrow{\pi_0[x]} \\ \xrightarrow{\pi_1[x]} \end{array} \bigoplus_{v \in T_x} \langle x^v \rangle \right)$$

where $\pi_i[x]$, on the component indexed by $v_0 \neq v_1 \in T_x$, is given by the inclusion $\langle x^{\max\{v_0, v_1\}} \rangle \hookrightarrow \langle x^{v_i} \rangle$. Thus the columns of the matrix representing $\pi_i[x]$ have all entries zero except one which is one.

- The module $\mathbf{R}F$ is then isomorphic to the cokernel of the difference $\pi_0 - \pi_1$. Note that the columns of the matrix $M(F_x)$ representing $\pi_0 - \pi_1$ are vectors of the form: one entry is 1, one entry is -1 , and all other entries are zero.

To summarize, with a multifiltration $F : \mathbb{N}^r \rightarrow \text{Sets}$ we have associated the following invariants:

- (1) a set $\text{colim } F$;
- (2) for any x in $\text{colim } F$, a finite subset $T_x := \text{gen}(\text{supp}(F[x]))$ of \mathbb{N}^r ;
- (3) for any x in $\text{colim } F$, a $|T_x| \times \binom{|T_x|}{2}$ matrix $M(F_x)$, representing the map $\pi_0[x] - \pi_1[x]$ whose columns are vectors of the form: one entry is 1, one entry is -1 , and all other entries are zero.

These invariants can be used to get the \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -module associated to the R -span $\mathbf{R}F$ as the cokernel of the map:

$$\bigoplus_{x \in \text{colim } F} \left(\bigoplus_{v_0 \neq v_1 \in T_x} \langle x^{\max\{v_0, v_1\}} \rangle \xrightarrow{M(F_x)} \bigoplus_{v \in T_x} \langle x^v \rangle \right)$$

5. FUNCTORS WITH VALUES IN Spaces

Let $F : \mathbb{N}^r \rightarrow \text{Spaces}$ be a multifiltration of simplicial complexes, $X := \text{colim } F$, and R a commutative ring with identity. Let us choose an ordering on the set of vertices of X . Since F is a multifiltration, we can restrict this ordering to the set of vertices of $F(v)$, for any v in \mathbb{N}^r . In this way the maps $F(v \leq w)$ are order preserving and we can form a functor of ordered n -simplices to get a multifiltration of sets $F_n : \mathbb{N}^r \rightarrow \text{Sets}$ (see 2.2) which assigns to any v in \mathbb{N}^r the set $F(v)_n$ of ordered n -simplices in $F(v)$. These functors, for various n 's, are connected via the natural transformations given by the maps $d_i : F_{n+1}(v) \rightarrow F_n(v)$ which forget the i -th element of an ordered simplex (see 2.2). By applying the R -span functor and

taking the alternating sum of the induced maps we obtain a diagram of natural transformations in $\text{Fun}(\mathbb{N}^r, R\text{-Mod})$:

$$RF_{n+1} \xrightarrow{\partial_{n+1} := \sum_{i=0}^{n+1} (-1)^i d_i} RF_n \xrightarrow{\partial_n := \sum_{i=0}^n (-1)^i d_i} RF_{n-1}$$

The composition of these maps is trivial and hence we can form a homology functor $H_n(F, R): \mathbb{N}^r \rightarrow R\text{-Mod}$ which in general may not be a multifiltration. This could be done in two stages. First we could take the cokernel of the first differential $\text{coker}(\partial_{n+1}: RF_{n+1} \rightarrow RF_n)$ and then the kernel of the induced map $\partial_n: \text{coker}(\partial_{n+1}) \rightarrow RF_{n-1}$ or we could take the kernel of the second differential $\text{ker}(\partial_n: RF_n \rightarrow RF_{n-1})$ and then the cokernel of the induced map $\partial_{n+1}: RF_{n+1} \rightarrow \text{ker}(\partial_n)$. Let us consider the case of $n = 0$. Recall that since RF_{-1} is assumed to be the trivial functor (see 2.2), $H_0(F, R)$ is given by the cokernel $\text{coker}(d_0 - d_1: RF_1 \rightarrow RF_0)$. This cokernel is simply the coequalizer of the two maps $d_0, d_1: RF_1 \rightarrow RF_0$. As the R -span functor commutes with colimits, we then get an isomorphism between $H_0(F, R)$ and the R -span of the following functor with values in the category of sets:

$$\text{colim} \left(\begin{array}{ccc} & & \\ & \xrightarrow{d_0} & \\ F_1 & & F_0 \\ & \xrightarrow{d_1} & \end{array} \right)$$

This is a special property of the 0-th homology. If $n \geq 1$, then it is not true in general that the functors $H_n(F, R)$, $\text{coker}(\partial_{n+1}: RF_{n+1} \rightarrow RF_n)$, and $\text{ker}(\partial_n: RF_n \rightarrow RF_{n-1})$ are R -spans of functors with values in the category of sets, even if R is a field as the following example illustrates:

5.1. Example. Consider the two multifiltrations of spaces $F, G: \mathbb{N}^2 \rightarrow \text{Spaces}$ which on the square $\{v \leq (2, 2)\} \subset \mathbb{N}^2$ are described in Figure 1 and for w in $\mathbb{N}^2 \setminus \{v \leq (2, 2)\}$, the maps induced by $(\min\{w_1, 2\}, \min\{w_2, 2\}) \leq w$ are the identities.




FIGURE 1. multifiltrations with values in Spaces

In the multifiltration $F: \mathbb{N}^r \rightarrow \text{Spaces}$ there are no 2-simplices and hence $H_1(F, R) = \text{ker}(\partial_1: RF_1 \rightarrow RF_0)$. On the square $\{v \leq (2, 2)\} \subset \mathbb{N}^2$, the functors $\text{ker}(\partial_1: RF_1 \rightarrow RF_0)$ and $\text{coker}(\partial_2: RG_2 \rightarrow RG_1)$ are given respectively by

the diagrams:

$$\begin{array}{ccc}
R \xrightarrow{\begin{pmatrix} 1 \\ 1 \end{pmatrix}} R \oplus R \xrightarrow{\text{id}} R \oplus R & R \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} R \oplus R \xrightarrow{\text{id}} R \oplus R \\
\uparrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \uparrow \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad \uparrow \text{id} & \uparrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \uparrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} \quad \uparrow \text{id} \\
0 \xrightarrow{\quad} R \xrightarrow{\begin{pmatrix} 1 \\ 0 \end{pmatrix}} R \oplus R & 0 \xrightarrow{\quad} R \xrightarrow{\begin{pmatrix} 0 \\ 1 \end{pmatrix}} R \oplus R \\
\uparrow \quad \uparrow \begin{pmatrix} 0 \\ 1 \end{pmatrix} & \uparrow \quad \uparrow \begin{pmatrix} -1 \\ -1 \end{pmatrix} \\
0 \xrightarrow{\quad} 0 \xrightarrow{\quad} R & 0 \xrightarrow{\quad} 0 \xrightarrow{\quad} R
\end{array} ,$$

By Example 4.3 both of the functors are not the R -span of any multifiltration of sets.

We now assume that $F : \mathbb{N}^r \rightarrow \text{Spaces}$ is a compact multifiltration of spaces. It follows that $X = \text{colim } F$ is a finite complex. Since in general the functor $H_n(F, R)$ is not the R -span of a multifiltration of sets we cannot directly use the construction in Section 4 to compute a free presentation of the module $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$. Instead our goal is to describe the module $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$ in such a way that one can use very efficiently standard commutative algebra software or an algorithm presented in [2] which often is faster. As it was pointed out in [2] this efficiency is a consequence of homogeneity and the fact that matrices involved are very simple. We proceed as follows:

- (1) Consider the decomposition $F_{n-1} = \coprod_{\sigma \in X_{n-1}} F_{n-1}[\sigma]$ (see 3.2). Define $D_{n-1} := \coprod_{\sigma \in X_{n-1}} \text{mor}_{\mathbb{N}^r}(0, -)$ and $\phi : F_{n-1} \rightarrow D_{n-1}$ to be the coproduct of the unique inclusions $\coprod_{\sigma \in X_{n-1}} (F_{n-1}[\sigma] \hookrightarrow \text{mor}_{\mathbb{N}^r}(0, -))$. Note that D_{n-1} is a free functor.
- (2) Information about F together with the presentations and natural transformations given at the end of Section 3 and in step (1) can be organized into the following commutative diagrams for any $0 \leq i \leq n+1$ and $0 \leq j \leq n$:

$$\begin{array}{ccccc}
\mathcal{G}F_{n+1} & \xrightarrow{p_{F_{n+1}}} & F_{n+1} & & \\
\downarrow \bar{d}_i & & \downarrow d_i & & \\
\mathcal{K}F_n \xrightarrow[\pi_1]{\pi_0} \mathcal{G}F_n & \xrightarrow{p_{F_n}} & F_n & & \\
\downarrow \alpha_j & & \downarrow d_j & & \\
D_{n-1} & \xleftarrow{\phi} & F_{n-1} & &
\end{array}$$

- (3) This leads to the following natural transformations:

$$\begin{array}{ccc}
\mathcal{G}F_{n+1} & & \\
\downarrow \bar{d}_0 \quad \cdots \quad \downarrow \bar{d}_{n+1} & & \\
\mathcal{K}F_n \xrightarrow[\pi_1]{\pi_0} \mathcal{G}F_n & & \\
\downarrow \alpha_0 \quad \cdots \quad \downarrow \alpha_n & & \\
D_{n-1} & &
\end{array}$$

- (4) By applying the R -span functor and additivity we get two homomorphisms of \mathbb{N}^r -graded free $R[x_1, \dots, x_r]$ -modules:

$$\mathbf{RK}F_n \oplus \mathbf{RG}F_{n+1} \xrightarrow{[\pi_0 - \pi_1 \quad \sum_{i=0}^{n+1} (-1)^i \bar{\mathbf{d}}_i]} \mathbf{RG}F_n \xrightarrow{\sum_{j=0}^n (-1)^j \alpha_j} \mathbf{RD}_{n-1}$$

5.2. Proposition. *The composition of the above homomorphisms is trivial and the homology of this complex is isomorphic to $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$.*

Proof. Consider the complex whose homology is $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$:

$$\mathbf{RF}_{n+1} \xrightarrow{\sum_{i=0}^{n+1} (-1)^i \mathbf{d}_i} \mathbf{RF}_n \xrightarrow{\sum_{i=0}^n (-1)^i \mathbf{d}_i} \mathbf{RF}_{n-1}$$

Since $\phi: F_{n-1} \hookrightarrow D_{n-1}$ is an inclusion and $p_{F_{n+1}}: \mathcal{G}F_{n+1} \rightarrow F_{n+1}$ is a surjection, the bottom row of the following commutative diagram is also a complex whose homology is isomorphic to $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$:

$$\begin{array}{ccccc} \mathbf{RK}F_n \oplus \mathbf{RG}F_{n+1} & \xrightarrow{[\pi_0 - \pi_1 \quad \sum_{i=0}^{n+1} (-1)^i \bar{\mathbf{d}}_i]} & \mathbf{RG}F_n & \xrightarrow{\sum_{j=0}^n (-1)^j \alpha_j} & \mathbf{RD}_{n-1} \\ \downarrow \text{projection} & & \downarrow p_{F_n} & & \parallel \\ \mathcal{G}F_{n+1} & \xrightarrow{\sum_{i=0}^{n+1} (-1)^i \mathbf{d}_i p_{F_{n+1}}} & \mathbf{RF}_n & \xrightarrow{\sum_{i=0}^n (-1)^i \phi \mathbf{d}_i} & \mathbf{RD}_{n-1} \end{array}$$

Recall that \mathbf{RF}_n is the cokernel of the map $\pi_0 - \pi_1: \mathbf{RK}F_n \rightarrow \mathbf{RG}F_n$. This implies the top row of the above diagram is also a complex whose homology is isomorphic to $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$ proving the proposition. \square

An important fact is that the above sequence of free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules that computes $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$ can be easily and explicitly described in terms of the original multifiltration of spaces. Here are the involved modules:

$$\begin{aligned} \mathbf{RK}F_n &= \bigoplus_{\sigma \in X_n} \bigoplus_{v_0 \neq v_1 \in \text{gen}(\sigma)} \langle x^{\max\{v_0, v_1\}} \rangle \\ \mathbf{RG}F_n &= \bigoplus_{\sigma \in X_n} \bigoplus_{v \in \text{gen}(\sigma)} \langle x^v \rangle \\ \mathbf{RD}_{n-1} &= \bigoplus_{\sigma \in X_{n-1}} R[x_1, \dots, x_r] \end{aligned}$$

and here is how to find the matrices associated to the maps in this sequence (see 2.3 for our convention to describe homomorphisms between free \mathbb{N}^r -graded $R[x_1, \dots, x_r]$ -modules).

- Let σ be a simplex in X_n or X_{n+1} . Consider the set $\{v \in \mathbb{N}^r \mid \sigma \in F(v)\}$. This is a saturated set and hence admits a finite minimal set of generators which we denote by $\text{gen}(\sigma)$. This set coincides with $\text{gen}(\text{supp}(F[\sigma]))$ and its elements are exactly the minimal elements of the set $\{v \in \mathbb{N}^r \mid \sigma \in F(v)\}$.
- The matrix $[\pi_0 - \pi_1 \quad \sum_{i=0}^{n+1} (-1)^i \bar{\mathbf{d}}_i]$ is a concatenation of two matrices one for $\pi_0 - \pi_1$ and one for $\sum_{i=0}^{n+1} (-1)^i \bar{\mathbf{d}}_i$.
- The matrix for $\pi_0 - \pi_1$ is a block diagonal. The blocks are indexed by simplices in X_n and the block corresponding to σ in X_n is of the size $|\text{gen}(\sigma)| \times \binom{|\text{gen}(\sigma)|}{2}$. The entry in this block indexed by v in $\text{gen}(\sigma)$ and $v_0 \neq v_1$ in $\binom{\text{gen}(\sigma)}{2}$ has row grade v and column grade $\max\{v_0, v_1\}$. Its value is 1 if $v = v_0$, -1 if $v = v_1$, and 0 otherwise.

- The rows of the matrix for $\sum_{i=0}^{n+1} (-1)^i \overline{\mathbf{d}}_i$ are indexed in the same way and have the same grades as the rows of the matrix for $\pi_0 - \pi_1$. The columns of the matrix for $\sum_{i=0}^{n+1} (-1)^i \overline{\mathbf{d}}_i$ are divided into blocks indexed by simplices in X_{n+1} . The columns in the block corresponding to σ in X_{n+1} are indexed by $\text{gen}(\sigma)$. The corresponding element in $\text{gen}(\sigma)$ is the grade of the column. Each column has exactly $n + 2$ non-zero entries which are either 1 or -1 . For a column indexed by v in $\text{gen}(\sigma)$, the non-zero entries occur in the row blocks corresponding to the simplices $d_i(\sigma)$. In each such block there is only one non-zero entry and is equal to $(-1)^i$ and occurs in the row corresponding to the minimal element with respect to the lexicographical order in the set $\{w \in \text{gen}(d_i(\sigma)) \mid w \leq v\}$.
- The matrix for $\sum_{j=0}^n (-1)^j \alpha_j$ has rows indexed by simplices in X_{n-1} . All the rows have grade 0. The columns are divided into blocks indexed by simplices in X_n . The columns in the block corresponding to σ in X_n are indexed by $\text{gen}(\sigma)$. The corresponding element in $\text{gen}(\sigma)$ is the grade of the column. The entry in this matrix in the row indexed by τ in X_{n-1} and the column indexed by v in $\text{gen}(\sigma)$ for σ in X_n has value $(-1)^i$ if $\tau = d_i(\sigma)$ and 0 otherwise. Note that in any row, the entries in the same column block have the same value but different grades.

We will now show our procedure to compute the module $H_1(F, R)$ with an example.

5.3. Example. Consider the multifiltration $F : \mathbb{N}^2 \rightarrow \text{Spaces}$ which on the square $\{v \leq (2, 2)\} \subset \mathbb{N}^2$ is described in Figure 2 and for w in $\mathbb{N}^2 \setminus \{v \leq (2, 2)\}$, the maps induced by $\min(w, (2, 2)) \leq w$ are the identities. The simplicial complex $X = \text{colim } F$ is given by the complex $F(2, 2)$ and we choose an ordering of its vertices as indicated also in Figure 2.




FIGURE 2

The functors $H_1(F, R) : \mathbb{N}^2 \rightarrow R\text{-Mod}$ on the square $\{v \leq (2, 2)\} \subset \mathbb{N}^2$ is given by the following commutative diagram:

$$\begin{array}{ccccc}
 R & \xrightarrow{1} & R & \xrightarrow{1} & R \\
 \uparrow & & \uparrow_0 & & \uparrow_1 \\
 0 & \longrightarrow & R & \xrightarrow{0} & R \\
 \uparrow & & \uparrow & & \uparrow_1 \\
 0 & \longrightarrow & 0 & \longrightarrow & R
 \end{array}$$

We will now go through the steps presented above and construct the elements needed in Proposition 3.11 to compute $H_1(\mathbf{F}, \mathbf{R})$:

- $X_0 = \{0, 1, 2, 3\}$, $X_1 = \{0 < 1, 0 < 2, 1 < 2, 1 < 3, 2 < 3\}$, $X_2 = \{1 < 2 < 3\}$, and $X_n = \emptyset$ for $n \geq 3$.
- For an ordered simplex σ in X , the minimal set of generators $\text{gen}(\sigma)$, ordered by the lexicographical order, is given by the tables:

σ	0	1	2	3
$\text{gen}(\sigma)$	(0,2) (2,0)	(0,2) (1,1) (2,0)	(0,2) (1,1) (2,0)	(0,2) (1,1)

σ	0 < 1	0 < 2	1 < 2	1 < 3	2 < 3
$\text{gen}(\sigma)$	(0,2) (2,0)	(0,2) (2,0)	(1,1) (2,0)	(0,2) (1,1)	(0,2) (1,1)

σ	1 < 2 < 3
$\text{gen}(\sigma)$	(1,2) (2,1)

- We thus have:

$\mathbf{RK}\mathbf{F}_1$	$2\langle x^{(1,2)} \rangle \oplus \langle x^{(2,1)} \rangle \oplus 2\langle x^{(2,2)} \rangle$
$\mathbf{RG}\mathbf{F}_1$	$4\langle x^{(0,2)} \rangle \oplus 3\langle x^{(1,1)} \rangle \oplus 3\langle x^{(2,0)} \rangle$
$\mathbf{RG}\mathbf{F}_2$	$\langle x^{(1,2)} \rangle \oplus \langle x^{(2,1)} \rangle$
\mathbf{RD}_0	$4R[x_1, x_2]$

- The matrix associated to $\pi_0 - \pi_1 : \mathbf{RK}\mathbf{F}_1 \rightarrow \mathbf{RG}\mathbf{F}_1$ with the block decomposition and the column and row grades is given by:

$$\begin{array}{c}
\begin{array}{c} 0 < 1 \\ 0 < 2 \\ 1 < 2 \\ 1 < 3 \\ 2 < 3 \end{array}
\begin{array}{c} (0,2) \\ (2,0) \\ (0,2) \\ (2,0) \\ (1,1) \\ (2,0) \\ (0,2) \\ (1,1) \\ (0,2) \\ (1,1) \end{array}
\left(\begin{array}{c|c|c|c|c}
\begin{array}{c} 0 < 1 \\ (2,2) \end{array} & \begin{array}{c} 0 < 2 \\ (2,2) \end{array} & \begin{array}{c} 1 < 2 \\ (2,1) \end{array} & \begin{array}{c} 1 < 3 \\ (1,2) \end{array} & \begin{array}{c} 2 < 3 \\ (1,2) \end{array} \\
\hline
\begin{array}{c} 1 \\ -1 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} \\
\hline
\begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ -1 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} \\
\hline
\begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ -1 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} \\
\hline
\begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ -1 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} \\
\hline
\begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 0 \\ 0 \end{array} & \begin{array}{c} 1 \\ -1 \end{array}
\end{array} \right)
\end{array}$$

- The matrix associated to $\sum_{j=0}^2 (-1)^j \overline{d}_i : \mathbf{RGF}_2 \rightarrow \mathbf{RGF}_1$ with the block decomposition and the column and row grades is given by:

$$\begin{array}{c}
\begin{array}{c} 0 < 1 \\ 0 < 2 \\ 1 < 2 \\ 1 < 3 \\ 2 < 3 \end{array} \begin{array}{cc} & \begin{array}{c} 1 < 2 < 3 \\ (1,2) \quad (2,1) \end{array} \\ \begin{array}{c} (0,2) \\ (2,0) \\ (0,2) \\ (2,0) \\ (1,1) \\ (2,0) \\ (0,2) \\ (1,1) \\ (0,2) \\ (1,1) \end{array} & \begin{pmatrix} 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 0 & 0 \\ 1 & 1 \\ 0 & 0 \\ -1 & 0 \\ 0 & -1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix} \end{array}
\end{array}$$

- The matrix associated to $\sum_{j=0}^1 (-1)^j \alpha_j : \mathbf{RGF}_1 \rightarrow \mathbf{RD}_0$ with the block decomposition and the column and row grades is given by:

$$\begin{array}{c}
\begin{array}{c} 0 \\ 1 \\ 2 \\ 3 \end{array} \begin{array}{c} 0 \\ 0 \end{array} \begin{array}{c} \begin{array}{c} 0 < 1 \\ (0,2) \quad (2,0) \end{array} \\ \begin{array}{c} 0 < 2 \\ (0,2) \quad (2,0) \end{array} \\ \begin{array}{c} 1 < 2 \\ (1,1) \quad (2,0) \end{array} \\ \begin{array}{c} 1 < 3 \\ (0,2) \quad (1,1) \end{array} \\ \begin{array}{c} 2 < 3 \\ (0,2) \quad (1,1) \end{array} \end{array} \begin{pmatrix} -1 & -1 & -1 & -1 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 0 & 0 & -1 & -1 & -1 & -1 & 0 & 0 \\ 0 & 0 & 1 & 1 & 1 & 1 & 0 & 0 & -1 & -1 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1 & 1 & 1 & 1 \end{pmatrix}
\end{array}$$

6. PRESENTATIONS OF BIFILTRATIONS

Assume that R is a field. For a general multifiltration of spaces, to get a presentation of its homology, one can apply a standard algebra software to the exact sequence given in 5.2. In the case of a bifiltration $F: \mathbb{N}^2 \rightarrow \text{Spaces}$ one can try to be more efficient. Instead of applying the software directly to the complex given in 5.2, one can first use the fact that the polynomial ring $R[x_1, x_2]$ has the projective dimension 2. This implies that the kernel of any map between free modules is free. In particular the kernel \mathbf{Z} of the map $\sum_{j=0}^n (-1)^j \alpha_j: \mathbf{RGF}_n \rightarrow \mathbf{RD}_{n-1}$ is free. Let $\phi: \mathbf{RKF}_n \oplus \mathbf{RGF}_{n+1} \rightarrow \mathbf{Z}$ be the map that fits into the following commutative diagram:

$$\begin{array}{ccc}
\mathbf{RKF}_n \oplus \mathbf{RGF}_{n+1} & \xrightarrow{\phi} & \mathbf{Z} \hookrightarrow \mathbf{RGF}_n \\
& \searrow & \uparrow \\
& & [\pi_0 - \pi_1 \quad \sum_{i=0}^{n+1} (-1)^i \overline{d}_i]
\end{array}$$

The map $\phi: \mathbf{RKF}_n \oplus \mathbf{RGF}_{n+1} \rightarrow \mathbf{Z}$ is a free presentation of $\mathbf{H}_n(\mathbf{F}, \mathbf{R})$. To take a full advantage of this idea, one would need to be able to describe in an efficient way a set of free generators of \mathbf{Z} . As of writing this paper, we have not found a method for doing it.

ACKNOWLEDGEMENTS

We would like to thank Bernd Sturmfels for suggesting the problem of finding an efficient way to compute a presentation of multidimensional persistence modules. We would also like to thank Sandra Di Rocco for her advice and support and Antonio Patriarca for stimulating discussions. This paper would not be the same without valuable contributions of the referee.

REFERENCES

- [1] G. Carlsson, A. Zomorodian, The Theory of Multidimensional Persistence, *Discr. Comput. Geomet.*,(1)(42)(2009) 71-93.
- [2] G. Carlsson, G. Singh, A. Zomorodian, Computing Multidimensional Persistence, *Journal of Computational Geometry*,(1)(2010) 72-100.
- [3] S. Mac Lane, *Categories for the Working Mathematician*, vol. 5 Graduate Texts in Mathematics, Springer (1998).
- [4] A. Adcock, D. Rubin, G. Carlsson, Classification of Hepatic Lesions using the Matching Metric, *Computer vision and image understanding*,(121)(2014), 36-42.

(Wojciech Chachólski) DEPARTMENT OF MATHEMATICS, KTH, S 10044 STOCKHOLM, SWEDEN
E-mail address: `wojtek@math.kth.se`

(Martina Scolamiero) DEPARTMENT OF MATHEMATICS, KTH, S 10044 STOCKHOLM, SWEDEN
E-mail address: `scola@kth.se`

(Francesco Vaccarino) DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO,
C.SO DUCA DEGLI ABRUZZI N.24,TORINO,10129, ITALY.
E-mail address: `francesco.vaccarino@polito.it`

(Francesco Vaccarino) ISI FOUNDATION, VIA ALASSIO 11/C,TORINO,10126, ITALY.
E-mail address: `vaccarino@isi.it`