

NLS ground states on graphs

Original

NLS ground states on graphs / Adami, Riccardo; Serra, Enrico; Tilli, Paolo. - In: CALCULUS OF VARIATIONS AND PARTIAL DIFFERENTIAL EQUATIONS. - ISSN 0944-2669. - STAMPA. - 54:1(2015), pp. 743-761. [10.1007/s00526-014-0804-z]

Availability:

This version is available at: 11583/2603580 since: 2020-01-31T15:40:39Z

Publisher:

Springer

Published

DOI:10.1007/s00526-014-0804-z

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

NLS ground states on graphs

Riccardo Adami*, Enrico Serra†, Paolo Tilli

Dipartimento di Scienze Matematiche “G.L. Lagrange”, Politecnico di Torino
Corso Duca degli Abruzzi, 24, 10129 Torino, Italy

Abstract

We investigate the existence of ground states for the subcritical NLS energy on metric graphs. In particular, we find out a topological assumption that guarantees the nonexistence of ground states, and give an example in which the assumption is not fulfilled and ground states actually exist. In order to obtain the result, we introduce a new rearrangement technique, adapted to the graph where it applies. Owing to such a technique, the energy level of the rearranged function is improved by conveniently mixing the symmetric and monotone rearrangement procedures.

AMS Subject Classification: 35R02, 35Q55, 81Q35, 49J40, 58E30.

Keywords: Minimization, metric graphs, rearrangement, nonlinear Schrödinger Equation.

1 Introduction

In this paper we investigate the existence of a ground state for the NLS energy functional

$$(1) \quad E(u, \mathcal{G}) = \frac{1}{2} \|u'\|_{L^2(\mathcal{G})}^2 - \frac{1}{p} \|u\|_{L^p(\mathcal{G})}^p$$

with the *mass constraint*

$$(2) \quad \|u\|_{L^2(\mathcal{G})}^2 = \mu,$$

where $\mu > 0$ and $p \in (2, 6)$ are given numbers and \mathcal{G} is a *connected metric graph*.

* Author partially supported by the FIRB 2012 project “Dispersive dynamics: Fourier Analysis and Variational Methods”.

† Author partially supported by the PRIN 2012 project “Aspetti variazionali e perturbativi nei problemi differenziali nonlineari”.




Figure 1: (a) Two half-lines with a double-bridge in between. (b) A straight line with one pendant attached to it.

Here we present a rather informal description of the problem and of the main results of the paper, whereas a precise setting and formal definitions are given in Section 2,

A metric graph \mathcal{G} ([6, 12, 16]) is essentially a one-dimensional singular variety, made up of several, possibly unbounded intervals (the edges of the graph) some of whose endpoints are glued together according to the topology of the graph. The spaces $L^p(\mathcal{G})$, $H^1(\mathcal{G})$ etc. are defined in the natural way. All the functions we consider are real valued: this is not restrictive, because $E(|u|, \mathcal{G}) \leq E(u, \mathcal{G})$ and any ground state is in fact real valued, up to multiplication by a constant phase $e^{i\theta}$.

When $\mathcal{G} = \mathbb{R}$ the minimization problem

$$(3) \quad \min E(u, \mathcal{G}), \quad u \in H^1(\mathcal{G}), \quad \|u\|_{L^2(\mathcal{G})}^2 = \mu$$

is well understood and the minimizers, called *solitons*, are known explicitly. The same is true when $\mathcal{G} = [0, +\infty)$ is a half-line (the minimizer being “half a soliton” of mass 2μ) and, to some extent, when $\mathcal{G} = [a, b]$ is a bounded interval. Much more interesting is the case when \mathcal{G} is non-compact and has a nontrivial topology with multiple junctions, loops and so on (see Figure 2). The aim of this paper is that of studying existence and qualitative properties of solutions to (3), under quite general assumptions on \mathcal{G} (for papers devoted to particular graphs, see [1, 3, 10]).

Since when \mathcal{G} is compact existence of minimizers for (3) is immediate, we focus on graphs where at least one edge is unbounded (a half-line), so that the embeddings $H^1(\mathcal{G}) \hookrightarrow L^r(\mathcal{G})$ are not compact and existence for (3) is non-trivial. In fact, even though the infimum of $E(u, \mathcal{G})$ is always trapped between two finite values (Theorem 2.2), it turns out that the existence of minimizers heavily depends on the topology of \mathcal{G} : if, for instance, \mathcal{G} consists of two half-lines with a “double bridge” in between (Figure 1.a) then (3) has *no solution*, while if \mathcal{G} is a straight line with one pendant attached to it (Figure 1.b) then minimizers *do exist*.

Our results extend in two directions. First, we prove a nonexistence result (Theorems 2.3 and 2.5) for a broad family of non-compact graphs (see condition (H) in Section 2): roughly speaking, if no cut-edge of \mathcal{G} segregates all the half-lines of \mathcal{G} in the same connected component, then (3) has no

solution, the only exceptions being certain graphs with a particular topology which we characterize completely (see Example 2.4). Moreover, for all these graphs the unattained infimum of $E(u, \mathcal{G})$ coincides with the NLS energy of a soliton on the real line, for the same values of μ and p .

The mentioned condition on \mathcal{G} , that prevents existence of minimizers in (3), is incompatible with the presence of a bounded pendant edge attached to \mathcal{G} . This motivates a case study when \mathcal{G} is the simplest non-compact graph with one pendant, namely the graph in Figure 1.b: we prove that for this particular \mathcal{G} problem (3) does have a solution (Theorem 2.6) and we establish some qualitative properties of the minimizers (Theorem 2.7). In fact, in this case the energy level of the minimizers is *strictly lower* than the energy level of a soliton on the real line, and any minimizer u —in order to reduce the energy level—does exploit the topology of \mathcal{G} , in that $\sup u$ is attained at the tip of the pendant: indeed, the fact that concentrating mass on the pendant is energetically convenient prevents an a priori possible loss of compactness of a minimizing sequence along a half-line, and existence can be proved. In this respect, a key role is played by a new rearrangement technique, introduced in the proof of Lemma 6.1. The interesting feature is that a *hybrid* rearrangement of a function $u \in H^1(\mathcal{G})$ is needed, adapted to the topology of \mathcal{G} : *high* values of u are rearranged *increasingly* on the pendant, while *small* values of u are rearranged *symmetrically* on the straight line of \mathcal{G} .

It should be pointed out, however, that the presence of a bounded pendant attached to \mathcal{G} is not enough, alone, to guarantee solutions to (3). The problem of characterizing all non-compact \mathcal{G} such that (3) has a solution is certainly a challenging one, since the topology of \mathcal{G} alone is not enough to answer this question and, in general, also the *metric* properties of \mathcal{G} (i.e. the lengths of its edges) play a relevant role. This issue will be discussed in more detail in a forthcoming paper.

Among the physical motivations for this problem (see [17] and references therein), nowadays the most topical is probably given by the Bose-Einstein condensation (see [11]). It is widely known that, under a critical temperature, a boson gas undergoes a phase transition that leads a large number N of particles into the same quantum state, represented by the *wave function* that minimizes the *Gross-Pitaevskii energy* functional

$$(4) \quad E_{GP}(\varphi, \Omega) = \|\varphi'\|_{L^2(\Omega)}^2 + 8\pi\alpha\|\varphi\|_{L^4(\Omega)}^4$$

under the *normalization condition* $\|\varphi\|_{L^2(\Omega)}^2 = N$. The real number α is the scattering length associated with the two-body interaction between the particles in the gas. The functional in (1) corresponds to the case of a negative scattering length, that is realized, for instance, by an attractive two-body interaction. Besides, in (1) we consider a general subcritical nonlinearity power.

In (4), the domain Ω corresponds to the shape of the (magnetic and/or optical) trap where the gas has been confined in order to induce the phase transition. Present technology allows various shapes for traps, like discs, cigars, and so on. Recently, the possibility of building ramified traps ([15, 19]) has been envisaged theoretically, even though, at least to our knowledge, they have not been experimentally realized so far.

To give a mathematical description of such an experimental setting, one should choose a spatial domain Ω that reproduces the shape of the trap and then minimize the energy in (4). One would expect that, for branched traps, the domain Ω may be replaced by a suitable graph \mathcal{G} . The possible ground state then provides the state of the condensate in the trap Ω , while the absence of a ground state would in principle signal an instable character of the system. For instance, in the situation depicted by Theorems 2.2, 2.3, the system would run away along an infinite edge, mimicking the shape of a soliton. For further results on nonlinear evolution on graphs, see e.g. [4, 8, 13, 18].

The paper is organized as follows: Section 2 contains a precise setting of the problem and the statements of the main results. In Sections 3 and 4 we discuss some preliminary facts and techniques (in particular, rearrangements on graphs) and some auxiliary statements that may have some interest in themselves. Finally, Sections 5 and 6 contain the proofs of the results stated in Section 2.

Remark on Figures: some figures have been included to better describe the topology of certain metric graphs. In these pictures, vertices “at infinity” are denoted by the symbol ∞ , while ordinary vertices (i.e. junctions of two or more edges) are denoted by a bullet.

2 Setting, notation and main results

Although we shall not need deep results from graph theory, the notion of graph is central to this paper: we refer the reader to [5, 7] for a modern account on the subject.

Throughout the paper a graph is always meant as a (connected) *multi-graph*, that is, we allow for multiple edges joining the same pair of vertices. Self-loops (i.e. edges starting and ending at the same vertex) are also allowed. More precisely, the central objects of the paper are *metric graphs* (see [16, 12]), i.e. (connected) graphs $\mathcal{G} = (V, E)$ where each edge $e \in E$ is associated with either a closed bounded interval $I_e = [0, \ell_e]$ of length $\ell_e > 0$, or a closed half-line $I_e = [0, +\infty)$, letting $\ell_e = +\infty$ in this case. Two edges $e, f \in E$ joining the same pair of vertices, if present, are *distinct objects* in all respects: in particular, the corresponding intervals I_e and I_f need not have the same length, and must be considered *distinct* even in the case where $\ell_e = \ell_f$. For every $e \in E$ joining two vertices $v_1, v_2 \in V$, a coordinate




Figure 2: A metric graph with 5 half-lines and 13 bounded edges, one of which forms a self-loop.

x_e is chosen along I_e , in such a way that v_1 corresponds to $x_e = 0$ and v_2 to $x_e = \ell_e$, or *viceversa*: if $\ell_e = +\infty$, however, we always assume that the half-line I_e is attached to the remaining part of the graph at $x_e = 0$, and the vertex of the graph corresponding to $x_e = +\infty$ is called a *vertex at infinity*. The subset of V consisting of all vertices at infinity will be denoted by V_∞ . With this respect, we shall always assume that

- (5) all vertices at infinity of \mathcal{G} (if any) have degree one

where, as usual, the degree of a vertex $v \in V$ is the number of edges incident at v (counting twice any self-loop at v , of course). Finally, the cardinalities of E and V are assumed to be finite. An example of a typical metric graph \mathcal{G} is given in Figure 2.

A connected metric graph \mathcal{G} has the natural structure of a locally compact metric space, the metric being given by the shortest distance measured along the edges of the graph. Observe that

- (6) \mathcal{G} is compact \iff no edge of \mathcal{G} is a half-line $\iff V_\infty = \emptyset$.

With some abuse of notation, we often identify an edge e with the corresponding interval I_e : thus, topologically, the *metric space* \mathcal{G} is the *disjoint union* $\bigsqcup I_e$ of its edges, with some of their endpoints *glued together* into a single point (corresponding to a vertex $v \in V \setminus V_\infty$), according to the topology of the *graph* \mathcal{G} (using the same symbol \mathcal{G} for both the metric graph and the induced metric space should cause no confusion). We point out that any vertex at infinity $v \in V_\infty$ is of course a vertex of the graph \mathcal{G} , but is *not* a point of the metric space \mathcal{G} (this is consistent with (5)).

With \mathcal{G} as above, a function $u : \mathcal{G} \rightarrow \mathbb{R}$ can be regarded as a bunch of functions $(u_e)_{e \in E}$, where $u_e : I_e \rightarrow \mathbb{R}$ is the restriction of u to the edge I_e . Endowing each edge I_e with Lebesgue measure, one can define L^p spaces over \mathcal{G} in the natural way, with norm

$$\|u\|_{L^p(\mathcal{G})}^p = \sum_{e \in E} \|u_e\|_{L^p(I_e)}^p, \quad u = (u_e).$$

Similarly, the Sobolev space $H^1(\mathcal{G})$ is defined as the set of those functions $u : \mathcal{G} \rightarrow \mathbb{R}$ such that

- (7) $u = (u_e)$ is continuous on \mathcal{G} , and $u_e \in H^1(I_e)$ for every edge $e \in E$,

with the natural norm

$$\|u\|_{H^1(\mathcal{G})}^2 = \int_{\mathcal{G}} (|u'(x)|^2 + |u(x)|^2) dx = \sum_{e \in E} \int_{I_e} (|u'_e(x_e)|^2 + |u_e(x_e)|^2) dx_e.$$

Note that $H^1(\mathcal{G})$ can be identified with a closed subspace (determined by the continuity of u at the vertices of \mathcal{G}) of the Cartesian product $\bigoplus_e H^1(I_e)$. In terms of the coordinate system $\{x_e\}$, continuity on \mathcal{G} means that, whenever two edges e, f meet at a vertex v of \mathcal{G} , the corresponding branches of u satisfy a no-jump condition of the kind $u_e(0) = u_f(0)$ (or $u_e(\ell_e) = u_f(\ell_f)$, or $u_e(\ell_e) = u_f(0)$ etc., depending on the orientation of I_e, I_f induced by the coordinates x_e, x_f). Notice that, according to (5), vertices at infinity are never involved in these continuity conditions: on the other hand, if $u \in H^1(\mathcal{G})$, then automatically

$$(8) \quad I_e = [0, +\infty) \quad \Rightarrow \quad \lim_{x_e \rightarrow +\infty} u_e(x_e) = 0,$$

because in particular $u_e \in H^1(I_e)$.

Within this framework, we are now in a position to state our main results. Fix \mathcal{G} as above, and numbers μ, p satisfying

$$(9) \quad \mu > 0 \quad \text{and} \quad 2 < p < 6.$$

For $u \in H^1(\mathcal{G})$, the NLS energy in (1) is finite and takes the concrete form

$$E(u, \mathcal{G}) = \frac{1}{2} \sum_e \int_{I_e} |u'_e(x_e)|^2 dx_e - \frac{1}{p} \sum_e \int_{I_e} |u_e(x_e)|^p dx_e.$$

If we let

$$(10) \quad H_\mu^1(\mathcal{G}) := \{u \in H^1(\mathcal{G}) \quad \text{such that} \quad \|u\|_{L^2(\mathcal{G})}^2 = \mu\},$$

the minimization problem (3) takes the compact form

$$(11) \quad \min_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}).$$

Remark 2.1. The classical instance of (11) where \mathcal{G} is the real line \mathbb{R} falls within our framework as a particular case, when \mathcal{G} is made up of *two* unbounded edges (half-lines), joined at their initial point (Figure 3.a).

In this case, the solutions to (11) are called *solitons*, and are known to be unique up to translations and a change of sign. In particular, there is a unique minimizer which is a *positive and even function*: we shall denote this function by ϕ_μ (the dependence on p being understood), and of course

$$(12) \quad E(\phi_\mu, \mathbb{R}) = \min_{\phi \in H_\mu^1(\mathbb{R})} E(\phi, \mathbb{R}) < 0.$$

It is well known that solitons obey the scaling rule

$$(13) \quad \phi_\mu(x) = \mu^\alpha \phi_1(\mu^\beta x), \quad \alpha = \frac{2}{6-p}, \quad \beta = \frac{p-2}{6-p}$$

where $\alpha, \beta > 0$ by (9), and $\phi_1(x) = C_p \operatorname{sech}(c_p x)^{\alpha/\beta}$ with $C_p, c_p > 0$.

The following is a general result for non-compact graphs.

Theorem 2.2. *If \mathcal{G} contains at least one half-line, then*

$$(14) \quad \inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) \leq \min_{\phi \in H_\mu^1(\mathbb{R})} E(\phi, \mathbb{R}) = E(\phi_\mu, \mathbb{R})$$

and, in the other direction,

$$(15) \quad \inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) \geq \min_{\phi \in H_\mu^1(\mathbb{R}^+)} E(\phi, \mathbb{R}^+) = \frac{1}{2} E(\phi_{2\mu}, \mathbb{R}).$$

In order to investigate whether the infimum in (14) is attained or not, the following structure assumption on the graph \mathcal{G} will play a crucial role:

- (H) After removal of any edge $e \in E$, *every* connected component of the graph $(V, E \setminus \{e\})$ contains at least one vertex $v \in V_\infty$.

Some remarks are in order. Firstly, (H) entails that \mathcal{G} has *at least* one vertex at infinity $v_1 \in V_\infty$, whence \mathcal{G} is not compact by (6). Secondly, the condition on e is relevant only when e is a *cut-edge* for \mathcal{G} (i.e. when the removal of e disconnects \mathcal{G}), because when $(V, E \setminus \{e\})$ is connected the presence of $v_1 \in V_\infty$ makes the condition trivial. On the other hand, the edge e (half-line) that has v_1 as vertex at infinity is necessarily a cut-edge, since by (5) its removal leaves vertex v_1 *isolated* in the graph $(V, E \setminus \{e\})$: therefore, the other connected component necessarily contains a vertex at infinity $v_2 \neq v_1$. Hence we see that, in particular,

$$(16) \quad (H) \Rightarrow \mathcal{G} \text{ has at least two vertices at infinity.}$$

Roughly speaking, assumption (H) says that there is always a vertex at infinity on *both sides* of any cut-edge. Any cut-edge e that violates (H), would therefore leave *all* the vertices at infinity on the same connected component thus forming a sort of “bottleneck”, as regards the location of V_∞ relative to e . Thus, in a sense, we may consider (H) as a no-bottleneck condition on \mathcal{G} . Finally, the fact that (H) concerns cut-edges only makes it easy to test algorithmically, when the topology of \mathcal{G} is intricate: for instance, one can easily check that the graph in Figure 2 satisfies (H).

Under assumption (H), the inequality in (14) is in fact an equality:

Theorem 2.3. *If \mathcal{G} satisfies (H), then*

$$(17) \quad \inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) = \min_{\phi \in H_\mu^1(\mathbb{R})} E(\phi, \mathbb{R}) = E(\phi_\mu, \mathbb{R}).$$

Remark. Hypothesis (H) and hence Theorem 2.3 apply to examples of graphs previously treated in the literature: star-graphs with unbounded edges ([1]) and general multiple bridges ([3]). Furthermore, they apply to any semi-eulerian graph with two vertices at infinity, as well as to more complicated networks like the one represented in Figure 2.

It is easy to construct examples of graphs \mathcal{G} satisfying (H), for which the infimum in (17) is achieved.




Figure 3: Graphs described in Example 2.4, for which (14) is an equality.

Example 2.4. (a) If \mathcal{G} is isometric to \mathbb{R} (see Remark 2.1 and Figure 3.a), then the soliton ϕ_μ can be seen as an element of $H_\mu^1(\mathcal{G})$, and by (12) the infimum in (17) is achieved.

(b) The symmetry of the soliton $\phi_\mu \in H^1(\mathbb{R})$ can be exploited to construct other examples. Given $a_1 > 0$, let \mathcal{G} be the quotient space $\mathbb{R}/\{\pm a_1\}$, obtained by gluing together the two points a_1 and $-a_1$ into a unique point x_1 . As a metric graph, \mathcal{G} is depicted in Figure 3.b, the length of the loop being $2a_1$. Since $\phi_\mu(a_1) = \phi_\mu(-a_1)$, ϕ_μ can be seen as an element of $H_\mu^1(\mathcal{G})$, letting $x = 0$ correspond to the north pole of the loop in Figure 3.b. As before, by (12) the infimum in (17) is achieved.

(c) More generally, for $n \geq 2$ fix $a_n > \dots > a_1 > 0$, and let \mathcal{G} be obtained from \mathbb{R} by gluing together each pair of points $\pm a_1, \dots, \pm a_n$, the corresponding new points being denoted $\{x_j\}$. As a metric graph, \mathcal{G} is as in Figure 3.d (the length of the loop at the top being $2a_1$, while the pairs of parallel edges have lengths $a_j - a_{j-1}$, $2 \leq j \leq n$). Since $\phi_\mu(a_i) = \phi_\mu(-a_i)$, reasoning as in (b) we see that the infimum in (17) is attained. \square

In fact, the graphs of the previous example are the *only ones* for which the infimum is attained.

Theorem 2.5. *If \mathcal{G} satisfies (H) then (17) holds true, but the infimum is never achieved unless \mathcal{G} is isometric to one of the graphs discussed in Example 2.4.*

Thus, with the only exception of the graphs of Example 2.4, assumption (H) rules out the existence of minimizers. Among metric graphs with at least two half-lines, the simplest one that violates (H) is the graph in Figure 1.b, made up of two half-lines and one bounded edge (of arbitrary length) joined at their initial point.

For this graph, we have the following result.

Theorem 2.6. *Let \mathcal{G} consist of two half-lines and one bounded edge (of arbitrary length $\ell > 0$) joined at their initial point. Then*

$$(18) \quad \inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) < \min_{\phi \in H_\mu^1(\mathbb{R})} E(\phi, \mathbb{R}) = E(\phi_\mu, \mathbb{R})$$

and the infimum is achieved.

As mentioned in the introduction, any minimizer exploits the peculiar topology of this graph, and tends to concentrate on the pendant. This is described in the following theorem.

Theorem 2.7. *Let \mathcal{G} be as in Theorem 2.6, and let $u \in H_\mu^1(\mathcal{G})$ be any minimizer that achieves the infimum in (18). Then, up to replacing u with $-u$, we have $u > 0$ and*

- (i) *u is strictly monotone along the pendant, with a maximum at the tip.*
- (ii) *If u_1, u_2 denote the restrictions of u to the two half-lines, with coordinates $x \geq 0$ starting both ways at the triple junction, then*

$$u_1(x) = u_2(x) = \phi_{\mu^*}(x + y) \quad \forall x \geq 0,$$

for suitable $y > 0$ and $\mu^ > \mu$ that depend on the mass μ and the length of the pendant ℓ . In particular, the restriction of u to the straight line is symmetric and radially decreasing, with a corner point at the origin.*

- (iii) *For fixed μ , the infimum in (18) is a strictly decreasing function of ℓ .*

From (i) and (ii) it follows that the minimum of u along the pendant coincides with its maximum on the straight line (in other words, u tends to concentrate on the pendant). Observe that, on each half-line, u coincides with a suitable portion of the soliton ϕ_{μ^*} .

3 Some preliminary results

The *decreasing rearrangement* u^* of a function $u \in H^1(\mathcal{G})$, where \mathcal{G} is a metric graph, was first used in [12] where it is proved that, as in the classical case where \mathcal{G} is an interval (see [14]), this kind of rearrangement does not increase the Dirichlet integral (see also [2]). Besides the increasing rearrangement u^* , we shall also need the *symmetric rearrangement* \widehat{u} , whose basic properties we now recall.

Given $u \in H^1(\mathcal{G})$, assume for simplicity that

$$(19) \quad m := \inf_{\mathcal{G}} u \geq 0, \quad M := \sup_{\mathcal{G}} u > 0$$

and, as in [12], let $\rho(t)$ denote the distribution function of u :

$$\rho(t) = \sum_{e \in E} \text{meas}(\{x_e \in I_e : u_e(x_e) > t\}), \quad t \geq 0,$$

where the u_e 's are the branches of u as in (7). Set

$$(20) \quad \omega := \sum_{e \in E} \text{meas}(I_e), \quad I^* := [0, \omega), \quad \widehat{I} := (-\omega/2, \omega/2)$$

where $\omega \in [0, \infty]$ is the total length of \mathcal{G} . As usual, one can define the following rearrangements of u :

(i) the *decreasing* rearrangement $u^* : I^* \rightarrow \mathbb{R}$ as the function

$$(21) \quad u^*(x) := \inf\{t \geq 0 : \rho(t) \leq x\}, \quad x \in I^*;$$

(ii) the *symmetric decreasing* rearrangement $\widehat{u} : \widehat{I} \rightarrow \mathbb{R}$ as the function

$$\widehat{u}(x) := \inf\{t \geq 0 : \rho(t) \leq 2|x|\}, \quad x \in \widehat{I}.$$

Since u , u^* and \widehat{u} are equimeasurable, one has

$$(22) \quad \int_{I^*} |u^*(x)|^r dx = \int_{\widehat{I}} |\widehat{u}(x)|^r dx = \int_{\mathcal{G}} |u(x)|^r dx \quad \forall r > 0$$

and

$$\inf_{I^*} u^* = \inf_{\widehat{I}} \widehat{u} = \inf_{\mathcal{G}} u = m, \quad \sup_{I^*} u^* = \sup_{\widehat{I}} \widehat{u} = \sup_{\mathcal{G}} u = M.$$

As in the classical case where \mathcal{G} is an interval (see [14]), when \mathcal{G} is a *connected* metric graph it turns out (see [12]) that $u^* \in H^1(I^*)$ and $\widehat{u} \in H^1(\widehat{I})$ respectively (connectedness of \mathcal{G} is not essential, as long as the image of u is connected). However, while the passage from u to u^* never increases the Dirichlet integral ([12]), this is not always true for \widehat{u} , a sufficient condition being that the number of preimages

$$N(t) := \#\{x \in \mathcal{G} : u(x) = t\}, \quad t \in (m, M)$$

is *at least two* (see Remark 2.7 in [14]). More precisely, we have

Proposition 3.1. *Let \mathcal{G} be a connected metric graph, and let $u \in H^1(\mathcal{G})$ satisfy (19). Then*

$$(23) \quad \int_{I^*} |(u^*)'|^2 dx \leq \int_{\mathcal{G}} |u'|^2 dx,$$

with strict inequality unless $N(t) = 1$ for a.e. $t \in (m, M)$. Finally,

$$(24) \quad N(t) \geq 2 \quad \text{for a.e. } t \in (m, M) \quad \Rightarrow \quad \int_{\widehat{I}} |(\widehat{u})'|^2 dx \leq \int_{\mathcal{G}} |u'|^2 dx,$$

where equality implies that $N(t) = 2$ for a.e. $t \in (m, M)$.

The part concerning u^* can be found in [12], while the corresponding statements for \widehat{u} can be proved in exactly the same way.

Remark 3.2. If \mathcal{G} is non-compact, i.e. if \mathcal{G} contains at least one half-line, then clearly $\omega = +\infty$ in (20), so that $I^* = \mathbb{R}^+$ and $\widehat{I} = \mathbb{R}$. Thus, in particular, $u^* \in H^1(\mathbb{R}^+)$ while $\widehat{u} \in H^1(\mathbb{R})$.

The following standard result deals with the optimality conditions satisfied by any solution to (11).

Proposition 3.3. *Let \mathcal{G} be a metric graph, and $u \in H_\mu^1(\mathcal{G})$ a solution to (11). Then*

(i) *there exists $\lambda \in \mathbb{R}$ such that*

$$(25) \quad u_e'' + u_e |u_e|^{p-2} = \lambda u_e \quad \text{for every edge } e;$$

(ii) *for every vertex v (that is not a vertex at infinity)*

$$(26) \quad \sum_{e \succ v} \frac{du_e}{dx_e}(v) = 0 \quad (\text{Kirchhoff conditions}),$$

where the condition $e \succ v$ means that edge e is incident at v ;

(iii) *up to replacing u with $-u$, one has that $u > 0$ on \mathcal{G} .*

The Kirchhoff condition (26) is well known (see [12, 16]) and is a natural form of continuity of u' at the vertices of \mathcal{G} . Observe that, by (25), $u_e \in H^2(I_e)$ for every edge e , so that u_e' is well defined at both endpoints of I_e : in (26), of course, the symbol $du_e/dx_e(v)$ is a shorthand notation for $u_e'(0)$ or $-u_e'(\ell_e)$, according to whether the coordinate x_e is equal to 0 or ℓ_e at v .

Proof. Since both the energy $E(u, \mathcal{G})$ and the L^2 constraint in (10) are differentiable in $H^1(\mathcal{G})$ and u is a constrained critical point, computing Gâteaux derivatives one has

$$(27) \quad \int_{\mathcal{G}} (u' \eta' - u |u|^{p-2} \eta) dx + \lambda \int_{\mathcal{G}} u \eta dx = 0 \quad \forall \eta \in H^1(\mathcal{G})$$

where λ is a Lagrange multiplier. Fixing an edge e , choosing $\eta \in C_0^\infty(I_e)$ and integrating by parts, one obtains (25).

Now fix a vertex v (not at infinity) and choose $\eta \in H^1(\mathcal{G})$, null at every vertex of \mathcal{G} except at v : integrating by parts in (27) and using (i), only the boundary terms at v are left, and one finds

$$-\sum_{e \succ v} \frac{du_e}{dx_e}(v) \eta(v) = 0,$$

and (26) follows since $\eta(v)$ is arbitrary.

To prove (iii), observe that if u is a minimizer so is $|u|$, hence we may assume that $u \geq 0$. First assume that u vanishes at a vertex v . Since $u \geq 0$ on \mathcal{G} , no term involved in (26) can be negative: since their sum is zero, every derivative in (26) is in fact zero. Then, by uniqueness for the ODE (25), we see that $u_e \equiv 0$ along every edge e such that $e \succ v$: since \mathcal{G} is connected, this argument can be iterated through neighboring vertices and one obtains that $u \equiv 0$ on \mathcal{G} , a contradiction since $u \in H_\mu^1(\mathcal{G})$. If, on the other hand, $u_e(x) = 0$ at some point x interior to some edge e , from $u \geq 0$ we see that also $u'_e(x) = 0$ and, as before, from (25) we deduce that $u_e \equiv 0$ along e . Thus, in particular, $u(v) = 0$ at a vertex $v \prec e$, and one can argue as above. \square

Another useful result, valid for any metric graph \mathcal{G} , is the following Gagliardo-Nirenberg inequality:

$$\|u\|_{L^p(\mathcal{G})}^p \leq C \|u\|_{L^2(\mathcal{G})}^{\frac{p}{2}+1} \|u\|_{H^1(\mathcal{G})}^{\frac{p}{2}-1} \quad \forall u \in H^1(\mathcal{G}),$$

where $C = C(\mathcal{G}, p)$. This is well known when \mathcal{G} is an interval (bounded or not, see [9]): for the general case, it suffices to write the inequality for each edge of \mathcal{G} , and take the sum.

In particular, when $\|u\|_{L^2(\mathcal{G})}^2 = \mu$ is fixed, we obtain

$$\|u\|_{L^p(\mathcal{G})}^p \leq C + C \|u'\|_{L^2(\mathcal{G})}^{\frac{p}{2}-1} \quad \forall u \in H_\mu^1(\mathcal{G}),$$

where now $C = C(\mathcal{G}, p, \mu)$. Since our p satisfies (9), this shows that the negative term in (1) grows *sublinearly*, at infinity, with respect to the positive one. As a consequence, Young's inequality gives $\|u'\|_{L^2(\mathcal{G})}^2 \leq C + CE(u, \mathcal{G})$ when $u \in H_\mu^1(\mathcal{G})$, and hence also

$$(28) \quad \|u\|_{H^1(\mathcal{G})}^2 \leq C + CE(u, \mathcal{G}) \quad \forall u \in H_\mu^1(\mathcal{G}), \quad C = C(\mathcal{G}, p, \mu).$$

4 Some auxiliary results

In this section we discuss two auxiliary *double-constrained* problems, on \mathbb{R}^+ and on \mathbb{R} respectively, that will be useful in Section 6 and may be of some interest in themselves.

We begin with the double-constrained problem on the half-line

$$(29) \quad \min E(\phi, \mathbb{R}^+), \quad \phi \in H^1(\mathbb{R}^+), \quad \int_0^\infty |\phi|^2 dx = \frac{m}{2}, \quad \phi(0) = a$$

for fixed $m, a > 0$. This corresponds to (3) when $\mathcal{G} = \mathbb{R}^+$ and $\mu = m/2$, with the *additional* Dirichlet condition

$$\phi(0) = a.$$

Theorem 4.1. *For every $a, m > 0$ there exist unique $M > 0$ and $y \in \mathbb{R}$ such that the soliton ϕ_M satisfies the two conditions*

$$(30) \quad \phi_M(y) = a \quad \text{and} \quad \int_0^{+\infty} \phi_M(y+x)^2 dx = \frac{m}{2}.$$

Moreover, the function $x \mapsto \phi_M(y+x)$ is the unique solution to (29).

Finally, there holds

$$(31) \quad a > \phi_m(0) \iff y > 0.$$

Proof. Recalling the scaling rule of solitons (13), let $z = \mu^\beta y$ (to be determined). Then, changing variable $x = M^{-\beta}t$ in the integral, the conditions in (30) become

$$(32) \quad M^\alpha \phi_1(z) = a, \quad \text{and} \quad M \int_0^{+\infty} \phi_1(z+t)^2 dt = \frac{m}{2}.$$

Using the first condition, we can eliminate M from the second and obtain

$$(33) \quad \phi_1(z)^{-\frac{1}{\alpha}} \int_0^{+\infty} \phi_1(z+t)^2 dt = \frac{ma^{-\frac{1}{\alpha}}}{2}.$$

Denoting by $g(z)$ the function on the left-hand side, we claim that

$$(34) \quad \lim_{z \rightarrow -\infty} g(z) = +\infty \quad \text{and} \quad \lim_{z \rightarrow +\infty} g(z) = 0.$$

The first limit is clear as $\phi_1(z) \rightarrow 0$ while the integral tends to $\|\phi_1\|_{L^2(\mathbb{R})}^2 = 1$. For the second, since ϕ_1 is decreasing on \mathbb{R}^+ one can estimate

$$\phi_1(z+t)^2 \leq \phi_1(z)^{\frac{1}{\alpha}} \phi_1(z+t)^{2-\frac{1}{\alpha}} \quad \forall z, t \geq 0$$

in the integral and observe that $2 - 1/\alpha > 0$ by (9).

Moreover, g is strictly decreasing: this is clear when $z < 0$, since in this case $g(z)$ is the product of two strictly decreasing functions. When $z > 0$, differentiation yields

$$\begin{aligned} g'(z) &= 2 \int_0^{+\infty} \frac{\phi_1(z+t)^2}{\phi_1(z)^{\frac{1}{\alpha}}} \left[\frac{\phi_1'(z+t)}{\phi_1(z+t)} - \frac{1}{2\alpha} \frac{\phi_1'(z)}{\phi_1(z)} \right] dt \\ &< 2 \int_0^{+\infty} \frac{\phi_1(z+t)^2}{\phi_1(z)^{\frac{1}{\alpha}}} \left[\frac{\phi_1'(z+t)}{\phi_1(z+t)} - \frac{\phi_1'(z)}{\phi_1(z)} \right] dt < 0, \end{aligned}$$

having used $1/2\alpha < 1$ and $\phi_1'(z) < 0$ in the first inequality, and the log-concavity of ϕ_1 (see Remark 2.1) in the second. This and (34) show that, given $a, m > 0$, there exists a unique $z \in \mathbb{R}$ (hence a unique y) satisfying (33), while M is uniquely determined by the first condition in (32).

To prove (31) observe that, by (13), $a > \phi_m(0)$ is equivalent to $a > m^\alpha \phi_1(0)$, which in turn is equivalent to $g(0) > ma^{-1/\alpha}/2$ since

$$g(0) = \phi_1(0)^{-\frac{1}{\alpha}} \int_0^{+\infty} \phi_1(t)^2 dt = \frac{\phi_1(0)^{-\frac{1}{\alpha}}}{2}.$$

But since $g(z) = ma^{-1/\alpha}/2$ by (33) and g is decreasing, the last inequality is equivalent to $z > 0$, which proves (31).

For the last part of the claim, by Remark 2.1 we see that the function $x \mapsto \phi_M(x+y)$ minimizes $E(\phi, \mathbb{R})$ under the *two constraints* $\|\phi\|_{L^2(\mathbb{R})}^2 = M$ and $\phi(0) = a$. If a competitor $\varphi(x)$ better than $x \mapsto \phi_M(x+y)$ could be found for (29), then the function equal to $\phi_M(x+y)$ for $x < 0$ and to $\varphi(x)$ for $x \geq 0$ would violate the mentioned optimality of $\phi_M(x+y)$. Finally, uniqueness follows from the uniqueness of M and y satisfying (30): indeed, any other solution $\varphi(x)$ to (29) not coinciding on \mathbb{R}^+ with any soliton, arguing as before would give rise to a *non-soliton* minimizer of $E(\phi, \mathbb{R})$ with mass constraint $\|\phi\|_{L^2(\mathbb{R})}^2 = M$. □

Now we consider the analogue problem on the *whole* real line, namely

$$(35) \quad \min E(\phi, \mathbb{R}), \quad \phi \in H^1(\mathbb{R}), \quad \int_{-\infty}^{\infty} |\phi|^2 dx = m, \quad \phi(0) = a$$

for fixed $m, a > 0$. Its solutions are characterized as follows, a special role being played by the soliton ϕ_m of mass m .

Theorem 4.2. *Let $a, m > 0$ be given.*

- (i) *If $a < \phi_m(0)$ then problem (35) has exactly two solutions, given by $x \mapsto \phi_m(x \pm y)$ for a suitable $y > 0$.*
- (ii) *If $a = \phi_m(0)$, then problem (35) has $\phi_m(x)$ as unique solution.*
- (iii) *if $a > \phi_m(0)$, then problem (35) has exactly one solution, namely $x \mapsto \phi_M(|x| + y)$ for suitable $M, y > 0$.*

Proof. Cases (i) and (ii) are immediate since a is in the range of ϕ_m : as ϕ_m and its translates are the only positive minimizers of $E(\cdot, \mathbb{R})$ in $H_m^1(\mathbb{R})$, the second constraint in (35) can be matched for free by a translation, with $y \geq 0$ such that $\phi_m(\pm y) = a$. Case (ii) is when $a = \max \phi_m$, and so $y = 0$.

Now consider (iii), where the value a is not in the range of ϕ_m . Let M, y be the numbers provided by Theorem 4.1, and observe that $y > 0$ according

to (31). Since $\phi_M(y) = a$ by (30), the soliton ϕ_M is clearly the unique solution of the constrained problem

$$(36) \quad \min E(\phi, \mathbb{R}), \quad \phi \in H_M^1(\mathbb{R}), \quad \phi(-y) = \phi(y) = a.$$

Moreover, as $y > 0$, the second condition in (30) implies that

$$\int_{\mathbb{R} \setminus (-y, y)} |\phi_M(x)|^2 dx = 2 \int_y^\infty |\phi_M(x)|^2 dx = m.$$

Therefore, the function $w(x) = \phi_M(|x| + y)$ is an admissible competitor for (35), and obviously

$$E(w, \mathbb{R}) = E(\phi_M, (-\infty, -y)) + E(\phi_M, (y, +\infty)).$$

The existence of a $v(x)$ admissible for (35) and such that $E(v, \mathbb{R}) < E(w, \mathbb{R})$, would allow the construction of a competitor better than ϕ_M in (36), by redefining $\phi_M(x)$ when $|x| \geq y$, setting it equal to $v(x - y)$ or $v(x + y)$, according to whether $x \geq y$ or $x \leq -y$. Thus w solves (35). Similarly, a competitor $v \neq w$ with $E(v, \mathbb{R}) = E(w, \mathbb{R})$ would violate the uniqueness of ϕ_M as a solution of (36), and therefore w is the unique solution of (35). \square

5 Proof of the nonexistence results

Proof of Theorem 2.2. Consider $u_\varepsilon \in H^1(\mathbb{R})$ with compact support, such that $\|u_\varepsilon\|_{L^2}^2 = \mu$ and $u_\varepsilon \rightarrow \phi_\mu$ strongly in $H^1(\mathbb{R})$ as $\varepsilon \rightarrow 0$. Since $u_\varepsilon \rightarrow \phi_\mu$ also in $L^p(\mathbb{R})$, we see that

$$E(u_\varepsilon, \mathbb{R}) \rightarrow E(\phi_\mu, \mathbb{R}) \quad \text{as } \varepsilon \rightarrow 0.$$

Now, by a translation, we may assume that u_ε is supported in $[0, +\infty)$: identifying this interval with one of the half-lines of \mathcal{G} , we may consider u_ε as a function in $H_\mu^1(\mathcal{G})$, by extending it to zero on any other edge of \mathcal{G} . Then we have from the previous equation

$$\inf_{u \in H_\mu^1(\mathcal{G})} E(u, \mathcal{G}) \leq \lim_{\varepsilon \rightarrow 0} E(u_\varepsilon, \mathcal{G}) = E(\phi_\mu, \mathbb{R}),$$

and (14) follows from (12). The inequality in (15) is immediate from (22) and (23) (see Remark 3.2), by rearranging an arbitrary $u \in H_\mu^1(\mathcal{G})$. Finally, the equality in (15) is well known. \square

To prove of Theorems 2.3 and 2.5 we need to investigate how assumption (H), which is purely graph-theoretical, reflects on \mathcal{G} as a *metric-space*.

Lemma 5.1. *Assume \mathcal{G} is connected and satisfies condition (H). Then \mathcal{G} , as a metric space, satisfies the following condition as well:*

(H') For every point $x_0 \in \mathcal{G}$, there exist two injective curves $\gamma_1, \gamma_2 : [0, +\infty) \rightarrow \mathcal{G}$ parameterized by arclength, with disjoint images except for finitely many points, and such that $\gamma_1(0) = \gamma_2(0) = x_0$.

Proof. We recall that a *trail* in a graph (see [7]) is a finite sequence of consecutive edges, in which no edge is repeated (note that a vertex, instead, may be repeated, if the trail contains cycles or self-loops). Let \mathcal{T} be the class of all those *trails* T in the graph \mathcal{G} , such that the initial edge and the final edge of T are half-lines (observe that, due to (5), no edge of T other than the initial and the final edge can be a half-line: thus, in a sense, every $T \in \mathcal{T}$ provides an immersion of \mathbb{R} in \mathcal{G} —not an embedding, however, as T may have cycles). Recalling (16), since \mathcal{G} is connected we see that $\mathcal{T} \neq \emptyset$. Moreover, any point $x_0 \in \mathcal{G}$ that is covered by at least one trail $T \in \mathcal{T}$, satisfies condition (H'). Indeed, to construct γ_1 and γ_2 , it suffices to start at x_0 and move along T both ways, according to arclength: the two obtained curves can then be made injective, by removing any useless loop that each of them, separately, may form. Moreover, since no edge is repeated in T , γ_1 and γ_2 may intersect only at finitely many vertices of \mathcal{G} . Since the initial and the final edges of T are half-lines, each $\gamma_i(t)$ is eventually trapped in a half-line and thus each γ_i has infinite length, which makes it possible to parameterize $\gamma_i(t)$ injectively by arclength with $t \in [0, +\infty)$. Therefore, it suffices to prove that the trails in \mathcal{T} cover \mathcal{G} .

Let $E_0 \subseteq E$ denote the set of those edges that do not belong to any trail $T \in \mathcal{T}$. Assuming $E_0 \neq \emptyset$, since \mathcal{G} is connected there must be some edge $e \in E_0$ with one vertex w on some trail $T \in \mathcal{T}$ ($w \in V \setminus V_\infty$, by (5)). If e were a cut-edge for \mathcal{G} , then by (H) there would be a vertex $v_1 \in V_\infty$ in the connected component of $(V, E \setminus \{e\})$ disjoint from T : then it would be possible to go from v_1 to w along a trail that crosses e , and then proceed over a portion of T up to another vertex $v_2 \in V_\infty$, thus constructing a path $T' \in \mathcal{T}$ that contains e , which is impossible since $e \in E_0$. Now, as e is not a cut-edge, it necessarily belongs to a cycle C (made up of bounded edges only, due to (5)): then, by inserting C in the middle of T at vertex w , we can construct a trail $T' \in \mathcal{T}$ that contains e . Since this is a contradiction, we see that $E_0 = \emptyset$ and therefore \mathcal{T} covers \mathcal{G} . \square

Proof of Theorem 2.3. We first prove that for every $u \in H_\mu^1(\mathcal{G})$,

$$(37) \quad E(u, \mathcal{G}) \geq E(\hat{u}, \mathbb{R}) \geq \min_{\phi \in H_\mu^1(\mathbb{R})} E(\phi, \mathbb{R})$$

where \hat{u} is the symmetric rearrangement of u as defined in (21).

As $E(u, \mathcal{G}) = E(|u|, \mathcal{G})$, we may assume that $u \geq 0$. In fact we have that $M := \max_{\mathcal{G}} u > 0$ by (2), and that $m := \inf_{\mathcal{G}} u = 0$ by (8), as \mathcal{G} contains at least two half-lines due to (16). From Remark 3.2 $\hat{u} \in H^1(\mathbb{R})$ and, in fact, $\hat{u} \in H_\mu^1(\mathbb{R})$ by (22) with $r = 2$, which proves the second inequality in (37).

The first inequality, due to (22) written with $r = p$, is in fact equivalent to the integral inequality in (24), which follows as soon as we show that

$$(38) \quad N(t) := \#\{x \in \mathcal{G} : u(x) = t\} \geq 2 \quad \text{for a.e. } t \in (0, M).$$

This, in turn, follows from (H') of Lemma 5.1. If γ_1, γ_2 are as in (H'), relative to a point $x_0 \in \mathcal{G}$ where $u(x_0) = M$, we may define the continuous function

$$(39) \quad v : \mathbb{R} \rightarrow \mathbb{R}, \quad v(z) = \begin{cases} u(\gamma_1(z)) & \text{if } z \geq 0, \\ u(\gamma_2(-z)) & \text{if } z < 0. \end{cases}$$

Clearly $v(0) = u(x_0) = M$. Moreover, as each $\gamma_i(z)$ parameterizes a half-line of \mathcal{G} for z large enough, from (8) we have that $v(z) \rightarrow 0$ as $|z| \rightarrow \infty$, hence v has at least two distinct preimages (in \mathbb{R}) for every value $t \in (0, M)$. But as the images of γ_1, γ_2 are disjoint except for finitely many points of \mathcal{G} , (38) is established and (37) follows. Now (37), combined with (14), proves Theorem 2.3. \square

Proof of Theorem 2.5. Carrying on with the previous proof, assume that some $u \in H_\mu^1(\mathcal{G})$ achieves the infimum in (17) (i.e. in (37)). Then, both inequalities in (37) are equalities. From them (combining the former with Proposition 3.1) we infer that

- (i) $N(t) = 2$ for a.e. $t \in (0, M)$,
- (ii) \hat{u} is a soliton of mass μ , i.e. $\hat{u} = \phi_\mu$.

Now, if Γ_i denotes the image of the curve γ_i defined above, we claim that

- (iii) the union $\Gamma = \Gamma_1 \cup \Gamma_2$ covers \mathcal{G} .

Indeed, in the proof of (38), we proved a slightly stronger statement, namely that $\#(u^{-1}(t) \cap \Gamma) \geq 2$ for a.e. t (while $N(t)$ counts preimages in the whole \mathcal{G}). Then, from (i), it follows that those values $t \in (0, M)$ attained by u on $\mathcal{G} \setminus \Gamma$ (if any) form a set of measure zero. Since u is continuous on \mathcal{G} and $\mathcal{G} \setminus \Gamma$ is open (Γ is, in fact, the trail mentioned in assumption (H)), this implies that u is *constant* on any edge e of \mathcal{G} not belonging to Γ (as Γ is a trail, we may regard it as a subgraph of \mathcal{G}). But u and \hat{u} are equimeasurable and, by (ii), every level set of \hat{u} has measure zero, hence the same is true for u : therefore, for any edge e of \mathcal{G} not in Γ , the only possibility is that $u \equiv 0$ on e (for the moment, this cannot be excluded: as $\text{meas}(\{\hat{u} > 0\}) = +\infty$, the quantity $\text{meas}(\{u = 0\})$ cannot be computed by complementation). However, since $u \equiv 0$ outside Γ , considering Γ as a subgraph, the restriction $u|_\Gamma$ satisfies

$$u|_\Gamma \in H_\mu^1(\Gamma), \quad E(u|_\Gamma, \Gamma) = E(u, \mathcal{G}) = E(\hat{u}, \mathbb{R}).$$

Since the $\gamma_i : [0, +\infty) \rightarrow \Gamma$ are injective and parameterized by arclength, it follows from (39) and the previous relations that

$$v \in H_\mu^1(\mathbb{R}), \quad E(v, \mathbb{R}) = E(u|_\Gamma, \Gamma) = E(\widehat{u}, \mathbb{R}).$$

Then, by (ii), v is necessarily the translate of a soliton of mass μ but, since $v(0) = M = \widehat{u}(0)$, v is centered at the origin and so $v = \widehat{u} = \phi_\mu$. In particular, $v > 0$ whence also $u|_\Gamma > 0$: then, by continuity, no edge of \mathcal{G} where $u \equiv 0$ can be attached to Γ , hence (iii) is proved since \mathcal{G} is connected.

Now, as $v = \phi_\mu$, v is injective if restricted to either \mathbb{R}^+ or \mathbb{R}^- , hence

$$\forall z_1, z_2 \geq 0, \quad \gamma_i(z_1) = \gamma_i(z_2) \quad \Rightarrow \quad v(z_1) = v(z_2) \quad \Rightarrow \quad z_1 = z_2,$$

which shows that each γ_i is a *simple* curve in \mathcal{G} . Recall that, by (H'), the images of the γ_i 's intersect at at their starting point x_0 and, possibly, also at finitely many other points x_1, \dots, x_n of \mathcal{G} . If $n = 0$ (i.e. if x_0 is the only intersection), then clearly Γ is isometric to the real line \mathbb{R} , and by (iii) we see that \mathcal{G} is necessarily the graph of Example 2.4 (a).

If $n = 1$, then Γ is a straight line with two points glued together, hence \mathcal{G} is necessarily the graph of Example 2.4 (b).

Finally, if $n > 1$, since $v = \varphi_\mu$ is an even and strictly radially decreasing function, we have

$$\forall z_1, z_2 > 0, \quad \gamma_1(z_1) = \gamma_2(z_2) = x_j \quad \Rightarrow \quad v(z_1) = v(-z_2) \quad \Rightarrow \quad z_1 = z_2.$$

Since the γ_i 's are parameterized by arclength, this shows that for every $j \in \{1, \dots, n\}$ the distance a_j of x_j from x_0 , measured along γ_1 , is the same as the distance measured along γ_2 . And this forces \mathcal{G} to be isometric to the graph in Example 2.4 (c). \square

6 Proof of the existence results

Throughout this section, \mathcal{G} is the graph described in Theorem 2.6 (see Figure 4). In dealing with (11), however, it is convenient to let $u = (\phi, \psi)$, and identify each $u \in H^1(\mathcal{G})$ with a pair of functions ϕ, ψ , with $\phi \in H^1(\mathbb{R})$, $\psi \in H^1(I)$ with $I = [0, \ell]$, satisfying the continuity condition $\phi(0) = \psi(0)$: here, of course, the interval $I = (0, \ell)$ represents the pendant of \mathcal{G} , while \mathbb{R} represents the union of its two half-lines. Then (11) is equivalent to the minimization problem

$$\min \left(E(\phi, \mathbb{R}) + E(\psi, I) \right), \quad \phi \in H^1(\mathbb{R}), \quad \psi \in H^1(I)$$

subject to the constraints

$$(40) \quad \phi(0) = \psi(0), \quad \int_{-\infty}^{\infty} |\phi(x)|^2 dx + \int_0^\ell |\psi(x)|^2 dx = \mu.$$




Figure 4: A straight line with a pendant of length $\ell > 0$ attached to it.

As mentioned in the Introduction, the key to the proofs of Theorems 2.6 and 2.7 is the following lemma, that shows how the value of $E(u, \mathcal{G})$ can be reduced, for any $u \in H_\mu^1(\mathcal{G})$, by a proper *hybrid rearrangement* operation, tailored to the topology of \mathcal{G} .

Lemma 6.1 (hybrid rearrangement). *Assume that $u \in H_\mu^1(\mathcal{G})$, with $u > 0$ and $\text{meas}(\{u = t\}) = 0$ for every $t > 0$. Then there exists $\tilde{u} \in H_\mu^1(\mathcal{G})$, $\tilde{u} = (\tilde{\phi}, \tilde{\psi})$, with the following properties:*

- (i) $\tilde{\phi} : \mathbb{R} \rightarrow \mathbb{R}$ is even and radially decreasing;
- (ii) $\tilde{\psi} : I \rightarrow \mathbb{R}$ is increasing, so that $\min \tilde{\psi} = \tilde{\psi}(0) = \tilde{\phi}(0) = \max \tilde{\phi}$;
- (iii) $E(\tilde{u}, \mathcal{G}) \leq E(u, \mathcal{G})$ and, if equality occurs, then letting $u = (\phi, \psi)$ we necessarily have that ψ is increasing on $[0, \ell]$ and $\min \psi = \max \phi$.

Proof. By the assumptions on u , it is easy to see that there exists $\tau > 0$ such that

$$(41) \quad \text{meas}(\{u > \tau\}) = \ell.$$

The idea of the proof is to rearrange the portion of u above τ increasingly on I , and the portion below τ symmetrically on \mathbb{R} . More precisely, let

$$f = u \wedge \tau, \quad g = (u - \tau)^+,$$

and observe that $f, g \in H^1(\mathcal{G})$. Then one can define \hat{f} and g^* , the symmetric and the decreasing rearrangements of f and g , respectively, in such a way that $\hat{f} \in H^1(\mathbb{R})$ and $g^* \in H^1(\mathbb{R}^+)$ by Remark 3.2. Note that: (i) since $\text{meas}(\{g > 0\}) = \ell$ by (41), g^* is supported in $[0, \ell]$; (ii) since $\text{meas}(\{f = \tau\}) = \ell$ by (41), $f^* \equiv \tau$ on $[-\ell/2, \ell/2]$. Therefore, if we let

$$\tilde{\psi}(x) = \tau + g^*(\ell - x) \quad \forall x \in [0, \ell], \quad \text{and} \quad \tilde{\phi}(x) = \begin{cases} \hat{f}(x + \ell/2) & \text{if } x \geq 0, \\ \hat{f}(x - \ell/2) & \text{if } x < 0, \end{cases}$$

we have $\tilde{\psi}(0) = \tau + g^*(\ell) = \tau$ and $\tilde{\phi}(0) = \hat{f}(\ell/2) = \tau$, and the first condition in (40) is satisfied. Moreover, as $\tilde{\phi} \in H^1(\mathbb{R})$ and $\tilde{\psi} \in H^1(I)$, letting $\tilde{u} =$

$(\tilde{\phi}, \tilde{\psi})$ we have that $\tilde{u} \in H^1(\mathcal{G})$. In fact, since by construction for $t > 0$

$$\begin{aligned} \text{meas}(\{\tilde{u} > t\}) &= \text{meas}(\{\tilde{\phi} > t\}) + \text{meas}(\{\tilde{\psi} > t\}) = \\ &= \text{meas}(\{\hat{f} > t\} \setminus [-\ell/2, \ell/2]) + \text{meas}(\{\tau + g^* > t\} \cap [0, \ell]) = \\ &= \text{meas}(\{\tau > u \wedge \tau > t\}) + \text{meas}(\{(u - \tau)^+ > (t - \tau)^+\}) = \text{meas}(\{u > t\}), \end{aligned}$$

\tilde{u} and u are equimeasurable and therefore also $\tilde{u} \in H_\mu^1(\mathcal{G})$ (i.e., (40) is fully satisfied). For the same reason,

$$(42) \quad \|\tilde{u}\|_{L^p(\mathcal{G})}^p = \int_{-\infty}^{\infty} |\tilde{\phi}(x)|^p dx + \int_0^\ell |\tilde{\psi}(x)|^p dx = \|u\|_{L^p(\mathcal{G})}^p.$$

Now we claim that

$$(43) \quad \int_{-\infty}^{\infty} |\tilde{\phi}'(x)|^2 dx = \int_{-\infty}^{\infty} |\hat{f}'(x)|^2 dx \leq \int_{\mathcal{G}} |(u \wedge \tau)'|^2 dx.$$

The equality is immediate from the definition of $\tilde{\phi}(x)$, since $\hat{f} \equiv \tau$ on $[-\ell/2, \ell/2]$, while the inequality follows from (24), as soon as we prove that

$$(44) \quad N(t) := \#\{x \in \mathcal{G} : u(x) \wedge \tau = t\} \geq 2 \quad \text{for a.e. } t \in (0, \tau).$$

Indeed, by (41), the closed set $\{u \geq \tau\}$ cannot be *strictly* contained inside the pendant edge of \mathcal{G} , since the latter has measure ℓ : therefore, we have that $u(x) \geq \tau$ for at least one point x_0 in the union of the two half-lines of \mathcal{G} (possibly, only at the triple junction). But this makes (44) obvious, even for *every* $t \in (0, \tau)$, since $u \rightarrow 0$ as one approaches the two points at infinity of \mathcal{G} , both ways from x_0 .

Similarly, since g^* is supported in $[0, \ell]$ and $\tilde{\psi}'(x) = (g^*)'(\ell - x)$,

$$(45) \quad \int_0^\ell |\tilde{\psi}'(x)|^2 dx = \int_0^\ell |(g^*)'(x)|^2 dx \leq \int_{\mathcal{G}} \left|((u - \tau)^+)' \right|^2 dx$$

where now (23) is used. Taking the sum with (43), we find

$$\int_{\mathcal{G}} |\tilde{u}'(x)|^2 dx \leq \int_{\mathcal{G}} |u'(x)|^2 dx,$$

which combined with (42) shows that $E(\tilde{u}, \mathcal{G}) \leq E(u, \mathcal{G})$. Finally, if equality holds, then it must hold in (45) as well: from the discussion after (23), then, we see that the function $(u - \tau)^+$ must have exactly *one preimage* $x_t \in \mathcal{G}$, for almost every value $t > 0$ that it achieves on \mathcal{G} . Now split $u = (\phi, \psi)$, with $\phi \in H^1(\mathbb{R})$ and $\psi \in H^1(I)$: if $M = \max \phi > \tau$ then, as $\phi(x) \rightarrow 0$ when $|x| \rightarrow \infty$, for *every* $t \in (\tau, M)$ we would have $\#\phi^{-1}(t) \geq 2$, hence $(u - \tau)^+$ would achieve *every* value $t \in (\tau, M)$ *at least twice* on \mathcal{G} , which would be a contradiction. We deduce that $\phi \leq \tau$, hence (41) implies that the set

$\{u > \tau\}$ is contained in the pendant edge of \mathcal{G} : indeed, since the pendant itself has measure ℓ , we obtain that $\psi \geq \tau$ on I , so that $\psi \geq \phi$ and $(u - \tau)^+$ is supported on the pendant edge of \mathcal{G} . Since $(u - \tau)^+$ has one preimage for almost every value, we deduce that ψ is monotone, and this completes the proof. \square

Proof of Theorem 2.6. To prove (18), consider the soliton ϕ_μ , and define

$$\phi(x) := \phi_\mu(|x| + \ell/2) \quad \forall x \in \mathbb{R}, \quad \psi(x) := \phi_\mu(x - \ell/2) \quad \forall x \in [0, \ell].$$

It is clear that $\phi \in H^1(\mathbb{R})$, $\psi \in H^1([0, \ell])$, and (40) is satisfied. Then the function $u = (\phi, \psi)$ belongs to $H_\mu^1(\mathcal{G})$, and clearly $E(u, \mathcal{G}) = E(\phi_\mu, \mathbb{R})$. Moreover, $u > 0$ and $\text{meas}(\{u = t\}) = 0$ for every t , so that we can apply Lemma 6.1 and obtain a new function $\tilde{u} \in H_\mu^1(\mathcal{G})$ as claimed there: from (iii), we see that $E(\tilde{u}, \mathcal{G}) < E(\phi, \mathcal{G})$, because our ψ is *not* increasing in $[0, \ell]$. Then we have

$$E(\tilde{u}, \mathcal{G}) < E(\phi, \mathcal{G}) = E(\phi_\mu, \mathbb{R}),$$

and this proves (18) since ϕ_μ achieves the minimum in (18).

To prove that the infimum in (18) is attained, let $u_n = (\phi_n, \psi_n)$ be a minimizing sequence, i.e.

$$(46) \quad \lim_n E(u_n, \mathcal{G}) = \lim_n (E(\phi_n, \mathbb{R}) + E(\psi_n, I)) = \inf_{v \in H_\mu^1(\mathcal{G})} E(v, \mathcal{G}).$$

As $E(u_n, \mathcal{G}) = E(|u_n|, \mathcal{G})$, we may assume that $u_n \geq 0$ and even more, by a density argument using e.g. piecewise linear functions, that each u_n satisfies the assumptions of Lemma 6.1. Thus, we may assume that ϕ_n, ψ_n satisfy properties as in (i) and (ii) therein, so that in particular

$$(47) \quad \phi_n \text{ is even and radially decreasing.}$$

Moreover from (28) we see that $\{u_n\}$ is bounded in $H^1(\mathcal{G})$, hence there exists a $u \in H^1(\mathcal{G})$, $u = (\phi, \psi)$, such that up to subsequences

$$(48) \quad \psi_n \rightharpoonup \psi \text{ weakly in } H^1(I), \quad \phi_n \rightharpoonup \phi \text{ weakly in } H^1(\mathbb{R}).$$

Since $I = [0, \ell]$ is bounded, $\psi_n \rightarrow \psi$ strongly in $L^p(I)$, so that

$$(49) \quad E(\psi, I) \leq \liminf_n E(\psi_n, I)$$

because the positive term in (1) is lower semicontinuous. Concerning ϕ_n , clearly $\phi_n \rightarrow \phi$ uniformly on compact sets: however, combining (47) with the fact that $\phi(x) \rightarrow 0$ when $|x| \rightarrow \infty$, one can easily check that $\phi_n \rightarrow \phi$ in $L^\infty(\mathbb{R})$. Since $\{\phi_n\}$ is bounded in $L^2(\mathbb{R})$ and $p > 2$, by interpolation we infer that also $\phi_n \rightarrow \phi$ strongly in $L^p(\mathbb{R})$, so that

$$E(\phi, \mathbb{R}) \leq \liminf_n E(\phi_n, \mathbb{R}).$$

Combining with (49) and (46), we have that

$$(50) \quad E(u, \mathcal{G}) = E(\phi, \mathbb{R}) + E(\psi, I) \leq \inf_{v \in H_\mu^1(\mathcal{G})} E(v, \mathcal{G}) < 0$$

(the last inequality follows from (18) and (12)). Observe that, since $u_n \in H_\mu^1(\mathcal{G})$, one has $\|u\|_{L^2(\mathcal{G})}^2 \leq \mu$ by semicontinuity: if the inequality were strict, one would have $\sigma u \in H_\mu^1$ for a proper constant $\sigma > 1$ (observe $u \not\equiv 0$, since $E(u, \mathcal{G}) < 0$ by (50)). But then, since $\sigma^p > \sigma^2$ and $E(u, \mathcal{G}) < 0$, we would find

$$E(\sigma u, \mathcal{G}) = \frac{\sigma^2}{2} \|u'\|_{L^2(\mathcal{G})}^2 - \frac{\sigma^p}{p} \|u\|_{L^p(\mathcal{G})}^p < \sigma^2 E(u, \mathcal{G}) < E(u, \mathcal{G})$$

which combined with (50) would be a contradiction, since now $\sigma u \in H_\mu^1(\mathcal{G})$. This shows that, $\|u\|_{L^2(\mathcal{G})}^2 = \mu$, so $u \in H_\mu^1(\mathcal{G})$ and u is a minimizer. \square

Proof of Theorem 2.7, (i)-(ii). Let $u \in H_\mu^1(\mathcal{G})$ be a minimizer. From (iii) in Proposition 3.3, we may assume that $u > 0$ on \mathcal{G} . Moreover, splitting $u = (\phi, \psi)$ and letting $a = \phi(0)$, $m = \|\phi\|_{L^2(\mathbb{R})}^2$, we see that ϕ necessarily solves the double-constrained problem (35), and therefore ϕ is one of the functions described in Theorem 4.2 (i)–(iii): in any case, one has $\text{meas}(\{\phi = t\}) = 0$ for every $t > 0$. And the same is true for ψ : indeed, as ψ solves an ODE like (25) in $(0, \ell)$, all its level sets have measure zero unless ψ is constant on $[0, \ell]$. But if ψ were constant then $\psi(0) = \psi(\ell)$, and the function

$$\phi : \mathbb{R} \rightarrow \mathbb{R}, \quad \phi(x) = \begin{cases} \phi(x) & \text{if } x < 0, \\ \psi(x) & \text{if } 0 \leq x \leq \ell, \\ \phi(x - \ell) & \text{if } x > \ell, \end{cases}$$

would therefore belong to $H_\mu^1(\mathbb{R})$, with $E(\phi, \mathbb{R}) = E(u, \mathcal{G})$. But this would contradict (18), since u achieves the infimum there. This shows that u satisfies the assumptions of Lemma 6.1: since $E(u, \mathcal{G})$ cannot be diminished within $H_\mu^1(\mathcal{G})$, an *equality* must occur in (iii) of Lemma 6.1, and this proves that ψ is increasing on $[0, \ell]$ and $\min \psi = \max \phi$. Since $\phi(0) = \psi(0)$, we see that $\phi(0) = \max \phi$, and this rules out the possibility that ϕ is as in (i) of Theorem 4.2. On the other hand, if ϕ were as in (ii), i.e. $\phi(x) = \phi_m(x)$, then the Kirchhoff condition (26) at the triple junction of \mathcal{G} , namely

$$-\phi'(0^-) + \phi'(0^+) + \psi'(0) = 0,$$

would reduce to $\psi'(0) = 0$, since $\phi = \phi_m$ would be of class $C^1(\mathbb{R})$ with $\phi'(0) = 0$. But then ϕ, ψ , other than solving the same ODE (25), would also satisfy the same initial conditions $\phi(0) = \psi(0)$, $\phi'(0) = \psi'(0)$: by uniqueness, ψ would then coincide with the restriction of $\phi = \phi_m$ to $[0, \ell]$, and since $\phi'_m(\ell) < 0$, this would violate the Kirchhoff condition (26) at the tip of the pendant, namely $\psi'(\ell) = 0$. Therefore, the only possibility is that ϕ is as in (iii) of Theorem 4.2, and this completes the proof. \square

Proof of Theorem 2.7, (iii). Given any two different lengths $\ell > \ell' > 0$ for the pendant, let $\mathcal{G}, \mathcal{G}'$ denote the corresponding graphs, and let $u \in H_\mu^1(\mathcal{G}')$, $u > 0$, be a minimizer on \mathcal{G}' : proceeding *exactly* as in the proof of Lemma 6.1, we shall construct a function $\tilde{u} \in H_\mu^1(\mathcal{G})$ such that $E(u, \mathcal{G}') > E(\tilde{u}, \mathcal{G})$. Since u has qualitative properties as in (i) and (ii) of Theorem 2.7, we can certainly find $\tau = \tau(\ell) > 0$ satisfying (41) (now $u \in H_\mu^1(\mathcal{G}')$ and ℓ is *unrelated* to \mathcal{G}' , but this is irrelevant to the computations following (41)). Then we proceed *verbatim* with the construction of $\tilde{u} \in H_\mu^1(\mathcal{G})$ satisfying (42) and (45), with the proviso that \mathcal{G} be replaced with \mathcal{G}' wherever u is involved (the same applies to (44) and (43), hereafter). Since the structure of u is as in (i)-(ii) of Theorem 2.7, and the pendant of \mathcal{G}' has length ℓ' , from $\ell > \ell'$ and (41) we see that $f = u \wedge \tau$ has exactly two preimages for every value $t \in (0, \tau)$ so that also (44) and hence (43) are satisfied, and the same is true for $g = (u - \tau)^+$ for all those values $t > 0$ *small enough*, so that the inequality in (45) is now *strict*. Therefore, we have $E(u, \mathcal{G}') > E(\tilde{u}, \mathcal{G})$ hence the infimum in (18), as a function of ℓ , is strictly decreasing. \square

References

- [1] R. Adami, C. Cacciapuoti, D. Finco, D. Noja. *On the structure of critical energy levels for the cubic focusing NLS on star graphs*. J. Phys. A **45** (2012), no. 19, 192001, 7pp.
- [2] R. Adami, C. Cacciapuoti, D. Finco, D. Noja. *Variational properties and orbital stability of standing waves for NLS equation on a star graph*. arXiv:1206.5201 (2012).
- [3] R. Adami, E. Serra, P. Tilli. *Lack of ground state for NLS on bridge-like graphs*. arXiv:1404.6973 (2014).
- [4] F. Ali Mehmeti. *Nonlinear waves in networks*. Akademie Verlag, Berlin, 1994.
- [5] V. K. Balakrishnan, K. Ranganathan. *A textbook of graph theory*. Second edition. Universitext. Springer, New York, 2012.
- [6] G. Berkolaiko, P. Kuchment. *Introduction to quantum graphs*. Mathematical Surveys and Monographs, 186. AMS, Providence, RI, 2013.
- [7] B. Bollobas. *Modern Graph Theory*. GTM 184, Springer Verlag, New York, 1998.
- [8] J. Bona, R. Cascaval. *Nonlinear dispersive waves on trees*. Can. J. App. Math. **16** (2008), no. 1, 1–18.
- [9] H. Brezis. *Functional Analysis, Sobolev Spaces and Partial Differential Equations*. Universitext, Springer Verlag, New York, 2011.

- [10] C. Cacciapuoti, D. Finco, D. Noja. *Topology induced bifurcations for the NLS on the tadpole graph*. arXiv:1405.3465 (2014).
- [11] F. Dalfovo, S. Giorgini, L.P. Pitaevskii, S. Stringari. *Theory of Bose-Einstein condensation in trapped gases*. Rev. Mod. Phys. **71** (1999) 463–511.
- [12] L. Friedlander. *Extremal properties of eigenvalues for a metric graph*. Ann. Inst. Fourier (Grenoble) **55** (2005), no. 1, 199–211
- [13] S. Gnutzman, U. Smilansky, S. Derevyanko. *Stationary scattering from a nonlinear network*. Phys. Rev. A **83** (2001), 033831, 6pp.
- [14] B. Kawohl. *Rearrangements and convexity of level sets in PDE*. Lect. Notes in Math. 1150, Springer Verlag, Berlin, 1985.
- [15] P.G. Kevrekidis, D.J. Frantzeskakis, G. Theocharis, I.G. Kevrekidis. *Guidance of matter waves through Y-junctions*. Phys. Lett. A **317** (2003), 513–522.
- [16] P. Kuchment. *Quantum graphs I. Some basic structures*. Waves in Random Media **14** (2004), no. 1, S107–S128.
- [17] D. Noja. *Nonlinear Schrödinger equation on graphs: recent results and open problems*. Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **372** (2007), 20130002, 20 pp.
- [18] Z. Sobirov, D. Matrasulov, K. Sabirov, S. Sawada, K. Nakamura. *Integrable nonlinear Schrödinger equation on simple networks: connection formula at vertices*. Phys. Rev. E **81** (2010), no. 6, 066602, 10 pp.
- [19] E.J.G. Vidal, R.P. Lima, M.L. Lyra. *Bose-Einstein condensation in the infinitely ramified star and wheel graphs*. Phys. Rev. E **83** (2011), 061137.