

Rock glaciers on Mars viewed with Google Earth

Original

Rock glaciers on Mars viewed with Google Earth / Sparavigna, Amelia Carolina. - In: PHILICA. - ISSN 1751-3030. - ELETTRONICO. - 2015:510(2015), pp. 1-5.

Availability:

This version is available at: 11583/2616677 since: 2015-09-08T14:57:26Z

Publisher:

PHILICA

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Rock glaciers on Mars viewed with Google Earth

Amelia Carolina Sparavigna

Department of Applied Science and Technology, Politecnico di Torino

Published in enviro.philica.com

Abstract

Several glacier-like landforms have been identified on Mars. We can view some of them with Google Earth and its virtual Mars planet. In the images of Ismeniae Fossae glaciers, the flow lines of these landforms are clearly visible too.

Keywords: Mars, Google Earth, Satellite Imagery

Article body


Ice is a substance which is common in the solar system. Even Mercury, the planet closest to the Sun, is hosting ice, as observed by a NASA spacecraft that provided the first optical images of it in the shadows of a crater [1]. Ice is also covering moons of Jupiter and Saturn and comets are made of it, mixed with rocks, dust and frozen gases such as carbon oxides, methane and ammonia [2]. Besides water ice, the solar system possesses many other types of frozen substances, not only in the comets. The poles of Mars for instance contain frozen carbon dioxide, the deposition of which, during a pole's winter, is caused by low temperatures. In fact, 25–30% of the Martian atmosphere is deposited into slabs of CO₂ ice (dry ice) [3].

Mars has two permanent polar ice caps, which consist primarily of water ice [4,5]. However, besides in its polar caps, the planet has ice in several glaciers [6-8]. A particular concentration of them had been found in the Ismenius Lacus region [6]. At mid-Martian latitudes, the current models of Martian atmosphere are showing that ice should not be stable if exposed [9]: it is supposed therefore, that most glaciers must be covered with a layer of rubble or dust that can prevent the transfer of water vapour in the sublimation of ice into atmosphere [10]. The atmospheric pressure on Mars is so low that ice simply evaporates, if it is not well protected under a thick layer of dust. As told in [6], because Martian glaciers are contain substantial proportions of debris, they could be better described as rock glaciers [11-14].


Satellites orbiting over the planet had been used to identify the Martian rock glaciers [15]: with the radar measurements made by the NASA Mars Reconnaissance Orbiter, the presence of ice had be determined and measured. In this manner, a research team of the University of Copenhagen has recently estimated the total volume of water ice present in massive belts of glaciers at the mid-latitudes of Mars, concluding that this volume is a very important water reservoir of this planet [16]. The belts of glaciers are located around Mars between the latitudes of 30 and 50 degrees, on both northern and southern hemispheres.

Some of the Martian glaciers can be viewed with Google Earth and its virtual Mars planet (maps of Google Earth are of a higher resolution than those on the browser version of Google Mars). On the Mars virtual globe, we can find some very-high-resolution images coming from the Mars Reconnaissance Orbiter's HiRISE camera (HiRISE means High Resolution Imaging Science Experiment). To see the glaciers, let us consider the region of Ismeniae Fossae. In the Figure 1A, we can see an elevation map (Google Maps) of this region and, in 1B, the same region in Google Earth. The Context Camera (CTX) on board NASA's Mars Reconnaissance Orbiter (MRO) spacecraft has taken the images we find in 1B (Location: 40.59°N 42.08°E, acquired on January 28, 2007; Location: 40.46°N 43.48°E, acquired on February 8, 2007; Location: 41.77°N 44.11°E, acquired on November 19, 2006). HiRISE images are also available. In the Figure 2, we can see details from the image taken by the Mars Reconnaissance Orbiter at location 40.59°N 42.08°E, on January 28, 2007. The visibility of flow lines is enhanced using Gimp Retinex [17,18]. In both panels of Figure 2, we have the merging of two valleys in which ice, according to the local slope, is flowing.

The streamlines we clearly see in the images are interpreted to represent ice-flow features, as told in [19]. In any case, using Google Earth, we can easily make comparison to glaciers on Earth: in the Figure 3 for instance, the flow lines of glaciers in Alaska are shown.



Courtesy NASA / JPL / GSFC / Arizona State University



Courtesy Google Earth

Figure 1: A) Elevation maps (Google Maps) of Ismeniae Fossae region. B) The same region in Google Earth. We can see images taken by the Context Camera (CTX) on board NASA's Mars Reconnaissance Orbiter (MRO) spacecraft. Location: 40.59°N 42.08°E, acquired on January 28, 2007. Location: 40.46°N 43.48°E, acquired on February 8, 2007. Location: 41.77°N 44.11°E, acquired on November 19, 2006. HiRISE (High Resolution Imaging Science Experiment) images are also available. Note the merging of two valleys in the image at location 40.59°N 42.08°E.




Figure 2: Two details from the image taken by the Context Camera (CTX) on board NASA's Mars Reconnaissance Orbiter (MRO) spacecraft. Location: 40.59°N 42.08°E, acquired on January 28, 2007. The visibility of flow lines is enhanced using Gimp Retinex [17,18]. Note the behaviour of glaciers at the merging of two valleys. Glaciers are flowing according to the local slope.




Figure 3: Two images of glaciers in Alaska for comparison. The visibility of flow lines is enhanced using Gimp Retinex [17,18].

References

1. N.L. Chabot, C.M. Ernst, B.W. Denevi, H. Nair, A.N. Deutsch, D.T. Blewett, S.L. Murchie, G.A. Neumann, E. Mazarico, D.A. Paige, J.K. Harmon, J.W. Head, S.C. Solomon, Images of surface volatiles in Mercury's polar craters acquired by the MESSENGER spacecraft, *Geology*, vol. 42, pp. 1051-1054, 2014. DOI: 10.1130/G35916.1
2. J.M. Greenberg, Making a comet nucleus, *Astronomy and Astrophysics*, vol. 330, pp. 375-380, 1998.
3. J.T. Mellon, W.C. Feldman, T.H. Prettyman, The presence and stability of ground ice in the southern hemisphere of Mars, *Icarus*, vol. 169, pp. 324-340, 2003. DOI: 10.1016/j.icarus.2003.10.022
4. H.H. Kieffer, T.Z. Martin, S.C. Chase, E.D. Miner, F. Don Palluconi, Martian north pole summer temperatures - Dirty water ice, *Science*, vol. 194, pp. 1341-1344, 1976. DOI: 10.1126/science.194.4271.1341
5. J.-P. Bibring, Y. Langevin, F. Poulet, A. Gendrin, B. Gondet, M. Berthé, A. Soufflot, P. Drossart, M. Combes, G. Bellucci, V. Moroz, N. Mangold, B. Schmitt, OMEGA team, Perennial water ice identified in the south polar cap of Mars, *Nature*, vol. 428, no. 6983, pp. 627-630, 2004. DOI: 10.1038/nature02461

6. VV. AA., Glaciers on Mars, Wikipedia and references therein.
7. S.W. Squyres, M.H. Carr, Geomorphic evidence for the distribution of ground ice on Mars Science, vol. 213, pp. 249–253, 1986. DOI: 10.1126/science.231.4735.249
8. J.W. Head, D.R. Marchant, M.C. Agnew, C.I. Fassett, M.A. Kreslavsky, Extensive valley glacier deposits in the northern mid-latitudes of Mars: Evidence for Late Amazonian obliquity-driven climate change. Earth and Planetary Science Letters, vol. 241, no.3, pp. 663-671, (2006). DOI: 10.1016/j.epsl.2005.11.016
9. K.E. Williams, O. Toon, J. Heldmann, C. McKay, M. Mellon, Stability of mid-latitude snowpacks on Mars, Icarus, vol. 196, no. 2, pp. 565-577, 2008. DOI: 10.1016/j.icarus.2008.03.017
10. J. Head, G. Neukum, R. Jaumann, H. Hiesinger, E. Hauber, M. Carr, P. Masson, B. Foing, H. Hoffmann, M. Kreslavsky, S. Werner, S. Milkovich, S. van Gasselt, HRSC Co-Investigator Team, Tropical to mid-latitude snow and ice accumulation, flow and glaciation on Mars, Nature, vol. 434, no. 7031, pp. 346–350, (2005). DOI: 10.1038/nature03359
11. J.W. Head, D.R. Marchant, Cold-based mountain glaciers on Mars: western Arsia Mons, Geology, vol. 31, no. 7, pp. 641-644, (2003). DOI: 10.1130/0091-7613(2003)031<0641:cmgomw>2.0.co;2
12. A. Colaprete, B.M. Jakosky, Ice flow and rock glaciers on Mars, Journal of Geophysical Research, vol. 103, no. E3, pp. 5897-5909, 1998. DOI: 10.1029/97je03371
13. W. Haeberli, B. Hallet, L. Arenson, R. Elconin, O. Humlum, A. Käab, V. Kaufmann, B. Ladanyi, N. Matsuoka, S. Springman, D. Vonder Mühll, Permafrost creep and rock glacier dynamics, Permafrost and Periglacial Processes, vol. 17, no. 3, pp. 189-214, 2006. DOI: 10.1002/ppp.561
14. Vv. Aa., Rock Glacier, Wikipedia.
15. W.B. Whalley, F. Azizi, Rock glaciers and protalus landforms: Analogous forms and ice sources on Earth and Mars, Journal of Geophysical Research, vol. 108, no. E4, pp. 2156-2202, 2003. DOI: 10.1029/2002JE001864
16. N.B. Karlsson, L.S. Schmidt, C.S. Hvidberg, Volume of Martian midlatitude glaciers from radar observations and ice flow modelling, Geophysical Research Letters, vol. 42, no. 8, pp. 2627–2633, 2015. DOI: 10.1002/2015GL063219
17. A.C. Sparavigna, Gimp Retinex for Enhancing Images from Microscopes, International Journal of Sciences, vol. 4, no. 6, pp. 72-79, 2015. DOI: 10.18483/ijsci.758
18. Vv. Aa., Retinex, at <http://docs.gimp.org/en/plugin-retinex.html>
19. D.M. Baker, J.W. Head, D.R. Marchant, Flow patterns of lobate debris aprons and lineated valley fill north of Ismeniae Fossae, Mars, 40th Lunar and Planetary Science Conference, 1822.pdf, (2009).

Amelia Carolina Sparavigna, Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

Information about this Article

This Article was published on 4th September, 2015 at 09:35:28



This work is licensed under a [Creative Commons Attribution 2.5 License](https://creativecommons.org/licenses/by/2.5/).

The full citation for this Article is:

Sparavigna, A. (2015). Rock glaciers on Mars viewed with Google Earth. *PHILICA.COM Article number 510*.