

A note on the Einstein-Hilbert action and Dirac operators on  $n$

*Original*

A note on the Einstein-Hilbert action and Dirac operators on  $n$  / Battisti, U., Coriasco, S.. - In: JOURNAL OF PSEUDO-DIFFERENTIAL OPERATORS AND APPLICATIONS. - ISSN 1662-9981. - 2:3(2011), pp. 303-315. [10.1007/s11868-011-0031-8]

*Availability:*

This version is available at: 11583/2624985 since: 2015-12-04T17:40:20Z

*Publisher:*

SP Birkhäuser Verlag Basel

*Published*

DOI:10.1007/s11868-011-0031-8

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

## A NOTE ON THE EINSTEIN-HILBERT ACTION AND DIRAC OPERATORS ON $\mathbb{R}^n$

UBERTINO BATTISTI AND SANDRO CORIASCO

ABSTRACT. We prove an extension to  $\mathbb{R}^n$ , endowed with a suitable metric, of the relation between the Einstein-Hilbert action and the Dirac operator which holds on closed spin manifolds. By means of complex powers, we first define the regularised Wodzicki Residue for a class of operators globally defined on  $\mathbb{R}^n$ . The result is then obtained by using the properties of heat kernels and generalised Laplacians.

### INTRODUCTION

In 1984 M. Wodzicki [40] introduced a trace on the algebra of classical pseudo-differential operators on a closed manifold  $M$ . A similar result was independently obtained by V. Guillemin [18], in order to give a *soft* proof of Weyl formula. Wodzicki residue became then a standard tool in Non-commutative Geometry. In 1988, A. Connes [10] proved that, for operator of order  $-\dim(M)$ , Dixmier Trace and Wodzicki residue are equivalent. Moreover, he conjectured that Wodzicki Residue could connect Dirac operators and Einstein-Hilbert actions on  $M$ . In 1995, D. Kastler [21], W. Kalau and M. Walze [19] proved this conjecture. Namely, let  $\mathcal{D}$  be the classical Atiyah-Singer operator defined on a closed spin manifold  $M = (M, g)$  of even dimension  $n \geq 4$ . Then

$$(0.1) \quad \text{wres}(\mathcal{D}^{-n+2}) = -\frac{(n-2)2^{\lfloor \frac{n}{2} \rfloor}}{\Gamma(\frac{n}{2})(4\pi)^{\frac{n}{2}}} \int_M \frac{1}{12} s(x) dx,$$

where  $s(x)$  is the scalar curvature and  $dx$  the measure on  $M$  induced by the Riemannian metric  $g$  (see, e.g., [4] for an overview about Wodzicki Residue and Non-commutative Geometry). T. Ackermann [1] gave a proof of (0.1), using the relationship between heat trace and  $\zeta$ -function and the properties of the second term in the asymptotic expansion of the heat trace of a generalised Laplacian. Y. Wang [37], [38], [39] suggested an extension of the result to a class of manifolds with boundary.

In the last years, the definition of Wodzicki residue has been extended to other settings: manifolds with boundary [14], manifolds with conical singularities [34], [15], SG-calculus on  $\mathbb{R}^n$  [28], anisotropic operators on  $\mathbb{R}^n$  [9], and manifolds with cylindrical ends [8]. In [26], in order to study critical metrics on  $\mathbb{R}^n$ , it has been introduced the regularised trace of operators, which leads to the definition of regularised  $\zeta$ -function. Wodzicki residue has been used also by R. Ponge [31] to introduce *lower dimensional volumes*.

In this paper, we make use of SG-operators, a class of pseudodifferential operators on  $\mathbb{R}^n$  whose symbols  $a(x, \xi)$ , for fixed  $\mu, m \in \mathbb{R}$  and all  $x, \xi \in \mathbb{R}^n$ , satisfy the estimates

$$(0.2) \quad |D_x^\alpha D_\xi^\beta a(x, \xi)| \leq C_{\alpha, \beta} (1 + |x|)^{m-|\alpha|} (1 + |\xi|)^{\mu-|\beta|}$$

---

2000 *Mathematics Subject Classification*. Primary: 58J40; Secondary: 58J42, 47A10, 47G30, 47L15.

*Key words and phrases*. Wodzicki Residue, Einstein-Hilbert action, Dirac operator.

for suitable constants  $C_{\alpha,\beta} \geq 0$ : the set of such symbols is denoted by  $SG^{\mu,m}(\mathbb{R}^n)$ . If one deals with operators acting on sections of a vector bundle, the symbols are matrices whose entries must satisfy the inequalities (0.2). The corresponding operators can be defined via the usual left-quantisation

$$Au(x) = \text{Op}(a) = \frac{1}{(2\pi)^{\frac{n}{2}}} \int e^{ix\xi} a(x, \xi) \hat{u}(\xi) d\xi,$$

$u \in \mathcal{S}(\mathbb{R}^n)$ , and the set  $\{\text{Op}(a) \mid a \in SG^{\mu,m}(\mathbb{R}^n)\}$  is denoted by  $L^{\mu,m}(\mathbb{R}^n)$ . Such operators form a graded algebra, that is  $L^{\mu,m}(\mathbb{R}^n) \circ L^{\mu',m'}(\mathbb{R}^n) \subseteq L^{\mu+\mu',m+m'}(\mathbb{R}^n)$ , map continuously  $\mathcal{S}(\mathbb{R}^n)$  to itself and can be extended as continuous operators from  $\mathcal{S}'(\mathbb{R}^n)$  to itself.  $L^{-\infty}(\mathbb{R}^n) = \bigcap_{\mu,m \in \mathbb{R}} L^{\mu,m}(\mathbb{R}^n)$ , the set of the so-called smoothing operators, coincides with the set of operators with kernel in  $\mathcal{S}(\mathbb{R}^{2n})$ : these constitute the residual elements of the SG-calculus. A main tool, in particular, is the subclass of SG-classical operators, whose symbols, belonging to  $SG_{\text{cl}}^{\mu,m}(\mathbb{R}^n) \subset SG^{\mu,m}(\mathbb{R}^n)$ , admit a double asymptotic expansion in terms which are homogeneous w.r.t. the  $\xi$ -variable or the  $x$ -variable, respectively. More precisely, for  $a \in SG_{\text{cl}}^{\mu,m}(\mathbb{R}^n)$  we denote

- (i) by  $a_{\mu-j}$ , the terms of order  $(\mu-j, m)$ , homogeneous w.r.t. the  $\xi$ -variable, such that, for a fixed 0-excision function  $\chi = \chi(\xi)$ ,  $a \sim \sum_j \chi a_{\mu-j} \text{ mod } SG^{-\infty,m}(\mathbb{R}^n)$ ;
- (ii) by  $a_{\cdot,m-k}$  the terms of order  $(\mu, m-k)$ , homogeneous w.r.t. the  $x$ -variable, such that, for a fixed 0-excision function  $\omega = \omega(x)$ ,  $a \sim \sum_k \omega a_{\cdot,m-k} \text{ mod } SG^{\mu,-\infty}(\mathbb{R}^n)$ ;
- (iii) by  $a_{\mu-j,m-k}$  the terms of order  $(\mu-j, m-k)$ , homogeneous w.r.t. the  $\xi$ -variable and the  $x$ -variable, such that, for the same 0-excision functions,  $\chi a_{\mu-j} \sim \sum_k \chi \omega a_{\mu-j,m-k} \text{ mod } SG^{\mu-j,-\infty}$  and  $\omega a_{\cdot,m-k} \sim \sum_j \chi \omega a_{\mu-j,m-k} \text{ mod } SG^{-\infty,m-k}$ .

The set  $\{\text{Op}(a) \mid a \in SG_{\text{cl}}^{\mu,m}(\mathbb{R}^n)\}$  is denoted by  $L_{\text{cl}}^{\mu,m}(\mathbb{R}^n)$ : typical examples are differential operators with polynomial coefficients. For details about the SG-calculus and its properties, see, e.g., [7], [11], [13], [26], [29], [32], and the references quoted therein.

Here we prove a *regularised version* of (0.1) on  $\mathbb{R}^n$ ,  $n \geq 4$ , endowed with a suitable Riemannian metric  $g$  and corresponding induced measure  $dx$ , see Sections 2 and 3 below. We consider a generalised positive Laplacian  $\Delta = \nabla^* \nabla + \mathcal{K}$ , defined on a Hermitian vector bundle  $E$  with connection  $\nabla$ , where  $\mathcal{K}$  is a symmetric endomorphism field. Moreover, in order to define and make use of its fractional powers, we assume that the spectrum  $\sigma(\Delta)$  of  $\Delta$  lies outside a sector of the complex plane, with at most the exception of the origin. Finally, we obtain

$$(0.3) \quad \widehat{\text{wres}}(\Delta^{-\frac{n}{2}+1}) = \frac{(n-2)}{\Gamma(\frac{n}{2})(4\pi)^{\frac{n}{2}}} \oint \left[ \frac{\text{Rk}(E)}{6} s(x) - \text{Trace}(\mathcal{K}_x) \right] dx,$$

where  $\widehat{\text{wres}}(\cdot)$  is a generalised Wodzicki residue, defined for SG-operators of order 0 w.r.t. the  $x$ -variable, and  $\oint$  denotes the *finite part integral*<sup>1</sup>.

The paper is organised as follows. Sections 1 and 2 are devoted to illustrate the definitions of the finite part integral of classical symbols, and of regularised trace and  $\zeta$ -function for SG-operators, respectively. In Section 3 we give the proof of (0.3) under the assumption that, if the origin belongs to  $\sigma(\Delta)$ , it is an isolated point of  $\sigma(\Delta)$ , and, finally, we obtain a regularised version of (0.1).

#### ACKNOWLEDGEMENTS

The authors would like to thank R. Mazzeo, L. Rodino, and E. Schrohe, for useful suggestions and discussions.

<sup>1</sup>A similar result was obtained in [7] by direct evaluation.

## 1. FINITE PART INTEGRAL

The *finite-part integral*, introduced in [25], gives a meaning to the integral of a classical symbol  $a$ , and coincides with the usual integral when  $a \in L^1(\mathbb{R}^n)$ .  $dS$  denotes the usual measure on  $|x| = 1$ , induced by the Euclidean metric on  $\mathbb{R}^n$ , while, in this Section,  $dx$  denotes the standard Lebesgue measure on  $\mathbb{R}^n$ .

**Definition 1.1.** Let  $a$  be an element of the classical Hörmander symbol class  $S_{\text{cl}}^m(\mathbb{R}^n)$ , that is,

- (i)  $a \in C^\infty(\mathbb{R}^n)$  and  $\forall x \in \mathbb{R}^n$   $|D^\alpha a(x)| \leq C_\alpha(1 + |x|)^{m-|\alpha|}$ ;
- (ii)  $a$  admits an asymptotic expansion in homogeneous terms  $a_{m-j}$  of order  $m-j$ : explicitly, for a fixed 0-excision function  $\omega$  and all  $N \in \mathbb{N}$ ,

$$a - \sum_{j=0}^{N-1} \omega a_{m-j} \in S^{m-N}(\mathbb{R}^n).$$

Then:

- if  $m \in \mathbb{Z}$ , set

$$\begin{aligned} \int a(x) dx &:= \lim_{\rho \rightarrow \infty} \left[ \int_{|x| \leq \rho} a(x) dx - \sum_{j=0}^{m-n} \int_{|x| \leq \rho} a_{m-j}(x) dx \right] \\ &= \lim_{\rho \rightarrow \infty} \left[ \int_{|x| \leq \rho} a(x) dx - \sum_{j=0}^{m-n} \frac{\beta_j}{n+m-j} \rho^{n+m-j} - \beta_{n+m} \log \rho \right] \end{aligned}$$

where

$$(1.4) \quad \beta_j := \int_{|x|=1} a_{m-j} dS;$$

- if  $m \notin \mathbb{Z}$ , set

$$(1.5) \quad \int a(x) dx := \lim_{\rho \rightarrow \infty} \left[ \int_{|x| \leq \rho} a(x) dx - \sum_{j=0}^{[m]-n-1} \int_{|x| \leq \rho} a_{m-j}(x) dx \right].$$

From the above Definition it is clear that if  $a \in L^1(\mathbb{R}^n)$  the finite part integral is equivalent to the standard integral. If  $m \notin \mathbb{Z}$  the finite part integral coincides with the Kontsevich-Vishik density [22], [23].

**Remark 1.** If one considers the radial compactification of  $\mathbb{R}^n$  to  $\mathbb{S}_+^n$ , namely

$$rc: \mathbb{R}^n \rightarrow \mathbb{S}_+^n : x = (x_1, \dots, x_n) \mapsto y = \left[ \frac{x_1}{(1+|x|^2)^{\frac{1}{2}}}, \dots, \frac{x_n}{(1+|x|^2)^{\frac{1}{2}}}, \frac{1}{(1+|x|^2)^{\frac{1}{2}}} \right],$$

and chooses  $y_{n+1}$  as boundary defining function on  $\mathbb{S}_+^n$ , its composition with  $rc$  coincides in the interior with  $\frac{1}{(1+|x|^2)^{\frac{1}{2}}}$ ,  $x = rc^{-1}(y) \in \mathbb{R}^n$ . Then

$$\int a(x) dx = \int_{\mathbb{S}_+^n} a(rc^{-1}(y)) dS(y)$$


where the right hand side is defined as the term of order  $\epsilon^0$  in the asymptotic expansion of

$$\int_{\mathbb{S}_+^n \cap \{y_{n+1} \geq \epsilon\}} a(rc^{-1}(y)) dS(y), \quad \epsilon \searrow 0.$$

$\int_{\mathbb{S}_+^n} f dS$  is called Renormalised integral, see [2] and the references quoted therein for its precise definition, properties and applications.

2. REGULARISED TRACE AND REGULARISED  $\zeta$ -FUNCTION

In the sequel we will often make reference to sectors of the complex plane with vertex at the origin, that is, subsets of  $\mathbb{C}$  given by  $\Lambda = \{z \in \mathbb{C} \mid -\pi + \theta \leq \arg(z) \leq \pi - \theta\}$ ,  $0 < \theta < \pi$ , as in the next picture.



The definition of  $\Lambda$ -elliptic operator is the standard one, here given for operators defined through matrix-valued symbols, whose spectrum we denote by  $\sigma(a(x, \xi))$ :

**Definition 2.1.** The operator  $A \in L^{\mu,0}(\mathbb{R}^n)$  is  $\Lambda$ -elliptic if there exists a constant  $R > 0$  such that

$$(2.6) \quad \sigma(a(x, \xi)) \cap \Lambda = \emptyset \quad \forall |\xi| \geq R, \quad \forall x \in \mathbb{R}^n$$

and

$$(2.7) \quad (a(x, \xi) - \lambda)^{-1} \in SG^{-\mu,0}(\mathbb{R}^n) \quad \forall |\xi| \geq R, \quad \forall x \in \mathbb{R}^n, \quad \forall \lambda \in \Lambda.$$

It is well known that, if an operator  $A$  is  $\Lambda$ -elliptic, we can build a *weak* parametrix  $B(\lambda)$  such that

$$(2.8) \quad \begin{aligned} B(\lambda) \circ (A - \lambda I) &= \text{Id} + R_1(\lambda), \\ (A - \lambda I) \circ B(\lambda) &= \text{Id} + R_2(\lambda), \quad R_1, R_2 \in L^{-\infty,0}(\mathbb{R}^n). \end{aligned}$$

Moreover

$$(2.9) \quad \begin{aligned} \lambda B(\lambda) &\in L^{-\mu,0}(\mathbb{R}^n), \\ \lambda^2 [(A - \lambda I)^{-1} - B(\lambda)] &\in L^{-\infty,0}(\mathbb{R}^n), \quad \forall \lambda \in \Lambda \setminus \{0\}. \end{aligned}$$

From now on,  $\mu > 0$  and  $A$  is considered as an unbounded operator with dense domain  $D(A) = H^\mu(\mathbb{R}^n) \hookrightarrow L^2(\mathbb{R}^n) \rightarrow L^2(\mathbb{R}^n)$ . To define the complex powers of a  $\Lambda$ -elliptic operator  $A$ , we assume that the following property holds for its spectrum  $\sigma(A)$ :

$$(A1) \quad \sigma(A) \cap \{\Lambda \setminus \{0\}\} = \emptyset \text{ and the origin is at most an isolated point of } \sigma(A).$$

**Proposition 2.1.** Let  $A \in L^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be a  $\Lambda$ -elliptic operator that satisfies Assumption (A1). The complex power  $A^z$ ,  $\Re(z) < 0$ , can be defined as

$$(2.10) \quad A^z := \frac{1}{2\pi i} \int_{\partial^+ \Lambda_\epsilon} \lambda^z (A - \lambda I)^{-1} d\lambda.$$

where  $\Lambda_\epsilon = \Lambda \cup \{z \in \mathbb{C} \mid |z| \leq \epsilon\}$ , with  $\epsilon > 0$  chosen such that  $\sigma(A) \cap \{\Lambda_\epsilon \setminus \{0\}\} = \emptyset$  and  $\partial^+ \Lambda_\epsilon$  is the (positively oriented) boundary of  $\Lambda_\epsilon$ .

*Proof.* By the definition of  $\Lambda$ -elliptic operator, we know that  $(A - \lambda I)^{-1}$  exists for all  $\lambda \in \Lambda \setminus \{0\}$ . Moreover, by (2.9) we have that  $A$  is sectorial, so the integral (2.10) converges in  $\mathcal{L}(L^2(\mathbb{R}^n))$ .  $\square$

**Remark 2.** The definition of  $A^z$  is then extended to arbitrary  $z \in \mathbb{C}$  in the standard way, that is  $A^z := A^{z-j} \circ A^j$ , where  $j \in \mathbb{Z}_+$  is chosen so that  $\Re(z) - j < 0$ , see, e.g., [8], [26], [35].

**Theorem 2.2.** Let  $A \in L^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be  $\Lambda$ -elliptic and satisfy Assumption (A1). Then,  $A^z \in L^{\mu z,0}(\mathbb{R}^n)$ . Moreover, if  $A$  is SG-classical then  $A^z$  is still SG-classical<sup>2</sup>.

**Remark 3.** In order to define the symbol of  $A^z$ , the resolvent  $(A - \lambda I)^{-1}$  can be approximated with the weak parametrix  $B(\lambda)$  defined in (2.8). In this way, a symbol for  $A^z$  can be computed, modulo smoothing operators w.r.t. the  $\xi$ -variable.  $A^z$  can then be considered as an element of the algebra  $\mathcal{A}$  given by

$$(2.11) \quad \mathcal{A} := \bigcup_{\mu \in \mathbb{Z}} L^{\mu,0}(\mathbb{R}^n) / L^{-\infty,0}(\mathbb{R}^n).$$

The proof of the Theorem 2.2 has been given in [26].

From here on,  $dx$  will denote the measure induced on  $\mathbb{R}^n$  by a smooth Riemannian metric  $g = (g_{jk})$ . In order to obtain a result similar to (0.1) we have to impose some condition on  $g$ , namely<sup>3</sup>

(A2)  $g$  is a matrix-valued SG-classical symbol of order  $(0,0)$ .

If  $A \in L^{\mu,m}(\mathbb{R}^n)$  is trace class, that is  $\mu < -n, m < -n$ , we can define its trace

$$TR(A) := \int K_A(x, x) dx,$$

where  $K_A(x, x)$  is the kernel of  $A$  restricted to the diagonal. The concept of regularised trace, valid for classical SG-operators under less restrictive hypotheses on the order, has been introduced in [25], using the finite part integral defined in the previous Section:

**Definition 2.2.** Let  $A \in L_{\text{cl}}^{\mu,m}(\mathbb{R}^n)$  be such that  $\mu < -n$ . We define the regularised trace of  $A$  as

$$(2.12) \quad \widehat{TR}(A) := \int K_A(x, x) dx.$$

**Remark 4.** Note that the condition  $\mu < -n$  implies that  $K_A(x, x)$  is indeed a function and that the finite part integral (2.12) is well defined.

Now, using the regularised integral, we can give the definition of regularised  $\zeta$ -function:

**Definition 2.3.** Let  $A \in L_{\text{cl}}^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be a  $\Lambda$ -elliptic operator that satisfies (A1); then we define

$$(2.13) \quad \hat{\zeta}(A, z) := \widehat{TR}(A^{-z}) = \int K_{A^{-z}}(x, x) dx, \quad \Re(z) > \frac{n}{\mu},$$

where  $K_{A^{-z}}(x, x)$  is the kernel of the operator  $A^z$ .

<sup>2</sup>To define the class of symbols  $SG^{z,w}(\mathbb{R}^n)$ ,  $z, w \in \mathbb{C}$ , we just have to substitute  $\Re(z)$  and  $\Re(w)$  in place of  $m, \mu$ , respectively, in the estimates (0.2).

<sup>3</sup>In the b-calculus setting, this condition implies that the underlying metric is polyhomogeneous: this is used, for instance, in [3].

It is simple to prove that  $\hat{\zeta}(A, z)$  is holomorphic for  $\Re(z) > \frac{n}{\mu}$ , in view of the fact that the hypotheses imply that the kernel  $K_{A^z}(x, x)$  is a function. As in the case treated in [35], we can look for meromorphic extensions of  $\hat{\zeta}(A, z)$ .

**Theorem 2.3.** *Let  $A \in L_{\text{cl}}^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be a SG-operator that admits complex powers. Then the function  $\hat{\zeta}(A, z)$  can be extended as a meromorphic function with, at most, poles at the points  $z_j = \frac{n-j}{\mu}$ ,  $j \in \mathbb{N}$ .*

Following the idea of M. Wodzicki [40], [20], we can now introduce a regularised version of non-commutative residue.

**Definition 2.4.** Let  $A \in L_{\text{cl}}^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be a  $\Lambda$ -elliptic operator that satisfies (A1). We define the regularised Wodzicki residue of  $A$  as

$$\widehat{\text{wres}}(A) := \mu \text{res}_{z=-1} \hat{\zeta}(A, z).$$

In the case  $\mu \in \mathbb{N}$ , using the explicit expression of the regularised integral and of the residues of  $\hat{\zeta}(A, z)$ , we get

$$(2.14) \quad \widehat{\text{wres}}(A) = \frac{1}{(2\pi)^n} \lim_{\rho \rightarrow \infty} \left[ \int_{|x| \leq \rho} \int_{|\xi|=1} a_{-n, \cdot}(x, \xi) dS(\xi) dx - \sum_{j=0}^{\mu+n-1} \frac{\beta_j}{n-j} \rho^{n-j} - \beta_{\mu+n} \log \rho \right]$$

where

$$\beta_j = \int_{|x|=1} \int_{|\xi|=1} a_{n-j, \cdot} dS(\xi) \widetilde{dS}(x),$$

$\widetilde{dS}(x)$  the metric induced by  $g$  on  $|x| = 1$ . The case  $\mu \notin \mathbb{Z}$  is not very interesting, since then  $\widehat{\text{wres}}(A)$  always vanishes, due to the fact that, in this case, the kernel  $K_{A^z}(x, x)$  has no poles at  $z = -1$ . The residue  $\widehat{\text{wres}}(\cdot)$  also vanishes on smoothing operators w.r.t. the  $\xi$ -variable, so it is well defined on the algebra  $\mathcal{A}$ . Incidentally, let us notice that the expression (2.14) is analogous to the functional  $\text{res}_\psi(A)$  defined by F. Nicola in [28], by means of holomorphic operator families.

### 3. A KASTLER-KALAU-WALZE TYPE THEOREM ON $\mathbb{R}^n$

First, we restrict to the case of  $\mathbb{R}^4$  and consider the classical Atiyah-Singer Dirac operator  $\mathcal{D}$  acting on the spinor bundle  $\Sigma \mathbb{R}^4$ . If the metric on  $\mathbb{R}^4$  satisfies Assumption (A2), it is immediate to verify that  $\mathcal{D} \in L_{\text{cl}}^{1,0}$ . Let  $\mathcal{D}^{-2}$  denote a weak parametrix of the square of the Dirac operator, that is  $\mathcal{D}^2 \circ \mathcal{D}^{-2} = I + R$ ,  $R \in L^{-\infty,0}$ . The calculus implies that  $\mathcal{D}^{-2} \in L_{\text{cl}}^{-2,0}$ . Via direct computation, following the idea of D. Kastler [21], it is possible to compute  $a_{-4, \cdot}(x, \xi)$ , the term of order  $-4$  in the asymptotic expansion w.r.t. the  $\xi$ -variable of the symbol of  $\mathcal{D}^{-2}$ . Evaluating the integral on the sphere w.r.t. the  $\xi$  variable one gets

$$\int_{|\xi|=1} a_{-4, \cdot}(x, \xi) dS(\xi) = -\frac{1}{24\pi^2} s(x).$$

So we have that

$$(3.15) \quad \widehat{\text{wres}}(\mathcal{D}^{-2}) = -\frac{1}{24\pi^2} \int s(x) dx.$$

The proof of (3.15) is contained in [7]. Let us notice the slight abuse of notation in (3.15), due to the fact that, in general,  $\mathcal{D}^{-2}$  does not satisfy Assumption (A1): anyway, we can use (2.14) as a definition of  $\widehat{\text{wres}}(\mathcal{D}^{-2})$  in this case.

In order to obtain a generalisation of (3.15) to higher dimensions and to more general operators, the direct approach seems to be rather cumbersome. For this

reason, we follow an idea of T. Ackermann [1] and exploit the properties of the asymptotic expansion of the heat kernel of generalised Laplacians.

As explained in the previous Section, if  $A \in L_{\text{cl}}^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , is  $\Lambda$ -elliptic and satisfies Assumption (A1), we can define the complex powers of  $A$  and the heat semigroup  $e^{-tA}$  as well:

$$e^{-tA} := \frac{i}{2\pi} \int_{\partial^+ \Lambda_c} e^{-t\lambda} (A - \lambda I)^{-1} d\lambda.$$

In [25] it has been proved that  $e^{-tA}$  is a SG-operator belonging to  $L^{-\infty,0}(\mathbb{R}^n)$ , so we can also consider the regularised heat trace  $\widehat{\text{TR}}(e^{-tA})$ . There is a deep link between regularised heat trace and  $\hat{\zeta}$ -function:

**Theorem 3.1.** *Let  $A \in L_{\text{cl}}^{\mu,0}(\mathbb{R}^n)$ ,  $\mu > 0$ , be an operator that admits complex powers. Then, for suitable constants  $c_{kl} = c_{kl}(A)$ , the following two asymptotic expansions hold:*

$$(3.16) \quad \Gamma(z) \hat{\zeta}(A, z) \sim \sum_{k=0}^{\infty} \sum_{l=0}^1 c_{kl} \left( z - \frac{n-k}{\mu} \right)^{-l+1}$$

$$(3.17) \quad \widehat{\text{TR}}(e^{-tA}) \sim \sum_{k=0}^{\infty} \sum_{l=0}^1 (-1)^l c_{kl} t^{-\frac{n-k}{\mu}} \log^l t, \quad t \searrow 0.$$

*Proof.* The statement follows by adapting the arguments given in [25] to the present situation. A main role in the proof is played by an abstract theorem by G. Grubb and R. Seeley [17], connecting  $\zeta$ -functions and heat traces.  $\square$

Let us now consider a generalised positive Laplacian  $\Delta \in L_{\text{cl}}^{-2,0}(E)$ , where  $E$  is a Hermitian vector bundle on  $\mathbb{R}^n$  with connection  $\nabla$ , that is

$$\Delta = \nabla^* \nabla + \mathcal{K}, \quad \mathcal{K} \in C^\infty(\text{End}(E)) \text{ symmetric endomorphism field.}$$

We require that  $\Delta$  satisfies Assumption (A1): in this way, we can define  $e^{-t\Delta}$  as above. In the case of closed manifolds, it is well known (see, e.g., [4]) that

$$(3.18) \quad K_{e^{-t\Delta}}(x, x) = k_t(x, x) \sim (4\pi t)^{-\frac{n}{2}} \left[ 1 \cdot \text{Id}_E + \left( \frac{1}{6} s(x) \text{Id}_E - \mathcal{K}_x \right) t + O(t^2) \right], \quad t \searrow 0.$$

where  $s(x)$  is the scalar curvature of the underlying manifold and the remainder term depends only on the connection and on the endomorphism field. The asymptotic expansion (3.18) also holds in the case of manifolds with cylindrical ends, since the computations are completely analogous and purely local, see [2]. The evaluation of the first term of the asymptotic expansion can be found in [24]: the expression of the second term then follows, using the properties of generalised Laplacians. In view of our hypotheses, the right hand side of (3.18) is a classical SG-symbol: then, we obtain

$$(3.19) \quad \widehat{\text{TR}}(e^{-t\Delta}) \sim (4\pi t)^{-\frac{n}{2}} \left\{ \int \text{Rk}(E) dx + t \int \left[ \frac{\text{Rk}(E)}{6} s(x) - \text{Trace}(\mathcal{K}_x) \right] dx + O(t^2) \right\}, \quad t \searrow 0.$$

Since, trivially, when  $h$  is a meromorphic function with a simple pole in  $z_0$ , the function  $\tilde{h}(z) = h(cz)$ ,  $c \in \mathbb{R}$ , is a meromorphic function with a simple pole in  $\frac{z_0}{c}$  and

$$\text{res}_{w=\frac{z_0}{c}} \tilde{h} = \frac{1}{c} \text{res}_{z=z_0} h,$$

we also have that

$$(3.20) \quad \begin{aligned} \widehat{\text{wres}}(\Delta^{-\frac{n}{2}+1}) &= (2-n) \text{res}_{z=-1} \hat{\zeta}(\Delta^{-\frac{n}{2}+1}, z) = 2 \text{res}_{z=\frac{n-2}{2}} \hat{\zeta}(\Delta, z) \\ &= 2\Gamma\left(\frac{n-2}{2}\right)^{-1} c_{2,0}(\Delta), \end{aligned}$$

where  $c_{2,0}(\Delta)$  is coefficient of the term of order  $t^{-\frac{n-2}{2}}$  in the asymptotic expansion (3.17). Finally, by (3.19) and the properties of  $\Gamma(z)$ ,

$$(3.21) \quad \widehat{\text{wres}}(\Delta^{-\frac{n}{2}+1}) = \frac{n-2}{\Gamma(\frac{n}{2})(4\pi)^{\frac{n}{2}}} \int \left[ \frac{\text{Rk}(E)}{6} s(x) - \text{Trace}(\mathcal{K}_x) \right] dx.$$

**Remark 5.** Assumption (A1) does not imply that  $\Delta$  is invertible, since we allow the origin to be an isolated point of  $\sigma(\Delta)$ . In view of this, the operator  $\Delta^{-\frac{n}{2}+1}$  has to be interpreted in the sense of the complex powers defined above.

If we consider a generalised Laplacian  $\Delta$ , then its principal homogeneous symbol is  $g^{jk}(x)\xi_j\xi_k = |\xi|^2 > 0$ ,  $\xi \neq 0$ .  $\Delta$  turns out to be always  $\Lambda$ -elliptic with respect to a suitable sector of the complex plane, while  $\sigma(\Delta)$  can admit the origin as an accumulation point. For example, it is well known that the classical Atiyah-Singer Dirac operator on  $\mathbb{R}^n$ , endowed with the canonical Euclidean metric, has no point spectrum, but the essential spectrum is the whole real line. In this case Assumption (A1) of course fails to be true<sup>4</sup>. A simple example such that Assumption (A1) is satisfied can be built in the following way. Let us consider a general Dirac operator  $D$ , defined on a Clifford bundle  $E$  over  $\mathbb{R}^n$ :  $D^2$  is then a generalised Laplacian and a non-negative operator. If we consider  $D_\epsilon^2 = D^2 + \epsilon I$ , we obtain an invertible generalised Laplacian, that clearly satisfies (A1). If we consider the classical Atiyah-Singer Dirac operator  $\mathcal{D}$ , formula (3.21) turns to

$$(3.22) \quad \widehat{\text{wres}}((D_\epsilon^2)^{-\frac{n}{2}+1}) = \frac{(n-2)2^{\lfloor \frac{n}{2} \rfloor}}{\Gamma(\frac{n}{2})(4\pi)^{\frac{n}{2}}} \left( -\frac{1}{12} \int s(x) dx - \epsilon \int dx \right).$$

On the other hand, a natural example of a metric on  $\mathbb{R}^n$  which can satisfy Assumption (A2) is an asymptotically flat one. In General Relativity, such a hypothesis on the metric is commonly assumed (e.g., in order to define the ADM-mass). Explicitly, we can consider a metric  $g$  such that, for a constant  $\alpha > 0$ ,

$$g_{jk}(x) - \delta_{jk} = O(|x|^{-\alpha}) \text{ outside a compact set } K \subset \mathbb{R}^n.$$

Moreover, restricting ourself to  $\mathbb{R}^4$ , if  $\alpha > 2$  the scalar curvature  $s(x)$  is integrable: in this case, (3.15) becomes

$$\widehat{\text{wres}}(\mathcal{D}^{-2}) = -\frac{1}{24\pi^2} \int s(x) dx.$$

The method above can be used to treat also the case of manifolds with cylindrical ends, using the contents of [8]: one defines in this setting a regularised Wodzicki Residue and exploits its connection with the zeta function. The asymptotic expansion of the heat kernel as  $t \searrow 0$  is locally defined, so, using suitable regularised integrals, see [8], the results can be generalised to those manifolds in this class which admit a spin structure. To keep this exposition at a reasonable length, we omit here the details.

<sup>4</sup>For further properties of the Dirac spectrum on open manifolds, the reader can refer, for instance, to the monograph by N. Ginoux [16].

## REFERENCES

- [1] T. Ackermann. A note on the Wodzicki residue. *J. Geom. Phys.*, 20(4):404–406, 1996.
- [2] P. Albin. A renormalized index theorem for some complete asymptotically regular metrics: The Gauss-Bonnet Theorem. *Advances in Mathematics*, Volume 213, Issue 1, 1 August 2007, Pages 1–52.
- [3] P. Albin and R. Mazzeo. Geometric construction of heat kernels: a users guide. In preparation, 2010.
- [4] B. Ammann and C. Bär. The Einstein-Hilbert action as a spectral action. In *Noncommutative geometry and the standard model of elementary particle physics (Hesselberg, 1999)*, volume 596 of *Lecture Notes in Phys.*, pages 75–108. Springer, Berlin, 2002.
- [5] C. Bär. The Dirac operator on hyperbolic manifolds of finite volume. *J. Diff. Geom.*, 54:439, 2000.
- [6] C. Bär. Spectral bounds for Dirac operators on open manifolds. *Ann. Global Anal. Geom.*, 36(1):67–79, 2009.
- [7] U. Battisti. Residuo di Wodzicki ed operatori di Dirac su  $\mathbb{R}^n$ , 2008. Tesi di Laurea Magistrale, Università di Torino (under the supervision of S. Coriasco).
- [8] U. Battisti and S. Coriasco. Wodzicki residue for operators on manifolds with cylindrical ends. Preprint, arXiv:1002.2804v1 [math.FA].
- [9] P. Boggiatto and F. Nicola. Non-commutative residues for anisotropic pseudo-differential operators in  $\mathbb{R}^n$ . *J. Funct. Anal.*, 203(2):305–320, 2003.
- [10] A. Connes. The action functional in noncommutative geometry. *Comm. Math. Phys.*, 117(4):673–683, 1988.
- [11] H.O. Cordes *The Technique of Pseudodifferential Operators*. Cambridge Univ. Press, 1995.
- [12] S. Coriasco, E. Schrohe, and J. Seiler. Bounded imaginary powers of differential operators on manifolds with conical singularities. *Math. Z.*, 244(2):235–269, 2003.
- [13] Y. Egorov and B.-W. Schulze. *Pseudo-Differential Operators, Singularities, Applications*. Operator Theory Advances and Application. Birkhäuser Verlag, Berlin, 1997.
- [14] B. V. Fedosov, F. Golse, E. Leichtnam, and E. Schrohe. The noncommutative residue for manifolds with boundary. *J. Funct. Anal.*, 142(1):1–31, 1996.
- [15] J. B. Gil and P. A. Loya. On the noncommutative residue and the heat trace expansion on conic manifolds. *Manuscripta Math.*, 109(3):309–327, 2002.
- [16] N. Ginoux. *The Dirac spectrum*, volume 1976 of *Lecture Notes in Mathematics*. Birkhäuser, 2009.
- [17] G. Grubb and R. T. Seeley. Zeta and eta functions for Atiyah-Patodi-Singer operators. *J. Geom. Anal.*, 6(1):31–77, 1996.
- [18] V. Guillemin. A new proof of Weyl’s formula on the asymptotic distribution of eigenvalues. *Adv. in Math.*, 55(2):131–160, 1985.
- [19] W. Kalau and M. Walze. Gravity, non-commutative geometry and the Wodzicki residue. *J. Geom. Phys.*, 16(4):327–344, 1995.
- [20] C. Kassel. Le résidu non commutatif (d’après M. Wodzicki). *Astérisque*, (177-178):Exp. No. 708, 199–229, 1989. Séminaire Bourbaki, Vol. 1988/89.
- [21] D. Kastler. The Dirac operator and gravitation. *Comm. Math. Phys.*, 166(3):633–643, 1995.
- [22] M. Kontsevich and S. Vishik. Determinants of elliptic pseudo-differential operators, 1994. <http://www.citebase.org/abstract?id=oai:arXiv.org:hep-th/9404046>.
- [23] M. Kontsevich and S. Vishik. Geometry of determinants of elliptic operators. In *Functional analysis on the eve of the 21st century, Vol. 1 (New Brunswick, NJ, 1993)*, volume 131 of *Progr. Math.*, pages 173–197. Birkhäuser Boston, Boston, MA, 1995.
- [24] L. Maniccia and P. Panarese. Eigenvalue asymptotics for a class of md elliptic Pdos on manifolds with cylindrical exits *Annali di Matematica Pura ed Applicata*, Volume 181, Number 3, 283–308.
- [25] L. Maniccia, E. Schrohe, and J. Seiler. Determinants of classical SG-pseudodifferential operators. Preprint <http://www.ifam.uni-hannover.de/seiler/artikel/ifam86.pdf>.
- [26] L. Maniccia, E. Schrohe, and J. Seiler. Complex powers of classical SG-pseudodifferential operators. *Ann. Univ. Ferrara Sez. VII Sci. Mat.*, 52(2):353–369, 2006.
- [27] R. Melrose. The Atiyah-Patodi-Singer index theorem. *Research Notes in Mathematics*, 4. A K Peters, Ltd., Wellesley, MA, 1993.
- [28] F. Nicola. Trace functionals for a class of pseudo-differential operators in  $\mathbb{R}^n$ . *Math. Phys. Anal. Geom.*, 6(1):89–105, 2003.
- [29] C. Parenti. Operatori pseudodifferenziali in  $\mathbb{R}^n$  e applicazioni. *Ann. Mat. Pura Appl.*, 93:359–389, 1972.
- [30] A. Pazy. Semigroups of operators in Banach spaces. In *Equadiff 82 (Würzburg, 1982)*, volume 1017 of *Lecture Notes in Math.*, pages 508–524. Springer, Berlin, 1983.
- [31] R. Ponge. Noncommutative geometry and lower dimensional volumes in Riemannian geometry. *Lett. Math. Phys.*, 83(1):19–32, 2008.

- [32] E. Schrohe. Spaces of weighted symbols and weighted Sobolev spaces on manifolds. In *Pseudodifferential operators (Oberwolfach, 1986)*, volume 1256 of *Lecture Notes in Math.*, pages 360–377. Springer, Berlin, 1987.
- [33] E. Schrohe. Complex powers on noncompact manifolds and manifolds with singularities. *Math. Ann.*, 281(3):393–409, 1988.
- [34] E. Schrohe. Wodzicki’s noncommutative residue and traces for operator algebras on manifolds with conical singularities. In *Microlocal analysis and spectral theory (Lucca, 1996)*, volume 490 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 227–250. Kluwer Acad. Publ., Dordrecht, 1997.
- [35] R. T. Seeley. Complex powers of an elliptic operator. In *Singular Integrals (Proc. Sympos. Pure Math., Chicago, Ill., 1966)*, pages 288–307. Amer. Math. Soc., Providence, R.I., 1967.
- [36] H. M. Srivastava and J. Choi. *Series associated with the zeta and related functions*. Kluwer Academic Publishers, Dordrecht, 2001.
- [37] Y. Wang. Differential forms and the noncommutative residue for manifolds with boundary in the non-product case. *Lett. Math. Phys.*, 77(1):41–51, 2006.
- [38] Y. Wang. Differential forms and the Wodzicki residue for manifolds with boundary. *Journal of Geometry and Physics*, 56:731, 2006.
- [39] Y. Wang. Lower dimensional volumes and the Kastler-Kalau-Walze type theorem for manifolds with boundary. <http://www.citebase.org/abstract?id=oai:arXiv.org:0907.3012>.
- [40] M. Wodzicki. Noncommutative residue. I. Fundamentals. In *K-theory, arithmetic and geometry (Moscow, 1984–1986)*, volume 1289 of *Lecture Notes in Math.*, pages 320–399. Springer, Berlin, 1987.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, ITALY  
E-mail address: ubertino.battisti@unito.it

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, ITALY  
E-mail address: sandro.coriasco@unito.it