

## SOLAR WIND SPECTRAL ANALYSIS IN HELIOSHEATH FROM VOYAGER 2 DATA

Federico Fraternali<sup>1</sup>, Luca Gallana<sup>1</sup>, Sophie M. Fosson<sup>2</sup>, Enrico Magli<sup>2</sup>, Merav Opher<sup>3</sup>, John D. Richardson<sup>4</sup>, Michele Iovieno<sup>1</sup> & Daniela Tordella<sup>1</sup>

<sup>1</sup>*Dipartimento di Ingegneria Meccanica ed Aerospaziale, Politecnico di Torino, Torino, Italy*

<sup>2</sup>*Dipartimento di Elettronica e Telecomunicazioni, Politecnico di Torino, Torino, Italy*

<sup>3</sup>*Astronomy Department, Boston University, Boston, USA*

<sup>4</sup>*Kavli Institute for Astrophysics and Space Research, Massachusetts Institute of Technology, Boston, USA*

**Abstract** The solar wind is a supersonic flow of magnetized plasma. It is time-dependent on all scales and expands with distance. The flow has fluctuations on a broad range of scales and frequencies. These fluctuations are not just convected outward but show energy cascades among the different scales. The solar wind turbulence peculiar phenomenology has been comprehensively reviewed by Tu and Marsch [11] and Bruno and Carbone [2]. As the distance from the sun increases, the available data on plasma and magnetic field become increasingly scarce. At distance of the order of 1 astronomical unit (AU), several measurements have been performed by various crafts, but, nowadays, only the Voyager spacecrafts can measure data in the heliosheath, the outermost layer in heliosphere where the solar wind is slowed by the pressure of the interstellar gas, and only the Voyager 2 craft can measure both plasma and magnetic fields (Voyager 1 can measure only the magnetic field, and Pioneer 10 and 11 has ceased communications). Taken together, the Voyager 1 and 2 probes have collected over 11 years of data in the heliosheath. The Voyager plasma experiment observes plasma currents in the energy/charge range 10 – 5950 eV/q using four modulated-grid Faraday cup detectors [1]. The observed currents are fit to convected isotropic proton Maxwellian distributions to derive the parameters (velocity, density, and temperature) used in this work. Magnetic field and plasma data are taken from the COHO web site and MIT Space Plasma Group repository.


Several studies have been done in order to extend the existing models to make them consistent with the energetic particles and magnetic fields measured in the heliosheath, but so far an exhaustive explanation has not yet been obtained. In particular, the differences between the energetic particle intensity variations seen by the two crafts are unexplained. The electron intensity measured by Voyager 2 varies steeper by a factor of 10 in a single year, while the same quantity from Voyager 1 changes gradually over time.[7] A possible explanation can be the presence of a bubble of turbulence that travels in the heliosheath. Therefore, a characterization of turbulence and its intermittency is necessary to explain this phenomenology. The aim of this work is to provide the first spectral analysis of heliosheath solar wind, trying to characterize the plasma turbulence in that region by estimating the spectral slopes. A first result is represented in figure 2, where it can be seen that the low frequency spectral slope is lower when the electron intensity is low.

In order to compute spectra, signal reconstruction techniques are mandatory: at distance over 80 AU, available data are very spotty. For the plasma velocity, there are 97% of missings due to unsteadiness in the signals, see 1, the most important of which are: tracking gaps due to the V2 location and due to limited deep space network availability; interference from other instruments; possible errors in the measurement chain (from the Faraday cups up to the data acquisition system and the signal shipping to Earth); the temporal sequence of the nuclear propulsion used to control the Voyager trajectory and to assist in several critical repositionings of the craft.


For data recovery, we mainly use two different methods. The first method used is based on the correlation computation [8] that allows to reconstruct correlations and use it to compute spectra. Better results can be achieved implementing the maximum likelihood reconstruction by Rybicki and Press [10] based on a minimum-variance recovery with a stochastic component. The second method comes from Compressed Sensing, a recent technique widely used in telecommunications, that provides the reconstruction of the signal from a sparse dataset [3], by using sparse Fourier matrices [9]. The methods used have been previously tested on 1979 data and on synthetic fluid turbulent fields. Results were in good agreement with the literature, and allow to compute largest spectra of solar wind at 5 AU, with frequencies ranging from  $10^{-7}$  to  $10^{-2}$  Hz [4, 5, 6].

## References

- [1] H. S. Bridge, J. W. Belcher, R. J. Butler, A. J. Lazarus, A. M. Mavretic, J. D. Sullivan, G. L. Siscoe, and V. M. Vasyliunas. The plasma experiment on the 1977 voyager mission. *Space Sci. Rev.*, **21**:259–287, 1977.
- [2] R. Bruno and V. Carbone. The solar wind as a turbulence laboratory. *Living Rev. Solar Phys.*, **10**(2), 2013.
- [3] D. L. Donoho. Compressed sensing. *IEEE T. Inform. Theory*, **52**:1289–1306, 1977.
- [4] F. Fraternali, L. Gallana, M. Opher, J. D. Richardson, M. Iovieno, and D. Tordella. Turbulence in the solar wind: spectra from voyager 2 data at 5 au. *Physica Scripta*, submitted.
- [5] F. Fraternali, L. Gallana, J. D. Richardson, M. Iovieno, and D. Tordella. Turbulence in the solar wind: spectra from voyager 2 data at 5 au. In *Proceedings of the Turbulent Mixing and Beyond Workshop*. ICTP.
- [6] L. Gallana, F. Fraternali, M. Opher, J. D. Richardson, M. Iovieno, and D. Tordella. Spectra and correlations in the solar wind from voyager 2 around 5 au. In *Proceedings of the 67th Annual Meeting of the APS Division of Fluid Dynamics*. APS.
- [7] M. E. Hill, R. B. Decker, L. E. Brown, J. F. Drake, D. C. Hamilton, S. M. Krimigis, and M. Opher. Dependence of energetic ion and electron intensities on proximity to the magnetically sectored heliosheath: Voyager 1 and 2 observations. *J. Plasma Phys.*, **781**:94–100, 2014.
- [8] W. H. Matthaeus and M. L. Goldstein. Measurement of the rugged invariants of magnetohydrodynamic turbulence in the solar wind. *J. Geophys. Res.*, **87**(16):6011–6028, 1982.
- [9] H. Rauhut. *Theoretical Foundations and Numerical Methods for Sparse Recovery*, **9** of *Radon Series Comp. Appl. Math.* M. Fornasier, 2010.
- [10] G. B. Rybicki and W. H. Press. Interpolation, realization, and reconstruction of noisy, irregularly sampled data. *The Astroph. J.*, **398**, 1992.
- [11] C.-Y. Tu and E. Marsch. Mhd structures, waves and turbulence in the solar wind: observations and theories. *Space Sci. Rev.*, **73**:1–210, 1995.



**Figure 1.** Plot of plasma velocity (red lines) and magnetic field components in Alfvénic units (blue lines) for the first 180 days in 2008 in the Heliocentric Radial Tangential Normal spherical reference frame. In the top panel, to highlight the gap presence, a magnification of a four days period is shown.



**Figure 2.** Kinetic energy spectra in the heliosheath computed through correlation method on hourly averages data in different periods after the termination shock. The periods of observation (indicated in years and fraction of year) are chosen in order to show the differences between different parts of the heliosheath, as seen by the Voyager 2 (unipolar and sectored heliosheath regions [7]), which are characterised by different electron intensity (measured as a flux of particles per unit energy). The low frequencies spectral slope module is 0.7, in presence of high electron intensity (sectored heliosheath). When the electron intensity decreases (unipolar heliosheath) the slope decreases to values around 0.3.