



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Monocentric survey about the use of Raltitrexed in the daily clinical practice for patients with metastatic colorectal cancer.

Original

Monocentric survey about the use of Raltitrexed in the daily clinical practice for patients with metastatic colorectal cancer. / Ponzetti, A.; Zaccaria, G.M.; Bustreo, S.; Mecca, C.; Spadi, R.; Fanchini, L.; Ritorto, G.; Zanini, M.; Ciuffreda, L.; Racca, P.. - In: ANNALS OF ONCOLOGY. - ISSN 0923-7534. - ELETTRONICO. - 27 Issue supplementary 4(2016), pp. 53-54. ((Intervento presentato al convegno XVIII National Congress of Medical Oncology, 28-30 October 2016, Rome, Italy tenutosi a Roma nel 28-30 October 2016.

Availability:

This version is available at: 11583/2655384 since: 2017-06-21T17:47:29Z

Publisher:

Oxford University

Published

DOI:10.1093/annonc/mdw335.50

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

XVIII CONGRESSO NAZIONALE AIOM

Roma, 28-30 ottobre 2016
Marriott Park Hotel



16/05/2016 13.14.35

Primo autore

Ponzetti Agostino (Medico)
Oncologia Medica 1, Città della Salute e della Scienza - Torino
agoponz@hotmail.com Tel.3482735282
Età uguale o inferiore a 40 anni
Sono socio AIOM

Area tematica

Gastrointestinal (colorectal) cancers

Titolo

Monocentric survey about the use of Raltitrexed in the daily clinical practice for patients with metastatic colorectal cancer

Testo

Background: The main setting of use of Raltitrexed for patients with colorectal cancer is when cardiac toxicity is feared or when a deficit of Dihydropyrimidine (DPD) dehydrogenase has been demonstrated. However, this drug is used as an alternative to fluororacil in patients with older age or in third or further line of treatment when exposure to fluoropyrimidines has been huge. However, data from recent clinical trials about these possible settings are scarce and case series can show the real clinical relevance of this drug. Here we describe our clinical experience in using raltitrexed in a tertiary care hospital with a specific Colorectal Cancer Unit.

Patients and methods: data about patients receiving Raltitrexed at our Institution since January 2015 were retrospectively collected by using electronic charts records. Patients' data and data about scheme of administration, response rate and toxicities were collected. Patients were analyzed according to the clinical setting of administration:

- A) 1st line treatment in patients with cardiac comorbidities;
- B) 1st line treatment in older patients;
- C) third or further line treatment after FOLFOX and FOLFIRI;
- D) Deficit of DPD

Results: globally since January 2015 thirty-two patients received Raltitrexed at our hospital, all for metastatic colorectal cancer (mCRC). Patients belonged to clinical setting A, B or C while none to group D.

The following Table resumes the main results of our survey:

Group	N- median age - % male	Characteristics	Benefits	G3 or G4 toxicities*
A) Cardiac comorbidities	6 pts – 70 years – 100% male	100% 1 st line, 50 % ECOG PS 1, 85,7% doublet	ORR 50% (2 pts not evaluable), CB 75%	33,3%
B) Older Age	2 pts – 82 years – 100% male	100% 1 st line, 100% monotherapy	ORR 50%, no SD	100%
C) Further line	24 pts – 64 years – 45,8% male	33,3% 4 th -6 th line, 54% ECOG PS 1, 50% doublet (45% TOMOX)	ORR 0% (3 pts not evaluable), CB 9,5% Relief from symptoms 25%	16,7%

Doublet = TOMOX or TOMIRI. ORR= overall response rate. CB = ORR + stable disease.

Globally 11 patients (45,8%) had G0 or G1 toxicities while none experienced G3 renal failure from Raltitrexed.

Conclusions: the administration of Raltitrexed is a part of daily clinical practice at our center in patients with mCRC and cardiac comorbidities or older age with relevant efficacy but frequent severe toxicities. The use in the very advanced setting can bring a clinical benefit/relief from symptoms in up to 25% of patients but with severe toxicities in up to 17%.

Parole Chiave

1. Raltitrexed
2. Metastatic Colorectal Cancer
3. Safety

Co-autori

1. **Zaccaria Gian Maria** (Altro)
Dipartimento di Elettronica e Telecomunicazioni, Politecnico - Torino
2. **Bustreo Sara** (Medico - Età uguale o inferiore a 40 anni)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
3. **Mecca Caterina** (Medico - Età uguale o inferiore a 40 anni)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
4. **Spadi Rosella** (Medico - Età uguale o inferiore a 40 anni)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
5. **Fanchini Laura** (Medico)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
6. **Ritorto Giuliana** (Medico)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
7. **Zanini Marcello** (Medico)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
8. **Ciuffreda Libero** (Medico)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino
9. **Racca Patrizia** (Medico)
Colorectal Cancer Unit, Oncologia Medica 1, Città della Salute e della Scienza - Torino