

Hydrogeochemical study of Bossea karst system

Original

Hydrogeochemical study of Bossea karst system / Fiorucci, A., Moitre, B., Vigna, B.. - STAMPA. - Unico:(2015), pp. 290-294. (Environmental safety and construction in karst areas Perm 26-29 maggio 2015).

Availability:

This version is available at: 11583/2667665 since: 2017-03-24T16:10:03Z

Publisher:

Perm State University

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Hydrogeochemical study of Bossea karst system

A. Fiorucci, B. Moitre & B. Vigna

Politecnico di Torino – Dipartimento di Ingegneria dell'Ambiente, del Territorio e delle Infrastrutture, C.so Duca degli Abruzzi 24, Torino, Italy, email: adriano.fiorucci@polito.it, barbara.moitre@tin.it, bartolomeo.vigna@polito.it




Figure 1. Study area.

ABSTRACT: The present work concerns the geochemical characterization of the water circulating in the Bossea karst system. In the cavity were sampled, in different hydrodynamic conditions, the waters of the main collector (Mora torrent) and numerous water supplies from drainage network of the unsaturated zone above the cavity and that flow into the main collector. Some seepages are located at the tectonic contact between the rocks of the metamorphic basement (metavolcanic) and carbonate coverage, others come directly from the network of cracks in the limestone and dolomite. The flow of these seepages generally is very low, less than $1 \text{ L}\cdot\text{s}^{-1}$. The characterization of the individual contributions and the main collector was performed through comparison of the major elements concentration using the Schöeller diagram and have also considered other parameters such as the lanthanides content, the calcite and dolomite saturation index. The data related to the lanthanides (REE - Rare Earth Elements) content normalized using the PAAS (Post-Archean Australian Shale) contribute to diversify the individual contributions that have different anomalies (especially cerium and europium) and/or trends in relation to the sampling period. Finally particularly interesting data have emerged from the correlation between the saturation indices of calcite and dolomite. For all the individual seepages is observed good correlation between the two indices, correlation is less marked when you consider all the values of the karst system under consideration. The data obtained shows, therefore, further differentiation between the individual seepages. From the overall analysis of the geochemical data obtained emerging a substantial complexity of the Bossea Karst system already highlighted by the monitoring data of flow rate, temperature and specific electrical conductivity of the water coming from the individual seepages and from the main collector.

KEYWORDS: hydrogeochemistry, saturation index, Bossea cave, rare earth elements.

1 STUDY AREA

The Bossea Karst system is located in the southern Piedmont, in the Ligurian Alps, at an altitude between 800 and 1700 m above sea level. The main absorption area is located between the Corsaglia Valley and the Maudagna Valley.

1.1 Geological and hydrogeological layout

The area is characterized by a complex tectonic history that brought the carbonate sediments, originally have built up over an ancient substrate of permocarboniferous age (represented by metavolcanics and quartzites characterized by a very low permeability), to instead be interspersed in large outcrops separated from each other by important surfaces of dislocation which border compartments hydro-

geologically independent, sometimes interconnected with each other in a particularly complex (figure 1).

The Bossea system is characterized by limestone and dolomitic limestone, laterally confined by the rocks of the metamorphic basement, metavolcanics and quartzites, through a series of sub-vertical tectonic contacts.

The carbonate aquifer is characterized by a relatively high permeability, with an underground circulation set to a main collector that receives the contributions coming from limestone-dolomitic storage and from rocks of the metamorphic basement. These rocks form a secondary aquifer, set along the discontinuity that border the carbonate structure and which, through a series of underground transfers, feeds the main aquifer (Peano et al. 2011; Banzato et al. 2011; Vigna & Doleatto 2008).

1.2 Groundwater flow description

The Bossea cave is a cavity collector, Mora Torrent, the main collector, is fed by numerous springs in proximity of the Mora and Saglia Torrent. The collector receives numerous secondary contributions by the circulation in the cavity also by the surface water infiltration. The rocks of the metamorphic basement water contributions are abundant. The concentrated infiltration in riverbeds and at tectonic contact (figure 1).

The seepages more important are composed of contributions from open and karstified, usually located between the carbonate aquifer and the basement (Polla delle Anate, Polletta). The secondary seepages "stillicidi" and distributed in the cavity, are located on the contact derived from fractures massive carbonate deposits (Stillicidio Milanese, Stillicidio Sacrestia). They have flow rates less than $0.008 \text{ L}\cdot\text{s}^{-1}$, with variations only in particular situations.

Near the catchment area there are numerous springs with moderate flow rates rather constant in time, recharged by the metamorphic basement. One of these, Sorgente dei Matti, was taken as representative of the underground water in the metamorphic rocks (figures 1 and 2).

2 HYDROCHEMISTRY OF INDIVIDUAL CONTRIBUTIONS

The water of Bossea Karst system shows a moderate mineralization between 100 and $200 \text{ mg}\cdot\text{L}^{-1}$ in terms of TDS. The mineralization in the waters of the main collector and those circulating in the




Figure 1. Study area.

1.2 Groundwater flow description

The Bossea cave is a cavity crossed by a mainkarst collector, Mora Torrent, that feeds a series of localized springs in proximity of the riverbed of the Corsaglia Torrent. The collector receives the waters by numerous secondary contributions and is fed in part by the circulation in the carbonate rocks and in part also by the surface water contributions flowing in the rocks of the metamorphic basement. The surface water contributions are absorbed by a series of concentrated infiltration in riverbed in proximity of the contact (figure 1).

The seepages more important, called “*polle*”, are composed of contributions from fractures relatively open and karstified, usually present at the contact between the carbonate aquifer and the underlying basement (*Polla delle Anatre*, *Polla dell’Orso* and *Polletta*). The secondary seepages, denominated “*stillicidi*” and distributed in different sections of the cavity, are located on the cave vault and are usually derived from fractures masked by abundant calcite deposits (*Stillicidio Milano*, *Stillicidio Torre*, *Stillicidio Sacrestia*). They have a very low flow, lower than $0.008 \text{ L}\cdot\text{s}^{-1}$, with variations closely related to external weather conditions and become inactive only in particular situations of drought.

Near the catchment area are present a number of springs with moderate flow, around $0.5\text{--}2 \text{ L}\cdot\text{s}^{-1}$, but rather constant in time, recharged by the rocks of the metamorphic basement. One of these sources, called *Sorgente dei Matti*, was taken as “*sample spring*” representative of the underground circulation set in metavolcanic rocks (figures 2, 3).

2 HYDROCHEMISTRY CHARACTERIZATION OF INDIVIDUAL CONTRIBUTIONS

The water of Bossea Karst system presents an average mineralization between 162.23 and $409.51 \text{ mg}\cdot\text{L}^{-1}$ in terms of *TDS*. The lowest values are found in the waters of the main collector (*Torrente Mora*) and those circulating in metavolcanic rocks (*Sorgente dei*

Matti) while the higher ones belong to the individual seepages.




Figure 2. Collector flow and electrical conductivity.




Figure 3. *Stillicidio Milano* level and electrical conductivity.

The hydrochemical facies found in several samples remain essentially constant over time but are substantial differences between the different sampling points (figure 4). *Bicarbonate-calcium-magnesium facies* are observed in the seepages called *Milano* and *Polletta* and *bicarbonate-calcium facies* in other seepages, in the main collector and in water circulating in the metavolcanic rocks. In the *bicarbonate-calcium facies* can be seen, however, differences in the $\text{Ca}^{2+}/\text{Mg}^{2+}$ ratio ranging from 4.11 (*Polla Anatre* – mean value) to 41.49 (*Sacrestia* – mean value). This ratio report presents average values of 6.99 in main collector and 21.96 in the waters circulating in metavolcanic rocks. The wide variability of the $\text{Ca}^{2+}/\text{Mg}^{2+}$ ratio is due to the different contents of the ion Mg^{2+} which is the only parameter, including the main ones, to differentiate the different seepages. The parameters with concentrations lower than $0.1 \text{ meq}\cdot\text{L}^{-1}$ (alkali, chlorides and sulfates) are subject to greater variations in time.

Sorgente dei Matti and *Polla delle Anatre* are characterized by alkali concentrations greater than those of the chlorides. This fact shows the presence of water coming mainly from metavolcanic rocks.

3 LANTHANIDES FOOTPRINT

The lanthanides are a family of 15 chemical elements with chemical properties very similar to each other that in the hydrogeological field can be used

for the characterization of the aquifers. The concentrations and distributions depend specifically by the different rocks with which the water came in contact, and despite the low concentrations present specific trends for different waters (Biddau et al. 2009; Banks et al. 1999; Fiorucci & Moitre 2012). Moreover, at present, do not have a full-blown human footprint, and for this reason can be used for the study of the natural geochemical fund.





Figure 4. Schöeller diagrams.

The rare earth elements concentration in the basal complex spring remains almost constant in the different sampling and seems quite high (average value equal to $765 \text{ ng}\cdot\text{L}^{-1}$). The seepages, instead, are

characterized by a different behavior and highlight remarkable changes in the course of the sampling.

The concentrations ranges of the seepages are quite large. *Polla delle Anatre* is the one with major variations, between $21 \text{ ng}\cdot\text{L}^{-1}$ (high flow condition) to $1481 \text{ ng}\cdot\text{L}^{-1}$ (low flow condition). The minimum and maximum concentrations are found in the same periods for *Stillicidio Milano*, the difference is in the values (40 and $180 \text{ ng}\cdot\text{L}^{-1}$). High values, above $400 \text{ ng}\cdot\text{L}^{-1}$, are found in two samples in the *Stillicidio Sacristia* and in a sampling of the main collector ($570 \text{ ng}\cdot\text{L}^{-1}$).

The concentrations of the REE were normalized with the *Post-Archean Australian Shale - PAAS* (McLennan 1989), to highlight the changes in time and the differences between the main collector, secondary seepages and the basal complex springs (figures 5-12).

Unlike Schöeller diagrams, the diagrams which characterize the relationship between the REE and PAAS not remain constant in time but show differences between the different sampling periods. The only exception is *Sorgente dei Matti*, whose trend remains unchanged in time. These differences may indicate that the Bossea Karst system is a complex system, in which is possible to highlight different catchment circuits between the various seepages and the main collector. Furthermore, important quantitative and trend changes are observed in the same seepages, in different hydrodynamic conditions.




Figure 5. Torrente Mora REE/PAAS concentration.




Figure 6. Stillicidio Milano REE/PAAS concentration.




Figure 7. Sorgente dei Matti REE/PAAS concentration.




Figure 8. Torre REE/PAAS concentration.




Figure 9. Polla Orso REE/PAAS concentration.




Figure 10. Polletta REE/PAAS concentration.




Figure 7. *Sorgente dei Matti* REE/PAAS concentration.




Figure 11. *Sacrestia* REE/PAAS concentration.




Figure 8. *Torre* REE/PAAS concentration.




Figure 12. *Polla Anatre* REE/PAAS concentration.




Figure 9. *Polla Orso* REE/PAAS concentration.




Figure 10. *Polletta* REE/PAAS concentration.

4 SATURATION INDICES

The calcite and dolomite saturation index were calculated using the ratio between the product of ionic activity (with the single ion activity determined with the equation of Debye-Hückel) and the solubility product.

The waters in almost all samples are to be super-saturated in respect of calcite, while recording a considerable number of cases in which they are under-saturated in respect of dolomite.

Considering all the samples there is a fairly good correlation (figure 13) between the two indices (r^2 equal to 0.7738), evaluating the correlation for each contribution is noted that these have values of correlation coefficient always greater than 96 % and angular coefficients of the correlation lines are very similar, varying from a minimum of 0.4330 (*Sorgente dei Matti*) to a maximum of 0.5036 (*Polla delle Anatre*).

5 CONCLUSIONS

The geochemical data show a substantial complexity of the Bossea Karst system already highlighted by the monitoring data of flow rate, temperature and specific electrical conductivity of the water of seep-ages and main collector.




Figure 13. Correlation calcite and dolomite saturation index.

The analysis of facies, performed considering only the major elements, highlights the differences between the individual contributions due substantially to the $\text{Ca}^{2+}/\text{Mg}^{2+}$ ratio and, in some cases, $(\text{Na}^+ + \text{K}^+)/\text{Cl}^-$ ratio. This last, when it assumes values much greater than 1 indicates a water circulation that involves the parts not carbonate of the system (*Polla delle Anatre*, *Sorgente dei Matti* and occasionally also the other seepages).

A clearer distinction between the individual seepages emerges from the data on the lanthanides content, which also seems to show the differences, relative to each seepages, linked to the different sampling periods.

The saturation indices calculated indicate waters almost always supersaturated in respect of calcite and also largely for the dolomite. The correlation between the two indices show high values when considering the individual seepages, while it is lower when comparing the overall data. This aspect seems to further highlight the complexity of the Bossea Karst system as already shown by other geochemical indices. Every circuit is independent of the others, but instead the collector receives the waters of all the karst and non-karst networks.

REFERENCES

- Banks, D., Reimann, C., Siewers, U. 1999. Distribution of rare earth elements in crystalline bedrock groundwaters: Oslo and Bergen region, Norway. *Appl. Geochem.* Vol. 14, pp. 27-39.
- Banzato, C., Dallagiovanna, G., Maino, M., Peano, G. & Vigna B. 2011. Correlation between the geological setting and groundwater flow: the Bossea karst underground laboratory. *Geitalia 2011, Epitome 4, Torino, 19-23 settembre 2011*. 14 p.
- Biddau, R., Bensimon, M., Cidu, R. & Parriaux, A. 2009. Rare earth elements in groundwaters from different Alpine aquifers. *Chemie der Erde-Geochemistry*. Vol. 69, pp. 327-339.
- Fiorucci, A. & Moitre, B. 2012. Hydrogeochemical study on the rare earth concentration in groundwater. *Romanian Journal of mineral Deposits*. Geological Institute of Romania, Society of Economic Geology of Romania. Vol. 85, pp. 56-59.

О зоне силикатного типа уралья

А.Я. Гаев

Оренбургский научный центр УрО РАН

АННОТАЦИЯ: Модель карстосферы сферно-мантийной средой. Гидросфера и флюиды в астеносфере и флюидными породами земной коры и в растворимости кристаллических силикатного карста и разработку

КЛЮЧЕВЫЕ СЛОВА: карстосфера

Г.А. Максимович выделил в разрезе ского фундамента платформ зону за, опираясь на терминологию (Максимович, 1964). Способность недр растворять даже вторичные этой зоне он отметил, анализируя мы в разрезах самых глубоких рудного золота Южной Африки. Он указал на наличие там силикатных пород занимают в разрезе толщи сотни километров в земной мантии. Это несоизмеримо, как и по объему с осадочной оболочкой соизмерима по масштабам области катного карста.

Однако, традиционно карстующими считаются только известняки, гели, терригенно-карбонатные породы гидриты и соли, которые занимают в разрезах континентов не менее осадочного чехла и только осадочная земной коры. Среди карстующих обладают карбонаты, составляющие 1.7% от веса земной коры. Они глубин порядка 5 км. Процессы глубже, почти не изучены.

Эксперименты со «сверхводоносными» породами, подверженными комбинационному рентгеновскому исследователи подвергли маленьким