

Composition law of -entropy for statistically independent systems

Original

Composition law of -entropy for statistically independent systems / Kaniadakis, Giorgio; Scarfone, ANTONIO MARIA; Sparavigna, Amelia Carolina; Wada, T.. - In: PHYSICAL REVIEW. E. - ISSN 2470-0045. - STAMPA. - 95:5(2017), p. 052112. [10.1103/PhysRevE.95.052112]

Availability:

This version is available at: 11583/2670532 since: 2017-06-06T12:10:32Z

Publisher:

American Physical Society

Published

DOI:10.1103/PhysRevE.95.052112

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Composition law of κ -entropy for statistically independent systems

G. Kaniadakis,^{1,*} A. M. Scarfone,² A. Sparavigna,¹ and T. Wada³

¹*Department of Applied Science and Technology, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

²*Istituto dei Sistemi Complessi (ISC-CNR) c/o Politecnico di Torino Corso Duca degli Abruzzi 24, 10129 Torino, Italy*

³*Department of Electrical and Electronic Engineering, Ibaraki University, Nakanarusawa, Hitachi, Ibaraki 316-8511, Japan*

(Received 2 March 2017; published 8 May 2017)

The intriguing and still open question concerning the composition law of κ -entropy $S_\kappa(f) = \frac{1}{2\kappa} \sum_i (f_i^{1-\kappa} - f_i^{1+\kappa})$ with $0 < \kappa < 1$ and $\sum_i f_i = 1$ is here reconsidered and solved. It is shown that, for a statistical system described by the probability distribution $f = \{f_{ij}\}$, made up of two statistically independent subsystems, described through the probability distributions $p = \{p_i\}$ and $q = \{q_j\}$, respectively, with $f_{ij} = p_i q_j$, the joint entropy $S_\kappa(pq)$ can be obtained starting from the $S_\kappa(p)$ and $S_\kappa(q)$ entropies, and additionally from the entropic functionals $S_\kappa(p/e_\kappa)$ and $S_\kappa(q/e_\kappa)$, e_κ being the κ -Napier number. The composition law of the κ -entropy is given in closed form and emerges as a one-parameter generalization of the ordinary additivity law of Boltzmann-Shannon entropy recovered in the $\kappa \rightarrow 0$ limit.

DOI: [10.1103/PhysRevE.95.052112](https://doi.org/10.1103/PhysRevE.95.052112)

I. INTRODUCTION

The one-parameter generalized entropy defined through

$$S_\kappa(f) = - \sum_i f_i \ln_\kappa(f_i), \quad (1.1)$$

with

$$\ln_\kappa(x) = \frac{x^\kappa - x^{-\kappa}}{2\kappa}, \quad (1.2)$$

and $0 < \kappa < 1$, $f = \{f_i; i = 1, \dots, N; \sum_i f_i = 1\}$, is considered to be a good candidate for the relativistic extension of Boltzmann-Shannon entropy, which is recovered in the $\kappa \rightarrow 0$ limit [1,2]. The κ -logarithm $\ln_\kappa(x)$ emerges as the relativistic generalization of the ordinary logarithm $\ln(x)$, which is obtained in the same limit. Its inverse function, namely, the κ -exponential, is given by

$$\exp_\kappa(x) = (\sqrt{1 + \kappa^2 x^2} + \kappa x)^{1/\kappa}, \quad (1.3)$$

and it is obtained within the special relativity, starting directly from the dynamical Lorentz transformations [3].

It has been shown that, in case of a particle system with energy levels E_i , the maximization of entropy (1.1) under the $\sum_i f_i = 1$ and $\sum_i f_i E_i = \langle E \rangle$ constraints, according to the Maximum Entropy Principle [4], leads to the probability distribution

$$f_i = \frac{1}{e_\kappa} \exp_\kappa(-\gamma_\kappa \beta E_i + \gamma_\kappa \beta \mu), \quad (1.4)$$

where β and $-\beta\mu$ are the usual Lagrange multipliers, while the constants γ_κ and e_κ , given by

$$\gamma_\kappa = \frac{1}{\sqrt{1 - \kappa^2}}, \quad (1.5)$$

$$e_\kappa = \left(\frac{1 + \kappa}{1 - \kappa} \right)^{1/2\kappa}, \quad (1.6)$$

linked through $e_\kappa = \exp_\kappa(\gamma_\kappa)$, represent a Lorentz factor and the relativistic generalization of the Napier number e , respectively.

The most interesting feature of the statistical distribution (1.4) is undoubtedly given by its asymptotic behavior. The low-energy limit of that distribution is equivalent to the $\kappa \rightarrow 0$ limit and yields the ordinary Boltzmann-Gibbs exponential distribution

$$f_i \approx \exp(-\beta E_i + \beta \mu - 1). \quad (1.7)$$

On the other hand, the distribution (1.4), in the high-energy limit shows a power-law tail

$$f_i \approx \frac{1}{e_\kappa} (2\kappa \gamma_\kappa \beta)^{-1/\kappa} E_i^{-1/\kappa}, \quad (1.8)$$

which has been observed experimentally in many physical, natural, and artificial systems [4].

Generalized statistical mechanics, based on κ -entropy, preserves the main features of ordinary Boltzmann-Gibbs statistical mechanics and is related to Boltzmann relativistic nonlinear kinetics [5–7]. Over the last 15 years this statistical theory has attracted the interest of many researchers, who have studied its foundation and mathematical aspects [8–15], the underlying thermodynamics [16–20], and, at the same time, specific applications of the theory to various fields. A nonexhaustive list of applications includes, among others, those in plasma physics [21–23], in astrophysics [23–29], in quantum statistics [30–32], in quantum entanglement [33,34], in genomics [35], in complex networks [36,37], in economy [38–41], and in finance [42–44].

The S_κ entropy, like Boltzmann-Shannon entropy, is thermodynamically and Lesche stable. On the other hand, while Boltzmann-Shannon entropy is additive for statistically independent systems, κ -entropy is superadditive [17], but its exact composition law has still not been completely established. The present paper deals with this intriguing and still open question regarding the composition law of S_κ , and the our main goal has just been its derivation.

The paper is organized as follows: First, in Sec. II, it is shown that $\ln_\kappa(xy)$ can be decomposed into the functions $\ln_\kappa(x)$ and $\ln_\kappa(y)$. In Sec. III, by employing this result, the

*giorgio.kaniadakis@polito.it

represents one of the main achievements of the present work.

IV. THE κ -PARENTROPY

In order to obtain a better understanding of the origin and meaning of the function $\ln_\kappa(x/e_\kappa)$ that appears in the composition law (2.4), the concept of scaled κ -logarithm introduced in Ref. [4], is briefly recalled. The scaled κ -logarithm is a two-parameter deformed logarithm that can be constructed, starting from the κ -logarithm and the scaling parameter ζ , according to

$$\ln_{\kappa\zeta}(x) = \frac{\ln_\kappa(\zeta x) - \ln_\kappa(\zeta)}{\sqrt{1 + \kappa^2 \ln_\kappa^2(\zeta)}}. \quad (4.1)$$

It is worth mentioning that the scaled κ -logarithm $\ln_{\kappa\zeta}(x)$ represents a family of deformed logarithms that contains the $\ln_\kappa(x)$ recovered when $\zeta = 1$, as a special case. This family also contains the functions $\ln_q(x)$ and $\ln_{2-q}(x)$ of the nonextensive statistical mechanics, obtained in the $\zeta \rightarrow 0^+$ and $\zeta \rightarrow +\infty$ limits, respectively, after posing $\kappa = q - 1$ [4]. In this sense, $\ln_{\kappa\zeta}(x)$ can also be viewed as a function that interpolates between $\ln_\kappa(x)$ and $\ln_q(x)$.

Let us focus attention on a particular scaled κ -logarithm that is obtained from the general formula (4.1) after fixing $\zeta = 1/e_\kappa$. This scaled κ -logarithm is hereafter called the κ -paralogarithm and is indicated as $\ln_\kappa^*(x)$. It is easy to verify that

$$\ln_\kappa^*(x) = \frac{1}{\gamma_\kappa} \ln_\kappa(x/e_\kappa) + 1. \quad (4.2)$$

After taking into account Eq. (2.3), it is possible to express $\ln_\kappa^*(x)$ in terms of $\ln_\kappa(x)$ according to

$$\ln_\kappa^*(x) = \ln_\kappa(x) - \sqrt{1 + \kappa^2 \ln_\kappa^2(x)} + 1, \quad (4.3)$$

from which it follows that $\ln_\kappa^*(x) \leq \ln_\kappa(x)$.

The $\ln_\kappa^*(x)$ function, defined in \mathbf{R}^+ , increases monotonically, $d \ln_\kappa^*(x)/dx > 0$, and is convex, $d^2 \ln_\kappa^*(x)/dx^2 > 0$. Furthermore, the following Taylor expansion holds

$$\ln_\kappa^*(1+x) \underset{x \rightarrow 0}{\sim} x - (1 + \kappa^2) \frac{x^2}{2}, \quad (4.4)$$

while the power-law asymptotic behaviors of $\ln_\kappa^*(x)$ are given by

$$\ln_\kappa^*(x) \underset{x \rightarrow 0^+}{\sim} -\frac{1 + \kappa}{2\kappa} x^{-\kappa}, \quad (4.5)$$

$$\ln_\kappa^*(x) \underset{x \rightarrow +\infty}{\sim} \frac{1 - \kappa}{2\kappa} x^\kappa, \quad (4.6)$$

and finally it holds that

$$\int_0^1 \ln_\kappa^*(x) dx = -\frac{1 + \kappa^2}{1 - \kappa^2}. \quad (4.7)$$

After introducing the κ -paralogarithm $\ln_\kappa^*(x)$, the composition law of $\ln_\kappa(xy)$, given in Eq. (2.4), assumes the following alternative form:

$$\begin{aligned} \ln_\kappa(xy) = & \ln_\kappa(x)[1 + \ln_\kappa(y) - \ln_\kappa^*(y)] \\ & + \ln_\kappa(y)[1 + \ln_\kappa(x) - \ln_\kappa^*(x)]. \end{aligned} \quad (4.8)$$

From the latter composition law, and after taking into account that $\ln_\kappa^*(x) \leq \ln_\kappa(x)$, it follows the property $\ln_\kappa(xy) \leq \ln_\kappa(x) + \ln_\kappa(y)$, holding for $0 < x, y \leq 1$.

Let us introduce the κ -parentropy of a normalized set of probability distributions $f = \{f_i\}$ defined by

$$S_\kappa^*(f) = -\langle \ln_\kappa^*(f_i) \rangle = -\sum_i f_i \ln_\kappa^*(f_i). \quad (4.9)$$

After taking into account the definition of the κ -paralogarithm (4.2), it is easy to verify that $S_\kappa^*(f)$ is related to the functional $S_\kappa(f/e_\kappa)$ through the linear transformation

$$S_\kappa^*(f) = \frac{1}{\gamma_\kappa} S_\kappa(f/e_\kappa) - 1. \quad (4.10)$$

In order to explain the meaning of the parentropy concept $S_\kappa^*(f)$ more clearly, let us consider a statistical system described by the ordered set of normalized probability distributions $f = \{f_i = \delta_{i1}; 1 \leq i \leq N\}$. This system occupies the state labeled by the index $i = 1$, and the entropy of the system is consequently equal to zero. It is easy to verify that $S_\kappa(f) = S_\kappa^*(f) = 0$, therefore entropy $S_\kappa(f)$ and parentropy $S_\kappa^*(f)$ can be adopted to describe the system. Otherwise it results that $S_\kappa(f/e_\kappa) = \gamma_\kappa \neq 0$, and the functional $S_\kappa(f/e_\kappa)$ therefore cannot represent the entropy of the system.

Finally, the composition law (3.6), after taking into account the relationship (4.10), assumes the following alternative form:

$$\begin{aligned} S_\kappa(pq) = & S_\kappa(p)[1 + S_\kappa^*(q) - S_\kappa(q)] \\ & + S_\kappa(q)[1 + S_\kappa^*(p) - S_\kappa(p)], \end{aligned} \quad (4.11)$$

while the functional $\mathcal{I}_\kappa(p)$, defined in (3.8), assumes the form

$$\mathcal{I}_\kappa(p) = 1 + S_\kappa^*(p) - S_\kappa(p). \quad (4.12)$$

After recalling that $\ln_\kappa^*(x) \leq \ln_\kappa(x)$, and then $S_\kappa^*(p) \geq S_\kappa(p)$ and $\mathcal{I}_\kappa(p) \geq 1$, it follows from (4.11) that the κ -entropy is superadditive, i.e., $S_\kappa(pq) \geq S_\kappa(p) + S_\kappa(q)$. Therefore, the difference between the entropy of the system obtained from the composition of two statistically independent subsystems and the sum of the entropies of the two individual subsystems,

$$\Delta S_\kappa = S_\kappa(pq) - S_\kappa(p) - S_\kappa(q), \quad (4.13)$$

results in $\Delta S_\kappa \geq 0$ and is given by

$$\Delta S_\kappa = S_\kappa(p)S_\kappa^*(q) + S_\kappa(q)S_\kappa^*(p) - 2S_\kappa(p)S_\kappa(q). \quad (4.14)$$

V. DISCUSSION

The question regarding the physical meaning of the entropy excess ΔS_κ , between the entropy of the composite system and the sum of the entropies of the two individual statistically independent subsystems is an important, and at the same time very complex, issue which should be considered within the context of the broader problem, which is still under intensive discussion, pertaining the origin of the generalized entropies, regardless of their particular form. It can be noted that if the classical Maxwell-Boltzmann statistics, corresponding to the additive Boltzmann-Shannon entropy, is excluded, all the other statistics considered in the literature are associated with particular types of nonadditive generalized entropies.

A first remark that deserves attention regards the thermodynamic stability of system described by a generalized

entropy. In the case of a system interacting with a bath in thermodynamic equilibrium, the particle density describing the state of the system in stationary conditions, is obtained starting from the maximum entropy principle. Alternatively the particle density function can be obtained as the asymptotic solution for $t \rightarrow \infty$, of a nonlinear Fokker-Planck evolution equation, obeying the second law of thermodynamics. On the other hand, this is also the case of an isolated system whose kinetics is governed by a nonlinear Boltzmann equation obeying the H-theorem that accounts the thermodynamics second law. This theorem states that the generalized entropy of the system grows continuously as it evolves towards equilibrium, where the entropy becomes maximum and reaches a finite value. The fact that κ -entropy is superadditive, and the entropy of the composite system is greater than the sum of the entropies of the individual subsystems, indicates that the composite system is thermodynamically more stable. This applies to all systems described by generalized superadditive entropies and was first noted by Landsberg and Vedral [45].

As far as the origin of the nonadditivity of generalized entropies is concerned, it can be observed that the systems described by these entropies are nonlinear. It is known that Boltzmann-Shannon entropy is associated with the ordinary Fokker-Planck, or Boltzmann, equations, depending on whether the system is in interaction with a bath or is isolated. In both cases, these equations describe linear kinetics. However, the generalized entropies are instead associated with nonlinear kinetics, governed by particular nonlinear evolution equations. In the case of systems subjected to κ -entropy, the corresponding nonlinear Fokker-Planck, or nonlinear Boltzmann, equations are known in the literature [1,7].

In Ref. [18], the thermodynamic potentials, i.e., the free energy and the Massieu function, have been obtained within the κ -statistical mechanics, and it has been shown that κ -thermodynamics preserves the Legendre structure of ordinary thermodynamics. Therefore, the nonadditivity of the κ -entropy implies the nonadditivity of the system internal energy. More in general and independently of the form of the generalized entropy, the positive sign of ΔS indicates the existence of an attractive force between the two statistically independent subsystems, and this interaction clearly originates from the interactions between the constituent particles of any subsystem. On the other hand, the interaction between the two subsystems is only negligible in the case of short-range particle interaction. For this reason, the hypothesis has been advanced in the literature that the systems described by generalized entropies could be composed of particles with long-range interactions.

Furthermore, another important question deserves to be addressed, i.e., the connection between nonadditive systems and ordinary additive thermodynamics. In Ref. [46], the authors claimed that a large ensemble of replicas of a nonadditive system obeys the standard additive thermodynamics principles, and the formalism of ordinary thermodynamics can be naturally applied, and the properties of nonadditive systems can be inferred.

Let us now briefly discuss the origin of κ -entropy. First, it should be noted that in the case Fermi-Dirac or Bose-Einstein statistics the nonlinear terms in the expression of the generalized fermionic or bosonic entropy are introduced precisely by the Pauli exclusion or inclusion principle, which accounts for the particle quantum nature. The related Fokker-Planck kinetics of fermions or bosons are nonlinear [47]. This is also the case of the Boltzmann kinetics of fermions or bosons described by the nonlinear Uehling-Uhlenbeck nonlinear equations. Analogously, the intermediate between the fermionic and bosonic nature of some many-body quantum systems leads to anyonic statistics, e.g., Haldane-Wu particles, quons, etc., which are described by generalized entropies.

Finally, let us consider κ -statistics. It should also be noted that κ -entropy emerges within the Einstein's special relativity. The fundamental law of relativistic dynamics governing the time evolution of any particle of the macroscopic statistical system, i.e., the Newton relativistic equation, is a nonlinear equation. As a consequence, according to the standard principles of molecular dynamics, a many-body system composed of relativistic particles necessarily is governed by nonlinear kinetics and thermodynamics. The related nonlinear kinetic equation of this statistical system is known in the literature, and it leads directly to κ -entropy [1–3]. Furthermore, the emergence of superadditive macroscopic quantities such as κ -entropy should not be surprising if it is taken into account that superadditive physical quantities can also be found at a microscopic level. In fact, it is well known, in relativistic dynamics, that the proper mass of an N -particle system is superadditive.

It has long been long known that the relativistic particles that form cosmic rays violate Maxwell-Boltzmann statistics, or, equivalently, Boltzmann-Shannon entropy. On the other hand, given the large extension of the cosmic ray spectrum, i.e., 33 decades in particle flux and 13 decades in energy, the excellent agreement observed over so many decades between these experimental data and the theoretical predictions of κ -statistical mechanics is quite remarkable. This result represents important experimental evidence of the relativistic origin of S_κ [1].

-
- [1] G. Kaniadakis, Statistical mechanics in the context of special relativity, *Phys. Rev. E* **66**, 056125 (2002).
 [2] G. Kaniadakis, Statistical mechanics in the context of special relativity II, *Phys. Rev. E* **72**, 036108 (2005).
 [3] G. Kaniadakis, Power-law tailed statistical distributions and Lorentz transformations, *Phys. Lett. A* **375**, 356 (2011).

- [4] G. Kaniadakis, Maximum entropy principle and power-law tailed distributions, *Eur. Phys. J. B* **70**, 3 (2009).
 [5] G. Kaniadakis, P. Quarati, and A. M. Scarfone, Kinetic foundations of non-conventional statistics, *Physica A (Amsterdam)* **305**, 76 (2002).
 [6] T. S. Biro and G. Kaniadakis, Two generalizations of the Boltzmann equation, *Eur. Phys. J. B* **50**, 3 (2006).

- [7] G. Kaniadakis, Relativistic kinetics and power-law-tailed distributions, *Europhys. Lett.* **92**, 35002 (2010).
- [8] R. Silva, The relativistic statistical theory and Kaniadakis entropy: An approach through a molecular chaos hypothesis, *Eur. Phys. J. B* **54**, 499 (2006).
- [9] J. Naudts, Deformed exponentials and logarithms in generalized thermostatics, *Physica A (Amsterdam)* **316**, 323 (2002).
- [10] F. Topsoe, Entropy and equilibrium via games of complexity, *Physica A (Amsterdam)* **340**, 11 (2004).
- [11] P. Tempesta, Group entropies, correlation laws, and zeta functions, *Phys. Rev. E* **84**, 021121 (2011).
- [12] A. M. Scarfone, Entropic forms and related algebras, *Entropy* **15**, 624 (2013).
- [13] N. T. C. M. Souza, D. H. A. L. Anselmo, R. Silva, M. S. Vasconcelos, and V. D. Mello, Analysis of fractal groups of the type d -(m , r)-Cantor within the framework of Kaniadakis statistics, *Phys. Lett. A* **378**, 1691 (2014).
- [14] A. M. Scarfone, On the κ -deformed cyclic functions and the generalized Fourier series in the framework of the κ -algebra, *Entropy* **17**, 2812 (2015).
- [15] A. M. Scarfone, κ -deformed Fourier transform, *Physica A (Amsterdam)* **480**, 63 (2017).
- [16] T. Wada, Thermodynamic stabilities of the generalized Boltzmann entropies, *Physica A (Amsterdam)* **340**, 126 (2004).
- [17] A. M. Scarfone and T. Wada, Canonical partition function for anomalous systems described by the κ -entropy, *Prog. Theor. Phys. Suppl.* **162**, 45 (2006).
- [18] A. M. Scarfone and T. Wada, Legendre structure of κ -thermostatistics revisited in the framework of information geometry, *J. Phys. A* **47**, 275002 (2014).
- [19] E. P. Bento, G. M. Viswanathan, M. G. E. da Luz, and R. Silva, Third law of thermodynamics as a key test of generalized entropies, *Phys. Rev. E* **91**, 022105 (2015).
- [20] T. Wada, H. Matsuzoe, and A. M. Scarfone, Dualistic Hessian structures among the thermodynamic potentials in the κ -thermostatistics, *Entropy* **17**, 7213 (2015).
- [21] I. Lourek and M. Tribeche, On the role of the κ -deformed Kaniadakis distribution in nonlinear plasma waves, *Physica A (Amsterdam)* **441**, 215 (2016).
- [22] L. A. Gougam and M. Tribeche, Electron-acoustic waves in a plasma with a κ -deformed Kaniadakis electron distribution, *Phys. Plasmas* **23**, 014501 (2016).
- [23] H. Chen, S. X. Zhang, and S. Q. Liu, The longitudinal plasmas modes of κ -deformed Kaniadakis distributed plasmas, *Phys. Plasmas* **24**, 022125 (2017).
- [24] J. C. Carvalho, R. Silva, J. D. do Nascimento, Jr., and J. R. De Medeiros, Power law statistics and stellar rotational velocities in the Pleiades, *Europhys. Lett.* **84**, 59001 (2008).
- [25] J. C. Carvalho, J. D. do Nascimento, Jr., R. Silva, and J. R. De Medeiros, Non-Gaussian statistics and stellar rotational velocities of main sequence field stars, *Astrophys. J. Lett.* **696**, L48 (2009).
- [26] J. C. Carvalho, R. Silva, J. D. do Nascimento, Jr., B. B. Soares, and J. R. De Medeiros, Observational measurement of open stellar clusters: A test of Kaniadakis and Tsallis statistics, *Europhys. Lett.* **91**, 69002 (2010).
- [27] M. Cure, D. F. Rial, A. Christen, and J. Casetti, A method to deconvolve stellar rotational velocities, *Astron. Astrophys.* **565**, A85 (2014).
- [28] E. M. C. Abreu, J. A. Neto, E. M. Barboza, and R. C. Nunes, Jeans instability criterion from the viewpoint of Kaniadakis' statistics, *Europhys. Lett.* **114**, 55001 (2016).
- [29] E. M. C. Abreu, J. A. Neto, E. M. Barboza, and R. C. Nunes, Tsallis and Kaniadakis statistics from the viewpoint of entropic gravity formalism, *Int. J. Mod. Phys. A* **32**, 1750028 (2017).
- [30] A. P. Santos, R. Silva, J. S. Alcaniz, and D. H. A. L. Anselmo, Kaniadakis statistics and the quantum H-theorem, *Phys. Lett. A* **375**, 352 (2011).
- [31] K. Ourabah and M. Tribeche, Planck radiation law and Einstein coefficients reexamined in Kaniadakis κ statistics, *Phys. Rev. E* **89**, 062130 (2014).
- [32] I. Lourek and M. Tribeche, Thermodynamic properties of the blackbody radiation: A Kaniadakis approach, *Phys. Lett. A* **381**, 452 (2017).
- [33] K. Ourabah, A. H. Hamici-Bendimerad, and M. Tribeche, Quantum Kaniadakis entropy under projective measurement, *Phys. Rev. E* **92**, 032114 (2015).
- [34] K. Ourabah, A. H. Hamici-Bendimerad, and M. Tribeche, Quantum entanglement and Kaniadakis entropy, *Phys. Scr.* **90**, 045101 (2015).
- [35] N. T. C. M. Souza, D. H. A. L. Anselmo, R. Silva, M. S. Vasconcelos, and V. D. Mello, A κ -statistical analysis of the Y-chromosome, *Europhys. Lett.* **108**, 38004 (2014).
- [36] A. Macedo-Filho, D. A. Moreira, R. Silva, and L. R. da Silva, Maximum entropy principle for Kaniadakis statistics and networks, *Phys. Lett. A* **377**, 842 (2013).
- [37] M. Stella and M. Brede, A κ -deformed model of growing complex networks with fitness, *Physica A (Amsterdam)* **407**, 360 (2014).
- [38] F. Clementi, M. Gallegati, and G. Kaniadakis, κ -generalized statistics in personal income distribution, *Eur. Phys. J. B* **57**, 187 (2007).
- [39] F. Clementi, M. Gallegati, and G. Kaniadakis, A κ -generalized statistical mechanics approach to income analysis, *J. Stat. Mech.* (2009) P02037.
- [40] M. L. Bertotti and G. Modenese, Exploiting the flexibility of a family of models for taxation and redistribution, *Eur. Phys. J. B* **85**, 261 (2012).
- [41] G. Modanese, Common origin of power-law tails in income distributions and relativistic gases, *Phys. Lett. A* **380**, 29 (2016).
- [42] B. Trivellato, The minimal κ -entropy martingale measure, *Int. J. Theor. Appl.* **15**, 1250038 (2012).
- [43] B. Trivellato, Deformed exponentials and applications to finance, *Entropy* **15**, 3471 (2013).
- [44] O. J. Tapiero, A maximum (non-extensive) entropy approach to equity options bid-ask spread, *Physica A (Amsterdam)* **392**, 3051 (2013).
- [45] P. T. Landsberg and V. Vedral, Distributions and channel capacities in generalized statistical mechanics, *Phys. Lett. A* **247**, 211 (1998).
- [46] I. Latella, A. Perez-Madrid, A. Campa, L. Casetti, and S. Ruffo, Thermodynamics of Nonadditive Systems, *Phys. Rev. Lett.* **114**, 230601 (2015).
- [47] G. Kaniadakis and P. Quarati, Classical model of bosons and fermions, *Phys. Rev. E* **49**, 5103 (1994).