

The Group Of The Fermat Numbers

*Original*

The Group Of The Fermat Numbers / Sparavigna, Amelia Carolina. - ELETTRONICO. - (2018).  
[10.5281/zenodo.1252422]

*Availability:*

This version is available at: 11583/2708600 since: 2018-05-25T08:04:01Z

*Publisher:*

Zenodo

*Published*

DOI:10.5281/zenodo.1252422

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

## The group of the Fermat Numbers

Amelia Carolina Sparavigna

Department of Applied Science and Technology, Politecnico di Torino, Torino, Italy

**Abstract:** In this work we are discussing the group that we can obtain if we consider the Fermat numbers with a generalized sum.

**Keywords:** generalized sum, groups, Abelian groups, transcendental functions, logarithmic and exponential functions, Fermat numbers.

**Turin, 24 May 2018. DOI: 10.5281/zenodo.1252422**

In [1] we find that there are two definitions of the Fermat numbers. We have a less common definition giving a Fermat number as  $F_n = 2^n + 1$ , which is obtained by setting  $x=1$  in a Fermat polynomial of  $x$ , and the commonly encounter definition  $F_n = 2^{2^n} + 1$ , which is a subset of the previous assembly of numbers. Here we will consider numbers  $F_n = 2^n + 1$  and - as we have recently proposed in [2] for  $q$ -integers and Mersenne numbers - investigate the set of them to find its generalized sum which defines the operation of the group.

Let us remember that a group is a set  $A$  having an operation  $\bullet$  which is combining the elements of  $A$ . That is, the operation combines any two elements  $a, b$  to form another element of the group denoted  $a \bullet b$ . To qualify  $(A, \bullet)$  as a group, the set and operation must satisfy the following requirements. *Closure:* For all  $a, b$  in  $A$ , the result of the operation  $a \bullet b$  is also in  $A$ . *Associativity:* For all  $a, b$  and  $c$  in  $A$ , it holds  $(a \bullet b) \bullet c = a \bullet (b \bullet c)$ . *Identity element:* An element  $e$  exists in  $A$ , such that for all elements  $a$  in  $A$ , it is  $e \bullet a = a \bullet e = a$ . *Inverse element:* For each  $a$  in  $A$ , there exists an element  $b$  in  $A$  such that  $a \bullet b = b \bullet a = e$ , where  $e$  is the identity (the notation is inherited from the multiplicative operation). A further requirement, is the *commutativity:* For all  $a, b$  in  $A$ ,  $a \bullet b = b \bullet a$ . In this case, the group is an Abelian group. For an Abelian group, one may choose to denote the *operation* by  $+$ , the *identity element* becomes the *neutral element* and the inverse element the *opposite element*. In this case, the group is called an additive group.

The generalized sum for the Fermat numbers  $F_n = 2^n + 1$  is:

$$F_m \oplus F_n = 2 - F_m - F_n + F_m F_n = (1 - F_m) + (1 - F_n) + F_m F_n \quad (1)$$

To have (1), let us evaluate:

$$\begin{aligned} F_{m+n} = 2^{m+n} + 1 &= F_m \oplus F_n = 2 - F_m - F_n + F_m F_n = 2 - (2^m + 1) - (2^n + 1) + (2^m + 1)(2^n + 1) \\ 2^{m+n} + 1 &= 2 - 2^m - 2^n - 2 + 2^m 2^n + 2^n + 2^m + 1 \end{aligned}$$

This gives also the *closure* of the group.

We can provide a recurrence relation as:  $F_{n+1} = 2^{n+1} + 1 = F_n \oplus F_1$

From (1), we can see that the *neutral element* is not 0. We have to use as a *neutral element* the integer 2, which is  $F_0 = 2^0 + 1 = 2$  and then an element of the group. We have:

$$F_n \oplus F_0 = 2 - F_n - F_0 + F_n F_0 = F_n$$

The *opposite element* is defined by  $F_n \oplus \text{Opposite}(F_n) = 2$ . We have:

$$\text{Opposite}(F_n) = \frac{F_n}{F_n - 1} = 1 + 2^{-n} = F_{-n} \quad (2)$$

Then, to have a group we need to add numbers (2) to the set of the Fermat numbers.

Therefore, we consider 2 as the *neutral element*, and the *opposite element* as given by (2).

Let us consider three Fermat numbers  $F_n, F_m, F_l$ ; to have a group we need the *associativity* of the generalized sum, so that  $(F_m \oplus F_n) \oplus F_l = F_m \oplus (F_n \oplus F_l)$ . Let us call  $x = F_n, y = F_m, z = F_l$  and evaluate:

$$\begin{aligned} (x \oplus y) \oplus z &= 2 - (x \oplus y) - z + (x \oplus y)z = 2 - 2 + x + y - xy - z + 2z - xz - yz + xyz \\ (x \oplus y) \oplus z &= x + y + z - xy - xz - yz + xyz \quad (3) \end{aligned}$$

And:

$$\begin{aligned} x \oplus (y \oplus z) &= 2 - x - (y \oplus z) + x(y \oplus z) = 2 - x - (2 - y - z + yz) + x(2 - y - z + yz) \\ x \oplus (y \oplus z) &= x + y + z - xy - xz - yz + xyz \quad (4) \end{aligned}$$

From (3) and (4), we have the *associativity*. The *commutativity* is evident.

We have already considered the generalized sum (1) in a recent work [3].

In [3], we consider some functions  $G(x)$ , having inverses so that  $G^{-1}(G(x))=x$ , which are *generators of group law* [4-6]:

$$\Phi(x, y) = G(G^{-1}(x) + G^{-1}(y))$$

The *group law* is giving the *generalized sum* of the group  $x \oplus y = G(G^{-1}(x) + G^{-1}(y))$ .

In [3] we considered the following generator and inverse:

$$G(x) = e^{-2x}(e^{2x} + 1) \quad G^{-1}(x) = \ln\left(\frac{1}{\sqrt{x-1}}\right) \quad (5)$$

and investigate a possible group from them. The *group law*  $\Phi(x, y)$  gives the generalized sum:

$$x \oplus y = G(G^{-1}(x) + G^{-1}(y)) = G\left(\ln\left(\frac{1}{\sqrt{x-1}}\right) + \ln\left(\frac{1}{\sqrt{y-1}}\right)\right) = G\left(\ln\left(\frac{1}{\sqrt{x-1}\sqrt{y-1}}\right)\right) = G\left(\ln\frac{1}{Z}\right)$$

$$G\left(\ln\frac{1}{Z}\right) = e^{-2\ln Z}(e^{2\ln Z} + 1) = (x-1)(y-1)\left(\frac{1}{(x-1)(y-1)} + 1\right)$$

$$x \oplus y = 2 - x - y + xy = (1-x) + (1-y) + xy \quad (6)$$

And (6) is the generalized sum (1) proposed for the Fermat numbers.

Let us also note that, if we use (5), we need  $x > 1$ . And this is a condition satisfied by the Fermat numbers and their opposites (2).

## References

1. Weisstein, Eric W. "Fermat Number." From MathWorld--A Wolfram Web Resource. <http://mathworld.wolfram.com/FermatNumber.html>
2. Sparavigna, A. C. (2018). The q-integers and the Mersenne numbers. Zenodo. DOI: 10.5281/zenodo.1251833
3. Sparavigna, A. C. (2018). Generalized Sums Based on Transcendental Functions. Zenodo. DOI: 10.5281/zenodo.1250020
4. Sicuro, G., & Tempesta, P. (2016). Groups, information theory, and Einstein's likelihood principle. Phys. Rev. E 93, 040101(R).
5. Curado, E. M., Tempesta, P., & Tsallis, C. (2016). A new entropy based on a group-theoretical structure. Annals of Physics, 366, 22-31.