

Some combinatorial properties of the Hurwitz series ring

Original

Some combinatorial properties of the Hurwitz series ring / Barbero, Stefano; Cerruti, Umberto; Murru, Nadir. - In: RICERCHE DI MATEMATICA. - ISSN 1827-3491. - 67:(2018), pp. 491-507.

Availability:

This version is available at: 11583/2719236 since: 2018-12-03T11:58:37Z

Publisher:

Springer

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Springer postprint/Author's Accepted Manuscript

(Article begins on next page)

Some combinatorial properties of the Hurwitz series ring

Stefano Barbero, Umberto Cerruti, Nadir Murru

Abstract We study some properties and perspectives of the Hurwitz series ring $H_R[[t]]$, for a commutative ring with identity R . Specifically, we provide a closed form for the invertible elements by means of the complete ordinary Bell polynomials, we highlight some connections with well-known transforms of sequences, and we see that the Stirling transforms are automorphisms of $H_R[[t]]$. Moreover, we focus the attention on some special subgroups studying their properties. Finally, we introduce a new transform of sequences that allows to see one of this subgroup as an ultrametric dynamic space.

Keywords Hurwitz series ring · binomial convolution · Bell polynomials · transforms of sequences

Mathematics Subject Classification (2000) 11B75 · 11B73

1 The ring of Hurwitz series, transformations of sequences and automorphisms

Given a commutative ring with identity R , let $H_R[[t]]$ denote the Hurwitz series ring whose elements are the formal series of the kind

$$A(t) := \sum_{n=0}^{+\infty} \frac{a_n}{n!} t^n,$$

equipped with the standard sum and the binomial convolution as product. Given two formal series $A(t)$ and $B(t)$, the binomial convolution is defined as follows:

$$A(t) \star B(t) := C(t),$$

where

$$c_n := \sum_{h=0}^n \binom{n}{h} a_h b_{n-h}.$$

The ring of Hurwitz series has been organizationally studied by Keigher [9] and in the recent years it has been extensively studied, see, e.g., [10], [11], [7], [3], [4], [5].

The Hurwitz series ring is trivially isomorphic to the ring H_R whose elements are infinite sequences of elements of R , with operations $+$ and \star . In the following, when we consider an element $a \in H_R$, we refer to a sequence $(a_n)_{n=0}^{+\infty} = (a_0, a_1, a_2, \dots)$, $a_i \in R$ for all $i \geq 0$, having exponential generating function (e.g.f.) denoted by $A(t)$. Clearly for two sequences $a, b \in H_R$ with exponential generating functions $A(t)$ and $B(t)$ respectively, the sequence $c = a \star b$ has e.g.f. $C(t) = A(t)B(t)$. Moreover, fixed any positive integer n , we can also consider the rings $H_R^{(n)}$ whose elements are sequences of elements of R with length n .

Remark 1 The binomial convolution is a commutative product and the identity in H_R is the sequence

$$(1, 0, 0, \dots).$$

Moreover, H_R can be also viewed as an R -algebra considering the map

$$\pi : R \rightarrow H_R, \quad \pi(r) := (r, 0, 0, \dots),$$

for any $r \in R$.

Proposition 1 *An element $a \in H_R$ is invertible if and only if $a_0 \in R$ is invertible, i.e.,*

$$H_R^* = \{a \in H_R : a_0 \in R^*\}.$$

Proof The proof is straightforward.

Given $a \in H_R^*$, we can recursively evaluate the terms of $b = a^{-1}$. Indeed, $b_0 = a_0^{-1}$ and for all $n \geq 1$

$$b_n = -a_0^{-1} \sum_{h=1}^n \binom{n}{h} a_h b_{n-h},$$

since the equality $a \star b = (1, 0, 0, \dots)$ implies

$$a_0 b_0 = 1, \quad \sum_{h=0}^n \binom{n}{h} a_h b_{n-h} = 0, \quad \forall n \geq 1.$$

On the other hand, we can find a nice closed form for the elements of b by means of the complete ordinary Bell polynomials [2]. First of all, we recall their definition as given in [13].

Definition 1 Let us consider the sequence $x = (x_1, x_2, \dots)$. The *complete ordinary Bell polynomials* are defined by

$$B_0(x) = 1, \quad \forall n \geq 1 \quad B_n(x) = B_n(x_1, x_2, \dots, x_n) = \sum_{k=1}^n B_{n,k}(x),$$

where $B_{n,k}(x)$ are the *partial ordinary Bell polynomials*, with

$$B_{0,0}(x) = 1, \quad \forall n \geq 1 \quad B_{n,0}(x) = 0, \quad \forall k \geq 1 \quad B_{0,k}(x) = 0,$$

$$B_{n,k}(x) = B_{n,k}(x_1, x_2, \dots, x_n) = \sum_{\substack{i_1+2i_2+\dots+n i_n=n \\ i_1+i_2+\dots+i_n=k}} \frac{k!}{i_1! i_2! \dots i_n!} x_1^{i_1} x_2^{i_2} \dots x_n^{i_n},$$

or, equivalently,

$$B_{n,k}(x) = B_{n,k}(x_1, x_2, \dots, x_{n-k+1}) = \sum_{\substack{i_1+2i_2+\dots+(n-k+1)i_{n-k+1}=n-k+1 \\ i_1+i_2+\dots+i_{n-k+1}=k}} \frac{k!}{i_1! i_2! \dots i_{n-k+1}!} x_1^{i_1} x_2^{i_2} \dots x_{n-k+1}^{i_{n-k+1}},$$

satisfying the equality

$$\left(\sum_{n \geq 1} x_n z^n \right)^k = \sum_{n \geq k} B_{n,k}(x) z^n.$$

Then, we introduce the Invert transform (see, e.g., [6] for a detailed survey).

The Invert transform

The Invert transform \mathcal{I} maps a sequence $a = (a_n)_{n=0}^{+\infty}$ into a sequence $\mathcal{I}(a) = b = (b_n)_{n=0}^{+\infty}$ whose ordinary generating function satisfies

$$\sum_{n=0}^{+\infty} b_n t^n = \frac{\sum_{n=0}^{+\infty} a_n t^n}{1 - t \sum_{n=0}^{+\infty} a_n t^n}.$$

Barbero et al. [1] highlighted the close relation between Invert transform and complete ordinary Bell polynomials: given $g \in H_R$ and $h = \mathcal{I}(g)$, we have, for all $n \geq 0$, that

$$h_n = B_{n+1}(g_0, g_1, g_2, \dots, g_n). \quad (1)$$

Now these tools allow us to explicitly find the terms of $b = a^{-1}$ for every $a \in H_R^*$.

Theorem 1 Let $a, b = a^{-1} \in H_R^*$ be sequences with e.g.f. $A(t)$ and $B(t)$, respectively. Then for all $n \geq 0$, we have

$$b_n = \frac{n!B_n(g_0, g_1, g_2, \dots, g_n)}{a_0},$$

where

$$g = (g_n)_{n=0}^{+\infty} = \left(-\frac{a_{n+1}}{a_0(n+1)!} \right)_{n=0}^{+\infty}.$$

Proof The ordinary generating function of the sequence g is

$$\bar{G}(t) = \frac{1}{t} \left(1 - \frac{A(t)}{a_0} \right)$$

since

$$\bar{G}(t) = \frac{1}{t} \left(1 - \frac{1}{a_0} \left(a_0 + \sum_{n=1}^{+\infty} \frac{a_n}{n!} t^n \right) \right) = -\frac{1}{t} \sum_{n=1}^{+\infty} \frac{a_n}{a_0 n!} t^n = \sum_{n=0}^{+\infty} \left(-\frac{a_{n+1}}{a_0(n+1)!} \right) t^n.$$

Moreover, considering $b = a^{-1}$ we have

$$B(t) = \frac{1}{A(t)} = \frac{1}{a_0(1 - t\bar{G}(t))} = \frac{1}{a_0} \left(1 + \frac{t\bar{G}(t)}{1 - t\bar{G}(t)} \right) = \frac{1}{a_0} (1 + t\bar{H}(t)),$$

where $\bar{H}(t)$ is the ordinary generating function of the sequence $h = \mathcal{I}(g)$. Thus relation (1) holds and, since $B_0(g) = 1$, we obtain

$$B(t) = \frac{1}{a_0} (1 + t\bar{H}(t)) = \frac{1}{a_0} \left(1 + \sum_{n=0}^{+\infty} B_{n+1}(g_0, g_1, g_2, \dots, g_n) t^{n+1} \right) = \sum_{n=0}^{+\infty} \frac{B_n(g_0, g_1, g_2, \dots, g_n)}{a_0} t^n$$

and the thesis easily follows.

We point out that some well-studied transforms acting on sequences can be viewed in $H_R[[t]]$ as the product (i.e., the binomial convolution) between a suitable fixed sequence and any sequence belonging to H_R . We present two enlightening and interesting examples.

The Binomial interpolated transform

The *Binomial interpolated transform* $\mathcal{L}^{(y)}$, with parameter $y \in R$, maps any sequence $a \in H_R$ into a sequence $b = \mathcal{L}^{(y)}(a) \in H_R$, whose terms are

$$b_n = \sum_{h=0}^n \binom{n}{h} y^{n-h} a_h.$$

For a survey and a detailed study of the action of $\mathcal{L}^{(y)}$ on recurrence sequences we refer the reader to [1]. The definition of this transform by means of the binomial convolution is straightforward. Indeed, considering the sequence

$$\lambda = (y^n)_{n=0}^{+\infty},$$

we have for any $a \in H_R$ with e.g.f. $A(t)$

$$\mathcal{L}^{(y)}(a) = \lambda \star a.$$

with the corresponding e.g.f. given by the product $e^{yt}A(t)$.

The Boustrophedon transform

The *Boustrophedon transform* \mathcal{B} , introduced and studied in [12], maps any sequence $a \in H_R$, with e.g.f. $A(t)$, into a sequence $b = \mathcal{B}(a) \in H_R$ with e.g.f.

$$B(t) = (\sec t + \tan t)A(t).$$

This transform is closely related to the sequence $\beta = (\beta_n)_{n=0}^{+\infty}$ of the Euler zigzag numbers (see [12]), with e.g.f.

$$\sum_{n=0}^{+\infty} \frac{\beta_n}{n!} t^n := \sec(t) + \tan(t),$$

since for any $a \in H_R$ clearly

$$\mathcal{B}(a) = \beta \star a.$$

The Hurwitz series ring is strictly connected to other well-known sequence transforms. We consider another two examples, the alternating sign transform, which is a little bit trivial, and the interesting Stirling transform. We also show that both are examples of H_R -automorphisms.

The alternating sign transform

The alternating sign transform \mathcal{E} maps any sequence $a \in H_R$ into a sequence $b = \mathcal{E}(a) \in H_R$, whose terms are

$$b_n = (-1)^n a_n.$$

The transform \mathcal{E} often appears in studying properties of integer sequences combined with other transforms. Clearly we have $\mathcal{E} = \mathcal{E}^{-1}$ and it is straightforward to see that, given any $a \in H_R$ with e.g.f. $A(t)$, then $\mathcal{E}(a)$ has e.g.f. $A(-t)$. Moreover it is easy to verify that for all sequences $a, b \in H_R$

$$\mathcal{E}(a + b) = \mathcal{E}(a) + \mathcal{E}(b),$$

and

$$\mathcal{E}(a \star b) = \mathcal{E}(a) \star \mathcal{E}(b),$$

showing that \mathcal{E} is an automorphism of H_R .

The Stirling transform

The *Stirling transform* \mathcal{S} maps any sequence $a \in H_R$ into a sequence $b = \mathcal{S}(a) \in H_R$, whose terms are

$$b_n = \sum_{h=0}^n \left\{ \begin{matrix} n \\ h \end{matrix} \right\} a_h,$$

where $\left\{ \begin{matrix} n \\ h \end{matrix} \right\}$ are the Stirling numbers of the second kind (see e.g. [8], chapter 6, for definition and properties of Stirling numbers of first and second kind). Some properties of this transform are exposed in [6], here we observe that \mathcal{S} is a bijection from H_R to itself. The inverse \mathcal{S}^{-1} maps any sequence $a \in H_R$ into a sequence $b = \mathcal{S}^{-1}(a) \in H_R$, whose terms are

$$b_n = \sum_{h=0}^n (-1)^{n-h} \left[\begin{matrix} n \\ h \end{matrix} \right] a_h,$$

where $\left[\begin{matrix} n \\ h \end{matrix} \right]$ are the (unsigned) Stirling numbers of the first kind. Moreover, we recall that for all $a \in H_R$ with e.g.f. $A(t)$, then $b = \mathcal{S}(a)$ has e.g.f. $B(t) = A(e^t - 1)$. It is very interesting to observe that, for all $a, b \in H_R$, \mathcal{S} obviously satisfies

$$\mathcal{S}(a + b) = \mathcal{S}(a) + \mathcal{S}(b),$$

but also

$$\mathcal{S}(a \star b) = \mathcal{S}(a) \star \mathcal{S}(b).$$

Indeed, remembering that, by definition, $\left\{ \begin{matrix} n \\ k \end{matrix} \right\} = 0$ when $n < k$, and that the e.g.f. of the Stirling number of the second kind is $\frac{(e^t - 1)^n}{n!}$ (see [8]), if we consider the e.g.f. $S(t)$ of $\mathcal{S}(a \star b)$, we have

$$\begin{aligned} S(t) &= \sum_{n=0}^{+\infty} \left(\sum_{h=0}^n \left\{ \begin{matrix} n \\ h \end{matrix} \right\} \left(\sum_{j=0}^h \binom{h}{j} a_j b_{h-j} \right) \right) \frac{t^n}{n!} = \sum_{h=0}^{+\infty} \left(\sum_{j=0}^h \binom{h}{j} a_j b_{h-j} \right) \sum_{n=h}^{+\infty} \left\{ \begin{matrix} n \\ h \end{matrix} \right\} \frac{t^n}{n!} = \\ &= \sum_{h=0}^{+\infty} \left(\sum_{j=0}^h \binom{h}{j} a_j b_{h-j} \right) \frac{(e^t - 1)^h}{h!} = \mathcal{A}(e^t - 1) \mathcal{B}(e^t - 1) \end{aligned}$$

and $\mathcal{A}(e^t - 1) \mathcal{B}(e^t - 1)$ clearly is the e.g.f. of $\mathcal{S}(a) \star \mathcal{S}(b)$. Hence \mathcal{S} is an automorphism of H_R .

2 Special subgroups of H_R^*

The purpose of this section is to highlight some properties of two interesting subgroups of H_R^* , with respect to the binomial convolution product \star operation. We also study their relationship with the transforms presented in the previous section and with other transforms which we will define in the next.

Definition 2 Let us denote U_R and B_R the subgroups of H_R^* defined as

$$U_R = \{a \in H_R^* : a_0 = 1\}, \quad B_R = \{a \in U_R : \mathcal{E}(a) = a^{-1}\}.$$

We start considering the subgroup U_R and observing that, for all $a \in H_R$, we can find sequences in U_R closely related with a , obtained by prepending to a a finite sequence of 1. So it is natural to consider these sequences as the images of a under the iteration of the following transform.

The *1-prepending* transform \mathcal{V} maps a sequence $a = (a_0, a_1, a_2, \dots) \in H_R$ into the sequence $b = \mathcal{V}(a) = (1, a_0, a_1, a_2, \dots) \in U_R$. We denote \mathcal{V}^k as the k -times iteration of \mathcal{V} and obviously $\mathcal{V}^k(a) \in U_R$, for all $k \geq 1$. We observe that the action of \mathcal{V}^k on a sequence $a \in H_R^*$ corresponds to the k -th iteration of the integral operator

$$\mathcal{J}(\cdot)(t) := 1 + \int_0^t (\cdot) du$$

on the e.g.f. $A(t)$ of a . In particular,

$$\mathcal{J}^k(A)(t) = \underbrace{\mathcal{J} \circ \dots \circ \mathcal{J}}_{k\text{-times}}(A)(t) = \sum_{h=0}^{k-1} \frac{t^h}{h!} + \sum_{h=k}^{+\infty} a_{h-k} \frac{t^h}{h!} = V(t),$$

where $V(t)$ is the e.g.f. of $\mathcal{V}^k(a)$, and clearly

$$V(0) = V'(0) = \dots = V^{(k-1)}(0) = 1, \quad V^{(k)}(t) = A(t),$$

being $V^{(k)}(t)$ the k -th derivative of $V(t)$.

Now, we explore some interesting properties related to the subgroup B_R . First of all we characterize all the elements in B_R .

Theorem 2 *All the elements in B_R corresponds to the sequences of U_R whose e.g.f. $A(t)$ is the solution of*

$$\begin{cases} A'(t) = g(t)A(t) \\ A(0) = 1 \end{cases} \quad (2)$$

where $g(t)$ is any fixed even function. Hence, if we consider the formal exponential operator \exp such that $\exp(f(t)) = \sum_{n=0}^{+\infty} \frac{(f(t))^n}{n!}$, the e.g.f. of $a \in B_R$ is $A(t) = \exp(h(t))$ where $h(t)$ is an odd function.

Proof It is immediate to see that $a \in B_R$ if and only if $A(t)A(-t) = 1$. If we differentiate this relation with respect to t we obtain

$$A'(t)A(-t) - A(t)A'(-t) = 0$$

which is equivalent to

$$\frac{A'(t)}{A(t)} = \frac{A'(-t)}{A(-t)}.$$

Thus we have $g(t) = \frac{A'(t)}{A(t)}$ where, from the previous relation, $g(t) = g(-t)$, i.e. $g(t)$ is an even function, and, obviously, we must have $A(0) = 1$, since for all sequences $a \in U_R$ $A(0) = a_0 = 1$. It is straightforward to verify that, given $g(t)$, a formal integration term by term of its power series corresponds to the series of an odd function $h(t)$, and clearly $A(t) = \exp(h(t))$ satisfies (2).

The transforms $\mathcal{L}^y, \mathcal{B}$, and \mathcal{E} act on B_R preserving the closure, as we point out in the following proposition.

Proposition 2 *The group B_R is closed with respect to the transforms \mathcal{E}, \mathcal{B} and $\mathcal{L}^{(y)}$, for any $y \in R$.*

Proof By definition of B_R , it is immediate to check that $\mathcal{E}(B_R) = B_R$ (with this notation, we say that given any $a \in B_R$, then $\mathcal{E}(a)$ is still in B_R). Given any $a \in B_R$, with e.g.f. $A(t)$, we have that $b = \mathcal{L}^{(y)}(a)$ has e.g.f. $e^{yt}A(t)$, and clearly $b \in B_R$ since

$$(e^{yt}A(t)) (e^{-yt}A(-t)) = A(t)A(-t) = 1.$$

Finally, let us recall that the Euler zig-zag numbers β have e.g.f. $B(t) = \sec(t) + \tan(t)$ which satisfies $B(t)B(-t) = 1$ as a simple calculation shows

$$B(t) - \frac{1}{B(-t)} = \frac{1 + \sin(t)}{\cos(t)} - \frac{\cos(-t)}{1 + \sin(-t)} = \frac{1 - \sin^2(t) - \cos^2(t)}{\cos(t)(1 - \sin(t))} = 0,$$

i.e., $\beta \in B_R$. Hence, given any $a \in B_R$, $\mathcal{B}(a) = \beta \star a \in B_R$.

Remark 2 The group B_R is not closed with respect to the transform \mathcal{S} . Indeed, if $A(t)$ is the e.g.f. of $a \in B_R$, the e.g.f. of $S(a)$ is $A(e^t - 1)$, while the e.g.f. of $\varepsilon(S(a))$ is $A(e^{-t} - 1)$ and in general $A(e^{-t} - 1)A(e^t - 1) \neq 1$. It would be interesting to characterize the group $\mathcal{S}(B_R)$.

Let us examine in depth the structure of a sequence $a \in B_R$. From the definition of B_R and from Theorem 2, we observe that the elements of a sequence $a \in B_R$ are constrained to severe restrictions, since the equality $\mathcal{E}(a) = a^{-1}$ must hold. If we pose $b = a^{-1}$ and $c = \mathcal{E}(a)$, we have, for instance,

$$b_0 = 1, \quad b_1 = -a_1, \quad b_2 = 2a_1^2 - a_2$$

and

$$c_0 = 1, \quad c_1 = -a_1, \quad c_2 = a_2,$$

i.e., the element a_1 of the sequence a can be arbitrary, while a_2 must satisfy

$$a_2 = 2a_1^2 - a_2,$$

i.e., $a_2 = a_1^2$. By continuing in this way, we can also see, e.g., that a_3 can be arbitrary, while $a_4 = -3a_1^4 + 4a_1a_3$. Thus, any sequence $a \in B_R$ is completely determined when we fix the values of $a_{2k-1}, k = 1, 2, \dots$. Indeed, the following theorem shows how to evaluate the terms with even positive index as functions of the ones with odd index, by means of the partial ordinary Bell polynomials described in Definition 1.

Theorem 3 *Given any $a \in B_R$, we have*

$$a_{2n} = (2n)! \sum_{k=0}^n \binom{\frac{1}{2}}{k} B_{n+k, 2k}(x_1, \dots, x_{n-k+1}), \quad \forall n \geq 1,$$

where $x_i = \frac{a_{2i-1}}{(2i-1)!}$, $\binom{\frac{1}{2}}{k} = \frac{\prod_{j=0}^{k-1} (\frac{1}{2}-j)}{k!}$.

Proof Let $A(t)$ be the e.g.f. of a . Clearly $A(t) = P(t) + D(t)$, where

$$P(t) = \sum_{n=0}^{+\infty} \frac{a_{2n}}{(2n)!} t^{2n} \quad (3)$$

and

$$D(t) = \sum_{n=1}^{+\infty} \frac{a_{2n-1}}{(2n-1)!} t^{2n-1} = \frac{1}{t} \sum_{n=1}^{+\infty} \frac{a_{2n-1}}{(2n-1)!} t^{2n}. \quad (4)$$

Moreover, we have

$$1 = A(t)A(-t) = (P(t) + D(t))(P(t) - D(t)) = (P(t))^2 - (D(t))^2,$$

since $A(-t) = P(-t) + D(-t) = P(t) - D(t)$ and $\mathcal{E}(a) = a^{-1}$. Now, observing that $P(0) = a_0 = 1$ and $D(0) = 0$, we obtain from the formal Maclaurin power series of $(1 + X)^{\frac{1}{2}}$ that

$$P(t) = \left(1 + (D(t))^2\right)^{\frac{1}{2}} = \sum_{k=0}^{+\infty} \binom{\frac{1}{2}}{k} (D(t))^{2k}.$$

By definition of partial ordinary Bell polynomials we have

$$(D(t))^{2k} = \sum_{m=2k}^{+\infty} B_{m, 2k}(x_1, \dots, x_{m-2k+1}) t^{2m-2k}.$$

If we set $n = m - k$, we get

$$P(t) = \sum_{k=0}^{+\infty} \binom{\frac{1}{2}}{k} \sum_{n=k}^{+\infty} B_{n+k, 2k}(x_1, \dots, x_{n-k+1}) t^{2n} = \sum_{n=0}^{+\infty} \left(\sum_{k=0}^n \binom{\frac{1}{2}}{k} B_{n+k, 2k}(x_1, \dots, x_{n-k+1}) \right) t^{2n}.$$

From this equality, comparing the coefficients of the respective even powers of t in (3) we finally obtain

$$a_{2n} = (2n)! \sum_{k=0}^n \binom{\frac{1}{2}}{k} B_{n+k, 2k}(x_1, \dots, x_{n-k+1}), \quad \forall n \geq 1.$$

By Definition 1 and observing that $B_{0,0} = 1$, $B_{h,0} = 0$ for $h \geq 1$, we immediately have the following corollary.

Corollary 1 Given $a \in B_R$, we have

$$a_{2n} = (2n)! \sum_{k=1}^n \binom{\frac{1}{2}}{k} \sum_{\substack{i_1+i_2+\dots+i_{n-k+1}=2k \\ i_1+2i_2+\dots+(n-k+1)i_{n-k+1}=n+k}} (2k)! \prod_{j=1}^{n-k+1} \frac{1}{i_j! ((2j-1)!)^{i_j}} \prod_{j=1}^{n-k+1} a_{2j-1}^{i_j}.$$

On the other hand, it is also possible to determine the sequences $a \in B_R$, with $a_2 \in R^*$ and such that a_2 is a square in R , by fixing the terms a_{2k} , $k = 1, 2, \dots$ and finding the terms with odd index as functions of the ones with even index.

Theorem 4 Given $a \in B_R$ such that $a_2 \in R^*$ and $x^2 = a_2$ is solvable in R , we have

$$a_{2n+1} = (2n+1)! \sum_{k=0}^n a_2^{\frac{1}{2}-k} \binom{\frac{1}{2}}{k} B_{n,k}(x_1, \dots, x_{n-k+1}), \quad \forall n \geq 0,$$

where $x_i = \frac{1}{(2i+2)!} \sum_{k=0}^{i+1} \binom{2i+2}{k} a_{2k} a_{2(n-k+1)}$, and $a_2^{\frac{1}{2}} \in R$ is a solution of $x^2 = a_2$.

Proof Let $A(t)$ be the e.g.f. of a , with the same notations used in the proof of Theorem 3, we have $(D(t))^2 = 1 + (P(t))^2$, where $D(t)$ and $P(t)$ as in (4) and in (3) respectively. Since the product $P(t) \cdot P(t)$ is equal to

$$(P(t))^2 = 1 + a_2 t^2 + \sum_{n=2}^{+\infty} \left(\sum_{k=0}^n \binom{2n}{2k} a_{2k} a_{2n-2k} \right) \frac{t^{2n}}{(2n)!},$$

we find

$$\begin{aligned} D(t) &= t a_2^{\frac{1}{2}} \left(1 + \sum_{n=2}^{+\infty} \frac{a_2^{-1}}{(2n)!} \left(\sum_{k=0}^n \binom{2n}{2k} a_{2k} a_{2n-2k} \right) t^{2n-2} \right)^{\frac{1}{2}} = \\ &= t a_2^{\frac{1}{2}} \left(1 + \sum_{n=1}^{+\infty} \frac{a_2^{-1}}{(2n+2)!} \left(\sum_{k=0}^{n+1} \binom{2n+2}{2k} a_{2k} a_{2n-2k+2} \right) t^{2n} \right)^{\frac{1}{2}} \end{aligned}$$

Then, considering the formal Maclaurin series expansion of $(1+X)^{\frac{1}{2}}$ and by Definition 1, we obtain

$$D(t) = \sum_{n=0}^{+\infty} \left(\sum_{k=0}^n a_2^{1/2-k} \binom{\frac{1}{2}}{k} B_{n,k}(x_1, \dots, x_{n-k+1}) \right) t^{2n+1}.$$

Now the thesis easily follows by a simple comparison of the corresponding coefficients of the odd powers of t in the expansion (4) of $D(t)$.

Remark 3 When $R = \mathbb{Z}$, $B_{\mathbb{Z}}$ contains many well-known and important integer sequences. We mention here some of them as interesting examples. We have seen that the Euler zigzag numbers belong to $B_{\mathbb{Z}}$. They are listed in OEIS [15] as A000111. Thus, all the sequences having as e.g.f. a power of $\sec(t) + \tan(t)$ are in $B_{\mathbb{Z}}$.

For instance the sequence A001250 in OEIS, whose n -th element is the number of alternating permutations of order n , has e.g.f. $(\sec(t) + \tan(t))^2$.

Moreover, the sequence A000667, which is the Boustrophedon transform of all-1's sequence, has e.g.f. $e^t(\sec(t) + \tan(t))$ and belongs to $B_{\mathbb{Z}}$.

Another sequence in $B_{\mathbb{Z}}$ is A000831, with e.g.f. $\frac{1+\tan(t)}{1-\tan(t)}$.

The sequences A006229 and A002017 also belong $B_{\mathbb{Z}}$ since they have exponential generating functions of the shape $\exp(f(t))$, with $f(t)$ odd function. Indeed, they have e.g.f. $e^{\tan(t)}$ and $e^{\sin(t)}$, respectively.

Thanks to Theorem 3 and Corollary 1, we have new interesting identities connecting many sequences in OEIS. Furthermore, it is quite surprising that all these (very different) sequences satisfy the same limiting conditions.

In the following, we will introduce a new transform of sequences that arises from the study of B_R , which will allow us to consider U_R as a dynamic ultrametric space.

Given $a, b = a^{-1} \in U_R$, we know that

$$\sum_{h=0}^n \binom{n}{h} a_h b_{n-h} = 0, \quad \forall n \geq 1,$$

from which it follows that

$$a_n = - \sum_{h=0}^{n-1} \binom{n}{h} a_h b_{n-h}, \quad b_n = - \sum_{h=0}^{n-1} \binom{n}{h} b_h a_{n-h}.$$

If $a \in B_R$, i.e. $A(-t) = A(t)^{-1}$, then, for all $n \geq 1$, we have

$$- \sum_{h=0}^{n-1} \binom{n}{h} (-1)^h a_h a_{n-h} = \begin{cases} 0 & \text{if } n \text{ odd} \\ 2a_n & \text{if } n \text{ even} \end{cases}.$$

Thus it is natural to define the following transform.

Definition 3 The *autoconvolution* transform \mathcal{A} maps a sequence $a \in H_R$ into a sequence $b = \mathcal{A}(a) \in H_R$, where

$$\begin{cases} b_0 = a_0 \\ b_{2n+1} = a_{2n+1}, \quad \forall n \geq 0 \\ b_{2n} = -\frac{1}{2} \sum_{h=1}^{n-1} \binom{n}{h} (-1)^h a_h a_{n-h}, \quad \forall n \geq 1 \end{cases}.$$

The following proposition is a straightforward consequence.

Proposition 3 Given any $a \in H_R$, we have $a \in B_R \Leftrightarrow \mathcal{A}(a) = a$.

Finally, we introduce another transform strictly related to \mathcal{A} .

Definition 4 The transform \mathcal{U} maps a sequence $a \in H_R$ into a sequence $\mathcal{U}(a) = b \in H_R$ as follows:

$$\begin{cases} b_0 = a_0 \\ b_{2n+1} = a_{2n+1}, \quad \forall n \geq 0 \\ b_{2n} = (2n)! \sum_{k=0}^n \binom{n}{k} B_{n+k, 2k}(x_1, \dots, x_{n-k+1}), \quad \forall n \geq 1 \end{cases},$$

where $x_i = \frac{a_{2i-1}}{(2i-1)!}$.

Remark 4 Given any sequence $a \in U_R$, the transform \mathcal{U} produces a sequence in B_R where the terms with odd index are the corresponding terms of a . Clearly, we have that a sequence $a \in U_R$ is in B_R if and only if $a = \mathcal{U}(a)$.

Proposition 4 Given $a \in H_R$, with e.g.f. $A(t)$, then $\mathcal{U}(a)$ has e.g.f.

$$U(t) = \left(1 + \left(\frac{A(t) - A(-t)}{2} \right)^2 \right)^{\frac{1}{2}} + \frac{A(t) - A(-t)}{2}.$$

Proof We can write $A(t) = P(t) + D(t)$, where $P(t)$ and $D(t)$ as in (3) and in (4) respectively. We have $A(-t) = P(t) - D(t)$ and consequently $D(t) = \frac{A(t) - A(-t)}{2}$. By Theorem 3, the terms in the even places of $\mathcal{U}(a)$ have e.g.f $P(t) = (1 + (D(t))^2)^{\frac{1}{2}}$. Thus, we have

$$U(t) = (1 + (D(t))^2)^{\frac{1}{2}} + D(t).$$

Given $a, b \in H_R$, let us define

$$\delta(a, b) := 2^{-k},$$

if $a_i = b_i$, for any $0 \leq i \leq k-1$. It is well-known that δ is an ultrametric in H_R . Indeed,

- $\delta(a, b) = 0 \Leftrightarrow a = b$,
- $\delta(a, b) = \delta(b, a)$,
- $\delta(a, c) \leq \max(\delta(a, b), \delta(b, c))$,

for any $a, b, c \in H_R$. Thus, (H_R, δ) is an ultrametric space.

Let us recall that we denote $H_R^{(n)}$ the ring whose elements are sequences of elements of R with length n . Similarly, $U_R^{(n)}$ and $B_R^{(n)}$ are the subgroups of $H_R^{(n)*}$ corresponding to the subgroups U_R and B_R of H_R^* , respectively.

Theorem 5 *Given any $a \in U_R$, we have*

$$\delta(\mathcal{A}^n(a), \mathcal{U}(a)) \leq \frac{1}{2^{2(n+1)}},$$

where $\mathcal{A}^n = \underbrace{\mathcal{A} \circ \dots \circ \mathcal{A}}_{n\text{-times}}$.

Proof We prove the thesis by induction.

Let us denote $a' = \mathcal{U}(a)$ and $b = \mathcal{A}(a)$. It is straightforward to check that

$$a' = (1, a_1, a_1^2, a_3, \dots), \quad b = (1, a_1, a_1^2, a_3, \dots).$$

Thus, a' and b coincide at least in the first 4 terms, i.e.,

$$\delta(\mathcal{A}(a), \mathcal{U}(a)) \leq \frac{1}{2^4}.$$

Now, let us suppose that given $b = \mathcal{A}^n(a)$, we have $\delta(\mathcal{A}^n(a), \mathcal{U}(a)) \leq \frac{1}{2^{2(n+1)}}$, i.e. $b_i = a'_i$ for all $i \leq 2n+1$ and consider $c = \mathcal{A}(b)$. Since $a' \in B_R$, we remember that for all $n \geq 1$ we have

$$-\sum_{h=0}^{n-1} \binom{n}{h} (-1)^h a'_h a'_{n-h} = \begin{cases} 0 & \text{if } n \text{ odd} \\ 2a'_n & \text{if } n \text{ even} \end{cases}.$$

Thus, by Definition 3, we obtain $c_i = a'_i$ for all $i \leq 2n+3$, since $(b_0, \dots, b_{2n+1}) = (a'_0, \dots, a'_{2n+1}) \in B_R^{(2n+2)}$ by inductive hypothesis. Hence, we have proved that

$$\delta(\mathcal{A}^{n+1}(a), \mathcal{U}(a)) \leq \frac{1}{2^{2(n+2)}}.$$

As a consequence of Theorem 5, we can observe that \mathcal{A} can be considered as an approximation of \mathcal{U} . Indeed, given a sequence $a \in U_R$, sequences $\mathcal{A}^n(a)$ have more elements equal to elements of $\mathcal{U}(a)$ for increasing values of n .

Example 1 Given $a = (a_0, a_1, a_2, a_3, a_4, a_5) \in U_R^{(6)}$, then

$$\mathcal{U}(a) = (1, a_1, a_1^2, a_3, 4a_1a_3 - 3a_1^4, a_5)$$

and

$$\mathcal{A}(a) = (1, a_1, a_1^2, a_3, 4a_1a_3 - 3a_2^2, a_5).$$

Considering \mathcal{A}^2 , we obtain

$$\mathcal{A}^2(a) = (1, a_1, a_1^2, a_3, 4a_1a_3 - 3a_1^4, a_5) = \mathcal{U}(a).$$

In other words, given any sequence $a \in U_R^{(6)}$, $\mathcal{A}^2(a) = \mathcal{U}(a) \in B_R^{(6)}$, i.e., in $U_R^{(6)}$ the transforms \mathcal{A}^2 and \mathcal{U} are identical.

From Theorem 5 easily follows the next corollary.

Corollary 2 *Given any $a \in U_R^{(2n)}$, we have*

$$\mathcal{U}(a) = \mathcal{A}^{n-1}(a).$$

Moreover, for any $a \in U_R$, we have

$$\mathcal{U}(a) = \lim_{n \rightarrow +\infty} \mathcal{A}^n(a).$$

Clearly, if two sequences $a, b \in H_R$ coincide in the first k terms, then $\mathcal{A}(a)$ and $\mathcal{A}(b)$ coincide at least in the first k terms. Thus, we have the following proposition.

Proposition 5 *Given any $a, b \in H_R$, then*

$$\delta(\mathcal{A}(a), \mathcal{A}(b)) \leq \delta(a, b).$$

By the previous proposition, we have that \mathcal{A} is a contraction mapping on the ultrametric space (H_R, δ) . As a first interesting consequence, we can observe that \mathcal{A} is a continuous function. Moreover, we have that the ultrametric group (U_R, \star, δ) with the contraction mapping \mathcal{A} is an ultrametric dynamic space, where the set of fixed points is the subgroup B_R . In this way, we have found a very interesting example of ultrametric dynamic space. Ultrametric dynamics are very studied in several fields, see [14] for a good reference about dynamics on ultrametric spaces.

References

1. S. Barbero, U. Cerruti, N. Murru, Transforming recurrent sequences by using the Binomial and Invert operators, *Journal of Integer Sequences*, Vol. **13**, Article **10.7.7**, (2010).
2. E. T. Bell, Partition Polynomials, *Annals of Mathematics*, Vol. **29**, 38–46, (1928).
3. A. Benhissi, Ideal structure of Hurwitz series rings, *Contributions to Algebra and Geometry*, Vol. **48**, No. **1**, 251–256, (2007).
4. A. Benhissi, PF and PP-properties in Hurwitz series ring, *Bull. Math. Soc. Sci. Math Roumanie*, Vol. **54**, No. **3**, 203–211, (2011).
5. A. Benhissi, F. Kojka, Basic properties of Hurwitz series rings, *Ricerche di Matematica*, Vol. **61**, No. **2**, 255–273, (2012).
6. M. Bernstein, N. J. A. Sloane, Some canonical sequences of integers, *Linear Algebra Appl.*, Vol. **226–228**, 57–72, (1995).
7. M. Ghanem, Some properties of Hurwitz series ring, *International Mathematical Forum*, Vol. **6**, No. **40**, 1973–1981, (2007).
8. R. Graham, D. Knuth, O. Patashnik, *Concrete mathematics: a foundation for computer science*. Addison–Wesley, 2nd Edition, (1994).
9. W. F. Keigher, On the ring of Hurwitz series, *Commun. Algebra*, Vol. **25**, 1845–1859, (1997).
10. W. F. Keigher, F. L. Pritchard, Hurwitz series as formal functions, *Journal of Pure and Applied Algebra*, Vol. **146**, 291–304, (2000).
11. Z. Liu, Hermite and PS-rings of Hurwitz series, *Communication in Algebra*, Vol. **28**, 299–305, (2000).
12. J. Millar, N. J. A. Sloane, N. E. Young, A new operation on sequences: the Boustrophedon transform, *J. Combinatorial Theory Series A*, Vol. **76**, 44–54, (1996).
13. D. Port, Polynomial maps with applications to combinatorics and probability theory. PhD Thesis, MIT, Available at <http://dspace.mit.edu/handle/1721.1/28041>, (1994).
14. S. Priess-Crampe, P. Ribenboim, Ultrametric dynamics, *Illinois Journal of Mathematics*, Vol. **55**, No. **1**, 287–303, (2011).
15. N. J. A. Sloane, The On-Line Encyclopedia of Integer Sequences, published electronically at <http://www.research.att.com/~njas/sequences>, (2010).