

Unconventional supersymmetry at the boundary of AdS4 supergravity

Original

Unconventional supersymmetry at the boundary of AdS4 supergravity / Andrianopoli, L.; Cerchiai, B. L.; D'Auria, R.; Trigiante, M.. - In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - ELETTRONICO. - 2018:4(2018). [10.1007/JHEP04(2018)007]

Availability:

This version is available at: 11583/2726638 since: 2021-11-21T19:23:44Z

Publisher:

Springer Verlag

Published

DOI:10.1007/JHEP04(2018)007

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Unconventional supersymmetry at the boundary of AdS_4 supergravity

L. Andrianopoli,^{a,b} B.L. Cerchiai,^{a,c} R. D'Auria^a and M. Trigiante^{a,b}

^a*DISAT, Politecnico di Torino,*

Corso Duca degli Abruzzi 24, Turin, I-10129 Italy

^b*Istituto Nazionale di Fisica Nucleare (INFN) Sezione di Torino,*

Via P. Giuria, 1, Torino, Italy

^c*Centro Fermi — Museo Storico della Fisica e Centro Studi e Ricerche “Enrico Fermi”,*

P.za del Viminale 1, Rome, Italy

E-mail: laura.andrianopoli@polito.it, bianca.cerchiai@polito.it,
riccardo.dauria@polito.it, mario.trigiante@polito.it

ABSTRACT: In this paper we perform, in the spirit of the holographic correspondence, a particular asymptotic limit of $\mathcal{N} = 2$, AdS_4 supergravity to $\mathcal{N} = 2$ supergravity on a locally AdS_3 boundary. Our boundary theory enjoys $\text{OSp}(2|2) \times \text{SO}(1, 2)$ invariance and is shown to contain the $D = 3$ super-Chern Simons $\text{OSp}(2|2)$ theory considered in [1] and featuring “unconventional local supersymmetry”. The model constructed in that reference describes the dynamics of a spin-1/2 Dirac field in the absence of spin 3/2 gravitini and was shown to be relevant for the description of graphene, near the Dirac points, for specific spatial geometries. Our construction yields the model in [1] with a specific prescription on the parameters. In this framework the Dirac spin-1/2 fermion originates from the radial components of the gravitini in $D = 4$.

KEYWORDS: Supergravity Models, AdS-CFT Correspondence

ARXIV EPRINT: [1801.08081](https://arxiv.org/abs/1801.08081)

Contents

1	Introduction	1
2	Boundary behavior of $\mathcal{N} = 2$, AdS_4 supergravity	3
2.1	Preliminaries	3
2.2	Supersymmetric boundary conditions	4
2.3	Explicit $D = 3$ description	5
2.4	Asymptotic limit	6
3	Comparison with “supersymmetry of a different kind”	8
4	New insights from the $D = 4$ approach	11
5	Conclusions and outlook	13
A	Notations and conventions	14
A.1	Conventions on spinors in $D = 4$ and $D = 3$	14
A.2	Map between our notations in $D = 3$ and the ones in [1]	15
B	The ultraspinning Kerr-AdS black hole	15
C	Proof of eq. (4.5)	16

1 Introduction

The AdS/CFT duality [2–5] has provided important insights into the properties of strongly coupled non-gravitational models from a classical supergravity theory in one dimension higher. At low energies, in the supergravity limit of string theory, it implies a one-to-one correspondence between quantum operators \mathcal{O} in the boundary conformal field theory and fields ϕ of the bulk supergravity and requires to supplement the supergravity action functional with appropriate boundary conditions $\phi^{(0)}$ for the supergravity fields, which act as sources for the operators of the CFT. As far as the metric field is concerned, in particular, the bulk metric is divergent near the boundary. These divergences can be however disposed of successfully by the so called holographic renormalization [6–14] through the inclusion of appropriate counterterms at the boundary.

In the spirit of the gauge-gravity correspondence the aim of this paper is to relate pure $\mathcal{N} = 2$ AdS_4 supergravity to a $D = 3$ Super-Chern-Simons theory developed in [1]. This latter model displays $\mathcal{N} = 2$ local supersymmetry in spite of the absence of gravitini. This special feature was named “supersymmetry of a different kind” and later “unconventional supersymmetry” by the authors. The only propagating field in the model is a spin-1/2 Dirac spinor with a possible mass term given in terms of the three-dimensional negative cosmological constant. The model discussed in [1] (denoted AVZ in the following) and further developed in [15, 16] was shown to have important applications for some topics in condensed matter physics, in particular the description of graphene near the Dirac points, in

agreement with the observation that the special properties of this material can be formally investigated using the techniques of 3D gravity and quantum field theory [19–24].

In this paper we succeed in reproducing the AVZ $D = 3$ model from $\mathcal{N} = 2$, $D = 4$ pure supergravity in the presence of a 3D boundary, using the results of [25] as a starting point. The supersymmetry of the $D = 4$ supergravity lagrangian constrains the boundary values of the super field strengths to have definite values (Neumann boundary conditions), as was already noticed in [26] in the $\mathcal{N} = 1$ case.

The precise correspondence with the Chern-Simons model of AVZ is found for a very specific choice of the $D = 3$ boundary, corresponding to a local AdS_3 geometry placed at spatial infinity of the $D = 4$ theory. Asymptotically AdS_4 solutions featuring this boundary geometry comprise the “ultraspinning limit” of AdS_4 -Kerr black hole considered in [27–31]. Once the curvatures and fields on the boundary are rewritten in terms of the 3D ones, in the appropriate limit they reproduce the field equations of [1], provided, following the prescription of that paper, the $D = 3$ gravitino only has a spin 1/2 part:

$$\psi_{A\mu} = i\gamma_\mu \chi_A, \quad (1.1)$$

where $A = 1, 2$, $\mu = 0, 1, 2$ and γ^μ are γ -matrices in $D = 3$. From our point of view, this assumption is naturally embedded in the $D = 4$ theory since it is compatible with the on shell condition on the $D = 4$ gravitino $\Psi_{A\hat{\mu}}$ (here, $\hat{\mu} = 0, 1, 2, 3 = (\mu, 3)$, $A = 1, 2$ and we denote by $\Gamma^{\hat{\mu}}$ the γ -matrices in $D = 4$):

$$\Gamma^{\hat{\mu}} \Psi_{A\hat{\mu}} = 0 \Rightarrow \Gamma^\mu \Psi_{A\mu} + \Gamma^3 \Psi_{A3} = 0, \quad (1.2)$$

which projects out its spin-1/2 component in $D = 4$ superspace, but still allows a non trivial solution for $\Gamma^\mu \Psi_{A\mu}$, if the contribution from the radial component Ψ_{A3} of the gravitino is not suppressed in the limit.

The paper is organized as follows: in section 2 we start by recalling, in subsections 2.1 and 2.2, the main facts about the general formulation of pure $\mathcal{N} = 2$ AdS_4 supergravity as developed in [25], where the boundary terms compatible with a full supersymmetry invariance of the supergravity action are computed in a geometric approach. In particular in subsections 2.3 and 2.4 the conditions on the field strengths of the various $D = 4$ fields at the boundary are written in a covariant form with respect to Lorentz group at the boundary. Referring to the Fefferman-Graham parametrization of an asymptotically AdS four-dimensional space-time \mathcal{M} , a suitable behavior of the fields near the boundary $\partial\mathcal{M}$ is devised in order to obtain the model in [1]. To this aim we use a description of the asymptotically AdS_4 geometry which is typical of the “ultraspinning” limit of the AdS -Kerr solution. It is shown that, in the boundary limit $r \rightarrow \infty$, the conditions of the field strengths, which are required by $D = 4$ supersymmetry, naturally yield the field equations of a $D = 3$ super gravity Lagrangian with symmetry $\text{OSp}(2|2) \times \text{SO}(1, 2)$ which, as it is known, can be formulated in terms of (super) Chern-Simons theories [32–39].

In section 3, which is the central part of our paper, we explicitly compare the Chern-Simons lagrangian of [1] with the $D = 3$ supergravity lagrangian whose field equations do coincide with those obtained from the asymptotic values on $\partial\mathcal{M}$ of the super field-strengths. We show that, by imposing the condition (1.1) on the limit values on $\partial\mathcal{M}$ of

the fields and on the $D = 3$ Lagrangian, we exactly reproduce the theory of ref. [1] as a part of the $\mathcal{N} = 2$ super-AdS₃ supergravity Lagrangian.

Section 4 contains a discussion of our results and the specific features of the $D = 3$ models that we obtain from its holographic relation with $D = 4$ supergravity. We deduce the explicit relations among the fields naturally associated with the boundary limit of the $D = 4$ super vielbein and spin connection, and the corresponding fields of ref. [1], where instead supersymmetry is an internal symmetry leaving invariant the space-time dreibein. In particular we find that the supersymmetry parameter of the $D = 3$ theory is proportional to the propagating spinor field of the AVZ model. Furthermore, from the correspondence with $D = 4$ supergravity, we gain an interpretation of the propagating spinor field of the AVZ model in terms of the radial component of the $D = 4$ gravitino.

Our results and possible further developments are presented in section 5.

We left to the appendices some technical material, including the map between our notations and the ones of [1], the details of the ultraspinning limit and the explicit relation between the space-time dreibein of [1] and the superdreibein on $D = 3$ superspace that we get in the limit.

2 Boundary behavior of $\mathcal{N} = 2$, AdS₄ supergravity

2.1 Preliminaries

In the presence of a space-time defect, such as a boundary of space-time, compatibility of the lagrangian with local supersymmetry invariance requires appropriate boundary conditions to be imposed. This was first pointed out in [40–42], in early attempts to study the quantization of gravity with a path integral approach, in order to have an action which depends only on the first derivatives of the metric. More recently, the addition of boundary terms was considered in [43] to cancel gauge and gravitational anomalies in the Hořava-Witten model in 11D. Inclusion of boundary terms and counterterms is also an essential tool for the study of the AdS/CFT duality [2–5] and, as far as the bosonic sector of AdS supergravity is concerned, has been extensively studied in many different contexts. In particular, interesting results were obtained in [44, 45], where it was shown that the addition of the topological Euler-Gauss-Bonnet term to the Einstein action of four dimensional AdS gravity leads to a background-independent definition of Noether charges, without imposing Dirichlet boundary conditions on the fields. Such boundary term indeed regularizes the action and the related (background independent) conserved charges.

At the full supergravity level, boundary contributions were considered from several authors, using different approaches, and in particular in [26, 46–56], and more recently in [57–59]. In [46–51] the minimal AdS supergravity theory in three dimensions with a boundary and with no boundary conditions on the fields was constructed from the requirement of local supersymmetry invariance of the supergravity action, *without imposing Dirichlet boundary conditions on the fields*, in contrast to the Gibbons-Hawking prescription [42]. Within that approach, it was shown in particular that $\mathcal{N} = 1$, $D = 3$ pure supergravity, including its appropriate boundary term, actually reproduces not only the Gibbons-Hawking-York boundary term, but also the counterterm which regularizes the

total action, in the language of holographic renormalization. The problem of boundary conditions in the holographic framework was discussed in generality in [26], in particular for the minimal, $\mathcal{N} = 1$, AdS₄ supergravity.

2.2 Supersymmetric boundary conditions

In [25], the results of [44, 60–63] on four dimensional gravity were extended to the supersymmetric case for $\mathcal{N} = 1$ and $\mathcal{N} = 2$ pure supergravity, in the geometric approach, by adding to the supergravity lagrangian in four dimensions, besides the Euler-Gauss-Bonnet contribution, other topological terms allowing full supersymmetry invariance of the supergravity action, in the bulk and on the boundary of four dimensional space-time. Supersymmetry invariance constrains the boundary values of the superfield-strengths (Newmann boundary conditions), without fixing however the superfields themselves on the boundary. In particular, in [25] a pure $D = 4$, $\mathcal{N} = 2$ theory was written in the presence of boundary. If we denote by \mathcal{M} the four-dimensional space-time and by $\partial\mathcal{M}$ its three-dimensional boundary, supersymmetry requires the following equations to be satisfied at the boundary $\partial\mathcal{M}$ (still written in terms of four dimensional fields):¹

$$R^{ab}|_{\partial\mathcal{M}} = \left[\frac{1}{\ell^2} V^a \wedge V^b + \frac{1}{2\ell} \bar{\Psi}_A \Gamma^{ab} \wedge \Psi_A \right]_{\partial\mathcal{M}}, \quad (2.2)$$

$$\nabla^{(4)} V^a|_{\partial\mathcal{M}} = \frac{i}{2} \bar{\Psi}_A \Gamma^a \wedge \Psi_A|_{\partial\mathcal{M}}, \quad (2.3)$$

$$dA^{(4)}|_{\partial\mathcal{M}} = \bar{\Psi}_A \wedge \Psi_B \epsilon_{AB}|_{\partial\mathcal{M}}, \quad (2.4)$$

$$\nabla^{(4)} \Psi_A|_{\partial\mathcal{M}} = \frac{i}{2\ell} \Gamma_a \Psi_A \wedge V^a|_{\partial\mathcal{M}}, \quad (2.5)$$

where $a, b, \dots = 0, 1, 2, 3$ are four dimensional flat indices, $\nabla^{(4)}$ denotes the SO(1, 3) covariant differential, and we have defined:

$$\begin{aligned} R^{ab} &\equiv d\omega^{ab} + \omega^a_c \wedge \omega^{cb}, \\ \nabla^{(4)} V^a &\equiv dV^a + \omega^a_b \wedge V^b, \\ \nabla^{(4)} \Psi_A &\equiv d\Psi_A + \frac{1}{4} \omega_{ab} \Gamma^{ab} \wedge \Psi_A - \frac{1}{2\ell} \epsilon_{AB} A^{(4)} \wedge \Psi_B. \end{aligned} \quad (2.6)$$

In what follows, for the sake of notational simplicity, we shall omit the wedge symbol.

Equations (2.2)–(2.5) express the condition of an asymptotic super-AdS₄ geometry.

¹We shall use the notations of [25], where in particular the metric is mostly minus.

With respect to that paper, however, here we made some changes which make the formulas more transparent and better adapted to match the notations in three dimensions. More precisely, the spin connection and the curvature are defined with an extra minus sign: $\omega^{ab} \rightarrow -\omega^{ab}$, $R^{ab} \rightarrow -R^{ab}$, and the 1-form with a prefactor $-\frac{1}{\sqrt{2}} : A \rightarrow -\frac{1}{\sqrt{2}} A$. We will use Majorana spinors both in four as well in three dimensions and redefine the constants appearing in the quoted paper as follows:

$$\frac{1}{\ell} = 2e = \frac{P}{\sqrt{2}} = \sqrt{-\frac{\Lambda}{3}}; \quad L = \frac{1}{\sqrt{2}}, \quad (2.1)$$

where Λ is the cosmological constant and ℓ is the AdS₄ radius.

2.3 Explicit $D = 3$ description

The explicit $D = 3$ description of the above supergravity equations depends on the general symmetry properties of the theory on the boundary $\partial\mathcal{M}$ which we wish to relate to.

Our aim in the present paper is to derive the model in [1], which features locally AdS_3 geometries, as the effective theory on an asymptotic boundary $\partial\mathcal{M}$, placed at $r \rightarrow \infty$. To this end we need to suitably choose the boundary behavior of the $D = 4$ fields, which relate them to the $D = 3$ ones.

We refer to a Fefferman-Graham parametrization of the four-dimensional geometry, where the coordinates $x^{\hat{\mu}}$ of the asymptotically- AdS_4 geometry split into the three coordinates x^μ , $\mu = 0, 1, 2$ on $\partial\mathcal{M}$ and the radial coordinate $x^3 = r$. The boundary is located at $r \rightarrow \infty$.

Let us start rewriting the curvatures in (2.2)–(2.5) in a $\text{SO}(1, 1) \times \text{SO}(1, 2)$ -covariant way, where $\text{SO}(1, 1)$ acts as a dilation on the radial variable r and $\text{SO}(1, 2)$ is identified with the Lorentz group on $\partial\mathcal{M}$. Given the extrinsic curvature 1-form $h^i = \omega^{3i}$, we introduce new 1-forms K_+^i , K_-^i , as follows:²

$$K_+^i \equiv \frac{1}{2} (V^i - \ell \omega^{3i}) , \quad K_-^i \equiv \frac{1}{2} (V^i + \ell \omega^{3i}) . \quad (2.7)$$

These 1-forms are 3-vectors with respect to $\text{SO}(1, 2)$ -Lorentz group on the boundary with different $\text{SO}(1, 1)$ -grading: K_+^i have grading +1 and K_-^i grading −1. Similarly we decompose the four-component gravitinos into two-component ones corresponding to different $\text{SO}(1, 1)$ -gradings:

$$\Psi_A = \Psi_{+A} + \Psi_{-A} , \quad \Gamma^3 \Psi_{\pm A} = \pm i \Psi_{\pm A} . \quad (2.8)$$

In terms of these new quantities we find that equations (2.2)–(2.5) on the boundary $\partial\mathcal{M}$ can be cast in the following form:

$$\begin{aligned} \mathcal{R}^{ij} &\equiv d\omega^{ij} + \omega^i_k \wedge \omega^{kj} = \frac{4}{\ell^2} K_+^{[i} \wedge K_-^{j]} + \frac{1}{\ell} \bar{\Psi}_{+A} \Gamma^{ij} \Psi_{-A} , \\ dV^3 &= -\omega^3_i \wedge V^i + \bar{\Psi}_{+A} \Psi_{-A} , \\ \mathcal{D}K_\pm^i &\equiv dK_\pm^i + \omega^i_j \wedge K_\pm^j = \frac{i}{2} \bar{\Psi}_{\pm A} \Gamma^i \Psi_{\pm A} \pm \frac{1}{\ell} K_\pm^i \wedge V^3 , \\ \mathcal{D}\Psi_{\pm A} &\equiv d\Psi_{\pm A} + \frac{1}{4} \omega_{ij} \Gamma^{ij} \wedge \Psi_{\pm A} = -\frac{i}{\ell} K_{\pm i} \wedge \Gamma^i \Psi_{\mp A} + \frac{1}{2\ell} \epsilon_{AB} A^{(4)} \wedge \Psi_{\pm|B} \pm \frac{1}{2\ell} \Psi_{\pm A} \wedge V^3 , \\ dA^{(4)} &= \epsilon_{AB} \bar{\Psi}_A \wedge \Psi_B = 2 \epsilon_{AB} \bar{\Psi}_{+A} \wedge \Psi_{-B} , \end{aligned} \quad (2.9)$$

where use has been made of equations (A.6) of appendix A and we have identified the rigid four-dimensional index $a = 3$ with the radial direction orthogonal to $\partial\mathcal{M}$ at the boundary: $V^3 = dV^3 = 0$. This in particular implies the conditions:³

$$\omega^3_i = h_{i|j} V^j \quad \text{with} \quad h_{i|j} = h_{j|i} ; \quad \bar{\Psi}_{+A} \Psi_{-A} = 0 . \quad (2.10)$$

²Since we use $a, b, \dots = 0, 1, 2, 3$ as rigid $D = 4$ Lorentz indices, we denote the three dimensional $\text{SO}(1, 2)$ ones by $i, j, k, \dots = 0, 1, 2$.

³One can easily show that the extrinsic curvature 1-form ω^3_i cannot have a component along the gravitino field of the superspace.

2.4 Asymptotic limit

Since the model in [1] which we want to relate to is defined on locally AdS_3 geometries, we shall use a description of the asymptotic behavior for the $D = 4$ metric and the spin-connection corresponding to an asymptotically locally AdS_4 space-time with a locally AdS_3 boundary. An example of such backgrounds is the “ultraspinning limit” of an AdS_4 -Kerr black-hole defined in [27, 28], see discussion in appendix B.⁴ In particular we choose the following boundary behavior for the $D = 4$ fields:⁵

$$\begin{aligned} K_+^i(x, r) &= \frac{r}{\ell} E^i(x) + \dots, & K_-^i(x, r) &= \frac{1}{4} \frac{\ell}{r} E^i(x) + \dots, & V^3(r) &= \frac{\ell}{r} dr + \dots, \\ \Psi_{+A\mu}(x, r) &= \sqrt{\frac{r}{\ell}} (\psi_{A\mu}, \mathbf{0}) + \dots, & \Psi_{-A\mu}(x, r) &= \sqrt{\frac{\ell}{r}} \left(\mathbf{0}, \frac{\varepsilon}{2} \psi_{A\mu} \right) + \dots, \\ \omega^{ij}(x, r) &= \omega^{ij}(x) + \dots, & A^{(4)}(x, r) &= \varepsilon A_\mu(x^\nu) dx^\mu + \dots, \end{aligned} \quad (2.11)$$

where $\psi_{A\mu} = (\psi_{A\mu\alpha})$, $\mu = 0, 1, 2$, $\alpha = 1, 2$, are gravitini in $D = 3$, associated with Majorana spinor 1-forms $\psi_A \equiv \psi_{A\mu} dx^\mu$ and $\varepsilon = \pm 1$. The ellipses refer to subleading terms in the $r \rightarrow \infty$ limit. The boundary behavior of the radial components of the $D = 4$ gravitini, $\Psi_{\pm A3}$, will be explicitly discussed later, in eq. (4.11) of section 4. In order to make contact with the model of [1], we shall assume that the $O\left(\frac{\ell^3}{r^3}\right)$ term in the expansion of V^i is absent. This further restriction is satisfied by an AdS_3 slicing of AdS_4 space-time, see appendix B, obtained by sending, in the ultraspinning AdS -Kerr solution, the mass of the black hole to zero.⁶ It implies the vanishing of the energy momentum tensor of the boundary theory [26, 28] (Neumann condition):

$$T_{ij} = 0,$$

which is a feature of the model of [1, 15] that we want to make contact with. By supersymmetry this also implies restrictions on the subleading terms in the expansion of the gravitinos [26, 28]. In order to reproduce the effect of deformations of this model one should consider more general boundary behaviors. We leave this to a future investigation.

The components ω^{ij} , restricted to $\partial\mathcal{M}$, provide the spin-connection on that space, and \mathcal{R}^{ij} the corresponding Riemann curvature 2-form. Let us emphasize that the fields E^i , ψ_A , ω^{ij} , A are to be intended, in the spirit of the geometric approach to supergravity, as superfields in the superspace extension of $\partial\mathcal{M}$, corresponding to the (non-dynamical) super-dreibein and gauge field on $\partial\mathcal{M}$.

⁴See also the solution found in [30] which is described, in a suitable limit, by the same metric.

⁵The ensuing boundary behavior is related to the that given in [26] by the change of coordinates given in appendix B.

⁶In fact it is satisfied by a more general space-time which is the BTZ [64, 65] slicing of a locally AdS_4 , known as ‘BTZ-black string’ and found in [66], see section 5.

If we now perform the $r \rightarrow \infty$ limit, the leading contributions to (2.9) correspond to the following equations on the boundary:

$$\begin{aligned}
 \mathcal{R}^{ij} &= \frac{1}{\ell^2} E^i E^j + \frac{\varepsilon}{2\ell} \bar{\psi}_A \gamma^{ij} \psi_A, \\
 \mathcal{D}E^i &= \frac{i}{2} \bar{\psi}_A \gamma^i \psi_A, \\
 \mathcal{D}\psi_A &= -\varepsilon \frac{i}{2\ell} E_i \gamma^i \psi_A + \frac{\varepsilon}{2\ell} \epsilon_{AB} A \psi_B, \\
 dA &= \epsilon_{AB} \bar{\psi}_A \psi_B,
 \end{aligned} \tag{2.12}$$

where \mathcal{D} denotes Lorentz covariant derivative in $D = 3$, as defined in (2.9). It is expressed in terms of the (super) torsionless spin connection ω^{ij} .⁷ They define the vanishing of the supercurvatures of $\mathcal{N} = 2$ super AdS_3 algebra.

The set of constraints (2.12) on the 3D boundary can be found as equations of motion from the following $D = 3$ supergravity lagrangian [32]:

$$\mathcal{L}^{(3)} = \left(\mathcal{R}^{ij} - \frac{1}{3\ell^2} E^i E^j - \frac{\varepsilon}{2\ell} \bar{\psi}_A \gamma^{ij} \psi_A \right) E^k \epsilon_{ijk} - \frac{\varepsilon}{2\ell} A dA + 2\bar{\psi}_A \left(\mathcal{D}\psi_A - \frac{\varepsilon}{2\ell} \epsilon_{AB} A \psi_B \right). \tag{2.13}$$

Eq. (2.13) collects in fact two inequivalent lagrangians on $D = 3$ superspace, depending on the sign $\varepsilon = \pm 1$. As found in [32], in $D = 3$ a family of inequivalent \mathcal{N} -extended supergravity lagrangians can be constructed, enjoying invariance under $\text{OSp}(p|2) \times \text{OSp}(q|2)$, with $p + q = \mathcal{N}$. In our case, where $\mathcal{N} = 2$, the family of inequivalent lagrangians consists of only two elements, that we labeled with the sign ε . They are invariant under $\text{OSp}(2|2)_{(\varepsilon)} \times \text{SO}(1,2)_{(-\varepsilon)}$. For the time being we keep both theories, which differ by a flipping of the gravitino gauge charge. In the next section we will show that one of the two assignments (in particular, $\varepsilon = -1$) reproduces the results of [1].

In the geometric formalism, the supersymmetry transformation of a generic field Φ , with supersymmetry parameter ϵ_A , can be found as a super-diffeomorphism in the fermionic directions of superspace:

$$\delta\Phi = \iota_K(d\Phi) + d(\iota_K\Phi), \tag{2.14}$$

generated by the tangent vector $K = \bar{\epsilon}_A D^A$, where D^A is the tangent vector dual to the gravitino field, such that $\psi_A(D^B) = \delta_A^B$. The lagrangian (2.13) is then invariant under the following supersymmetry transformations:

$$\begin{aligned}
 \delta\omega^{ij} &= \frac{\varepsilon}{\ell} \bar{\epsilon}_A \gamma^{ij} \psi_A, \\
 \delta E^i &= i \bar{\epsilon}_A \gamma^i \psi_A, \\
 \delta\psi_A &= \mathcal{D}\epsilon_A + \varepsilon \frac{i}{2\ell} E_i \gamma^i \epsilon_A - \frac{\varepsilon}{2\ell} \epsilon_{AB} A \epsilon_B, \\
 \delta A &= 2\epsilon_{AB} \bar{\epsilon}_A \psi_B.
 \end{aligned} \tag{2.15}$$

⁷As it is usual in supergravity and evident from the second equation in (2.12), the spin connection ω^{ij} has instead a non-zero space-time torsion, which is proportional to a gravitino bilinear.

The full $\mathfrak{osp}(2|2)_{(\varepsilon)} \times \mathfrak{so}(1,2)_{(-\varepsilon)}$ can be made manifest by introducing the torsionful spin connections [33]:

$$\omega_{(\pm\varepsilon)}^{ij} = \omega^{ij} \pm \frac{\varepsilon}{\ell} E_k \epsilon^{ijk} = \epsilon^{ijk} \omega_{\pm(\varepsilon)k}, \quad (2.16)$$

in terms of which the constraints (2.12) read:

$$\mathcal{R}_{(\varepsilon)}^i = i \frac{\varepsilon}{\ell} \bar{\psi}_A \gamma^i \psi_A, \quad (2.17)$$

$$\mathcal{R}_{(-\varepsilon)}^i = 0, \quad (2.18)$$

$$\mathcal{D}_{(\varepsilon)} \psi_A = \frac{\varepsilon}{2\ell} \epsilon_{AB} A \psi_B, \quad (2.19)$$

$$dA = \epsilon_{AB} \bar{\psi}_A \psi_B, \quad (2.20)$$

where we defined:

$$R_{(\pm)}^i \equiv d\omega_{(\pm)}^i - \frac{1}{2} \omega_{(\pm)}^j \omega_{(\pm)}^k \epsilon^i{}_{jk}. \quad (2.21)$$

In terms of the new connections, the supersymmetry transformations (2.15) become:

$$\begin{aligned} \delta\omega_{(-\varepsilon)}^i &= 0, & \delta\omega_{(\varepsilon)}^i &= \varepsilon \frac{2i}{\ell} \bar{\epsilon}_A \gamma^i \psi_A, & \delta A &= 2\epsilon_{AB} \bar{\epsilon}_A \psi_B, \\ \delta\psi_A &= \mathcal{D}_{(\varepsilon)} \epsilon_A - \frac{\varepsilon}{2\ell} \epsilon_{AB} A \epsilon_B \equiv \nabla^{(\varepsilon)} \epsilon_A. \end{aligned} \quad (2.22)$$

As it is evident from eqs. (2.22), in this theory the superalgebra $\mathfrak{osp}(2|2) \times \mathfrak{so}(1,2)$ (and in particular supersymmetry) is realized as a gauge symmetry, since eqs. (2.17)–(2.20) can be interpreted as the vanishing of the supercurvatures of the above superalgebra (isomorphic to $\mathcal{N} = 2$ super AdS₃).

As shown in [33], using (2.16) the supergravity lagrangian (2.13) can be written as:

$$\mathcal{L}^{(3)} = \varepsilon (\mathcal{L}_{(\varepsilon)} - \mathcal{L}_{(-\varepsilon)}) \equiv \mathcal{L}_+^{(3)} - \mathcal{L}_-^{(3)}, \quad (2.23)$$

where:

$$\mathcal{L}_{(\varepsilon)} = \frac{\ell}{2} \left(\omega_{(\varepsilon)}^i d\omega_{(\varepsilon)|i} - \frac{1}{3} \omega_{(\varepsilon)}^i \omega_{(\varepsilon)}^j \omega_{(\varepsilon)}^k \epsilon_{ijk} \right) + 2\varepsilon \bar{\psi}_A \nabla^{(\varepsilon)} \psi_A - \frac{\varepsilon}{2\ell} A dA, \quad (2.24)$$

$$\nabla^{(\varepsilon)} \psi_A \equiv \left(d + \frac{1}{4} \omega_{(\varepsilon)}^{ij} \gamma_{ij} \right) \psi_A - \frac{\varepsilon}{2\ell} A \psi_B \epsilon_{AB}, \quad (2.25)$$

$$\mathcal{L}_{(-\varepsilon)} = \frac{\ell}{2} \left(\omega_{(-\varepsilon)}^i d\omega_{(-\varepsilon)|i} - \frac{1}{3} \omega_{(-\varepsilon)}^i \omega_{(-\varepsilon)}^j \omega_{(-\varepsilon)}^k \epsilon_{ijk} \right). \quad (2.26)$$

The component $\mathcal{L}_{(\varepsilon)}$ is the super-Chern Simons lagrangian of the superalgebra $\mathfrak{osp}(2|2)_{(\varepsilon)}$, while $\mathcal{L}_{(-\varepsilon)}$ is the Chern Simons lagrangian of the algebra $\mathfrak{so}(1,2)_{(-\varepsilon)}$.

3 Comparison with “supersymmetry of a different kind”

In ref. [1], the $D = 3$ Chern-Simons action for a $\text{OSp}(2|2)$ connection was considered, and shown to be invariant off-shell under gauge supersymmetry. We aim here to compare

it with the boundary theory discussed above. The explicit map between our notations and conventions on the space-time signature and fields with the ones of [1] is given in appendix A.2.

A peculiar feature of the $\text{OSp}(2|2)$ 1-form connection in [1] is that the spinor 1-form associated with the odd generator of the superalgebra is not a spin-3/2 gravitino, but is instead given in terms of a ($\mathcal{N} = 2$) spin 1/2 field that we name χ_A ,⁸ defined by the following condition:

$$\psi_A = i e_i \gamma^i \chi_A, \quad (3.1)$$

where e_i is a 1-form space-time dreibein, *invariant* under supersymmetry transformations:

$$\delta e^i = 0, \quad (3.2)$$

and satisfying:

$$\hat{\mathcal{D}} e^i = \frac{1}{\ell} \epsilon^{ijk} e_j e_k, \quad (3.3)$$

where $\hat{\mathcal{D}}$ denotes a covariant derivative with respect to an appropriate torsionful connection ω^i .

Once expressed in terms of the same notations, the super Chern-Simons lagrangian introduced in section 3 of [1] does coincide, modulo an overall scaling, with our lagrangian (2.24) *for the choice* $\varepsilon = -1$, if one identifies the spin-connection ω^i of [1] with our $\omega_{(\varepsilon)}^i = \omega_{(-)}^i$. For later convenience, in the following we shall write the lagrangian and the ensuing equations of motion in terms of a generic choice of $\varepsilon = \pm 1$. Indeed, if we plug the Ansatz (3.1) in the lagrangian (2.24) we obtain:

$$\begin{aligned} \mathcal{L}_{(\varepsilon)} = & \frac{\ell}{2} \left(\omega_{(\varepsilon)}^i d\omega_{(\varepsilon)|i} - \frac{1}{3} \omega_{(\varepsilon)}^i \omega_{(\varepsilon)}^j \omega_{(\varepsilon)}^k \varepsilon_{ijk} \right) + 2\varepsilon e_i \mathcal{D}^{(\varepsilon)} e^i \bar{\chi}_A \chi_A + \\ & - 4i \varepsilon \bar{\chi}_A \nabla^{(\varepsilon)} \chi_A e d^3x - \frac{\varepsilon}{2\ell} A dA, \end{aligned} \quad (3.4)$$

where $e \equiv \det(e_\mu^i)$ and we have defined:

$$\nabla^{(\varepsilon)} \chi_A \equiv \gamma^i \nabla_i^{(\varepsilon)} \chi_A = \mathcal{D}^{(\varepsilon)} \chi_A - \frac{\varepsilon}{2\ell} A_{i\epsilon AB} \gamma^i \chi_B. \quad (3.5)$$

In deriving the above lagrangian we have used the following properties:

$$\bar{\psi}_A \nabla^{(\varepsilon)} \psi_A = e_i \mathcal{D}^{(\varepsilon)} e^i \bar{\chi}_A \chi_A - 2i e \bar{\chi}_A \nabla^{(\varepsilon)} \chi_A d^3x, \quad (3.6)$$

being $e^i \wedge e^j \wedge e^k = \epsilon^{ijk} e d^3x$. Note that, defining the Dirac spinor $\chi = \chi_1 + i\chi_2$, we can also rewrite the kinetic term for the two spinors χ_A as:

$$\bar{\chi}_A \mathcal{D}^{(\varepsilon)} \chi_A = \frac{1}{2} \bar{\chi} \left(\overrightarrow{\not{\partial}} - \overleftarrow{\not{\partial}} - \frac{1}{2} \gamma^i \phi_{ij} \gamma^j \right) \chi, \quad (3.7)$$

⁸In ref. [1] the spin-1/2 Dirac field is called ψ . Since in the supergravity literature the symbol ψ is generally reserved to the gravitino, for the sake of clarity we preferred to adhere to the latter convention. Note that the extra factor i with respect to the conventions of ref. [1] ensures that χ_A is Majorana, being related to the Majorana gravitino 1-form ψ_A . Referring to the notation in [1], the spinors $i\chi_A$ introduced here should therefore be identified with the real and imaginary part of the spin- $\frac{1}{2}$ fermion ψ of [1]: $\psi = \chi_1 + i\chi_2$.

which is useful for comparison with [1]. The equation of motion for the spinors χ_A is the Dirac equation:⁹

$$\nabla^{(\varepsilon)} \chi_A - \frac{i}{2} \kappa \chi_A = 0, \quad (3.9)$$

where κ is defined as $e_i \mathcal{D}^{(\varepsilon)} e^i = -\kappa e d^3 x$, as in [1], and in our case $\kappa = \varepsilon 6/\ell$. Eq. (3.9) matches with the equation obtained from (2.19) by means of (3.1). Indeed, from (2.19), using (3.1) and (3.3), we easily derive the following equation:

$$\nabla_i^{(\varepsilon)} \chi_A = \frac{i\varepsilon}{\ell} \gamma_i \chi_A, \quad (3.10)$$

which, when contracted with γ^i , yields the Dirac equation (3.9). As shown in [15], equation (3.10) ensures that no local spin-3/2 components are generated through parallel transport of the fermions.

We remark that, as explicitly shown in appendix C, eq. (3.10) implies:

$$d(\bar{\chi}_A \chi_A) = 0, \quad (3.11)$$

so that $\bar{\chi}\chi$ is a free constant of the model. This is a consequence of having fixed the Weyl invariance hidden in (3.1) in such a way that, in agreement with ref. [1], the torsion tensor in (3.3) only has its fully antisymmetric component.

The supersymmetry transformation of the spinor χ_A can be found from (2.15) and (3.2), given the relation (3.1), and reads:

$$\delta \chi_A = -\frac{i}{3} \gamma^i \nabla_i^{(\varepsilon)} \epsilon_A. \quad (3.12)$$

Consistency requires $\nabla_i^{(\varepsilon)} \epsilon_A$ to have no spin-3/2 component, see [1]:

$$P_i^j \nabla_j^{(\varepsilon)} \epsilon_A = 0, \quad (3.13)$$

where $P_i^j = \delta_i^j - \frac{1}{3} \gamma_i \gamma^j$ is the projector on the spin-3/2 component, which is a condition on ϵ_A . In next section we shall find that our derivation of the model implies a relation between ϵ_A and χ_A which is consistent with (3.13) by virtue of eq. (3.10), in accordance with the fact that supersymmetry is a global symmetry of the $D = 3$ model, which does not affect the counting of the degrees of freedom.¹⁰

The equations of motion for $\omega_{(\varepsilon)}^i$ and A following from the lagrangian (3.4) are given by:

$$\mathcal{R}_{(\varepsilon)}^i = -\frac{\varepsilon}{\ell} \bar{\chi}_A \chi_A \epsilon^{ijk} e_j e_k, \quad (3.14)$$

$$dA = i\epsilon_{AB} \bar{\chi}_A \gamma^k \chi_B e^i e^j \epsilon_{ijk}. \quad (3.15)$$

⁹In terms of the Dirac spinor χ we can write the above equation in the same form as [1]:

$$\left(i\not{\partial} + \frac{k}{2} - \frac{i}{4} \gamma^i \phi_{ij} \gamma^j - \frac{\varepsilon}{2\ell} \not{A} \right) \chi + \frac{i}{2e} \partial_\mu \left(e_i^\mu \gamma^i \right) \chi = 0, \quad (3.8)$$

where $e_i^\mu e_\mu^j = \delta_i^j$.

¹⁰We thank Pedro Alvarez and Jorge Zanelli for a relevant comment on this point.

They reproduce exactly those obtained respectively from the equations (2.17), (2.20), substituting the Ansatz (3.1).

The equation of motion of the e^i field implies the vanishing of the energy momentum tensor $T_{ij} = 0$ [15].¹¹ This is consistent with the assumption we made in section 2.4 when defining the asymptotic limit of the $D = 4$ fields. Indeed, as shown in [26, 28], the absence of $\mathcal{O}\left(\frac{\ell^3}{r^3}\right)$ terms in the expansion of the vielbein is related to the Neumann condition $T_{ij} = 0$.

4 New insights from the $D = 4$ approach

After having derived the model in [1] as a holographic limit of AdS_4 supergravity, let us now emphasize some specific features of our construction of the three-dimensional theory within the four-dimensional supergravity.

- Our boundary lagrangian is $\mathcal{L}^{(3)}$ in (2.13) and is built from the difference of $\mathcal{L}_{(\varepsilon)}$ and $\mathcal{L}_{(-\varepsilon)}$, as in (2.23). This expresses the fact that our $D = 3$ boundary model always has an $\text{OSp}(2|2)_{(\varepsilon)} \times \text{SO}(1, 2)_{(-\varepsilon)}$ symmetry. In particular the cosmological constant of the three dimensional space-time, which defines the mass $|\kappa|$ of the Dirac spinor χ , as observed in [1], in our case is always negative (local AdS_3 geometry).
- The space-time dreibein e^i introduced in [1] for the definition (3.1) satisfies:

$$\mathcal{D}_{(\varepsilon)} e^i \equiv de^i - \epsilon^{ijk} \omega_{(\varepsilon)j} e_k = -\frac{\varepsilon}{\ell} \epsilon^{ijk} e_j e_k. \quad (4.1)$$

It should not to be confused with the superfield dreibein E^i , appearing in (2.12). The presence of two sets of dreibein, e^i and E^i , satisfying different differential equations, seems at first sight a puzzling feature of our model. To clarify this point, let us explicitly compare the torsion equation for the (non dynamical) bosonic dreibein e^i , eq. (4.1), with the super-torsion equation in (2.12) for the super-dreibein E^i entering the definition of $\omega_{(\pm)}^i$, after using (3.1). If written in terms of the spin connection $\omega_{(\varepsilon)}^i$, the second and third equation in (2.12) read:

$$\mathcal{D}_{(\varepsilon)} E^i \equiv dE^i - \epsilon^{ijk} \omega_{(\varepsilon)j} E_k = -\frac{\varepsilon}{\ell} \epsilon^{ijk} E_j E_k + \frac{1}{2} \bar{\chi}_A \chi_A \epsilon^{ijk} e_j e_k, \quad (4.2)$$

$$\gamma_i (\mathcal{D}_{(\varepsilon)} \chi_A) e^i + \gamma_i \chi_A \mathcal{D}_{(\varepsilon)} e^i = \frac{\varepsilon}{2\ell} A \epsilon_{AB} \gamma_i \chi_B e^i. \quad (4.3)$$

Eq. (4.2) contains in its right hand side a redundant set of vielbein, and hints to a relation expressing the E^i in terms of the basis of space-time dreibein e^i and of the complex spinor $\chi \equiv \chi_1 + i\chi_2$.

¹¹Care must be taken to obtain this result from (2.23) since it contains a torsionful connection, so that the contribution from the contorsion tensor to the matter Lagrangian should be included.

We consider the following Ansatz:

$$E^i = (1 + a \bar{\chi}\chi + b(\bar{\chi}\chi)^2) e^i. \quad (4.4)$$

Plugging (4.4) in (4.2), and using (4.1), (4.3), one can determine E^i to be:

$$E^i = \left(1 + \varepsilon \frac{\ell}{2} \bar{\chi}\chi - \frac{\ell^2}{4} (\bar{\chi}\chi)^2\right) e^i \equiv M(\bar{\chi}\chi) e^i. \quad (4.5)$$

The details of the calculation are given in appendix C.

Let us now check compatibility of the supersymmetry transformation of E^i , in eq. (2.15), with the one obtained by its expression (4.5), using (3.12). The direct comparison yields:

$$\bar{\epsilon}_A (\gamma^{ij} + \eta^{ij}) \chi_A = \frac{2}{3} i \eta^{ij} (a + 2b \bar{\chi}\chi) \bar{\chi}_A \gamma^k \nabla_k^{(\varepsilon)} \epsilon_A, \quad (4.6)$$

that in particular requires $\bar{\epsilon}_A \gamma^{ij} \chi_A = 0$. This condition can be non trivially satisfied if:

$$\epsilon_A = iN(\bar{\chi}\chi) \chi_A, \quad N(\bar{\chi}\chi) = \alpha(x) + \beta(x) \bar{\chi}\chi \quad (4.7)$$

with α, β real functions of $D = 3$ space-time. Substituting this Ansatz in (4.6), we find the solution:

$$\alpha = 0, \quad \epsilon_A = i\beta \bar{\chi}_B \chi_B \chi_A, \quad \beta \text{ constant}, \quad (4.8)$$

which still depends on the free parameter β .

- The reinterpretation of the model of [1] in the framework of AdS₄ supergravity gives a new perspective on the meaning of the spinor χ . Indeed, the condition (3.1) identifies the spinor χ_A as the spin- $\frac{1}{2}$ projection of the $D = 3$ gravitino ψ_A :

$$\chi_A = -\frac{i}{3} \gamma_i e^{i|\mu} \psi_{A\mu}, \quad (4.9)$$

where $\mu = 0, 1, 2$ labels curved space-time indices. In our framework, however, we are expressing ψ_A in terms of the $\mathcal{N} = 2$ gravitino in $D = 4$, $\Psi_{A\hat{\mu}}$, where $\hat{\mu} = (\mu, 3)$. The gravitino in $D = 4$ obeys the condition:

$$\Gamma_a V^{a|\hat{\mu}} \Psi_{A\hat{\mu}} = 0, \quad (4.10)$$

which projects out its spin- $\frac{1}{2}$ component all over $D = 4$ superspace, including its boundary $\partial\mathcal{M}$.

Expanding eq. (4.10) in the limit $r \rightarrow \infty$ and using the boundary behavior (2.11) for the supergravity fields, eq. (4.10) implies the following behavior for the radial component Ψ_{A3} of the gravitino fields:

$$\psi_{+A3} = \frac{3i\varepsilon}{2M(\bar{\chi}\chi)} \left(\frac{\ell}{r}\right)^{\frac{5}{2}} (\chi_A, \mathbf{0}), \quad \psi_{-A3} = \frac{3i}{M(\bar{\chi}\chi)} \left(\frac{\ell}{r}\right)^{\frac{3}{2}} (\mathbf{0}, \chi_A). \quad (4.11)$$

In this way we give an interpretation of the spin 1/2 fields χ_A as originating from the radial component of the gravitino field on $\partial\mathcal{M}$. These components, although subleading with respect to the others, are related to them in a non-trivial way through eq. (4.10).

5 Conclusions and outlook

In this paper we have found a holographic relation, at the full supersymmetric level, between AdS_4 , $\mathcal{N} = 2$ pure supergravity and a $D = 3$ Chern-Simons theory, found in [1], exhibiting unconventional supersymmetry, which was shown to describe the behavior of graphene near the Dirac points [17, 18]. It is well known that there exists a close analogy between the properties of some condensed matter special systems, in particular of graphene, and the theoretical framework of quantum field theory and general relativity [19–24]. The formalisms of these theories can and have been used in the past years because the equations of motion of the quasiparticles in graphene (at the Dirac points) have formally the same expression as the Dirac equation of the free electrons both in special and general relativity. In particular, the powerful methods of the relativistic theories turn out to be a very useful tool for exploring the special properties of graphene possessing a two-dimensional spatially curved surface. Viceversa graphene has been proposed as a simple laboratory to check gravitational cosmic phenomena like Hawking-Unruh radiation etc.

In our paper the model of reference [1] was reproduced, after projecting out the spin $3/2$ part of the $D = 3$ gravitino, as a theory living at the boundary of $\mathcal{N} = 2$, $D = 4$ Supergravity, in the spirit of $\text{AdS}_4/\text{CFT}_3$ correspondence (for a recent review on a top-down approach to the subject, see [67]). This hints to conjecture a more physical meaning for such relationship. Supersymmetry plays a crucial role in establishing the above correspondence, and the results presented in this paper reinforce the holographic relationship between $(2+1)$ -dimensional condensed matter models and supergravity at the full supersymmetric level, already in the absence of interactions.

Consistently with [1], the $D = 3$ model that we derive has vanishing energy-momentum tensor, as it follows from the equations of motion for $e_\mu{}^i$. The analyses of [26, 28] imply that the dual supergravity background is locally AdS. The duality correspondence discussed in the present paper refers to “vacuum” theories. Reproducing the effects of deformations of the CFT on $\partial\mathcal{M}$ calls for a generalization of boundary behaviors (2.11). Already in [1] it was observed that the solutions of the model include the Bañados-Teitelboim-Zanelli black holes [64, 65]. In light of our duality with $D = 4$ supergravity, these solutions correspond to the so-called ‘BTZ-black string’ found in [66]:

$$ds^2 = \left(1 + \frac{\rho^2}{\ell^2}\right) ds_{\text{BTZ}}^2 - \frac{d\rho^2}{1 + \frac{\rho^2}{\ell^2}}, \quad (5.1)$$

where ds_{BTZ}^2 is the metric of a generic BTZ black hole and we can identify $\rho = \ell \sinh(\alpha)$. The above issues are left to future work.

Finally, let us observe that the Supergravity approach gives, as an extra bonus, the possibility to obtain directly a gauge supersymmetry in $(2+1)$ -dimensional models, even if the Lagrangian of the mother theory only enjoys a supersymmetric invariance which is not gauge since its algebra only closes on-shell.

Acknowledgments

The motivation of this paper arose from a stimulating and fruitful discussion with Jorge Zanelli, to whom we express all our gratitude.

We also acknowledge interesting discussions with Pedro Alvarez, Andrés Anabalon, Matteo Baggioli, Antonio Gallerati, Dietmar Klemm and Lucrezia Ravera.

A Notations and conventions

Throughout the paper we used a mostly minus signature, both in four and in three dimensions. World indices are denoted with greek letters: $\hat{\mu} \dots = 0, 1, 2, 3 = (\mu, 3)$; $\mu, \dots = 0, 1, 2$; $x^3 = r$ denoting the radial coordinate, while to label flat indices we used latin indices: $a, \dots = 0, 1, 2, 3 = (i, 3)$; $i, \dots = 0, 1, 2$; the flat direction orthogonal to $\partial\mathcal{M}$ being labeled with 3.

A.1 Conventions on spinors in $D = 4$ and $D = 3$

It is convenient to use the following representation of the Clifford algebra:

$$\begin{aligned}\Gamma^i &= \begin{pmatrix} \mathbf{0} & \gamma^i \\ \gamma^i & \mathbf{0} \end{pmatrix} = \sigma_1 \otimes \gamma^i, & \gamma^0 &= \sigma_2, & \gamma^1 &= i\sigma_1, & \gamma^2 &= i\sigma_3, \\ \Gamma^3 &= \begin{pmatrix} i & 0 & 0 & 0 \\ 0 & i & 0 & 0 \\ 0 & 0 & -i & 0 \\ 0 & 0 & 0 & -i \end{pmatrix} = i\sigma_3 \otimes \mathbf{1}, \\ \Gamma^5 &= i\Gamma^0\Gamma^1\Gamma^2\Gamma^3 = \begin{pmatrix} 0 & 0 & i & 0 \\ 0 & 0 & 0 & i \\ -i & 0 & 0 & 0 \\ 0 & -i & 0 & 0 \end{pmatrix} = -\sigma_2 \otimes \mathbf{1}.\end{aligned}\tag{A.1}$$

The charge conjugation matrix is taken to be:

$$C = \Gamma^0, \quad C^{-1}\Gamma^a C = -(\Gamma^a)^T.\tag{A.2}$$

Let us define Ψ_{\pm} as the following projections:

$$\Gamma^3\Psi_{\pm} = \pm i\Psi_{\pm}.\tag{A.3}$$

A Majorana spinor λ satisfies the condition:

$$\bar{\lambda} \equiv \lambda^\dagger \Gamma^0 = \lambda^T C \implies \lambda = \lambda^*,\tag{A.4}$$

namely it is real. In the new basis the projected Majorana spinor 1-forms Ψ_{\pm} satisfying the condition $\Gamma^3\Psi_{\pm} = \pm i\Psi_{\pm}$ have the following forms:

$$\Psi_+ = (\xi_1, \xi_2, 0, 0), \quad \Psi_- = (0, 0, \xi_3, \xi_4), \quad \xi_{\alpha}^* = \xi_{\alpha}.\tag{A.5}$$

The following bilinears vanish:

$$\overline{\Psi_{\pm}}\Gamma^{ij}\Psi_{\pm} = \overline{\Psi_{\pm}}\Gamma^3\Psi_{\pm} = \overline{\Psi_{\pm}}\Psi_{\pm} = 0.\tag{A.6}$$

A.2 Map between our notations in $D = 3$ and the ones in [1]

The conventions used here are quite different from the ones of [1] in various respects. For the sake of clarity, we are providing here the relations between the notions used throughout this paper and the ones of [1]. To make the comparison, the items referring to [1] are labeled with a tilde.

- i) We are using a mostly minus signature for the space-time metric, $\eta = (+, -, -)$, while $\tilde{\eta} = (-, +, +)$ is mostly plus. they are related by

$$\tilde{\eta}_{ij} = -\eta_{ij},$$

from which it descends:

$$\begin{aligned}\tilde{\gamma}^i &= i\gamma^i, & \tilde{\gamma}_i &= -i\gamma_i, \\ \tilde{\epsilon}_{ijk} &= \epsilon_{ijk}, & \tilde{\epsilon}^{ijk} &= -\epsilon^{ijk}, \\ \tilde{\omega}^i_j &= \omega^i_j, & \tilde{\omega}_{ij} &= -\omega_{ij}, & \tilde{\omega}^i &= \frac{1}{2}\tilde{\epsilon}^{ijk}\tilde{\omega}_{jk} = \frac{1}{2}\epsilon^{ijk}\omega_{jk} = \omega^i, \\ \tilde{J}_i &= -\frac{\tilde{\gamma}_i}{2} = i\frac{\gamma_i}{2} = J_i.\end{aligned}\tag{A.7}$$

- ii) In [1] all fields are part of a dimensionless connection. The dimensions of the $D = 3$ supergravity fields considered here are:

$$[\psi_A] = [L^{\frac{1}{2}}], \quad [E^i] = [e^i] = [L], \quad [\chi_A] = [L^{-\frac{1}{2}}], \quad [A] = [L], \quad [\omega^{ij}] = [L^0].\tag{A.8}$$

Let us denote here by $\tilde{\chi}$ the spin-1/2 field in [1], denoted there by ψ . The relation between the fields used here and in [1] is:

$$\tilde{e}^i = e^i, \quad \tilde{\chi} = -\frac{1}{\sqrt{\ell}}\chi = -\frac{1}{\sqrt{\ell}}(\chi_1 + i\chi_2), \quad \tilde{\bar{\chi}} = -\frac{1}{\sqrt{\ell}}\bar{\chi}, \quad \tilde{A} = \frac{A}{2\ell}.\tag{A.9}$$

B The ultraspinning Kerr-AdS black hole

In this appendix we give a specific four-dimensional background which satisfies the boundary conditions (2.11). It is the ultraspinning limit of the four-dimensional Kerr-AdS geometry [27–31]. This space-time is obtained by performing the limit $a \rightarrow \ell$ (ℓ being the AdS radius) of the rotation parameter a of a Kerr-AdS black hole, keeping the horizon area fixed and, at the same time, zooming into the pole. The metric reads:

$$ds^2 = V(\rho) \left[dt - \ell \sinh^2\left(\frac{\sigma}{2}\right) d\phi \right]^2 - \frac{\ell^2 + \rho^2}{4} (d\sigma^2 + \sinh^2(\sigma)d\phi^2) - \frac{d\rho^2}{V(\rho)},\tag{B.1}$$

where:

$$V(\rho) = -\frac{2M\rho}{\rho^2 + \ell^2} + \frac{\rho^2}{\ell^2} + 1.\tag{B.2}$$

The boundary is located at $\rho \rightarrow \infty$ and its metric ds_{bdry}^2 is:

$$ds_{\text{bdry}}^2 = \lim_{\rho \rightarrow \infty} \frac{\ell^2}{\rho^2} ds^2 = \left[dt - \ell \sinh^2 \left(\frac{\sigma}{2} \right) d\phi \right]^2 - \frac{\ell^2}{4} (d\sigma^2 + \sinh^2(\sigma) d\phi^2) . \quad (\text{B.3})$$

It describes an AdS_3 spacetime with radius ℓ . If we set $M = 0$ the metric (B.1) provides a parametrization of the AdS_4 vacuum with an AdS_3 boundary geometry. The asymptotic behaviors (2.11) are referred to the Fefferman-Graham radial coordinate r , which is related to ρ by the condition:

$$\frac{d\rho^2}{1 + \frac{\rho^2}{\ell^2}} = \ell^2 \frac{dr^2}{r^2} \Rightarrow \rho = \frac{r}{2} \left(1 - \frac{\ell^2}{r^2} \right) . \quad (\text{B.4})$$

The reader can verify that, for the vacuum solution $M = 0$, the following asymptotic expansions hold:

$$\begin{aligned} V^i &= \frac{r}{2\ell} E^i \left[1 + \frac{\ell^2}{r^2} + O\left(\frac{\ell^6}{r^6}\right) \right] , \\ \omega^{3i} &= -\frac{r}{2\ell} E^i \left[1 - \frac{\ell^2}{r^2} + O\left(\frac{\ell^6}{r^6}\right) \right] , \end{aligned} \quad (\text{B.5})$$

which yield asymptotic behaviors of K_{\pm}^i in (2.11). Note that there is no $O\left(\frac{\ell^3}{r^3}\right)$ term in the expansion of V^i , consistently with our assumptions in section 2.4. This, together with the absence of $O\left(\left(\frac{\ell}{r}\right)^{\frac{3}{2}}\right)$ -terms in the expansion of the gravitinos implies the vanishing of the energy momentum tensor and the supercurrent [26, 28]:

$$T_{ij} = 0 , \quad J_i = 0 , \quad (\text{B.6})$$

in the boundary CFT.

C Proof of eq. (4.5)

We are going to show here how the parametrization of the super-dreibein E^i on the 3D space-time spanned by the e^i :

$$E^i = \left(1 + \varepsilon \frac{\ell}{2} \bar{\chi} \chi - \frac{\ell^2}{4} (\bar{\chi} \chi)^2 \right) e^i \equiv M(\bar{\chi} \chi) e^i \quad (\text{C.1})$$

comes out. Let us start from eqs. (4.1). By plugging the ansatz (4.4) into eq. (4.1), we get:

$$2(a + 2b\bar{\chi}\chi)\bar{\chi}\mathcal{D}_{(\varepsilon)}\chi e^i - \frac{\varepsilon}{\ell} \epsilon^{ijk} M(\bar{\chi}\chi) e_j e_k = \epsilon^{ijk} e_j e_k \left[\frac{1}{2} \bar{\chi}\chi - \frac{\varepsilon}{\ell} (M(\bar{\chi}\chi))^2 \right] . \quad (\text{C.2})$$

To solve eq. (C.2) requires to first determine $\bar{\chi}\mathcal{D}_{(\varepsilon)}\chi$. This is found by looking at (4.3), that can be written:

$$\gamma_i(\mathcal{D}_{(\varepsilon)}\chi_A)e^i - \frac{\varepsilon}{\ell} \epsilon_{ijk} \gamma_k \chi_A e^j - \frac{\varepsilon}{2\ell} A \epsilon_{AB} \gamma_i \chi_B e^i = 0 , \quad (\text{C.3})$$

implying:

$$\gamma_{[i}\mathcal{D}_{(\varepsilon)j]}\chi_A + \frac{\varepsilon}{\ell}\epsilon_{ijk}\gamma_k\chi_A - \frac{\varepsilon}{2\ell}A_{[j}\epsilon_{AB}\gamma_{i]}\chi_B = 0. \quad (\text{C.4})$$

Multiplication of (C.4) by γ^{ij} gives the Dirac equation for χ_A :

$$\gamma^i\mathcal{D}_{(\varepsilon)i}\chi_A - \varepsilon\frac{3}{\ell}\mathbf{i}\chi_A - \frac{\varepsilon}{2\ell}A_i\epsilon_{AB}\gamma^i\chi_B = 0, \quad (\text{C.5})$$

while its multiplication for γ^i :

$$\mathcal{D}_{(\varepsilon)j}\chi_A - 4\mathbf{i}\frac{\varepsilon}{\ell}\gamma_j\chi_A - \frac{\varepsilon}{2\ell}A_j\epsilon_{AB}\chi_B + \gamma_j\left(\gamma^i\mathcal{D}_{(\varepsilon)i}\chi_A - \frac{\varepsilon}{2\ell}A_i\epsilon_{AB}\gamma^i\chi_B\right) = 0. \quad (\text{C.6})$$

Combining (C.5) with (C.6) we finally get:

$$\mathcal{D}_{(\varepsilon)j}\chi_A - \mathbf{i}\frac{\varepsilon}{\ell}\gamma_j\chi_A - \frac{\varepsilon}{2\ell}A_j\epsilon_{AB}\chi_B = 0, \quad (\text{C.7})$$

so that, if we now multiply on the left eq. (C.7) with $\bar{\chi}_A$, we get:

$$\bar{\chi}\mathcal{D}_{(\varepsilon)j}\chi = 0. \quad (\text{C.8})$$

Eq. (C.2) has thus turned into an algebraic equation:¹²

$$\begin{aligned} -\frac{\varepsilon}{\ell}M(\bar{\chi}\chi) &= \left[\frac{1}{2}\bar{\chi}\chi - \frac{\varepsilon}{\ell}(M(\bar{\chi}\chi))^2\right], \\ (1 + a\bar{\chi}\chi + b(\bar{\chi}\chi)^2) &= \left[1 + \left(2a - \varepsilon\frac{\ell}{2}\right)\bar{\chi}\chi + (a^2 + 2b)(\bar{\chi}\chi)^2\right], \end{aligned} \quad (\text{C.9})$$

which is solved for $a = \varepsilon\frac{\ell}{2}$, $b = -a^2$, so that (4.5) follows.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] P.D. Alvarez, M. Valenzuela and J. Zanelli, *Supersymmetry of a different kind*, *JHEP* **04** (2012) 058 [[arXiv:1109.3944](https://arxiv.org/abs/1109.3944)] [[INSPIRE](#)].
- [2] J.M. Maldacena, *The Large N limit of superconformal field theories and supergravity*, *Int. J. Theor. Phys.* **38** (1999) 1113 [*Adv. Theor. Math. Phys.* **2** (1998) 231] [[hep-th/9711200](https://arxiv.org/abs/hep-th/9711200)] [[INSPIRE](#)].
- [3] S.S. Gubser, I.R. Klebanov and A.M. Polyakov, *Gauge theory correlators from noncritical string theory*, *Phys. Lett. B* **428** (1998) 105 [[hep-th/9802109](https://arxiv.org/abs/hep-th/9802109)] [[INSPIRE](#)].
- [4] E. Witten, *Anti-de Sitter space and holography*, *Adv. Theor. Math. Phys.* **2** (1998) 253 [[hep-th/9802150](https://arxiv.org/abs/hep-th/9802150)] [[INSPIRE](#)].

¹²Due to the Grassman nature of χ_A , it is in fact a polynomial equation quadratic in $\bar{\chi}\chi$.

- [5] O. Aharony, S.S. Gubser, J.M. Maldacena, H. Ooguri and Y. Oz, *Large N field theories, string theory and gravity*, *Phys. Rept.* **323** (2000) 183 [[hep-th/9905111](#)] [[INSPIRE](#)].
- [6] V. Balasubramanian and P. Kraus, *A Stress tensor for Anti-de Sitter gravity*, *Commun. Math. Phys.* **208** (1999) 413 [[hep-th/9902121](#)] [[INSPIRE](#)].
- [7] J. de Boer, E.P. Verlinde and H.L. Verlinde, *On the holographic renormalization group*, *JHEP* **08** (2000) 003 [[hep-th/9912012](#)] [[INSPIRE](#)].
- [8] E.P. Verlinde and H.L. Verlinde, *RG flow, gravity and the cosmological constant*, *JHEP* **05** (2000) 034 [[hep-th/9912018](#)] [[INSPIRE](#)].
- [9] J. de Boer, *The Holographic renormalization group*, *Fortsch. Phys.* **49** (2001) 339 [[hep-th/0101026](#)] [[INSPIRE](#)].
- [10] S. de Haro, S.N. Solodukhin and K. Skenderis, *Holographic reconstruction of space-time and renormalization in the AdS/CFT correspondence*, *Commun. Math. Phys.* **217** (2001) 595 [[hep-th/0002230](#)] [[INSPIRE](#)].
- [11] K. Skenderis, *Lecture notes on holographic renormalization*, *Class. Quant. Grav.* **19** (2002) 5849 [[hep-th/0209067](#)] [[INSPIRE](#)].
- [12] I. Papadimitriou and K. Skenderis, *AdS/CFT correspondence and geometry*, *IRMA Lect. Math. Theor. Phys.* **8** (2005) 73 [[hep-th/0404176](#)] [[INSPIRE](#)].
- [13] A. Lawrence and A. Sever, *Holography and renormalization in Lorentzian signature*, *JHEP* **10** (2006) 013 [[hep-th/0606022](#)] [[INSPIRE](#)].
- [14] I. Heemskerk and J. Polchinski, *Holographic and Wilsonian Renormalization Groups*, *JHEP* **06** (2011) 031 [[arXiv:1010.1264](#)] [[INSPIRE](#)].
- [15] A. Guevara, P. Pais and J. Zanelli, *Dynamical Contents of Unconventional Supersymmetry*, *JHEP* **08** (2016) 085 [[arXiv:1606.05239](#)] [[INSPIRE](#)].
- [16] P.D. Alvarez, P. Pais and J. Zanelli, *Unconventional supersymmetry and its breaking*, *Phys. Lett. B* **735** (2014) 314 [[arXiv:1306.1247](#)] [[INSPIRE](#)].
- [17] A.H. Castro Neto, F. Guinea, N.M.R. Peres, K.S. Novoselov and A.K. Geim, *The electronic properties of graphene*, *Rev. Mod. Phys.* **81** (2009) 109 [[INSPIRE](#)].
- [18] A. Lucas and K.C. Fong, *Hydrodynamics of electrons in graphene*, *J. Phys. Condens. Matter* **30** (2018) 053001 [[arXiv:1710.08425](#)] [[INSPIRE](#)].
- [19] A. Cortijo and M.A.H. Vozmediano, *A Cosmological model for corrugated graphene sheets*, *Eur. Phys. J. ST* **148** (2007) 83 [[cond-mat/0612623](#)] [[INSPIRE](#)].
- [20] A. Cortijo and M.A.H. Vozmediano, *Effects of topological defects and local curvature on the electronic properties of planar graphene*, *Nucl. Phys. B* **763** (2007) 293 [Erratum *ibid.* **B 807** (2009) 659] [[cond-mat/0612374](#)] [[INSPIRE](#)].
- [21] A. Cortijo, F. Guinea and M.A.H. Vozmediano, *Geometrical and topological aspects of graphene and related materials*, *J. Phys. A* **45** (2012) 383001 [[arXiv:1112.2054](#)] [[INSPIRE](#)].
- [22] A. Iorio and G. Lambiase, *The Hawking-Unruh phenomenon on graphene*, *Phys. Lett. B* **716** (2012) 334 [[arXiv:1108.2340](#)] [[INSPIRE](#)].
- [23] A. Iorio and G. Lambiase, *Quantum field theory in curved graphene spacetimes, Lobachevsky geometry, Weyl symmetry, Hawking effect and all that*, *Phys. Rev. D* **90** (2014) 025006 [[arXiv:1308.0265](#)] [[INSPIRE](#)].

- [24] M. Cvetič and G.W. Gibbons, *Graphene and the Zermelo Optical Metric of the BTZ Black Hole*, *Annals Phys.* **327** (2012) 2617 [[arXiv:1202.2938](#)] [[INSPIRE](#)].
- [25] L. Andrianopoli and R. D'Auria, *$N = 1$ and $N = 2$ pure supergravities on a manifold with boundary*, *JHEP* **08** (2014) 012 [[arXiv:1405.2010](#)] [[INSPIRE](#)].
- [26] A.J. Amsel and G. Compere, *Supergravity at the boundary of AdS supergravity*, *Phys. Rev. D* **79** (2009) 085006 [[arXiv:0901.3609](#)] [[INSPIRE](#)].
- [27] M.M. Caldarelli, R. Emparan and M.J. Rodriguez, *Black Rings in (Anti)-deSitter space*, *JHEP* **11** (2008) 011 [[arXiv:0806.1954](#)] [[INSPIRE](#)].
- [28] M.M. Caldarelli, R.G. Leigh, A.C. Petkou, P.M. Petropoulos, V. Pozzoli and K. Siampos, *Vorticity in holographic fluids*, *PoS(CORFU2011)076* [[arXiv:1206.4351](#)] [[INSPIRE](#)].
- [29] A. Gnechchi, K. Hristov, D. Klemm, C. Toldo and O. Vaughan, *Rotating black holes in 4d gauged supergravity*, *JHEP* **01** (2014) 127 [[arXiv:1311.1795](#)] [[INSPIRE](#)].
- [30] D. Klemm, *Four-dimensional black holes with unusual horizons*, *Phys. Rev. D* **89** (2014) 084007 [[arXiv:1401.3107](#)] [[INSPIRE](#)].
- [31] R.A. Hennigar, D. Kubizňák, R.B. Mann and N. Musoke, *Ultraspinning limits and super-entropic black holes*, *JHEP* **06** (2015) 096 [[arXiv:1504.07529](#)] [[INSPIRE](#)].
- [32] A. Achúcarro and P.K. Townsend, *A Chern-Simons Action for Three-Dimensional anti-de Sitter Supergravity Theories*, *Phys. Lett. B* **180** (1986) 89 [[INSPIRE](#)].
- [33] A. Achúcarro and P.K. Townsend, *Extended Supergravities in $d = (2 + 1)$ as Chern-Simons Theories*, *Phys. Lett. B* **229** (1989) 383 [[INSPIRE](#)].
- [34] E. Witten, *$(2 + 1)$ -Dimensional Gravity as an Exactly Soluble System*, *Nucl. Phys. B* **311** (1988) 46 [[INSPIRE](#)].
- [35] E. Witten, *Three-Dimensional Gravity Revisited*, [arXiv:0706.3359](#) [[INSPIRE](#)].
- [36] E. Witten, *SL(2, Z) action on three-dimensional conformal field theories with Abelian symmetry*, in *From fields to strings: circumnavigating theoretical physics: Ian Kogan Memorial Collection. Volume 2*, M. Shifman, A. Vainshtein and J. Wheeler eds., World Scientific, Singapore (2005), pp. 1173–1200 [[hep-th/0307041](#)] [[INSPIRE](#)].
- [37] S.L. Cacciatori, M.M. Caldarelli, A. Giacomini, D. Klemm and D.S. Mansi, *Chern-Simons formulation of three-dimensional gravity with torsion and nonmetricity*, *J. Geom. Phys.* **56** (2006) 2523 [[hep-th/0507200](#)] [[INSPIRE](#)].
- [38] R.G. Leigh, N.N. Hoang and A.C. Petkou, *Torsion and the Gravity Dual of Parity Symmetry Breaking in AdS_4/CFT_3 Holography*, *JHEP* **03** (2009) 033 [[arXiv:0809.5258](#)] [[INSPIRE](#)].
- [39] P.K. Townsend and B. Zhang, *Thermodynamics of “Exotic” Bañados-Teitelboim-Zanelli Black Holes*, *Phys. Rev. Lett.* **110** (2013) 241302 [[arXiv:1302.3874](#)] [[INSPIRE](#)].
- [40] J.W. York Jr., *Role of conformal three geometry in the dynamics of gravitation*, *Phys. Rev. Lett.* **28** (1972) 1082 [[INSPIRE](#)].
- [41] J.D. Brown and J.W. York Jr., *Quasilocal energy and conserved charges derived from the gravitational action*, *Phys. Rev. D* **47** (1993) 1407 [[gr-qc/9209012](#)] [[INSPIRE](#)].
- [42] G.W. Gibbons and S.W. Hawking, *Action Integrals and Partition Functions in Quantum Gravity*, *Phys. Rev. D* **15** (1977) 2752 [[INSPIRE](#)].

- [43] P. Hořava and E. Witten, *Eleven-dimensional supergravity on a manifold with boundary*, *Nucl. Phys. B* **475** (1996) 94 [[hep-th/9603142](#)] [[INSPIRE](#)].
- [44] R. Aros, M. Contreras, R. Olea, R. Troncoso and J. Zanelli, *Conserved charges for gravity with locally AdS asymptotics*, *Phys. Rev. Lett.* **84** (2000) 1647 [[gr-qc/9909015](#)] [[INSPIRE](#)].
- [45] O. Mišković and R. Olea, *Topological regularization and self-duality in four-dimensional anti-de Sitter gravity*, *Phys. Rev. D* **79** (2009) 124020 [[arXiv:0902.2082](#)] [[INSPIRE](#)].
- [46] P. van Nieuwenhuizen and D.V. Vassilevich, *Consistent boundary conditions for supergravity*, *Class. Quant. Grav.* **22** (2005) 5029 [[hep-th/0507172](#)] [[INSPIRE](#)].
- [47] D.V. Belyaev, *Boundary conditions in supergravity on a manifold with boundary*, *JHEP* **01** (2006) 047 [[hep-th/0509172](#)] [[INSPIRE](#)].
- [48] D.V. Belyaev and P. van Nieuwenhuizen, *Tensor calculus for supergravity on a manifold with boundary*, *JHEP* **02** (2008) 047 [[arXiv:0711.2272](#)] [[INSPIRE](#)].
- [49] D.V. Belyaev and P. van Nieuwenhuizen, *Simple $d = 4$ supergravity with a boundary*, *JHEP* **09** (2008) 069 [[arXiv:0806.4723](#)] [[INSPIRE](#)].
- [50] D. Grumiller and P. van Nieuwenhuizen, *Holographic counterterms from local supersymmetry without boundary conditions*, *Phys. Lett. B* **682** (2010) 462 [[arXiv:0908.3486](#)] [[INSPIRE](#)].
- [51] D.V. Belyaev and T.G. Pugh, *The Supermultiplet of boundary conditions in supergravity*, *JHEP* **10** (2010) 031 [[arXiv:1008.1574](#)] [[INSPIRE](#)].
- [52] G. Esposito, A.Y. Kamenshchik and K. Kirsten, *One loop effective action for Euclidean Maxwell theory on manifolds with boundary*, *Phys. Rev. D* **54** (1996) 7328 [[hep-th/9606132](#)] [[INSPIRE](#)].
- [53] I.G. Avramidi and G. Esposito, *Gauge theories on manifolds with boundary*, *Commun. Math. Phys.* **200** (1999) 495 [[hep-th/9710048](#)] [[INSPIRE](#)].
- [54] I.G. Moss, *Boundary terms for eleven-dimensional supergravity and M-theory*, *Phys. Lett. B* **577** (2003) 71 [[hep-th/0308159](#)] [[INSPIRE](#)].
- [55] I.G. Moss, *Boundary terms for supergravity and heterotic M-theory*, *Nucl. Phys. B* **729** (2005) 179 [[hep-th/0403106](#)] [[INSPIRE](#)].
- [56] P.S. Howe, T.G. Pugh, K.S. Stelle and C. Strickland-Constable, *Ectoplasm with an Edge*, *JHEP* **08** (2011) 081 [[arXiv:1104.4387](#)] [[INSPIRE](#)].
- [57] D.Z. Freedman, K. Pilch, S.S. Pufu and N.P. Warner, *Boundary Terms and Three-Point Functions: An AdS/CFT Puzzle Resolved*, *JHEP* **06** (2017) 053 [[arXiv:1611.01888](#)] [[INSPIRE](#)].
- [58] P. Benetti Genolini, D. Cassani, D. Martelli and J. Sparks, *Holographic renormalization and supersymmetry*, *JHEP* **02** (2017) 132 [[arXiv:1612.06761](#)] [[INSPIRE](#)].
- [59] N. Drukker, D. Martelli and I. Shamir, *The energy-momentum multiplet of supersymmetric defect field theories*, *JHEP* **08** (2017) 010 [[arXiv:1701.04323](#)] [[INSPIRE](#)].
- [60] R. Aros, M. Contreras, R. Olea, R. Troncoso and J. Zanelli, *Conserved charges for even dimensional asymptotically AdS gravity theories*, *Phys. Rev. D* **62** (2000) 044002 [[hep-th/9912045](#)] [[INSPIRE](#)].
- [61] P. Mora, R. Olea, R. Troncoso and J. Zanelli, *Finite action principle for Chern-Simons AdS gravity*, *JHEP* **06** (2004) 036 [[hep-th/0405267](#)] [[INSPIRE](#)].

- [62] R. Olea, *Mass, angular momentum and thermodynamics in four-dimensional Kerr-AdS black holes*, *JHEP* **06** (2005) 023 [[hep-th/0504233](#)] [[INSPIRE](#)].
- [63] D.P. Jatkar, G. Kofinas, O. Mišković and R. Olea, *Conformal Mass in AdS gravity*, *Phys. Rev. D* **89** (2014) 124010 [[arXiv:1404.1411](#)] [[INSPIRE](#)].
- [64] M. Bañados, C. Teitelboim and J. Zanelli, *The Black hole in three-dimensional space-time*, *Phys. Rev. Lett.* **69** (1992) 1849 [[hep-th/9204099](#)] [[INSPIRE](#)].
- [65] M. Bañados, M. Henneaux, C. Teitelboim and J. Zanelli, *Geometry of the $(2+1)$ black hole*, *Phys. Rev. D* **48** (1993) 1506 [*Erratum ibid.* **D 88** (2013) 069902] [[gr-qc/9302012](#)] [[INSPIRE](#)].
- [66] R. Emparan, G.T. Horowitz and R.C. Myers, *Exact description of black holes on branes II: comparison with BTZ black holes and black strings*, *JHEP* **01** (2000) 021 [[hep-th/9912135](#)] [[INSPIRE](#)].
- [67] G.W. Semenoff, *Engineering holographic graphene*, *AIP Conf. Proc.* **1483** (2012) 305 [[INSPIRE](#)].