

POLITECNICO DI TORINO  
Repository ISTITUZIONALE

On Repunits

*Original*

On Repunits / Sparavigna, Amelia Carolina. - ELETTRONICO. - (2019). [10.5281/zenodo.2639620]

*Availability:*

This version is available at: 11583/2734816 since: 2019-06-07T09:16:28Z

*Publisher:*

Zenodo

*Published*

DOI:10.5281/zenodo.2639620

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

## On Repunits

Amelia Carolina Sparavigna

Politecnico di Torino

**Abstract** Here we discuss the repunits. An operation of addition of these numbers is proposed. A recursive formula is given accordingly. Symmetric repunits are also defined.

**Keywords** Repunits

**DOI:** 10.5281/zenodo.2639620

Written in Turin, on April 14, 2019

As explained in [1], the term “repunit” was coined by Beiler in a book of 1966 [2], for the numbers defined as:

$$R_n = \frac{10^n - 1}{10 - 1}$$

The sequence of repunits starts with 1, 11, 111, 1111, 11111, 111111, ... (sequence A002275 in the OEIS, <https://oeis.org/A002275>). As we can easily see, these numbers are linked to q-integers and Mersenne numbers [3-7]. A q-integer is defined as [3]:

$$[n] = \frac{q^n - 1}{q - 1}$$

so we have the Mersenne numbers for  $q=2$ . The repunits are the q-integers for  $q=10$  :

$$[n]_{q=10} = \frac{10^n - 1}{10 - 1}$$

We can use the same approach for the repunits of that proposed in [4-6]. Let us consider the following operation (generalized sum):

$$R_{m+n} = R_m \oplus R_n$$

defined in the following manner:

$$(1) \quad R_m \oplus R_n = R_m + R_n + (10 - 1) R_m R_n$$

This is the addition of the q-units as given in [4,5]. The neutral element for (1) is  $R_0 = 0$ , so that:  $R_m \oplus R_0 = R_m + R_0 + (10 - 1) R_m R_0 = R_m$ .

The recursive relation for the repunits, given according to (1) and starting from  $R_1 = 1$ , is:

$$R_m \oplus R_1 = R_m + R_1 + (10 - 1) R_m R_1$$

That is: 11, 111, 1111, 11111, 111111, 1111111, 11111111, and so on.

In [8], we have discussed the symmetric q-integers, which are defined as [3]:

$$[n]_s = \frac{q^n - q^{-n}}{q - q^{-1}}$$

We can define the “symmetric” repunits as:

$$R_{n,s} = \frac{10^n - 10^{-n}}{10 - 10^{-1}} = 2 \frac{\sinh(n \ln 10)}{10 - 10^{-1}}$$

The sequence is: 1, 10.1, 101.01, 1010.101, 10101.0101, 101010.10101, etc.

In this case, the addition is defined [8]:

$$R_{m,s} \oplus R_{n,s} = R_{m,s} \cosh(n \ln 10) + R_{n,s} \cosh(m \ln 10)$$

or

$$R_{m,s} \oplus R_{n,s} = R_{m,s} \sqrt{1 + k^2 (R_{n,s})^2} + R_{n,s} \sqrt{1 + k^2 (R_{m,s})^2}$$

where  $k = \frac{1}{2} \left(10 - \frac{1}{10}\right)$ . Let us note that  $R_{1,s} = \frac{10 - 10^{-1}}{10 - 10^{-1}} = 1$ .

The recursive formula for the symmetric repunits is:

$$R_{n+1,s} = R_{n,s} \oplus R_{1,s} = R_{n,s} \sqrt{1 + k^2} + \sqrt{1 + k^2 (R_{n,s})^2}$$

## References

- [1] Weisstein, Eric W. "Repunit." From MathWorld--A Wolfram Web Resource.  
<http://mathworld.wolfram.com/Repunit.html>
- [2] Beiler, A. H. (1966). "11111...111." Ch. 11 in *Recreations in the Theory of Numbers: The Queen of Mathematics Entertains*. New York: Dover.
- [3] Kac, V., & Cheung, P. (2001). *Quantum calculus*. Springer Science & Business Media.
- [4] Sparavigna, A. C. (2018). The q-integers and the Mersenne numbers. Zenodo.  
<http://doi.org/10.5281/zenodo.1251833>
- [5] Sparavigna, A. C. (2018). ON THE ADDITIVE GROUP OF q-INTEGERS. Zenodo . DOI:  
10.5281/zenodo.1245849
- [6] Sparavigna, A. C. (2018). On a generalized sum of the Mersenne Numbers. Zenodo.  
<http://doi.org/10.5281/zenodo.1250048>
- [7] Weisstein, Eric W. "Mersenne Number." From MathWorld--A Wolfram Web Resource.  
<http://mathworld.wolfram.com/MersenneNumber.html>
- [8] Sparavigna, A. C. (2018). On the generalized sum of the symmetric q-integers. Zenodo.  
DOI: 10.5281/zenodo.1248959