

TENOGENIC DIFFERENTIATION PROTOCOL IN XENOGENIC-FREE MEDIA ENHANCES TENDON-RELATED MARKER EXPRESSION IN ADIPOSE-DERIVED STEM CELLS

Original

TENOGENIC DIFFERENTIATION PROTOCOL IN XENOGENIC-FREE MEDIA ENHANCES TENDON-RELATED MARKER EXPRESSION IN ADIPOSE-DERIVED STEM CELLS / Stanco, D.; Soldati, G.; Ciardelli, G.. - ELETTRONICO. - Vol. 100-B, No. SUPP_15:(2018), pp. 128-128. (Intervento presentato al convegno The European Orthopaedic Research Society (EORS) 2018 Meeting tenutosi a Galway, Ireland nel September 2018).

Availability:

This version is available at: 11583/2747792 since: 2019-08-19T09:58:14Z

Publisher:

Bone & Joint Publishing

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

TENOGENIC DIFFERENTIATION PROTOCOL IN XENOGENIC-FREE MEDIA ENHANCES TENDON-RELATED MARKER EXPRESSION IN ADIPOSE-DERIVED STEM CELLS

D. Stanco, G. Soldati, G. Ciardelli

Published Online: 19 Nov 2018

Tendon injuries are common and current therapies often are unsuccessful. Cell-based therapy using mesenchymal stem cells (MSCs) seems to be the most promising approach to heal tendon. Moreover, providing safe and regulated cell therapy products to patients requires adherence to good manufacturing practices (GMP). Adipose-derived stem cells (n=4) were cultured in 6-well plates coated with type-I collagen in a chemically defined serum-free medium (SF) or a xenogenic-free human pooled platelet lysate medium (hPL). At passage 4, ASCs were induced to tendon lineage for 14 days using 100ng/ml CTGF, 10ng/ml TGF β 3, 50ng/ml BMP12 and 50 μ g/ml ascorbic acid in the SF (SF-TENO) or in the hPL (hPL-TENO) medium. Cells cultured without any supplements are used as control. Morphological appearance, cell viability and FACS were performed in undifferentiated cells to evaluate the xenogenic-free culture conditions; the gene and protein expression were performed by RT-PCR and immunofluorescence to evaluate to expression of stem cell- and tendon-related markers upon cell differentiation. SF-CTRL and hPL-CTRL showed similar viability and MSC's surface proteins and expressed the stemness markers NANOG, OCT4 and Ki67. Moreover, both SF-TENO and hPL-TENO expressed significant higher levels of SCX, COL1A1, COL3A1, COMP, MMP3 and MMP13 genes already at 3d (p<0.05) respect to CTRLs. Scleraxis and collagen were also detected in both SF-TENO and hPL-TENO at protein level in higher amount than CTRLs. In conclusion, ASCs exposed to CTGF, BMP12, TGF β 3 and AA in both serum and xenogenic-free media possess similar tenogenic differentiation ability moving forward the GMP-compliant approaches for the clinical use of ASCs.

Orthopaedic Proceedings contains abstracts of papers presented at recent scientific meetings organised by orthopaedic associations and specialist societies. *Orthopaedic Proceedings* only publishes the abstracts and does not publish full-text articles.

Re-using blood products as an alternative supplement in the optimisation of clinical-grade adipose-derived mesenchymal stem cell culture

J. Phetfong, *Bone & Joint Research*, 2017

Existence of mesenchymal stem cells in sites of atrophic nonunion

H. D. Ismail, *Bone & Joint Research*, 2013

Biological analysis of flexor tendon repair-failure stump tissue: a potential recycling of tissue for tendon regeneration

C-C. Lu et al., *Bone & Joint Research*, 2019

Angelica polysaccharide promotes proliferation and osteoblast differentiation of mesenchymal stem cells by regulation of long non-coding RNA H19

Xiaoyan Xie et al., *Bone & Joint Research*, 2019

Effects of platelet-rich plasma and triamcinolone acetonide on interleukin-1 β -stimulated human rotator cuff-derived cells

T. Muto, *Bone & Joint Research*, 2016

AB0094 DIFFERENTIATION OF ADIPOSE DERIVED MESENCHYMAL STEM CELLS OBTAINED FROM PATIENTS WITH SYSTEMIC LUPUS ERYTHEMATOSUS ANKYLOSING SPONDYLITIS AND SYSTEMIC SCLEROSIS

Ewa Kuca-Warnawin et al., *Ann Rheum Dis*, 2019

In vitro effect of direct current electrical stimulation on rat mesenchymal stem cells.

PeerJ

Preservation media, durations and cell concentrations of short-term storage affect key features of human adipose-derived mesenchymal stem cells for therapeutic application

PeerJ

P116 Basic characteristics of adipose-derived mesenchymal stem cells of ankylosing spondylitis patients

E Kuca-Warnawin et al., *Ann Rheum Dis*, 2019

New research: Nintedanib plus docetaxel after progression on immune checkpoint inhibitor therapy - a VARGADO study

Future Oncology

Powered by **TREND MD**

I consent to the use of Google Analytics and related cookies across the TrendMD network (widget, website, blog). [Learn more](#)

Yes

No



The British Editorial Society of Bone & Joint Surgery, 22 Buckingham Street, London WC2N 6ET +44 (0) 20 7782 0010

[Terms and conditions](#) [Privacy policy](#) [Refunds policy](#)

© 2019 The British Editorial Society of Bone & Joint Surgery. Registered charity no: 209299.

Powered by Atypon® Literatum