

Lipoplexes for miRNA delivery in myocardial regeneration

Original

Lipoplexes for miRNA delivery in myocardial regeneration / Nicoletti, Letizia; Tarricone, Giulia; Paoletti, Camilla; Divieto, Carla; Andreana, Ilaria; Stella, Barbara; Arpicco, Silvia; Mattu, Clara; Chiono, Valeria. - ELETTRONICO. - (2019), pp. 136-136. (Intervento presentato al convegno Nano-Day IV tenutosi a Milan nel December, 11-14).

Availability:

This version is available at: 11583/2776917 since: 2020-01-03T15:13:11Z

Publisher:

Cinsa and University of Parma

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Lipoplexes for miRNA delivery in myocardial regeneration

Letizia Nicoletti¹, Giulia Tarricone¹, Camilla Paoletti¹, Carla Divieto², Ilaria Andreana³,
Barbara Stella³, Silvia Arpicco³, Clara Mattu¹, Valeria Chiono^{1*}

¹*Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Corso Duca degli Abruzzi 24, Turin, Italy*

²*Division of Metrology for Quality of Life, Istituto Nazionale di Ricerca Metrologica, Strada delle Cacce 91, Turin, Italy*

³*Department of Drug Science and Technology, University of Turin, via P. Giuria 9, 10125, Turin, Italy*

**valeria.chiono@polito.it*

Myocardial infarction is one of the major causes of mortality in industrialized countries. Several strategies have been studied to regenerate myocardial infarction, reaching a limited success [1].

Direct reprogramming of cardiac fibroblasts populating post-infarct scar into cardiomyocytes could represent an alternative promising approach [1-3]. In 2012, mouse fibroblasts transfected with a combination of four microRNAs (miRNAs), termed “miRcombo”, were transdifferentiated into cardiomyocytes [1-3].

In this work, nanotechnology-based approaches were explored to efficiently deliver miRNAs to human cardiac fibroblasts in the perspective of their direct reprogramming. Lipoplexes containing negmiR or miR-1 (miRVana™ miRNA Mimic, Life Technologies) were prepared at different N:P ratios (3.0; 1.75; 0.70; 0.35). Lipoplexes showed an average size between 405 nm and 560 nm and an average zeta potential between +21 mV and -29 mV. Encapsulation efficiency and morphology (cryoTEM) were also analyzed.

In vitro tests with human cardiac fibroblasts showed a significantly higher miR-1 expression in the case of transfection with lipoplexes compared to a commercial agent, as evaluated by digital droplet PCR.

As a conclusion, in this work new lipoplexes were developed showing efficient delivery of miR-1 to human cardiac fibroblasts, for future use in direct reprogramming.

BIORECAR project is acknowledged: it has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme grant agreement No 772168

[1] C. Paoletti et al. Cells 2018;7: 114.

[2] T. M. Jayawardena et al. Circ. Res. 2012; 110: 1465.

[3] T. M. Jayawardena et al. Circ. Res. 2015; 116: 418.