

The protective role of forests to reduce rockfall risks and impacts in the Alps under a climate change perspective

Original

The protective role of forests to reduce rockfall risks and impacts in the Alps under a climate change perspective /
Lingua, Emanuele; Bettella, Francesco; Pividori, Mario; Marzano, Raffaella; Garbarino, Matteo; Piras, Marco; Kobal,
Milan; Berger, Frédéric - In: Climate Change, Hazards and Adaptation Options - Handling the Impacts of a Changing
ClimateSTAMPA. - [s.l.] : Springer International Publishing, 2020. - ISBN 978-3-030-37424-2. - pp. 333-347
[10.1007/978-3-030-37425-9_18]

Availability:

This version is available at: 11583/2780892 since: 2020-01-16T09:31:19Z

Publisher:

Springer International Publishing

Published

DOI:10.1007/978-3-030-37425-9_18

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Metadata of the chapter that will be visualized in SpringerLink

Book Title	Climate Change, Hazards and Adaptation Options	
Series Title		
Chapter Title	The Protective Role of Forests to Reduce Rockfall Risks and Impacts in the Alps Under a Climate Change Perspective	
Copyright Year	2020	
Copyright HolderName	Springer Nature Switzerland AG	
Corresponding Author	Family Name	Lingua
	Particle	
	Given Name	Emanuele
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	TESAF, University of Padova
	Address	Viale dell'Università, 16, 35020, Legnaro, Italy
	Email	emanuele.lingua@unipd.it
Author	Family Name	Bettella
	Particle	
	Given Name	Francesco
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	TESAF, University of Padova
	Address	Viale dell'Università, 16, 35020, Legnaro, Italy
	Email	francesco.bettella@unipd.it
Author	Family Name	Pividori
	Particle	
	Given Name	Mario
	Prefix	
	Suffix	
	Role	
	Division	
	Organization	TESAF, University of Padova
	Address	Viale dell'Università, 16, 35020, Legnaro, Italy
	Email	mario.pividori@unipd.it
Author	Family Name	Marzano
	Particle	
	Given Name	Raffaella
	Prefix	
	Suffix	

Role
Division
Organization DISAFA, University of Torino
Address L.go Braccini, 2, 10095, Grugliasco, Italy
Email raffaella.marzano@unito.it

Author Family Name **Garbarino**
Particle
Given Name **Matteo**
Prefix
Suffix
Role
Division
Organization DISAFA, University of Torino
Address L.go Braccini, 2, 10095, Grugliasco, Italy
Email matteo.garbarino@unito.it

Author Family Name **Piras**
Particle
Given Name **Marco**
Prefix
Suffix
Role
Division
Organization DIATI, Politecnico di Torino
Address C.so Duca degli Abruzzi, 24, 10129, Turin, Italy
Email marco.piras@polito.it

Author Family Name **Kobal**
Particle
Given Name **Milan**
Prefix
Suffix
Role
Division Department of Forestry and Renewable Forest Resources
Organization University of Ljubljana
Address Večna pot 83, 1000, Ljubljana, Slovenia
Email milan.kobal@bf.uni-lj.si

Author Family Name **Berger**
Particle
Given Name **Frédéric**
Prefix
Suffix
Role
Division Mountain Ecosystems Research Unit
Organization IRSTEA—National Research Institute of Science and Technology for Environment and Agriculture
Address rue de la Papeterie, 2, BP 76, 38402, Saint-Martin-d'Hères Cedex, France

Abstract

Among the functions provided by forests, protection has always been considered one of the preeminent in mountain areas. In order to fulfil, maximize, and sustain this function, specific forest structures should be obtained and maintained through properly designed forest management. A specific management goal should be defined with a well-defined forest target against each natural hazard, based on the protection potentially provided by the forest stands, in either an active (e.g. against avalanches) or passive way (e.g. against rockfall). Climate change is forecast to affect both disturbance regimes and forest ecosystems, leading to new challenging issues concerning protection forest management. This paper describes how a forest stand exerts its protective role against rockfalls and the target profile to be reached for sustaining this function. Potential consequences of climate change on forest ecosystems that management will have to face in the near future are also addressed. New perspectives are provided taking into account the knowledge coming from recent research studies and specifically the results obtained in the RockTheAlps project (ASP462), dealing with the assessment of protection forests against rockfall in the Alps.

Keywords
(separated by '-')

Protection forests - Alps - Rockfall - Forest management - Climate change

The Protective Role of Forests to Reduce Rockfall Risks and Impacts in the Alps Under a Climate Change Perspective



Emanuele Lingua, Francesco Bettella, Mario Pividori, Raffaella Marzano, Matteo Garbarino, Marco Piras, Milan Kobal and Frédéric Berger

1 **Abstract** Among the functions provided by forests, protection has always been
2 considered one of the preeminent in mountain areas. In order to fulfil, maximize, and
3 sustain this function, specific forest structures should be obtained and maintained
4 through properly designed forest management. A specific management goal should
5 be defined with a well-defined forest target against each natural hazard, based on the
6 protection potentially provided by the forest stands, in either an active (e.g. against
7 avalanches) or passive way (e.g. against rockfall). Climate change is forecast to
8 affect both disturbance regimes and forest ecosystems, leading to new challenging
9 issues concerning protection forest management. This paper describes how a forest
10 stand exerts its protective role against rockfalls and the target profile to be reached

E. Lingua (✉) · F. Bettella · M. Pividori
TESAF, University of Padova, Viale dell'Università, 16, 35020 Legnaro, Italy
e-mail: emanuele.lingua@unipd.it

F. Bettella
e-mail: francesco.bettella@unipd.it

M. Pividori
e-mail: mario.pividori@unipd.it

R. Marzano · M. Garbarino
DISAFA, University of Torino, L.go Braccini, 2, 10095 Grugliasco, Italy
e-mail: raffaella.marzano@unito.it

M. Garbarino
e-mail: matteo.garbarino@unito.it

M. Piras
DIATI, Politecnico di Torino, C.so Duca degli Abruzzi, 24, 10129 Turin, Italy
e-mail: marco.piras@polito.it

M. Kobal
Department of Forestry and Renewable Forest Resources, University of Ljubljana, Večna pot 83,
1000 Ljubljana, Slovenia
e-mail: milan.kobal@bf.uni-lj.si

F. Berger
Mountain Ecosystems Research Unit, IRSTEA—National Research Institute of Science and
Technology for Environment and Agriculture, rue de la Papeterie, 2, BP 76, 38402
Saint-Martin-d'Hères Cedex, France

© Springer Nature Switzerland AG 2020

W. Leal Filho et al. (eds.), *Climate Change, Hazards and Adaptation Options*,
Climate Change Management, https://doi.org/10.1007/978-3-030-37425-9_18

1

11 for sustaining this function. Potential consequences of climate change on forest
12 ecosystems that management will have to face in the near future are also addressed.
13 New perspectives are provided taking into account the knowledge coming from
14 recent research studies and specifically the results obtained in the RockTheAlps
15 project (ASP462), dealing with the assessment of protection forests against rockfall
16 in the Alps.

17 **Protection Forests**

18 In mountainous areas, forest stands have always played an important role in main-
19 taining valleys safe for living in and transiting. This importance is increasing, since
20 in the last decades settlements have also been built in remote mountain areas and
21 there is the need to access them throughout the year, mostly for tourism (Wehrli and
22 Dorren 2013; Lingua et al. 2017).

23 The protective role is considered among one of the most important functions pro-
24 vided by a forest stand. All forests, regardless of their location, can accomplish this,
25 for instance protecting the soil from surface erosion and taking part directly and indi-
26 rectly in the hydrological cycle. However, despite this general protective role, we do
27 not define all forests as protection forests. A protection forest is instead characterized
28 and defined due to the existence of a specific natural hazard against which it offers
29 protection in an active or passive way. It might also fulfil other functions, but its pro-
30 tective role is preeminent. Protection forests are particularly effective against some
31 types of abiotic disturbances, like snow avalanches, rockfalls, debris flows, shallow
32 landslides, surface erosion and floods. Since these are mostly gravity-driven haz-
33 ards, it is clear that the protective role provided by forests can become fundamental
34 in mountain areas, where steep slopes increase the risk of occurrence (Lingua et al.
35 2017). By definition, active protection is exerted when the forest helps to prevent the
36 occurrence of a natural hazard. Passive protection occurs instead in those situations
37 in which the presence of the forest contributes to mitigating the effects produced by
38 the disturbance. The distinction between these two kinds of protective roles mostly
39 depends on the characteristics of the considered process. For instance, forests play
40 an important protective role against snow avalanches in the starting zone, preventing
41 their release. For rocks falling down a slope, forests can instead reduce the speed and
42 number of rocks reaching the bottom of the slope.

43 We can further classify stands based on the presence of a specific object to be
44 protected into direct and indirect protection forests (Brang et al. 2006). A direct
45 protection forest is defined as such when it grows in close proximity to an endangered
46 asset to which it offers protection against natural hazards. This direct protection is
47 generally offered to people, buildings and any other infrastructure that might be
48 exposed to a specific hazard in a mountain area. The direct protective function is
49 usually provided over an area, which is limited in size and located below, and close
50 to the protection forest. A typical example of a direct protection forest is represented

51 by a stand located above a group of houses or a road, which are threatened by snow
52 avalanches.

53 The indirect protective function is instead exerted by forests, independently of
54 their exact location, simply by their presence at a broader scale (e.g. the landscape
55 level). This could be the case for stands in mountain catchments where they can
56 potentially reduce soil erosion and flooding at the closing section.

57 Almost any forest can offer some indirect protection, for instance through its
58 effect in intercepting precipitation or affecting the local climate. Only some forests
59 affected by gravitational hazards have a direct protective function.

60 The distinction between direct and indirect protection forests is particularly impor-
61 tant in the context of forest planning, specifically in the definition of intervention
62 priorities and management targets (Berger and Rey 2004).

63 A stand with a direct protective function should be permanently effective (Lingua
64 et al. 2017). This can be achieved if the stand has high resistance to natural hazards
65 and high persistency. The only way to maintain a stand in this efficiency window is
66 with active forest management.

67 All forests are anyway subject to stand dynamics, which may modify or limit
68 their protection effectiveness. Most of these dynamics are driven by the occurrence
69 of natural disturbances, which often coincide with the hazards they are meant to
70 prevent or mitigate.

71 Protection forests might be affected by different kinds of disturbances, including
72 wildfires, storms, snow break, bark beetle outbreaks, which can act at both small and
73 large scale. This last category of events can deeply alter the capacity of the stand to
74 maintain its protective function. The degree and temporal extent in the impairment
75 of the protective function produced after the disturbance is strongly related to the
76 severity and spatial extent of the disturbance itself, as well as the recovery process
77 undertaken naturally by the disturbed system.

78 At this stage, the quality and quantity of biological legacies (e.g. logs, root plates)
79 can affect both the recovery process, for instance favouring the establishment of
80 natural regeneration, and the residual protection function offered by the disturbed
81 stand (Lingua et al. 2017).

82 A forest can offer a certain degree of protection, being more or less effective in
83 its protective function, based on two main aspects: (1) the type and characteristics
84 of the natural hazards involved; (2) the main features of the stand, together with its
85 conditions when a damaging event occurs (Brang et al. 2008).

86 Concerning the disturbance regime, intensity and frequency of occurrence are the
87 two most influential attributes affecting the capacity of a forest stand to provide an
88 effective protective function.

89 In this context, proper management of protection forests should be oriented
90 towards the maintenance of forest structures with a high degree of resistance and
91 resilience to disturbances, as well as capable of providing effective protection to
92 people and structures at risk. To reach this objective may sometimes require making
93 compromises, since the characteristics needed to increase resistance to a certain dis-
94 turbance might not necessarily be those that maximise the protective function of the
95 same stand against some specific hazard. For instance, to decrease the probability

96 of a crown fire spreading in the forest canopy, we could decide to reduce the stand
97 density, but this management decision might consequently diminish the protection
98 effectiveness of that same forest against rockfall.

99 Structure and composition of protection forests should thus be designed based on
100 the type of disturbance that might affect the stand, the disturbance regime, the required
101 protection function and the possibility of guaranteeing stand renewal. Obtaining the
102 stand structures required to reach these management goals is a long-term process
103 requiring decades or even centuries.

104 **Managing Protection Forests Against Rockfall Hazard**

105 In natural forest stand dynamics, some stages will not provide a protective effect, since
106 trees are in turn too small, too sparse, or too few (Motta and Haudemand 2000). To be
107 effective, a protection forest should have defined characteristics, especially concern-
108 ing tree density and average stem size. In order to guarantee the protective function,
109 a certain structure (i.e., combination of tree density and average size) has therefore
110 to be achieved and maintained over time. As previously stated, appropriate silvicultural
111 management can guide a forest towards its highest protection effectiveness and
112 it should thus be designed and pursued to reach this aim.

113 Current management of protection forests adopts the term target profile to describe
114 the characteristics of the stand that we want to obtain through silvicultural interven-
115 tions, according to the natural hazards involved and local site conditions (Brang et al.
116 2006). A target profile thus describes the state of the forest that is expected to have
117 an effective protective action against natural hazards and that can be permanently
118 maintained with minimum effort.

119 Depending on the hazard type, different attributes of the forest have to be taken
120 into account in terms of their protective effect (Lingua et al. 2017). Considering snow
121 avalanches, for instance, the forest performs an active protective role in the starting
122 zone impeding the release of an avalanche. In this situation, the canopy cover is
123 the key parameter to consider since it plays the important role of snow interception
124 (Frehner et al. 2007). When dealing with a rockfall hazard, forests mainly act in
125 a passive way, limiting its impact by reducing the runout length. In this case, tree
126 density and size are the main stand parameters of interest.

127 A rockfall process is the movement of falling rocks and their interaction with
128 the environment (Frehner et al. 2007). On a slope characterized by rockfalls, three
129 distinct areas (that often overlap) can be found: the zone of origin, the transit zone
130 and the runout and deposition zone.

131 In the zone of origin, where rocks are released (generally corresponding to steep
132 slopes exceeding 30°), the forest does not play a relevant protective role, other than
133 general water regimentation and root holding functions. In some cases trees can
134 actually act as triggering factors. Roots can penetrate into cracks, increasing frost
135 wedging effects. Moreover, roots can produce acid exudates that, together with litter
136 (in the case of conifers), can corrode and weather rocks, and they can release rocks

137 they were previously holding when trees sway or are uprooted by strong winds.
138 Forest management in the zone of origin should aim to avoid the presence of unstable
139 trees potentially prone to uprooting, and perhaps even tall trees, since they are more
140 subjected to wind load (Table 1). If suitable for the site and species, coppicing can
141 be applied, to maintain stands at lower height.

142 When rocks start moving down the slope, they enter the transit zone. In this zone,
143 rocks can slide, roll and bounce (if the incline exceeds 35°). The potential contribution
144 of the forest in mitigating the impact of rockfall (considering blocks up to a size of
145 5 m^3) can be significant. Trees act as energy-dissipating elements, making falling
146 rocks lose part of their kinetic energy with each impact on stems, lying deadwood
147 logs and root plates. In addition to dissipating energy, the collision against a tree can
148 cause a falling rock to deviate its trajectory or even to stop (Fig. 1).

149 The diameter of the stems in relation to the size of the rocks strongly determines
150 the energy reduction efficiency. The larger the trees, the more effective they can be
151 in dissipating energy or stopping rocks. However, big trees usually belong to older
152 age classes, being more prone to falling as a consequence of senescence dynamics,
153 with subsequent short-lived protection effects.

154 A high density of trees is supposedly desirable since it increases the probability
155 for a falling rock to hit a stem. However, from an ecological point of view, a stand
156 with a high density of large trees is not sustainable. The self-thinning rule explains
157 why in the presence of a large number of stems, their average size will be smaller
158 compared to a stand with lower tree density where trees can grow bigger.

159 The desired target profile against rockfall can be obtained by silvicultural manage-
160 ment, focusing on prominent stand parameters such as basal area and mean diameter
161 at breast height (DBH; corresponding to the diameter at 1.30 m) or stem density.

162 It has been proven that, to be effective against rockfall, a protection forest should
163 have a minimal length of 250 m along the slope. Moreover, attention should be given
164 not to create gaps between stems larger than 40 m along the maximum slope. In this
165 short distance, rocks can in fact regain high speed, depleting all the protective role
166 of the upslope forest.

167 Rocks start decreasing their speed when the slope diminishes to less than 30° ;
168 they then stop rapidly if the slope reaches less than 25° . In this last area, known as
169 the runout and deposition zone, the forest can play an important role in reducing the
170 length of the rock path, contributing to slowing down the rocks in the same manner
171 as in the transit zone. Here, however, the forest stand can be even more effective,
172 since the rock energy (i.e. its speed) is already reducing because of the diminishing
173 slope steepness. Consequently, in the runout and deposition zone even small trees
174 can stop big rocks.

175 To increase the protection function against rockfall, silvicultural management
176 can also act on species composition, where possible, favouring the presence of
177 broadleaves (hardwood species), since they are usually more resistant than conifers.

178 The importance of the protective function of a forest stand against rockfall should
179 be considered looking at the residual hazard, considering the risk reduction (Dorren
180 and Berger 2007). The ability to dissipate the rocks kinetic energy and thus their
181 speed can effectively reduce the intensity of the hazard, leading, if the element

Table 1 Silvicultural guidelines for rockfall mitigation by forest stands. Modified from Berger et al. (2017), Frehner et al. (2007)

	Recommendations	Thresholds
Zone of origin (release zone)	Remove unstable trees (leverage effect due to the wind) at the top of cliffs or outcrops and in the release area	Coefficient of stability value (height/diameter at breast height = H/DBH) Conifers: $H/DBH \leq 65$ Broadleaves: $H/DBH \leq 80$
	Maintain a high basal area compatible with stand sustainability at the foot of the release area	
	Whenever possible, limit boulder's distance to the beginning of the stand	
	Promote broadleaved trees that are more resistant than conifers with equivalent diameter Maintain more than 30% of broadleaved trees among the largest trees. Depending on site conditions, a certain amount of conifers is needed for stand stability	
	Limit the size of gaps (same thresholds as for transit and runout zones)	
	Harvest trees leaving stumps with a min. height of 1.30 m or, if rockfall can occur, completely remove stump to ground-level (or screed in order to avoid a trampoline effect)	
	Fell/cut trees at an oblique angle to the slope leaving felled trees on the ground in a position from which they cannot be easily moved	
Transit and run out zones	If possible, increase the planimetric length of the forested slope	Recommended horizontal length of forested slope >200 m (ideal >250 m)
	Limit the size of gaps	Length of gap along the steepest slope: High forest <40 m Coppice <20 m, In all cases, recommended value (H = average height of trees): length $\leq 1.3H$ with a wooded strip below the gap >2H (recommended >4H)
	Promote broadleaved trees, which are more resistant than conifers with equivalent diameter Maintain more than 30% of broadleaved trees among the largest trees. Depending on the site conditions, a certain amount of conifers is needed for increasing the stand stability	
	Maintain an appropriate basal area for the efficient trees	In the transit zone: the basal area of trees with a DBH ≥ 15 cm is required to be ≥ 25 m ² /ha

(continued)

Table 1 (continued)

	Recommendations	Thresholds
		In the run out zone, the basal area of trees with a DBH ≥ 15 cm is required to be ≥ 20 m ² /ha
	Maintain an appropriate stem density for the efficient trees Maintain a high density in a band of 25 m on either side of a corridor	In all cases the stem density for trees with a diameter of ≥ 20 cm is required to be ≥ 350 stems/ha
	Remove unstable trees along corridors	Value of the coefficient of stability (height/diameter at breast height = H/DBH) Conifers: H/DBH ≤ 65 Broadleaves: H/DBH ≤ 80




Fig. 1 Rock stopped by trees in a protection forest (Auronzo di Cadore, BL, Italy). The stems show evidence of recent wounds produced by bouncing rocks

182 to be protected requires no marginal risk, to fewer and less impacting permanent
183 infrastructures being built. We should remember that a forest stand provides several
184 ecosystem services besides the protective function (i.e. habitat provision, recreation,
185 aesthetic value), while a rockfall net is only a passive protection.

186 Managing stands to perform a protective function requires accurate knowledge
187 of their spatial location. The availability of accurate maps with a good resolution is
188 thus highly desirable. When rock sources are known, the area affected by the rockfall
189 processes can be defined by modelling the path of the rocks. Rockfall trajectories
190 modelling is frequently implemented via simulation models with different spatial
191 dimensions (2D, 2.5D, or 3D), normally using topographic layers that are created in
192 most GIS programs (Volkwein et al. 2011; Pradhan and Fanos 2017). The reliability of
193 the results mostly depends on the input data; the accuracy of Digital Elevation Models
194 (DEM) is therefore crucial for a correct rockfall assessment (Žabota et al. 2019). The
195 availability of new remote sensing tools, particularly LiDAR (Light Detection and
196 Ranging), and platforms (i.e. UAV—Unmanned Aerial Vehicle), provides enhanced
197 information that renders the simulation closer to reality, at least concerning slope
198 profile. Forest parameters can be also extracted from remote sensing data with good
199 accuracy (Eysn et al. 2015), but the goodness of the results is still strongly affected
200 by the forest stand complexity (e.g. structural diversity, species composition).

201 In the Alps, several studies provided detailed maps of protection forests in single
202 municipalities or valleys (see for example Motta and Haudemand 2000), and some
203 maps are also available at the regional or country level. However, the coverage is not
204 complete and the methodologies and spatial scales adopted are not harmonized. The
205 project RockTheAlps (ASP462), carried out in the framework of the Interreg Alpine
206 Space (2014–2020), aimed at filling this gap, identifying and applying innovative
207 methodologies to detect and map rockfall risk and protection forests, providing deci-
208 sion makers and policy makers with harmonized information for the whole Alpine
209 Space. At the broader scale, a model (ROCK-EU) based on more than 10,000 real
210 cases collected from the Alps has been implemented to map the potential runout
211 zone (Fig. 2). The forests located in this area are thus considered protection forests.
212 In order to define protection forests with direct protective function, the presence of
213 endangered assets has been taken into account and added to the model. The effective-
214 ness of the identified protection forests will be assessed by means of TORRID
215 (Toolbox for assessing the protective effect of forests against rockfall and expressing
216 the protective role in a Risk Reduction InDex), a new toolbox developed for the entire
217 Alpine Space. The rockfall risk reduction provided by forest stands will be defined
218 and the gap between optimal characteristics and current situation will be identified,
219 guiding forest managers in the crucial phase of prioritization according to an adap-
220 tive management approach. The preliminary map of the protective function obtained
221 after the first version of the model shows that around 20% of forests in the Alpine
222 Space are potentially providing this important ecosystem service. The analyses have
223 been conducted on a 25 m DEM, since this is the resolution available for the entire
224 Alpine Space, but the methodologies applied are flexible and can easily be re-run as
225 soon as a better resolution becomes available for the whole area. Furthermore the
226 results of the simulation at 25 m spatial resolution have proven to be comparable




Fig. 2 Simulation of rockfall in the Colcuc study case (Colle Santa Lucia, BL, Italy). The number of passages (size 1.2 m^3 ; 1000 replication from each source cell) is shown for the scenario without forest (a) and with forest (b). The reduction in runout length provided by the protection forest is evident

227 with the results obtained at a finer scale and can therefore be used effectively at the
 228 regional level, where only a general overview of the potential risk is needed, and
 229 location accuracy is not the main purpose (Žabota et al. 2019).

230 Several case studies distributed over the Alps have been used to test the large-
 231 scale model locally, as well as to evaluate the economic value of the protective
 232 function (Bianchi et al. 2018). Some of the case studies were also selected among
 233 forest stands recently affected by high severity disturbances, which consequently
 234 altered their ability to sustain the provision of ecosystem services and particularly
 235 their potential protective function. In the western Alps, during autumn 2017, almost
 236 10,000 ha of forests were affected by large wildfires, and some of these stands were
 237 protection forests. In the post-disturbance silvicultural intervention plan, priorities
 238 were assigned to the protection forests characterized by high burn severity. The
 239 recent storm Vaia, in late October 2018, which affected more than 40,000 ha of
 240 forests in the central and eastern Italian Alps, resulted in more than 8.6 M cubic
 241 meters of windthrown trees. Several of these forest stands had provided, among
 242 other ecosystem services, a relevant protective function.

243 **Managing Protection Forests Under Climate Change**

244 Climate change is predicted to produce important consequences, both direct and
245 indirect, on forest ecosystems and on the disturbance regimes possibly affecting
246 them (Seidl et al. 2017; Seidl and Rammer 2017; Thom et al. 2017).

247 Forest ecosystems in mountain regions will be greatly affected by climate change.
248 Tree species compositions, forest cover and growth rate are among the parameters of
249 great importance for the fulfilment of the protective function that could be altered in
250 the mid- and long-term perspective (Albrich et al. 2018). Not only rising mean tem-
251 peratures and changes in precipitation patterns will shape future forests by selecting
252 species more adapted to the new conditions, but also the alteration of natural dis-
253 turbance regimes will have a crucial role. Disturbance regimes are forecast to be
254 altered by climate change in several parts of the world (Dale et al. 2001; Seidl et al.
255 2017). Both abiotic and biotic disturbance agents will be affected and their cumula-
256 tive impacts can result in unprecedented negative effects on forests (Temperli et al.
257 2013). Appropriate disturbance regimes are in equilibrium with species composition
258 and forest structure that are thus able to resist or persist after disturbances since they
259 are adapted to their occurrence with a specific intensity and return interval. Altering
260 the disturbance regime can instead lead to changes in species arrangement, and the
261 creation of degraded structures.

262 Extreme events have the potential to affect forests much more than gradual changes
263 in temperature (Lindner et al. 2014). Indeed natural disturbances are discrete events
264 that could produce sudden alterations to forests, while changes in climatic conditions
265 can increase stress resulting in a stand decline that can take several years before
266 causing tree death.

267 From the perspective of forest management, focusing specifically on protection
268 forests, there are two main topics to tackle: the non-permanence issue and species
269 fitness (Dyderski et al. 2019).

270 When a high severity disturbance affects a protection forest, forest cover will be
271 partially or totally removed, with consequences on the level of protection offered
272 by the disturbed forest, based on the residual structure and composition. In the case
273 of wildfires or insect outbreaks standing dead trees will dominate the landscape
274 immediately after the event; they will persist standing for a variable amount of time,
275 depending mostly on tree species and size (Marzano et al. 2012). In the case of a
276 windthrow there will instead be mostly logs and uprooted trees covering the ground.
277 The rise in temperatures will result in increasing fire risk, particularly in the driest
278 valleys of the Alps. Rather than the gradual increase in temperature, extreme and
279 prolonged droughts may increase the probability of large stand-replacing fires (Zum-
280 brunnen et al. 2009; Eelkin et al. 2013). Biotic agents can also be highly favoured
281 by the presence of trees stressed by water limitation, and even a moderate increase
282 in temperature or drought length has been found to potentially raise the risk of forest
283 pathogens outbreaks (Bentz et al. 2010). Natural disturbances are often interdepend-
284 ent, and the occurrence of one type can promote the occurrence of another one
285 and increase its intensity. Windthrows or severe droughts are generally followed by

286 insect outbreaks, and standing dead trees killed by pests and diseases can potentially
287 increase fire risk.

288 Forest regeneration will be the most susceptible stage. Mature trees show a sort
289 of biological inertia; they already have a well-developed and larger root system
290 compared to seedlings, and are therefore able to counteract increasing stresses up to
291 a certain threshold.

292 Concerning rockfall, one of the most probable consequences of climate change
293 will be an increase in occurrence (Berger et al. 2017). Although rockfall is an almost
294 unpredictable phenomenon, triggering factors are generally related to weather con-
295 ditions, such as freeze-thaw processes (frost wedging) and precipitations. A high
296 variation of temperature over a short period, and long duration and/or high intensity
297 of precipitations generally precede rockfall events. Indeed, some studies already indi-
298 cated an increase in rockfall occurrences for years with weather anomalies (Berger
299 et al. 2017). Climate change will not only directly affect rocks release, but also and
300 probably to a greater extent the forests that should mitigate this hazard.

301 Forest management should take into account climate change effect on forest
302 ecosystems, identifying and mitigating potential threats to resistance and resilience
303 traits of the stands.

304 Thinning can be a correct silvicultural treatment in order to enhance resistance to
305 increasing water stress. Reducing inter-tree competition and providing more growing
306 space for roots in search of water produced encouraging results also in post-drought
307 recovery (Hlásny et al. 2014). However, concerning rockfall mitigation, reducing tree
308 density is not always a good option, since we need to maintain the high probability
309 of impacts between rocks and stems to dissipate the energy, deviate the path, and
310 eventually stop the rolling rocks. In this case, it could be better to focus on manipu-
311 lating species composition towards the creation of a stand including a mixture more
312 adapted to the new conditions.

313 Given the expected alteration of disturbance regimes within mountain areas, par-
314 ticular attention should be paid to the assessment of the most suitable post-disturbance
315 management decisions. After a disturbance affecting a protection forest, management
316 practices to be adopted should be carefully considered. In the Alps the most common
317 post-disturbance management practice is salvage logging (i.e. the felling and removal
318 of affected trees), followed or not by plantation. This practice has proven to act as an
319 additional disturbance on the already disturbed environment, with several negative
320 effects on the ecosystem processes and services (Leverkus et al. 2018; Marzano et al.
321 2013). Furthermore, removing these deadwood elements from forests that had a pro-
322 tective function can further deplete the residual protection they can still offer after the
323 disturbance. Lying deadwood, especially if formed of large logs, greatly increases
324 the surface roughness of the forest floor, acting as additional obstacles for the rocks
325 (Fuhr et al. 2015). Nevertheless, its role is still generally underestimated, and not con-
326 sidered reliable. After a stand replacing disturbance, if the area is not salvaged, lying
327 logs can exert a protective function for a very long time, despite decaying processes.
328 The long-lasting and beneficial effect produced by the presence of large amounts of
329 lying deadwood resulting in greater ground roughness has already been proven after
330 past high-severity disturbance events. In forests affected by the storm Vivian in 1990,

331 for instance, very few follow-up rockfall events have occurred (Wohlgemuth et al.
332 2017).

333 Moreover, deadwood elements can result in enhanced microsite conditions faci-
334 litating regeneration establishment and survival, particularly in sites affected by severe
335 disturbances, where the disturbance might have exacerbated already harsh conditions
336 (Marzano et al. 2013; Leverkus et al. 2018). Preferential recruitment is in fact par-
337 ticularly evident in climatically stressed sites. Within areas characterized by high
338 insolation and low precipitation, fallen trees and uprooted stumps create microsites
339 where more shade and moisture are available for seedlings, stabilizing microclimatic
340 conditions (Beghin et al. 2010). The sheltering effects of deadwood elements directly
341 protecting tree regeneration against high radiation, high temperature and high tran-
342 spiration rates were found to be determinant in arid environments (Callaway 2007;
343 Marzano et al. 2013). These nurse objects can also act as traps for wind-dispersed
344 seeds. In areas with low winter temperatures, the beneficial effect of deadwood
345 material could also result in maintaining higher soil temperatures during the night,
346 positively affecting winter seedling survival, as found by Castro et al. (2011). At the
347 same time, similar results can be obtained in cold environments where deadwood
348 protects seedlings from snow gliding and favours snow melting, locally increasing
349 the length of the growing season.

350 Leaving standing and lying deadwood after a disturbance can thus produce a dou-
351 ble positive effect in the management of protection forests, providing both enhanced
352 microsites for regeneration, and eco-engineering structures. Even if deadwood under-
353 goes natural decaying processes, reducing its protective effects over time, during its
354 permanence standing or lying on the ground it may still perform an important func-
355 tion for the time lapse required by the natural or planted regeneration to establish
356 (Wohlgemuth et al. 2017).

357 With specific reference to rockfall activities, it could be even more appropriate to
358 discuss global change rather than just climate change. Including land-use and land
359 cover changes will in fact provide a better overview on the foreseen scenarios for this
360 natural hazard (Lopez-Saez et al. 2016) for the Alpine area. Since temperature is ris-
361 ing and land use is shifting from crops or pastures to forests due to the abandonment
362 of marginal lands, forest cover in mountain areas will increase, theoretically provid-
363 ing increased protection against rockfall propagations (Berger et al. 2017). Forests
364 will potentially cover more land at the upper elevations, tree growth will increase,
365 broadleaves will gain a higher share. The increase in lengths of forested slopes,
366 basal area, and broadleaves percentage are all consequences leading to a generic
367 increase in the protective effect against rockfall. In two French Alpine departments
368 (Haute-Savoie and Isère), this expansion of protection forests has been forecast to
369 reach around 20% of the current area (Berger et al. 2017). However, rockfall is a
370 site-specific phenomenon, involving local lithological, geomorphological, climate
371 and forest issues, so we cannot excessively generalize the possible effects of global
372 change.

Conclusions

The management of protection forests under climate change is a challenging task. Unfortunately, there is no “one size fits all” management approach, but it is advisable to adopt a site and case specific tailor-made solution. Forests in the Alps will probably grow better and increase their range in the next decades, but uncertainty related to natural disturbances calls for a more careful assessment of their potential protective role. Since site conditions are changing, and therefore so are species performances, we should also change the forest management approach. Adaptive management, considering both climate and land-use changes, should be promoted. The inclusion of natural hazards and protection function assessment in forest management plans should be supported by guidelines providing factual information derived from scientific knowledge acquired in the recent research projects. Specific attention should be paid to finding evidence from research studies focusing on monitoring protection forests affected by severe natural disturbances in order to define proper post-disturbance management considering their preeminent protection function.

Acknowledgements This work was funded by the European Commission (project ASP 462 RockTheAlps) within the European Territorial Cooperation program “Alpine Space”.

References

- Albrich K, Rammer W, Thom D, Seidl R (2018) Trade-offs between temporal stability and level of forest ecosystem services provisioning under climate change. *Ecol Appl* 28:1884–1896
- Beghin R, Lingua E, Garbarino M, Lonati M, Bovio G, Motta R, Marzano R (2010) *Pinus sylvestris* forest regeneration under different post-fire restoration practices in the northwestern Italian Alps. *Ecol Eng* 36:1365–1372
- Bentz BJ, Régnière J, Fettig CJ, Hansen EM, Hayes JL, Hicke JA, Kelsey RG, Negron JF, Seybold SJ (2010) Climate change and bark beetles of the western United States and Canada: direct and indirect effects. *Bioscience* 60:602–613
- Berger F, Dorren L, Kleemayr K, Maier B, Planinsek S, Bigot C, Bourrier F, Jancke O, Toe D, Cerbu G (2017). Eco-engineering and protection forests against rockfalls and snow avalanches. Management strategies to adapt alpine space forests to climate change risk, pp 191–210
- Berger F, Rey F (2004) Mountain protection forests against natural hazards and risks: new French developments by integrating forests in risk zoning. *Nat Hazards* 33:395–404
- Bianchi E, Accastello C, Trappmann D, Blanc S, Brun F (2018) The economic evaluation of forest protection service against rockfall: a review of experiences and approaches. *Ecol Econ* 154:409–418
- Brang P, Schönenberger W, Frehner M, Schwitter R, Thormann JJ, Wasser B (2006) Management of protection forests in the European Alps: an overview. *For Snow Landsc Res* 80:23–24
- Brang P, Schönenberger W, Ott E, Gardiner B (2008) Forests as protection from natural hazards. In: Evans J (ed) *The forest handbook*, vol II. Blackwell Science, London, pp 53–81
- Callaway RM (2007) Positive interactions and interdependence in plant communities. Springer, Dordrecht, The Netherlands

- 413 Castro J, Allen CD, Molina-Morales M, Marañón-Jiménez S, Sánchez-Miranda Á, Zamora R (2011)
 414 Salvage logging versus the use of burnt wood as a nurse object to promote post-fire tree seedling
 415 establishment. *Restor Ecol* 19:537–544
- 416 Dale VH, Joyce LA, McNulty S, Neilson RP, Ayres MP, Flannigan MD, Hanson PJ, Irland LC,
 417 Lugo AE, Peterson CJ, Simberloff D, Swanson FJ, Stocks BJ, Wotton BM (2001) Climate change
 418 and forest disturbances. *Bioscience* 51:723–734
- 419 Dyderski MK, Paž S, Frelich LE, Jagodziński AM (2019) How much does climate change threaten
 420 European forest tree species distributions? *Glob Change Biol* 24:1150–1163
- 421 Dorren LKA, Berger F (2007) Principles of the tool Rockfor.net for quantifying the rockfall hazard
 422 below a protection forest. *Schweizerische Zeitschrift für Forstwesen* 6:157–165
- 423 Eelkin C, Gutierrez AG, Leuzinger S, Manusch C, Temperli C, Rasche L, Bugmann H (2013) A
 424 2 °C warmer world is not safe for ecosystem services in the European Alps. *Glob Change Biol*
 425 19:1827–1840
- 426 Eysn L, Hollaus M, Lindberg E, Berger F, Monnet JM, Dalponte M, Kobal M, Pellegrini M, Lingua
 427 E, Mongus D, Pfeifer N (2015) A benchmark of LiDAR-based single tree detection methods
 428 using heterogeneous forest data from the alpine space. *Forests* 6:1721–1747
- 429 Frehner M, Wasser B, Schwitter R (2007) Sustainability and success monitoring in protection
 430 forests. *Guidelines for Silvicultural Interventions in Forests with Protective Functions*, FOEN,
 431 Bern
- 432 Fuhr M, Bourrier F, Cordonnier T (2015) Protection against rockfall along a maturity gradient in
 433 mountain forests. *For Ecol Manage* 354:224–231
- 434 Hlásny T, Mátyás C, Seidl R, Kulla L, Merganičová K, Trombik J, Dobor L, Barcza Z, Konôpka
 435 B (2014) Climate change increases the drought risk in central European forests: what are the
 436 options for adaptation? *Lesn Časopis—For J* 60:5–18
- 437 Lindner M, Fitzgerald JB, Zimmermann NE, Reyher C, Delzon S, van der Maaten E, Schelhaas MJ,
 438 Lasch P, Eggers J, van der Maaten-Theunissen M, Suckow F, Psomas A, Poulter B, Hanewinkel
 439 M (2014) Climate change and European forests: what do we know, what are the uncertainties,
 440 and what are the implications for forest management? *J Environ Manage* 146:69–83
- 441 Lingua E, Marzano R, Berger F (2017) Protection forests and their management: an overview. *EFI*
 442 *Tech Rep* 101:115–122
- 443 Leverkus A, Rey Benayas J, Castro J, Boucher D, Brewer S, Collins B, Donato D, Fraver S, Kishchuk
 444 B, Lee E-J, Lindenmayer DB, Lingua E, Macdonald E, Marzano R, Rhoades CC, Royo A, Thorn
 445 S, Wagenbrenner JW, Waldron K, Wohlgemuth T, Gustafsson L (2018) Salvage logging effects
 446 on regulating and supporting ecosystem services—a systematic map. *Can J For Res* 48:983–1000
- 447 Lopez-Saez J, Corona C, Eckert N, Stoffel M, Bourrier F, Berger F (2016) Impacts of land-use and
 448 land-cover changes on rockfall propagation: insights from the Grenoble conurbation. *Sci Total*
 449 *Environ* 547:345–355
- 450 Marzano R, Lingua E, Garbarino M (2012) Post-fire effects and short-term regeneration dynamics
 451 following high-severity crown fires in a Mediterranean forest. *iForest* 5:93–100
- 452 Marzano R, Garbarino M, Marcolin E, Pividori M, Lingua E (2013) Deadwood anisotropic faci-
 453 litation on seedling establishment after a stand-replacing wildfire in Aosta valley (NW Italy). *Ecol*
 454 *Eng* 51:117–122
- 455 Motta R, Haudemand JC (2000) Protective forests and silvicultural stability. An example of planning
 456 in the Aosta valley. *Mt Res Dev* 20:180–187
- 457 Pradhan B, Fanos AM (2017). Rockfall hazard assessment: an overview. in laser scanning
 458 applications in landslide assessment. Springer, pp 299–322
- 459 Seidl R, Rammer W (2017) Climate change amplifies the interactions between wind and bark beetle
 460 disturbances in forest landscapes. *Landscape Ecol* 32:1485–1498
- 461 Seidl R, Thom D, Kautz M, Martin-Benito D, Peltoniemi M, Vacchiano G, Wild J, Ascoli D, Petr
 462 M, Honkaniemi J, Lexer MJ, Trotsiuk V, Mairota P, Svoboda M, Fabrika M, Nagel TA, Reyher
 463 CPO (2017) Forest disturbances under climate change. *Nat Clim Chang* 7:395–402
- 464 Temperli C, Bugmann H, Elkin C (2013) Cross-scale interactions among bark beetles, climate
 465 change, and wind disturbances: a landscape modeling approach. *Ecol Monogr* 83:383–402

- 466 Thom D, Rammer W, Seidl R (2017) The impact of future forest dynamics on climate: interactive
467 effects of changing vegetation and disturbance regimes. *Ecol Monogr* 87:665–684
- 468 Volkwein A, Schellenberg K, Labiouse V, Agliardi F, Berger F, Bourrier F, Dorren LKA, Gerber W,
469 Jaboyedoff M (2011) Rockfall characterisation and structural protection—a review. *Nat Hazards*
470 *Earth Syst Sci* 11:2617–2651
- 471 Wehrli A, Dorren L (2013) Protection forests: a key factor in integrated risk management in the
472 Alps. In: Renaud FG, Sudmeier-Rieux K, Estrella M (eds) *The role of ecosystems in disaster risk*
473 *reduction*. UNU Press, Tokyo, pp 321–342
- 474 Wohlgemuth T, Schwitter R, Bebi P, Sutter F, Brang P (2017) Post-windthrow management in
475 protection forests of the Swiss Alps. *Eur J Forest Res* 136:1029–1040
- 476 Žabota B, Repe B, Kobal (2019). Influence of digital elevation model resolution on rockfall
477 modelling. *Geomorphology* 328: 183–195
- 478 Zumbrunnen T, Bugmann H, Conedera M, Burgi M (2009) Linking forest fire regimes and climate-a
479 historical analysis in a dry inner alpine valley. *Ecosystems* 12:73–86

Author Queries

Chapter 18

Query Refs.	Details Required	Author's response
AQ1	Kindly note that, reference 'Dyderski et al. (2018)' has been changed to 'Dyderski et al. (2019)', so that this citation match the list.	

UNCORRECTED PROOF

MARKED PROOF

Please correct and return this set

Please use the proof correction marks shown below for all alterations and corrections. If you wish to return your proof by fax you should ensure that all amendments are written clearly in dark ink and are made well within the page margins.

<i>Instruction to printer</i>	<i>Textual mark</i>	<i>Marginal mark</i>
Leave unchanged	... under matter to remain	Ⓟ
Insert in text the matter indicated in the margin	∧	New matter followed by ∧ or ∧ [Ⓢ]
Delete	/ through single character, rule or underline or ┌───┐ through all characters to be deleted	Ⓞ or Ⓞ [Ⓢ]
Substitute character or substitute part of one or more word(s)	/ through letter or ┌───┐ through characters	new character / or new characters /
Change to italics	— under matter to be changed	↙
Change to capitals	≡ under matter to be changed	≡
Change to small capitals	≡ under matter to be changed	≡
Change to bold type	~ under matter to be changed	~
Change to bold italic	≈ under matter to be changed	≈
Change to lower case	Encircle matter to be changed	≡
Change italic to upright type	(As above)	⊕
Change bold to non-bold type	(As above)	⊖
Insert 'superior' character	/ through character or ∧ where required	Υ or Υ under character e.g. Υ or Υ
Insert 'inferior' character	(As above)	∧ over character e.g. ∧
Insert full stop	(As above)	⊙
Insert comma	(As above)	,
Insert single quotation marks	(As above)	ʹ or ʸ and/or ʹ or ʸ
Insert double quotation marks	(As above)	“ or ” and/or ” or ”
Insert hyphen	(As above)	⊞
Start new paragraph	┌	┌
No new paragraph	┐	┐
Transpose	└┐	└┐
Close up	linking ○ characters	Ⓞ
Insert or substitute space between characters or words	/ through character or ∧ where required	Υ
Reduce space between characters or words		↑