



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Direct reprogramming of human cardiac fibroblasts to cardiomyocytes using microRNA mimics

Original

Direct reprogramming of human cardiac fibroblasts to cardiomyocytes using microRNA mimics / PAOLETTI, CAMILLA; DIVIETO, CARLA; DI MEGLIO, FRANCA; NURZYNSKA, DARIA; CHIONO, VALERIA. - (2019). ((Intervento presentato al convegno JRC Summer School 2019: Non-Animal Approaches in Science. Challenges & Future Directions.

Availability:

This version is available at: 11583/2798494 since: 2020-02-28T12:27:52Z

Publisher:

European Commission, Joint Research Center (JRC) Ispra

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Poster Abstract

JRC Summer School on Non-Animal Approaches in Science – Challenges and Future Directions

21-24 May 2019, Ispra (IT)

Contact details	
Name	Camilla
Surname	Paoletti
Affiliation	Politecnico di Torino, Dep. of Mechanical and Aerospace Engineering
Email address	camilla.paoletti@polito.it

Abstract details	
Title	Direct reprogramming of human cardiac fibroblasts to cardiomyocytes using microRNA mimics
Authors	C. Paoletti ¹ , C. Divieto ² , F. Di Meglio ³ , D. Nurzynska ³ , V. Chiono ¹
Author's affiliations	¹ Department of Mechanical and Aerospace Engineering, Politecnico di Torino, Turin, Italy, ² INRIM, Metrology for Life Quality, Turin, Italy, ³ Department of Public Health, University of Naples "Federico II", Naples, Italy.
Keywords	Cardiomyocytes, Myocardial infarction, microRNAs, cardiac fibroblasts

The combination of four different microRNAs (miR-1, 133, 208 and 499), named "miRcombo", has been used for the direct reprogramming of murine fibroblasts into cardiomyocytes (CMs) for myocardial infarction (MI) treatments.[1,2] Here, we evaluated miRcombo mediated reprogramming of human adult cardiac fibroblasts (AHCfs) into CMs in 2D and 3D culture. After 4 days in 2D culture, ddPCR analysis showed significantly increased expression of early cardiac transcription factors (TFs) Hand2 and Mef2c ($p < 0.005$), slightly increased expression of Tbx5 and Nkx2.5 although non-significant ($p > 0.05$), and reduced Vimentin expression ($p < 0.05$) in miRcombo-transfected AHCfs compared to controls. ICC analysis showed increased expression of late cardiac markers α -sarcomeric actinin and cTnT in miRcombo-transfected AHCfs after 10 and 20 days of culture in 2D. However, ddPCR showed no significant differences of late cardiac markers Myh6 and cTnI expression between the groups after 15 days in 2D culture. On the other hand, non-transfected AHCfs cultured in 3D fibrin-based hydrogels showed enhanced cardiac TFs expression compared to 2D cultures after 4 days, while, miRcombo transfection did not significantly increase cardiac gene expression of AHCfs cultured in 3D hydrogels for 4 days. After 15 days, AHCfs cultured in 3D hydrogels showed a strongly enhanced expression of cardiac genes such as cTnI and Myh6 compared to 2D cultures. In conclusion, results showed that a 3D environment plays a key role in enhancing direct reprogramming of AHCfs into CMs.

References:

- [1] Jayawardena TM et al. *Circ Res.* **2013**, *110*, 1465–1473
- [2] Li Y et al. *Sci. Rep.* **2016**, *6*, 1–11

This project has received funding from the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation programme (grant agreement No 772168).