

On the notion of weak isometry for finite metric spaces

Original

On the notion of weak isometry for finite metric spaces / DE GREGORIO, Alessandro; Fugacci, Ulderico; Memoli, Facundo; Vaccarino, Francesco. - ELETTRONICO. - (2020).

Availability:

This version is available at: 11583/2821495 since: 2020-05-09T15:51:47Z

Publisher:

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

On the notion of weak isometry for finite metric spaces

Alessandro De Gregorio^{*1}, Ulderico Fugacci^{†1}, Facundo Memoli^{‡2},
and Francesco Vaccarino^{§1}

¹Department of Mathematical Sciences, Politecnico di Torino

²Department of Mathematics and Department of Computer
Science and Engineering, The Ohio State University

May 2020

Abstract

Finite metric spaces are the object of study in many data analysis problems. We examine the concept of *weak isometry* between finite metric spaces, in order to analyse properties of the spaces that are invariant under strictly increasing rescaling of the distance functions. In this paper, we analyse some of the possible complete and incomplete invariants for weak isometry and we introduce a dissimilarity measure that asses how far two spaces are from being weakly isometric. Furthermore, we compare these ideas with the theory of persistent homology, to study how the two are related.

1 Introduction

Finite metric spaces arise in many applicative problems, whenever we have a set of objects on which it is defined a measure of dissimilarity. One of the main purposes of Topological Data Analysis (TDA) is to gather topological and geometric information from these datasets [4], in order to be able to perform comparisons between the phenomena that generated the measurements. The main input, in the TDA pipeline, is a nested sequence of simplicial complexes called *filtration*. Such a sequence is usually obtained from a weighted undirected network, or, in case the weights of the network satisfy the triangular inequality, from a finite metric space. In many problems, it is interesting to obtain

^{*}alessandro.degregorio@polito.it

[†]ulderico.fugacci@polito.it

[‡]memoli@math.osu.edu

[§]francesco.vaccarino@polito.it

information which is invariant under “rescalings” of the metric space. If, for example, the distances are physical quantities, we may want that the results that we get are independent from the system used for measurements. Sometimes, our observations undergo transformations that are not linear, but that preserve the order of distance between points. For example, in [11], Giusti et al. showed that such an approach is useful to study data from neural activity and connectivity. In the recent theory of monoid equivariant operators [5], an interesting family of operators is that of the so called *change of units*, that are nothing but functions that transform a dataset via a *rescaling* of the observations. In this work, we want to give a definition of weak isometry that let us identify two finite metric spaces if one of the distance functions can be obtained from the other by mean of a composition with a strictly increasing real valued function. We compare our approach to the work of Ganyushkin and Tsvirkunov [10], where they perform a classification of finite metric spaces and introduce a notion of isomorphism between them. We will show that the problem of determining if two metric spaces are weakly isometric or not can be reduced to the classical problem of isometry. Then, we will see how curvature sets introduced by Gromov [12] and Vietoris-Rips filtration can serve as complete and incomplete invariants for weak isometry.

2 Weakly isometric finite metric spaces

Henceforth we will consider all the metric spaces taken into account to be finite. We will write FMS for finite metric space. Whenever we write \mathbb{R}^+ we are considering the set $\{x \in \mathbb{R} \mid x \geq 0\}$.

Definition 1 (weak isometry). Let us consider two metric spaces (X, d_X) and (Y, d_Y) . We say that they are weakly isometric if there exist a bijection $\varphi : X \longrightarrow Y$ and a strictly increasing function $\psi : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ such that, for all $x_1, x_2 \in X$,

$$\psi(d_X(x_1, x_2)) = d_Y(\varphi(x_1), \varphi(x_2)). \quad (1)$$

If (X, d_X) and (Y, d_Y) are weakly isometric, we will write

$$(X, d_X) \cong^w (Y, d_Y).$$

This concept is also defined in [7], called by the authors *weak similarity*, where it is applied to semi-metric spaces of possibly infinite cardinality.

Theorem 1. *The relation of weak isometry is an equivalence relation.*

Proof. We check the three properties of equivalence relations.

- Reflexivity: using the identity, $X \cong^w X$ for all finite metric spaces X .
- Symmetry: if $X \cong^w Y$ we have a bijection $\varphi : X \longrightarrow Y$ and a strictly increasing function $\psi : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ with $\psi(d_X(x_1, x_2)) = d_Y(\varphi(x_1), \varphi(x_2))$ for all $x_1, x_2 \in X$. We have that ψ is an invertible function since it is

strictly monotone. For each $y_1, y_2 \in Y$ consider $x_1, x_2 \in X$ with $y_i = \varphi(x_i)$, where $i = 1, 2$. Then,

$$d_X(\varphi^{-1}(y_1), \varphi^{-1}(y_2)) = \psi^{-1}(d_Y(y_1, y_2)).$$

and so, by definition, $Y \cong^w X$.

- Transitivity: if $X \cong^w Y$ and $Y \cong^w Z$ consider the functions $\varphi_1, \varphi_2, \psi_1, \psi_2$ such that

$$\begin{aligned} \psi_1(d_X(x_1, x_2)) &= d_Y(\varphi_1(x_1), \varphi_1(x_2)) \quad \forall x_1, x_2 \in X, \\ \psi_2(d_Y(y_1, y_2)) &= d_Z(\varphi_2(y_1), \varphi_2(y_2)) \quad \forall y_1, y_2 \in Y. \end{aligned}$$

Then,

$$\psi_1(d_X(x_1, x_2)) = d_Y(\varphi_1(x_1), \varphi_1(x_2)) = \psi_2^{-1}(d_Z(\varphi_2(\varphi_1(x_1)), \varphi_2(\varphi_1(x_2))))$$

hence, considering the functions $\varphi_2 \circ \varphi_1$ and $\psi_2 \circ \psi_1$, we have $X \cong^w Z$. □

Ganyushkin and Tsvirkunov [10] introduced a notion of isomorphism for finite metric spaces which we will compare to weak isometry below.

Definition 2 (isomorphism - [10]). We say that two metric spaces (X, d_X) and (Y, d_Y) are isomorphic if there is a bijection $\varphi : X \rightarrow Y$ such that for all $x_1, x_2, x'_1, x'_2 \in X$ we have

$$d_X(x_1, x_2) = d_X(x'_1, x'_2) \Rightarrow d_Y(\varphi(x_1), \varphi(x_2)) = d_Y(\varphi(x'_1), \varphi(x'_2)) \quad (2)$$

$$d_X(x_1, x_2) < d_X(x'_1, x'_2) \Rightarrow d_Y(\varphi(x_1), \varphi(x_2)) < d_Y(\varphi(x'_1), \varphi(x'_2)). \quad (3)$$

If (X, d_X) and (Y, d_Y) are isomorphic we will write $(X, d_X) \simeq (Y, d_Y)$.

Now, we see how the two concepts are related.

Theorem 2. *Two finite metric spaces are weakly isometric if and only if they are isomorphic.*

Proof. Assume that $X \cong^w Y$. Then, we have two functions φ and ψ such that, for all $x_1, x_2, x'_1, x'_2 \in X$,

$$\psi(d_X(x_1, x_2)) = d_Y(\varphi(x_1), \varphi(x_2)), \quad (4)$$

$$\psi(d_X(x'_1, x'_2)) = d_Y(\varphi(x'_1), \varphi(x'_2)). \quad (5)$$

Then, if $d_X(x_1, x_2) = d_X(x'_1, x'_2)$, we have

$$d_Y(\varphi(x_1), \varphi(x_2)) = \psi(d_X(x_1, x_2)) = \psi(d_X(x'_1, x'_2)) = d_Y(\varphi(x'_1), \varphi(x'_2)). \quad (6)$$

If $d_X(x_1, x_2) < d_X(x'_1, x'_2)$, since ψ is a strictly increasing function,

$$d_Y(\varphi(x_1), \varphi(x_2)) = \psi(d_X(x_1, x_2)) < \psi(d_X(x'_1, x'_2)) = d_Y(\varphi(x'_1), \varphi(x'_2)). \quad (7)$$

Therefore, $X \cong^w Y \Rightarrow X \simeq Y$.

On the other hand, assume $X \simeq Y$ and consider the bijection φ given by Definition 2. It is possible to order all the pairs $(x_i, x_j) \in X \times X$ so that

$$d_X(x_{i_1}, x_{j_1}) \leq d_X(x_{i_2}, x_{j_2}) \leq \dots \leq d_X(x_{i_n}, x_{j_n}), \quad (8)$$

where n is the number of points of X and Y . Because of implications (2) and (3), we have that

$$d_Y(\varphi(x_{i_1}), \varphi(x_{j_1})) \leq d_Y(\varphi(x_{i_2}), \varphi(x_{j_2})) \leq \dots \leq d_Y(\varphi(x_{i_n}), \varphi(x_{j_n})), \quad (9)$$

hence, we can define an increasing function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ with

$$\psi(d_X(x_{i_r}, x_{j_r})) = d_Y(\varphi(x_{i_r}), \varphi(x_{j_r})) \quad \forall (x_{i_r}, x_{j_r}) \in X \times X \quad (10)$$

and then $X \simeq Y \Rightarrow X \cong^w Y$. \square

We have seen that the two concepts are the same, but weak isometry explicitly shows the rescaling that has to be performed to obtain one space from the other. Now, we want to associate to each equivalence class of weak isometry a good representative.

Definition 3 (distance set). Given a metric space (X, d) , we define its distance set $D(X, d)$ as the set of all pairwise distances between different points of X .

$$D(X, d) := \{d(x_1, x_2) \mid x_1, x_2 \in X, x_1 \neq x_2\}. \quad (11)$$

When there is no ambiguity for the metric d defined on the space X , we will simply write $D(X)$.

Lemma 1. *Two weakly isometric finite metric spaces have distance sets of the same cardinality.*

Proof. If $X \cong^w Y$, there is a strictly increasing function ψ such that

$$\psi(D(X)) := \{\psi(l) \mid l \in D(X)\} = D(Y). \quad (12)$$

So, since $\psi|_{D(X)}$ is injective by definition and surjective on $D(Y)$, then $|D(X)| = |D(Y)|$. \square

Remark 1. The reciprocal statement does not hold. Two spaces can have the same distance set but not be weakly isometric. For example, Boutin and Kemper in [1] study the problem of reconstruction of a metric space given the distribution of distances between points.

Example 1. The two metric spaces in Fig. 1, $X = \{a, b, c\}$ with $d_X(a, c) = d_X(b, c) = 6$, $d_X(a, b) = 5$ and $Y = \{d, e, f\}$ with $d_Y(d, f) = d_X(e, f) = 5$, $d_Y(d, e) = 6$ have the same distance set, but they are not weakly isometric.




Figure 1: FMS that are not weakly isometric.

Definition 4 (natural-valued metric space). We say that the metric d defined on the space X is natural-valued if $D(X) \subset \mathbb{N}$.

Definition 5 (dense distance set). Given a finite metric space (X, d) with natural-valued matrix d we say that its distance set is dense if it is a list of consecutive natural numbers,

$$D(X) = \{a + 1, a + 2, \dots, a + |D(X)|\}. \quad (13)$$

Definition 6 (canonical). A finite metric space of cardinality n is called k -canonical if it has a natural valued and dense distance set of the form

$$D_{n,k} := \left\{ 2\binom{n}{2}, 2\binom{n}{2} - 1, \dots, 2\binom{n}{2} - k + 1 \right\}. \quad (14)$$

Canonical metric spaces are interesting because for them the notion of weak isometry is equivalent to that of isometry, as we show in the next Lemma.

Lemma 2. *Let \mathcal{C}_1 and \mathcal{C}_2 be canonical finite metric spaces. Then, \mathcal{C}_1 is weakly isometric to \mathcal{C}_2 if and only if \mathcal{C}_1 is isometric to \mathcal{C}_2 .*

Proof. If \mathcal{C}_1 is isometric to \mathcal{C}_2 , then \mathcal{C}_1 is weakly isometric to \mathcal{C}_2 . On the other hand assume $\mathcal{C}_1 \cong^w \mathcal{C}_2$. By Lemma 1, we know that the two spaces have distance sets of the same cardinality. On the other hand, the distance set of a canonical space is defined by its cardinality, so

$$\begin{aligned} D(\mathcal{C}_1) &= \left\{ 2\binom{n}{2}, 2\binom{n}{2} - 1, \dots, 2\binom{n}{2} - |D(\mathcal{C}_1)| + 1 \right\} = \\ &= \left\{ 2\binom{n}{2}, 2\binom{n}{2} - 1, \dots, 2\binom{n}{2} - |D(\mathcal{C}_2)| + 1 \right\} = D(\mathcal{C}_2). \end{aligned} \quad (15)$$

By hypothesis, we have a bijection $\varphi : \mathcal{C}_1 \longrightarrow \mathcal{C}_2$ and a strictly increasing function $\psi : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ such that $\psi(D(\mathcal{C}_1)) = D(\mathcal{C}_2)$. Since the two distance sets are equal, such a function can only be the identity and, therefore,

$$\psi(d_{\mathcal{C}_1}(x_1, x_2)) = d_{\mathcal{C}_1}(x_1, x_2) = d_{\mathcal{C}_2}(\varphi(x_1), \varphi(x_2)) \quad (16)$$

and the two spaces are isometric. \square

In Proposition 2 of [10], the authors proved the following useful result.

Theorem 3. *Each finite metric space is isomorphic to a canonical metric space.*

Remark 2. We will say that a finite metric space \mathcal{C} is canonical for the FMS X if they are weakly isometric and \mathcal{C} is canonical.

We can now define a map called *canonicalization* which assigns to each finite metric space X a canonical metric space \mathcal{C}_X .

Definition 7 (canonicalization). Let X be a finite metric space of cardinality n and distance set $D(X) = \{a_1, a_2, \dots, a_k \mid a_i < a_j \text{ if } i < j\}$. Define $\psi : \mathbb{R}^+ \longrightarrow \mathbb{N}$ as

$$\psi(a_i) = 2 \binom{n}{2} - k + i. \quad (17)$$

The canonicalization of X is the space $(\mathcal{C}_X, d_{\mathcal{C}_X})$ with:

- $\mathcal{C}_X = X$
- $d_{\mathcal{C}_X}(x_1, x_2) = \begin{cases} \psi(d_X(x_1, x_2)) & \text{if } x_1 \neq x_2 \\ 0 & \text{if } x_1 = x_2. \end{cases}$

Canonical metric spaces are important because, as we will see with the following corollary, they allow us to associate to each weak isometry equivalence class a unique representative.

Corollary 1 (uniqueness of canonical representations). *Let X be a finite metric space with \mathcal{C} and \mathcal{C}' canonical for X . Then, \mathcal{C} and \mathcal{C}' are isometric, that is, there is a unique metric space canonical for X up to isometry.*

Proof. By the transitivity of weak isometry, we have that $\mathcal{C} \simeq X \simeq \mathcal{C}'$ and, by Lemma 2, they are isometric. \square

For a given FMS X , Corollary 1 allows us to identify a canonical representative of a X in the form of its canonicalization, as introduced in Definition 7.

In this way, we can now reformulate the problem of weak isometry to that of classical isometry between canonical spaces.

Theorem 4. *Two finite metric spaces are weakly isometric if and only if their canonicalizations are isometric.*

Proof. If \mathcal{C} is canonical for X and Y , then $X \cong^w \mathcal{C} \cong^w Y$ and, hence, X and Y are weakly isometric. On the other hand, assume that X is weakly isometric to Y . By the transitivity of weak isometry, we have

$$\mathcal{C}_X \cong^w X \cong^w Y \cong^w \mathcal{C}_Y,$$

therefore, \mathcal{C}_X and \mathcal{C}_Y are weakly isometric. By Lemma 2, we have that they have to be isometric. \square

3 A dissimilarity measure for weak isometry

We would like to define a pseudo-distance between finite metric spaces that measures how far are two metric spaces from being weakly isometric. We are looking for a pseudo-distance $d : \text{FMS} \times \text{FMS} \rightarrow \mathbb{R}$ such that:

1. $d((X, d_X), (Y, d_Y)) = 0$ if and only if (X, d_X) and (Y, d_Y) are weakly isometric,
2. d is “continuous”, that means that d is not discrete and takes into account the actual values assumed by the metrics d_X and d_Y , and does not discriminate them only because of their “combinatorial properties”.

Proposition 1. *There is no pseudo-distance that satisfies the two conditions above.*

Proof. Let us assume that the pseudo-distance d is such that it is zero if and only if two spaces are weakly isometric.

Since $d((X, d_X), (Y, d_Y)) = 0$ and d satisfies the triangle inequality, for any three spaces (X, d_X) , (Y, d_Y) and (Z, d_Z) with (X, d_X) weakly isometric to (Y, d_Y) we have that

$$d((X, d_X), (Z, d_Z)) = d((Y, d_Y), (Z, d_Z)). \quad (18)$$

Therefore, the function d is constant on the equivalence classes given by weak isometry, hence it is discrete. \square

Since it is not possible to find such a pseudo-distance, we will weaken our demands by dropping the assumption that the function d satisfies the triangle inequality.

Definition 8 (dissimilarity measure). A dissimilarity measure d over a set S is a function $d : S \times S \rightarrow \mathbb{R}$ such that:

1. there exists a number $d_0 \in \mathbb{R}$, with $-\infty < d_0 \leq d(s_1, s_2) < +\infty$, for all $s_1, s_2 \in S$, and $d(s, s) = d_0$, for all $s \in S$,
2. $d(s_1, s_2) = d(s_2, s_1)$, for all $s_1, s_2 \in S$.

Since we will make use of the Gromov-Hausdorff distance, we recall its definition.

Definition 9 (Gromov-Hausdorff distance). Given two metric spaces (X, d_X) and (Y, d_Y) , a *correspondence* between them is a set $C \subseteq X \times Y$ such that $\pi_X(C) = X$ and $\pi_Y(C) = Y$, where π_X and π_Y are the canonical projections of the product space. We denote with $\mathfrak{C}(X, Y)$ the set of all correspondences between X and Y . We define the *distortion* of a correspondence C , with respect to the metrics d_X and d_Y as

$$\text{dis}(C, d_X, d_Y) = \sup_{(x, y), (x', y') \in C} |d_X(x, x') - d_Y(y, y')|. \quad (19)$$

The Gromov-Hausdorff distance between (X, d_X) and (Y, d_Y) is

$$d_{GH}((X, d_X), (Y, d_Y)) = \frac{1}{2} \inf_{C \in \mathfrak{C}(X, Y)} \text{dis}(C, d_X, d_Y). \quad (20)$$

When (X, d_X) and (Y, d_Y) are finite metric spaces, the supremum in Eq. (19) is actually a maximum, and the infimum in Eq. (20) is a minimum. We refer the interested reader to [3] for further details.

Remark 3. From now on, we will denote by \mathcal{J} the set of strictly increasing functions $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $\psi(0) = 0$.

Consider now the map $\hat{d} : \text{FMS} \times \text{FMS} \rightarrow \mathbb{R}^+$ given by

$$\begin{aligned} \hat{d}((X, d_X), (Y, d_Y)) &= \inf_{\psi \in \mathcal{J}} d_{GH}((X, \psi \circ d_X), (Y, d_Y)) + \\ &\quad + \inf_{\psi \in \mathcal{J}} d_{GH}((X, d_X), (Y, \psi \circ d_Y)). \end{aligned} \quad (21)$$

We have the following result

Proposition 2. *The map \hat{d} is a dissimilarity on the collection of FMS.*

Proof. Since d_{GH} is a distance, for all (X, d_X) and (Y, d_Y) it holds

$$0 \leq \inf_{\psi \in \mathcal{J}} d_{GH}((X, \psi \circ d_X), (Y, d_Y)) < \infty$$

and

$$0 \leq \inf_{\psi \in \mathcal{J}} d_{GH}((X, d_X), (Y, \psi \circ d_Y)) < \infty.$$

Therefore there exists $d_0 = 0$ such that $d_0 \leq \hat{d}((X, d_X), (Y, d_Y)) < \infty$ for all (X, d_X) and (Y, d_Y) . It is also possible to notice that \hat{d} is symmetric by its very definition. Hence, \hat{d} is a dissimilarity. \square

Now, we show that that dissimilarity \hat{d} satisfies our initial requests, especially the fact that is 0 if and only if two spaces are weakly isometric.

Proposition 3. *Given two FMSs (X, d_X) , (Y, d_Y) , we have*

$$\hat{d}((X, d_X), (Y, d_Y)) = 0 \iff (X, d_X) \cong^w (Y, d_Y).$$

Proof. If $(X, d_X) \cong^w (Y, d_Y)$, there exists a strictly increasing function ψ such that $(X, \psi \circ d_X)$ is isometric to (Y, d_Y) and $(Y, \psi^{-1} \circ d_Y)$ is isometric to (X, d_X) . Since the Gromov-Hausdorff distance between two metric spaces is zero if and only if they are isometric, both the infima on the right hand side of Eq. (21) are 0, hence $\hat{d}((X, d_X), (Y, d_Y)) = 0$.

On the other hand, suppose that $\hat{d}((X, d_X), (Y, d_Y)) = 0$, so that

$$\inf_{\psi \in \mathcal{J}} d_{GH}((X, \psi \circ d_X), (Y, d_Y)) = 0 \text{ and } \inf_{\psi \in \mathcal{J}} d_{GH}((X, d_X), (Y, \psi \circ d_Y)) = 0.$$

Therefore, by the definition of infimum, there exist two sequences $(\psi_n)_{n \in \mathbb{N}} \subseteq \mathcal{J}$ and $(\tilde{\psi}_n)_{n \in \mathbb{N}} \subseteq \mathcal{J}$ such that

$$\begin{aligned} \lim_{n \rightarrow \infty} d_{GH}((X, \psi_n \circ d_X), (Y, d_Y)) &= 0 \\ \lim_{n \rightarrow \infty} d_{GH}((X, d_X), (Y, \tilde{\psi}_n \circ d_Y)) &= 0. \end{aligned} \tag{22}$$

Let us focus our attention on the first sequence, $(\psi_n)_{n \in \mathbb{N}} \subseteq \mathcal{J}$. By the finiteness of $\mathfrak{C}(X, Y)$, for every n in \mathbb{N} there exists a, possibly non-unique, correspondence R_n in $\mathfrak{C}(X, Y)$ such that $\text{dis}(R_n, \psi_n \circ d_X, d_Y) = d_{GH}((X, \psi_n \circ d_X), (Y, d_Y))$. By the axiom of choice, it is possible to construct a sequence $(\psi_n, R_n)_{n \in \mathbb{N}} \subseteq \mathcal{J} \times \mathfrak{C}(X, Y)$ such that

$$\lim_{n \rightarrow \infty} \text{dis}(R_n, \psi_n \circ d_X, d_Y) = 0.$$

The set $\mathfrak{C}(X, Y)$ is finite, henceforth we can find a subsequence $(\hat{\psi}_n, \hat{R}_n)_{n \in \mathbb{N}}$ of $(\psi_n, R_n)_{n \in \mathbb{N}}$ such that there exists a R_1 in $\mathfrak{C}(X, Y)$ and a \bar{n} in \mathbb{N} with $\hat{R}_n = R_1$, for all $n \geq \bar{n}$.

For this correspondence R_1 , it holds

$$\lim_{n \rightarrow \infty} |\hat{\psi}_n(d_X(x, x')) - d_Y(y, y')| = 0 \quad \forall (x, y), (x', y') \in R_1. \tag{23}$$

Hence, the restriction of the sequence $(\hat{\psi}_n)$ on the distance set $D(X)$ converges to a function $\psi_X : D(X) \rightarrow D(Y)$, such that $d_{GH}((X, \psi_X \circ d_X), (Y, d_Y)) = 0$. In the same way, we can prove the existence of a function $\psi_Y : D(Y) \rightarrow D(X)$ such that $d_{GH}((X, d_X), (Y, \psi_Y \circ d_Y)) = 0$.

We observe that

$$\begin{aligned} d_{GH}((X, \psi_Y \circ \psi_X \circ d_X), (X, d_X)) &\leq d_{GH}((X, \psi_Y \circ \psi_X \circ d_X), (Y, \psi_Y \circ d_Y)) + \\ &\quad + d_{GH}((Y, \psi_Y \circ d_Y), (X, d_X)). \end{aligned} \tag{24}$$

We already know that the second summand on the right hand side of the inequality is 0. For the first one it is easy to see that since $d_{GH}((X, \psi_X \circ d_X), (Y, d_Y)) = 0$, then also $d_{GH}((X, \psi_Y \circ \psi_X \circ d_X), (Y, \psi_Y \circ d_Y)) = 0$. Therefore, $d_{GH}((X, \psi_Y \circ \psi_X \circ d_X), (X, d_X)) = 0$, and $(X, \psi_Y \circ \psi_X \circ d_X)$ and (X, d_X) are isometric. Since ψ_X and ψ_Y are both non decreasing functions, also their

composition is non decreasing. Moreover, $\psi_Y \circ \psi_X$ has to be a bijective function from $D(X)$ to itself, otherwise the two spaces fail to be isometric. Then, it has to be $\psi_Y \circ \psi_X = \text{id}|_{D(X)}$, therefore ψ_X and ψ_Y are invertible. It is possible to extend, by linear interpolation, the domain and codomain of ψ_X to \mathbb{R}^+ , thus we have a strictly increasing function ψ_X such that $d_{GH}((X, \psi_X \circ d_X), (Y, d_Y)) = 0$ and this is equivalent to saying that (X, d_X) and (Y, d_Y) are weakly isometric. \square

Example 2. In this example, we show the computation of \tilde{d} between three finite metric spaces. Let us consider the spaces (X, d_X) , (Y, d_Y) and (Z, d_Z) depicted in Fig. 2. We can see that $\inf_{\psi \in \mathcal{S}} d_{GH}((X, \psi \circ d_X), (Y, d_Y)) = 0$. In fact, if we take a sequence $(\psi_n)_{n \in \mathbb{N}}$ such that

$$\psi_n(3) = 3, \psi_n(4) = 4, \psi_n(5) = 4 + \frac{1}{n},$$

clearly $\lim_{n \rightarrow \infty} d_{GH}((X, \psi_n \circ d_X), (Y, d_Y)) = 0$. On the other hand, for $\hat{\psi}$ with

$$\hat{\psi}(3) = 3, \hat{\psi}(4) = 4.5,$$

it holds $\inf_{\psi \in \mathcal{S}} d_{GH}((X, d_X), (Y, \psi \circ d_Y)) = d_{GH}((X, d_X), (Y, \hat{\psi} \circ d_Y)) = 0.5$. Therefore, $\tilde{d}((X, d_X), (Y, d_Y)) = 0.5$. Reasoning in a similar way it is possible to show that $\tilde{d}((Z, d_Z), (Y, d_Y)) = 1$. We also know by Proposition 3 that since X and Z are weakly isometric it has to be $\tilde{d}((X, d_X), (Z, d_Z)) = 0$. Hence,

$$\tilde{d}((Y, d_Y), (Z, d_Z)) = 1 > 0.5 = \tilde{d}((Y, d_Y), (X, d_X)) + \tilde{d}((X, d_X), (Z, d_Z))$$

and the triangular inequality does not hold.




Figure 2: Examples for the computation of \hat{d} .

4 Curvature sets of finite metric spaces

Establishing whether two spaces are isometric or not is in general a computationally intensive problem. Hence, we would like to have a set of complete or

incomplete invariants to study this problem. We will focus our attention firstly on the concept of curvature set introduced by Gromov in [12]. They have already been used by Mémoli in [14], they can in fact be used to obtain a lower bound for the Modified Gromov-Hausdorff distance between two metric spaces.

Definition 10 (curvature set - [12]). Given a, non-necessarily finite, metric space (X, d_X) we can consider the function

$$\begin{aligned} \Psi_X^m : X^m &\longrightarrow \mathbb{R}^{m \times m} \\ (x_1, \dots, x_m) &\longmapsto M \text{ s.t. } M_{i,j} = d_X(x_i, x_j). \end{aligned} \quad (25)$$

We call m -th curvature set of (X, d_X) the set

$$K_m(X) := \text{im} \Psi_X^m. \quad (26)$$

Let us see an example of some curvature sets for a finite metric space.

Example 3. Consider the set $X = \{x_1, x_2, x_3\}$ and endow it with the metric d such that $d(x_1, x_2) = 3$, $d(x_1, x_3) = 5$, $d(x_2, x_3) = 4$. Then, $K_2(X)$ and $K_3(X)$ are

$$\begin{aligned} K_2(X) &= \left\{ \begin{bmatrix} 0 & 3 \\ 3 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 4 \\ 4 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 5 \\ 5 & 0 \end{bmatrix}, \begin{bmatrix} 0 & 0 \\ 0 & 0 \end{bmatrix} \right\}. \\ K_3(X) &= \left\{ P^T \begin{bmatrix} 0 & 3 & 5 \\ 3 & 0 & 4 \\ 5 & 4 & 0 \end{bmatrix} P, P^T \begin{bmatrix} 0 & 0 & 3 \\ 0 & 0 & 3 \\ 3 & 3 & 0 \end{bmatrix} P, P^T \begin{bmatrix} 0 & 0 & 4 \\ 0 & 0 & 4 \\ 4 & 4 & 0 \end{bmatrix} P, \right. \\ &\quad \left. P^T \begin{bmatrix} 0 & 0 & 5 \\ 0 & 0 & 5 \\ 5 & 5 & 0 \end{bmatrix} P, \begin{bmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{bmatrix} \mid P \text{ runs in } 3 \times 3 \text{ permutation matrices} \right\}. \end{aligned}$$

Curvature sets encode information about finite metric subspaces of a given metric space. A similar, but coarser, invariant is the isometric sequence of a space, introduced by Hirasaka and Shinohara [13]. Curvature sets are important in virtue of the next theorem.

Theorem 5 (isometry of compact metric spaces - [12]). *Two compact metric spaces X and Y are isometric if and only if $K_m(X) = K_m(Y)$ for all $m \in \mathbb{N}$.*

Remark 4. Notice that every FMS is compact and then satisfies the hypothesis of the above theorem.

Therefore, checking the equality of curvature sets is a way to find out whether two metric spaces are isometric or not. In the general case, in which a metric space is not finite, we have to prove the equality of all curvature sets to ensure the isometry between metric spaces, but for the finite case the problem becomes easier. We can see that the r -th curvature set carries all the information included in all the l -th curvature sets for $l < r$.

Lemma 3. *For any two, possibly infinite, metric spaces (X, d_X) , (Y, d_Y) , if $K_r(X) = K_r(Y)$ for a certain $r \in \mathbb{N}$ then $K_l(X) = K_l(Y)$ for all $l \leq r$.*

Proof. Each matrix of $K_l(X)$ can be obtained from a matrix of $K_r(X)$ removing $r - l$ rows and columns. Given a set of indices $I := \{i_1, \dots, i_k\} \subseteq \{1, \dots, n\}$ and a matrix $M \in \mathbb{R}^{n \times n}$ we can define M_I as the matrix obtained by M removing the columns and the rows whose indices are in I . Then

$$\begin{aligned} K_l(X) &= \{M_I \mid M \in K_r(X), I \subseteq \{1, \dots, n\}, |I| = r - l\} = \\ &= \{M_I \mid M \in K_r(Y), I \subseteq \{1, \dots, n\}, |I| = r - l\} = K_l(Y). \end{aligned} \quad (27)$$

□

For finite metric spaces we can further improve this result. In the following theorem, we show that given a finite metric space X of cardinality n , all the curvature sets are determined by $K_n(X)$.

Theorem 6. *Let (X, d_X) and (Y, d_Y) be two finite metric space of cardinality n . Then*

$$K_m(X) = K_m(Y) \quad \forall m \in \mathbb{N} \iff K_n(X) = K_n(Y). \quad (28)$$

Proof. The forward implication is given by the hypothesis. We have to prove the other direction \Leftarrow . We already know, by Lemma 3, that $K_m(X) = K_m(Y)$ for all $m \leq n$. We have only to prove the case in which $m > n$. For every matrix $M \in K_m(X)$, we can find m points x_1, \dots, x_m of X such that $\Psi_X^m(x_1, \dots, x_m) = M$. Of these m points at most $k \leq n$ of them can be different, let them be x_{i_1}, \dots, x_{i_k} . Then, the matrix $\Psi_X^k(x_{i_1}, \dots, x_{i_k})$ is in $K_k(X) = K_k(Y)$ by Lemma 3. Then, we have $y_1, \dots, y_k \in Y$ such that $\Psi_X^k(x_{i_1}, \dots, x_{i_k}) = \Psi_Y^k(y_1, \dots, y_k)$, and it means that

$$d_X(x_{i_a}, x_{i_b}) = d_Y(y_a, y_b). \quad (29)$$

Consider the bijection $\phi : \{x_{i_1}, \dots, x_{i_k}\} \longrightarrow \{y_1, \dots, y_k\}$ with $\phi(x_{i_j}) = y_j$, $j = 1, \dots, k$. Thanks to Eq. (29) we have that

$$\Psi_X^m(x_1, \dots, x_m) = \Psi_Y^m(\phi(x_1), \dots, \phi(x_m)) \in K_m(Y).$$

Therefore, $K_m(X) \subseteq K_m(Y)$. Analogously, we can see that $K_m(Y) \subseteq K_m(X)$, hence they must be equal. □

We have seen in Example 3 that when we compute the m -th curvature set of a metric spaces we take m -tuples of points of X with repetitions. This means computing n^m matrices. We would like to reduce such a computational cost, and we try to do so introducing the concept of reduced curvature set.

Definition 11 (reduced curvature set). Consider a metric space (X, d_X) and the associated function $\Psi_X^m : X^m \longrightarrow \mathbb{R}^{m \times m}$ defined in Eq. (25). We call m -th reduced curvature set of (X, d_X) the set

$$\tilde{K}_m(X) = \{\Psi_X^m(x_1, \dots, x_m) \mid x_1, \dots, x_m \in X \text{ and } x_i \neq x_j \text{ if } i \neq j\}. \quad (30)$$

As we can see from the definition, to obtain the m -th reduced curvature set we need to compute $\frac{n!}{(n-m)!}$ matrices. We want to show that we do not lose any information with this reduction.

Lemma 4. *For any two metric spaces (X, d_X) , (Y, d_Y) , if $\tilde{K}_r(X) = \tilde{K}_r(Y)$ for a certain $r \in \mathbb{N}$, then $\tilde{K}_l(X) = \tilde{K}_l(Y)$ for all $l \leq r$.*

Proof. The proof is analogous to that of Lemma 3. \square

Theorem 7. *Let (X, d_X) and (Y, d_Y) be two finite metric spaces of cardinality n . For any $m \leq n$, we have*

$$K_m(X) = K_m(Y) \iff \tilde{K}_m(X) = \tilde{K}_m(Y). \quad (31)$$

Proof. Assume $K_m(X) = K_m(Y)$. Then a matrix $M \in K_m(X)$ is also an element of $\tilde{K}_m(X)$ if and only if $(M_{i,j} = 0 \iff i = j)$. The same argument holds also for $K_m(Y)$, therefore, given $M \in \tilde{K}_m(X)$, we have $M \in K_m(X) = K_m(Y)$ and $M \in K_m(Y)$. Since M has null entries only in its diagonal, $M \in \tilde{K}_m(Y)$. In this way, we can see that $\tilde{K}_m(X) \subseteq \tilde{K}_m(Y)$ and $\tilde{K}_m(Y) \subseteq \tilde{K}_m(X)$, hence they are equal. Assume now that $\tilde{K}_m(X) = \tilde{K}_m(Y)$. We want to prove that, for any $M \in K_m(X) \setminus \tilde{K}_m(X)$, we have $M \in K_m(Y)$. We know there are $x_1, \dots, x_m \in X$ such that $M = \Psi_X^m(x_1, \dots, x_m)$, where at most $k < m$ of the points are different. Suppose these points are x_{i_1}, \dots, x_{i_k} . For Lemma 4, we have that $\tilde{K}_{m-k}(X) = \tilde{K}_{m-k}(Y)$, then we have y_1, \dots, y_k points of Y such that

$$\Psi_X^{m-k}(x_{i_1}, \dots, x_{i_k}) = \Psi_Y^{m-k}(y_1, \dots, y_k).$$

Hence, given the bijection $\phi : \{x_{i_1}, \dots, x_{i_k}\} \longrightarrow \{y_1, \dots, y_k\}$ with $\phi(x_{i_j}) = y_j$, $j = 1, \dots, k$, we have

$$M = \Psi_X^m(x_1, \dots, x_m) = \Psi_Y^m(\phi(x_1), \dots, \phi(x_m)) \in K_m(Y). \quad (32)$$

Then, $K_m(X) \subseteq K_m(Y)$ and reasoning in the same way we have $K_m(Y) \subseteq K_m(X)$, therefore they are equal. \square

Thanks to the last theorem, we can see that reduced curvature sets carry the same information given by the non reduced version. Notice that for finite metric spaces we do not have m -th reduced curvature sets, with m greater than the number of points of the space. In the following theorem, we observe that isometry of finite metric spaces is characterised by the n -th reduced curvature set.

Theorem 8. *Two finite metric spaces (X, d_X) and (Y, d_Y) of cardinality n are isometric if and only if $\tilde{K}_n(X) = \tilde{K}_n(Y)$.*

Proof. We need to prove the “ \Leftarrow ” implication only. Since X and Y are finite they are compact and by Theorem 5 we know that they are isometric if and only if $K_m(X) = K_m(Y)$ for all $m \in \mathbb{N}$. By Theorem 6, since X and Y are finite, we know that this is true if and only if $K_n(X) = K_n(Y)$ and for Theorem 7 this holds if and only if $\tilde{K}_n(X) = \tilde{K}_n(Y)$. \square

We recall that, thanks to Theorem 4, two spaces are weakly isometric if and only if their canonicalizations are isometric, and this condition can now be checked using the above theorem. Hence, we have the following corollary.

Corollary 2. *Two finite metric spaces (X, d_X) , (Y, d_Y) of cardinality n with respective canonicalizations $(\mathcal{C}_X, d_{\mathcal{C}_X})$, $(\mathcal{C}_Y, d_{\mathcal{C}_Y})$ are weakly isometric if and only if $\tilde{K}_n(\mathcal{C}_X) = \tilde{K}_n(\mathcal{C}_Y)$.*

5 Vietoris-Rips filtration and finite metric spaces

We will provide a categorification of the concept of weak isometry of finite metric spaces, introduced in the work [10], in order to obtain another complete invariant for weak isometry.

Definition 12 (monotone map between FMS). Given two finite metric spaces (X, d_X) and (Y, d_Y) , we say the a map $f : X \longrightarrow Y$ is monotone if, for all $x_1, x_2, x'_1, x'_2 \in X$, we have:

$$d_X(x_1, x_2) \leq d_X(x'_1, x'_2) \Rightarrow d_Y(f(x_1), f(x_2)) \leq d_Y(f(x'_1), f(x'_2)). \quad (33)$$

Remark 5 (Proposition 3 - [10]). If $f : X \longrightarrow Y$ is a monotone map, then

$$d_X(x_1, x_2) = d_X(x'_1, x'_2) \Rightarrow d_Y(f(x_1), f(x_2)) = d_Y(f(x'_1), f(x'_2)). \quad (34)$$

The converse is not true.

Lemma 5. *A monotone map $f : X \longrightarrow Y$ between two finite metric spaces X and Y induces a non-decreasing function between the distance sets $\tilde{f} : D(X) \longrightarrow D(Y)$ given by*

$$\tilde{f}(a) = d_Y(f(x_i), f(x_j)) \quad \text{where } d_X(x_i, x_j) = a. \quad (35)$$

Proof. Thanks to Remark 5, we have that the function \tilde{f} is well defined. In fact, for any $a \in D(X)$, we have that, if $d_X(x_i, x_j) = d_X(x'_i, x'_j) = a$, we can write $\tilde{f}(a) = d_Y(f(x_i), f(x_j)) = d_Y(f(x'_i), f(x'_j))$. The function is non-decreasing because if $a = d_X(x_1, x_2) \leq b = d_X(x_3, x_4)$, by Definition 12, $\tilde{f}(a) = d_Y(f(x_1), f(x_2)) \leq d_Y(f(x_3), f(x_4)) = \tilde{f}(b)$. \square

Lemma 6. *A monotone map $f : X \longrightarrow Y$ between two finite metric spaces X and Y induces a non-decreasing function $\hat{f} : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ whose restriction is \tilde{f} as in Lemma 5.*

Proof. Thanks to Lemma 5, we have that f induces a non-decreasing function $\tilde{f} : D(X) \longrightarrow D(Y)$. Such a function can be extended to a non decreasing function $\hat{f} : \mathbb{R}^+ \longrightarrow \mathbb{R}^+$ in the following way. If $D(X) = \{a_1, \dots, a_k \mid a_i < a_j \text{ if } i < j\}$,

we define \hat{f} as

$$\hat{f}(x) = \begin{cases} \frac{\tilde{f}(a_1)}{a_1}x & \text{if } x \in [0, a_1] \\ \frac{\tilde{f}(a_{i+1}) - \tilde{f}(a_i)}{a_{i+1} - a_i}(x - a_i) + \tilde{f}(a_i) & \text{if } x \in [a_i, a_{i+1}] \\ (x - a_k) + \tilde{f}(a_k) & \text{if } x \in (a_k, \infty). \end{cases} \quad (36)$$

By definition, it follows that $\hat{f}|_{D(X)} = \tilde{f}$. \square

Definition 13 (category of FMS - [10]). We can define a category **FMS** of finite metric space whose objects are finite metric spaces and whose morphisms are monotone maps.

Remark 6. It is possible to observe that two finite metric spaces are isomorphic in the category **FMS** if and only if they are weakly isometric.

We recall some concepts of algebraic topology that we will use in the rest of this section. For more detailed information, we refer the reader to [15].

An *abstract simplicial complex* is a collection K of finite non-empty sets, called *simplices*, such that for any σ in K every τ , non-empty subset of σ is in K . If a simplex σ has elements x_0, \dots, x_k , we will say that is generated by the points x_0, \dots, x_k and we will write $\sigma = \{x_0, \dots, x_k\}$. The *dimension* of a simplex is defined as the number of its elements minus 1, so

$$\dim \sigma = \dim \{x_0, \dots, x_k\} = k.$$

The 0-dimensional simplices are also called *vertices*. Given two abstract simplicial complexes K and L , a *simplicial map* $s : K \rightarrow L$ is a function that sends the vertices of K to vertices of L , such that, if $\sigma = \{x_0, \dots, x_k\}$ is a k -simplex of K , then $s(\sigma) = \{s(\{x_0\}), \dots, s(\{x_k\})\}$ is a simplex of L . Beware that the function s does not need to be injective on the set of vertices. It may be possible that the image of a simplex σ is a simplex $s(\sigma)$ of lower dimension.

Example 4. Consider the simplicial complex

$$K = \{\{a\}, \{b\}, \{c\}, \{a, b\}, \{a, c\}, \{b, c\}, \{a, b, c\}\}.$$

An example of simplicial map is the function $s : K \rightarrow K$, that on the set of vertices is equal to

$$\begin{aligned} s(\{a\}) &= \{a\}, \\ s(\{b\}) &= \{b\}, \\ s(\{c\}) &= \{b\}. \end{aligned} \quad (37)$$

It is possible to see that some simplices degenerate, through s , to simplices of lower dimension. For example $s(\{a, b, c\}) = \{a, b\}$.

Definition 14 (category of simplicial complexes). We denote with **Simp** the category whose objects are finite abstract simplicial complexes, and morphisms are simplicial maps.

An example of abstract simplicial complex that we will use is the Vietoris-Rips complex.

Definition 15 (Vietoris-Rips complex). Given a finite metric space (X, d_X) and a positive real number ε , the Vietoris-Rips complex $\text{VR}_\varepsilon(X)$ is the abstract simplicial complex whose elements are subsets $\sigma = \{x_0, \dots, x_k\}$ of X such that

$$\forall x_i, x_j \in \sigma, \quad d_X(x_i, x_j) \leq \varepsilon.$$

We recall that we can associate to any partially ordered set (P, \leq_P) the category whose objects are the elements of P , and whose morphisms are the relations $a \leq_P b$. In this section, we will consider in this way the category (\mathbb{R}^+, \leq) .

Definition 16 (Vietoris-Rips filtration). Given a finite metric space (X, d_X) , we can consider the functor $\text{VR}_\bullet(X) : (\mathbb{R}^+, \leq) \rightarrow \mathbf{Simp}$ that assigns to each $a \in \mathbb{R}^+$ the Vietoris-Rips complex $\text{VR}_a(X)$ and to each morphism $a \leq b$ the inclusion $\iota_{a \leq b}^X : \text{VR}_a(X) \hookrightarrow \text{VR}_b(X)$.

Definition 17 (rescaling). Given a non-decreasing function $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ we call ψ -rescaling the functor $R_\psi : (\mathbb{R}^+, \leq) \rightarrow (\mathbb{R}^+, \leq)$ with

$$\begin{aligned} R_\psi(a) &= \psi(a) \\ R_\psi(a \leq b) &= \psi(a) \leq \psi(b). \end{aligned} \tag{38}$$

Remark 7 (notation). Given two functors $F : \mathbf{B} \rightarrow \mathbf{C}$ and $G : \mathbf{C} \rightarrow \mathbf{D}$ we denote their composition as $GF : \mathbf{B} \rightarrow \mathbf{D}$. If we have another functor $F' : \mathbf{B} \rightarrow \mathbf{C}$, a natural transformation η between F and F' is a family of morphisms $\{\eta_a : F(a) \rightarrow F'(a) \mid a \in \text{ob}(\mathbf{B})\}$ such that, for every morphism $m : a \rightarrow b$ in \mathbf{B} , the following diagram commute

$$\begin{array}{ccc} F(a) & \xrightarrow{F(m)} & F(b) \\ \downarrow \eta_a & & \downarrow \eta_b \\ F'(a) & \xrightarrow{F'(m)} & F'(b). \end{array} \tag{39}$$

In this case, we will write $\eta : F \Rightarrow F'$.

Lemma 7. *A morphism $f : X \rightarrow Y$ in **FMS** induces a rescaling $R_{\hat{f}}$ and a natural transformation $\eta^f : \text{VR}_\bullet(X) \Rightarrow \text{VR}_\bullet(Y)R_{\hat{f}}$.*

Proof. Thanks to Lemma 6, we have a non-decreasing function \hat{f} that induces a rescaling $R_{\hat{f}}$. We want to see that, for any $a \in \mathbb{R}^+$, we have a simplicial map

$\eta_a^f : \text{VR}_a(X) \longrightarrow \text{VR}_\bullet(Y)R_{\hat{f}}(a) = \text{VR}_{\hat{f}(a)}(Y)$ such that, for all $a, b \in \mathbb{R}^+$ with $a \leq b$, we have a commutative diagram

$$\begin{array}{ccc} \text{VR}_a(X) & \xleftarrow{\iota^X} & \text{VR}_b(X) \\ \downarrow \eta_a^f & & \downarrow \eta_b^f \\ \text{VR}_{\hat{f}(a)}(Y) & \xleftarrow{\iota^Y} & \text{VR}_{\hat{f}(b)}(Y). \end{array} \quad (40)$$

We define η_a^f as

$$\eta_a^f(\{x_{i_0}, \dots, x_{i_k}\}) = \{f(x_{i_0}), \dots, f(x_{i_k})\}. \quad (41)$$

By the very definition of Vietoris-Rips complex and \hat{f} , we have that, for every simplex σ of $\text{VR}_a(X)$, $\eta_a^f(\sigma)$ is a simplex of $\text{VR}_{\hat{f}(a)}(Y)$ and η_a^f is a well-defined simplicial map. We only need to prove that $\iota^Y \circ \eta_a^f = \eta_b^f \circ \iota^X$. Indeed, for any $\sigma \in \text{VR}_a(X)$ with $\sigma = \{x_{i_0}, \dots, x_{i_k}\}$ we have

$$\begin{aligned} \iota^Y \circ \eta_a^f(\sigma) &= \iota^Y(\{f(x_{i_0}), \dots, f(x_{i_k})\}) = \{f(x_{i_0}), \dots, f(x_{i_k})\} \\ \eta_b^f \circ \iota^X(\sigma) &= \eta_b^f(\{x_{i_0}, \dots, x_{i_k}\}) = \{f(x_{i_0}), \dots, f(x_{i_k})\}. \end{aligned} \quad (42)$$

Hence η^f is a natural transformation. \square

Now, we want to show that the Vietoris-Rips filtration can be used as a complete invariant for weak isometry.

Theorem 9. *Given two finite metric spaces X and Y , the following statements are equivalent:*

1. X and Y are isomorphic in **FMS**.
2. There exist a rescaling R_ψ and a natural isomorphism

$$\eta : \text{VR}_\bullet(X) \Longrightarrow \text{VR}_\bullet(Y)R_\psi.$$

Proof. Suppose that X and Y are isomorphic in **FMS**. Then we have a monotone map $f : X \longrightarrow Y$ that is a bijection. Thanks to Lemma 7, we have a rescaling $R_{\hat{f}}$ and a natural transformation $\eta^f : \text{VR}_\bullet(X) \Longrightarrow \text{VR}_\bullet(Y)R_{\hat{f}}$. We want to see that for all $a \in \mathbb{R}^+$, η_a^f is an isomorphism of simplicial complexes. Since f is a bijection, we know that η_a^f is injective. In fact, given $\sigma = \{x_{i_1}, \dots, x_{i_k}\}$ and $\tau = \{x_{j_1}, \dots, x_{j_l}\}$ with $\sigma \neq \tau$, we can assume without loss of generality that there is a $x_j \in \tau$ with $x_j \notin \sigma$. Then if $\eta_a^f(\sigma) = \eta_a^f(\tau)$, there is a $x_i \in \sigma$ with $f(x_j) = f(x_i)$ and this is absurd for the injectivity of f . On the other hand η_a^f is also surjective. Suppose that there is a $\rho \in \text{VR}_{\hat{f}(a)}(Y)$ with $\rho = \{y_{i_1}, \dots, y_{i_k}\}$ that is not in the image of η_a^f . We can consider the points $f^{-1}(y_{i_1}), \dots, f^{-1}(y_{i_k})$ of X and see that they form a simplex of $\text{VR}_a(X)$. In fact, since $\rho \in \text{VR}_{\hat{f}(a)}(Y)$, for all $u, v \in \rho$, we have $d_Y(u, v) \leq \hat{f}(a)$. It can be seen that, for all $x_1, x_2 \in X$,

we have $\hat{f}(d_X(x_1, x_2)) = d_Y(f(x_1), f(x_2))$. Therefore, for all $u, v \in \rho$, since \hat{f}^{-1} is also a strictly increasing function

$$d_X(f^{-1}(u), f^{-1}(v)) = \hat{f}^{-1}(d_Y(u, v)) \leq \hat{f}^{-1}(\hat{f}(a)) = a. \quad (43)$$

Then points $f^{-1}(y_{i_1}), \dots, f^{-1}(y_{i_k})$ span a simplex of $\text{VR}_a(X)$ whose image under η_a^f is ρ . Therefore η_a^f is also bijective and is an isomorphism of simplicial complexes. Hence, η is a natural isomorphism.

Now, suppose that we have a rescaling R_ψ and a natural isomorphism $\eta : \text{VR}_\bullet(X) \Rightarrow \text{VR}_\bullet(Y)R_\psi$. For each $a \in \mathbb{R}^+$, the restriction of the isomorphism η_a to the vertices of $\text{VR}_a(X)$ yields a bijection $f_a : X \rightarrow Y$. Moreover, these bijections are all the same because of the commutativity of diagrams that define the natural isomorphism. Hence we have a unique bijection $f : X \rightarrow Y$ associated with η . We claim that this bijection is an isomorphism of finite metric spaces. In fact, for each couple of points $x_1, x_2 \in X$, call $\bar{a} = d_X(x_1, x_2)$. Since η is a natural isomorphism, we have that $\sigma = \{x_1, x_2\} \in \text{VR}_{\bar{a}}(X)$ and that $\eta_{\bar{a}}(\sigma) = \{f(x_1), f(x_2)\}$ is a simplex of $\text{VR}_{\psi(\bar{a})}(Y)$, that is not present in any $\text{VR}_b(Y)$, for $b \leq \psi(\bar{a})$. This means that $d_Y(f(x_1), f(x_2)) = \psi(\bar{a}) = \psi(d_X(x_1, x_2))$, for all $x_1, x_2 \in X$, and therefore X and Y are weakly isometric and also isomorphic in **FMS**. \square

Remark 8. Given two Vietoris-Rips filtrations $\text{VR}_\bullet(X)$ and $\text{VR}_\bullet(Y)$, the existence of an isomorphism between each $\text{VR}_a(X)$ and $\text{VR}_a(Y)$ is not enough to ensure that X and Y are isometric. Indeed, all these isomorphism have to commute with the inclusions given by the filtrations. For example, consider the two metric spaces, depicted in Fig. 3, given by the distance matrices

$$d_X = (d_X(x_i, x_j)) = \begin{pmatrix} 0 & 7 & 9 & 10 \\ 7 & 0 & 8 & 11 \\ 9 & 8 & 0 & 12 \\ 10 & 11 & 12 & 0 \end{pmatrix}, \quad (44)$$

$$d_Y = (d_Y(y_i, y_j)) = \begin{pmatrix} 0 & 7 & 9 & 10 \\ 7 & 0 & 8 & 12 \\ 9 & 8 & 0 & 11 \\ 10 & 12 & 11 & 0 \end{pmatrix}.$$

They are not isometric, but, for each $a \in \mathbb{R}^+$, we can find an isomorphism between $\text{VR}_a(X)$ and $\text{VR}_a(Y)$. Notice that, on the other hand, if there exists an a in \mathbb{R}^+ such that there is no isomorphism between $\text{VR}_a(X)$ and $\text{VR}_a(Y)$, then the two spaces are for sure not isometric.

5.1 Persistent homology as an incomplete invariant for weak isometry

Topological data analysis (TDA) is a branch of applied mathematics developed in the last 30 years in order to have a set of tools, based on topological and




Figure 3: Embedding of the spaces of Remark 8 in \mathbb{R}^3 .

geometrical concepts, to analyse data [9, 8]. The main tool used in TDA is *persistent homology*, an algebraic topology technique that consider the evolution and relations of homology groups of a sequence of nested topological spaces. In the usual pipeline of topological data analysis, the Vietoris-Rips filtration is used to compute the so called *persistence module*, with which it is possible to keep track of the changes of homological features of the simplicial complexes at different scales of the filtration. Let us recall the definition of the k -th *degree simplicial homology functor*. Given a field \mathbb{F} , with $\mathbf{Vect}_{\mathbb{F}}$ we denote the category whose objects are finite dimensional vectors spaces over \mathbb{F} and with morphisms the linear maps between these spaces. Let us consider an abstract simplicial complex K . To simplify the definitions we will consider a total order on the set of vertices of K . Every k -simplex of K from now on will be considered as an ordered tuple $[x_0, \dots, x_k]$ of $k+1$ vertices of K . The group of k -chains of K , denoted with $C_k(K)$ is the vectors space whose elements are formal sums

$$\sum_i \lambda_i \sigma_i \quad (45)$$

where λ_i is an element of \mathbb{F} and σ_i is a k -simplex of K . It is possible to see that a basis is given by the formal sums $\{1_{\mathbb{F}}\sigma \mid \dim \sigma = k\}$. Between every $C_k(K)$ and $C_{k-1}(K)$, there is a linear map $\partial_k : C_k(K) \rightarrow C_{k-1}(K)$ called *boundary operator*. It is defined on the basis of C_k given by the set of k -simplices as

$$\partial_k([x_0, \dots, x_k]) = \sum_{i=0}^k (-1)^i [x_0, \dots, \hat{x}_i, \dots, x_k], \quad (46)$$

where with $[x_0, \dots, \hat{x}_i, \dots, x_k]$ we denote the $(k-1)$ -simplex obtained removing the vertex x_i . It is possible to show that for every $k > 0$ it holds $\partial_k \circ \partial_{k+1} = 0$, meaning that the set $\text{im } \partial_{k+1}$ is a vector subspace of $\ker \partial_k$. Then, the k -th

homology group of K with coefficients in \mathbb{F} is the quotient vector space

$$H_k(K) := \frac{\ker \partial_k}{\text{im } \partial_{k+1}}. \quad (47)$$

The k -th simplicial homology functor is a functor from the category **Simp** to the category **Vect** $_{\mathbb{F}}$, that assigns to every object K of **Simp** the homology group $H_k(K)$ in **Vect** $_{\mathbb{F}}$, and to every simplicial map $s : K \rightarrow L$ the induced map $H_k(s) : H_k(K) \rightarrow H_k(L)$ between the homology groups.

In general terms, a *filtration* is a functor $F : (\mathbb{R}_+, \leq) \rightarrow \mathbf{Simp}$. Given a filtration F , the composition $H_k F$ is called the k -th degree *persistence module*. It is possible to define an *interleaving distance* between persistence modules in the following way, as described in detail in [2]. Given a positive real number ε , it is possible to define a translation functor $T_\varepsilon : (\mathbb{R}^+, \leq) \rightarrow (\mathbb{R}^+, \leq)$, with $T_\varepsilon(x) = x + \varepsilon$ and a natural transformation $\nu_\varepsilon : \text{Id}_{(\mathbb{R}^+, \leq)} \Rightarrow T_\varepsilon$, with $\nu_\varepsilon(x) : x \rightarrow x + \varepsilon$ defined as $x \leq x + \varepsilon$. Two persistence modules $H_k F$ and $H_k G$ are said to be ε -interleaved if there exist two natural transformation $\eta_F : H_k F \Rightarrow H_k G T_\varepsilon$ and $\eta_G : H_k G \Rightarrow H_k F T_\varepsilon$ such that

$$(\eta_G T_\varepsilon) \eta_F = H_k F \nu_{2\varepsilon} \quad \text{and} \quad (\eta_F T_\varepsilon) \eta_G = H_k G \nu_{2\varepsilon}. \quad (48)$$

The interleaving distance d_I between two persistence modules $H_k F$ and $H_k G$ is defined as

$$d_I(H_k F, H_k G) = \inf \{ \varepsilon \geq 0 \mid H_k F \text{ and } H_k G \text{ are } \varepsilon\text{-interleaved} \}. \quad (49)$$

The reader familiar with persistent homology will find the following corollary to be natural.

Corollary 3. *The persistent homology of the Vietoris-Rips filtration is an incomplete invariant for isometry.*

Proof. The two spaces in Remark 8 have persistence modules with interleaving distance 0, yet they are not isometric. \square

Even if persistent homology is only an incomplete invariant, we can still use it to study the problem of weak isometry. Given two finite metric spaces, it is possible to compute their canonicalizations, from them their associated Vietoris-Rips filtrations and then the persistence modules. If the obtained persistence modules have interleaving distance greater than 0, then the two spaces cannot be weakly isometric.

5.2 A dissimilarity measure for persistence modules

In a spirit similar to Section 3, we can define a dissimilarity between persistence modules. Recall that we defined \mathcal{J} as the set of strictly increasing functions $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $\psi(0) = 0$.

Definition 18. Given two persistence modules HF_1 and HF_2 , the dissimilarity \tilde{d} between them is

$$\tilde{d}(HF_1, HF_2) = \inf_{\psi \in \mathcal{J}} d_I(HF_1, HF_2 R_\psi) + \inf_{\psi \in \mathcal{J}} d_I(HF_1 R_\psi, HF_2), \quad (50)$$

where d_I is the interleaving distance between persistence modules.

One of the key properties desired for distances between persistence modules is that they satisfy a form of stability theorem, that is, there must be a distance between the original metric spaces, that bounds from above the distance between the obtained persistence modules. We have a stability theorem of this kind for the Vietoris-Rips filtration and the interleaving distance [6], in fact

$$d_I(H_k \text{VR}_\bullet(X, d_X), H_k \text{VR}_\bullet(Y, d_Y)) \leq 2d_{GH}((X, d_X), (Y, d_Y)).$$

We will provide a similar stability theorem also for the distances introduced in this paper.

Theorem 10 (stability theorem for weak isometry). *Let $(X, d_X), (Y, d_Y)$ be two finite metric spaces. We denote with $H_k \text{VR}_\bullet(X, d_X)$ and $H_k \text{VR}_\bullet(Y, d_Y)$ the k -th persistence modules obtained from the Vietoris-Rips filtration associated with the two spaces. Then for all $k \in \mathbb{N}$,*

$$\tilde{d}(H_k \text{VR}_\bullet(X, d_X), H_k \text{VR}_\bullet(Y, d_Y)) \leq 2\hat{d}((X, d_X), (Y, d_Y)). \quad (51)$$

We can see that thanks to this theorem and Proposition 3, if the persistence modules obtained by two Vietoris-Rips filtration have distance \tilde{d} greater than 0, then the corresponding finite metric spaces cannot be weakly isometric.

Example 5. Consider the finite metric spaces (X, d_X) and (Y, d_Y) with

$$d_X = (d_X(x_i, x_j)) = \begin{pmatrix} 0 & 7 & 12 & 8 \\ 7 & 0 & 10 & 11 \\ 12 & 10 & 0 & 9 \\ 8 & 11 & 9 & 0 \end{pmatrix}, \quad (52)$$

$$d_Y = (d_Y(y_i, y_j)) = \begin{pmatrix} 0 & 7 & 12 & 8 \\ 7 & 0 & 10 & 9 \\ 12 & 10 & 0 & 11 \\ 8 & 9 & 11 & 0 \end{pmatrix}.$$

We will use \mathbb{Z}_2 as the field of coefficients with which we will compute homology. We can see that the persistence module $H_1 \text{VR}_\bullet(X, d_X)$ is the functor

$$\begin{aligned} H_1 \text{VR}_\varepsilon(X, d_X) &= \begin{cases} \mathbb{Z}_2 & \text{if } 10 \leq \varepsilon \leq 11 \\ 0 & \text{otherwise,} \end{cases} \\ H_1 \text{VR}_\bullet(X, d_X)(a \leq b) &= \\ &= \begin{cases} \text{id} : \mathbb{Z}_2 \rightarrow \mathbb{Z}_2 & \text{if } 10 \leq a \leq b \leq 11 \\ 0 : H_1 \text{VR}_a(X, d_X) \rightarrow H_1 \text{VR}_b(X, d_X) & \text{otherwise.} \end{cases} \end{aligned}$$




Figure 4: Embedding of the spaces of Example 5 in \mathbb{R}^3 .

The persistence module $H_1 \text{VR}_\bullet(Y, d_Y)$ is

$$\begin{aligned} H_1 \text{VR}_\varepsilon(Y, d_Y) &= 0 \quad \forall \varepsilon \geq 0 \\ H_1 \text{VR}_\bullet(Y, d_Y)(a \leq b) &= 0 : H_1 \text{VR}_a(Y, d_Y) \rightarrow H_1 \text{VR}_b(Y, d_Y) \quad \forall a \leq b. \end{aligned}$$

Let us consider the sequence $(\psi_n)_{n \in \mathbb{N}}$ with

$$\psi_n(x) = \begin{cases} x & \text{if } 0 \leq x \leq 10 \\ n(x - 10) + 10 & \text{if } 10 < x \leq 10 + \frac{1}{n} \\ x + 1 - \frac{1}{n} & \text{if } 10 + \frac{1}{n} < x. \end{cases}$$

It is easy to see that

$$\lim_{n \rightarrow \infty} d_I(H_1 \text{VR}_\bullet(Y, d_Y), H_1 \text{VR}_\bullet(X, d_X) R_{\psi_n}) = 0.$$

On the other hand, for all strictly increasing functions $\psi : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ it holds

$$d_I(H_1 \text{VR}_\bullet(X, d_X), H_1 \text{VR}_\bullet(Y, d_Y) R_\psi) = \frac{1}{2}.$$

Therefore, $\tilde{d}(H_1 \text{VR}_\bullet(X, d_X), H_1 \text{VR}_\bullet(Y, d_Y)) = \frac{1}{2} > 0$, and the two spaces are not weakly isometric.

Proof of Theorem 10. We have that, for every $\psi \in \mathcal{J}$,

$$H_k \text{VR}_\bullet(X, \psi \circ d_X) R_\psi = H_k \text{VR}_\bullet(X, d_X).$$

This is true because $(X, \psi \circ d_X)$ and (X, d_X) are weakly isometric and, as in the proof of Theorem 9, there are a natural isomorphism η and a rescaling R_ψ such that $\eta : \text{VR}_\bullet(X, d_X) \Rightarrow \text{VR}_\bullet(X, \psi \circ d_X) R_\psi$. In this case, the natural isomorphism is simply the identity between every $\text{VR}_a(X, d_X)$ and $\text{VR}_{\psi(a)}(X, \psi \circ d_X)$. Therefore, the two functors $\text{VR}_\bullet(X, d_X)$ and $\text{VR}_\bullet(X, \psi \circ d_X) R_\psi$ are equal.

Now, let us take a sequence $(\psi_n)_{n \in \mathbb{N}}$ in \mathcal{J} with

$$\lim_{n \rightarrow \infty} d_{GH}((X, \psi_n \circ d_X), (Y, d_Y)) = \inf_{\psi \in \mathcal{J}} d_{GH}((X, \psi \circ d_X), (Y, d_Y))$$

and a sequence $(\bar{\psi}_n)_{n \in \mathbb{N}}$ with

$$\lim_{n \rightarrow \infty} d_{GH}((X, d_X), (Y, \bar{\psi}_n \circ d_Y)) = \inf_{\psi \in \mathcal{J}} d_{GH}((X, d_X), (Y, \psi \circ d_Y)).$$

By the classical stability theorem [6], for all n in \mathbb{N} , we have

$$\begin{aligned} d_I(H_k \text{VR} \bullet (X, \psi_n \circ d_X), H_k \text{VR} \bullet (Y, d_Y)) &\leq 2d_{GH}((X, \psi_n \circ d_X), (Y, d_Y)), \\ d_I(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, \bar{\psi}_n \circ d_Y)) &\leq 2d_{GH}((X, d_X), (Y, \bar{\psi}_n \circ d_Y)). \end{aligned} \quad (53)$$

Recall that, for all n in \mathbb{N} by the definition of infimum

$$\begin{aligned} \inf_{\psi \in \mathcal{J}} d_I(H_k \text{VR} \bullet (X, \psi \circ d_X), H_k \text{VR} \bullet (Y, d_Y)) &\leq \\ &\leq d_I(H_k \text{VR} \bullet (X, \psi_n \circ d_X), H_k \text{VR} \bullet (Y, d_Y)), \\ &\quad \text{and} \\ \inf_{\psi \in \mathcal{J}} d_I(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, \psi \circ d_Y)) &\leq \\ &\leq d_I(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, \bar{\psi}_n \circ d_Y)). \end{aligned} \quad (54)$$

By the definition of \tilde{d} and because of the previous inequalities it holds

$$\begin{aligned} \tilde{d}(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, d_Y)) &\leq \\ \lim_{n \rightarrow \infty} (d_I(H_k \text{VR} \bullet (X, \psi_n \circ d_X), H_k \text{VR} \bullet (Y, d_Y)) + \\ + d_I(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, \bar{\psi}_n \circ d_Y))). \end{aligned} \quad (55)$$

Because of the inequalities in Eq. (53) and the definition of \hat{d} , the following is true

$$\begin{aligned} &\lim_{n \rightarrow \infty} (d_I(H_k \text{VR} \bullet (X, \psi_n \circ d_X), H_k \text{VR} \bullet (Y, d_Y)) + \\ &\quad + d_I(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, \bar{\psi}_n \circ d_Y))) \leq \\ &\leq \lim_{n \rightarrow \infty} (2d_{GH}((X, \psi_n \circ d_X), (Y, d_Y)) + 2d_{GH}((X, d_X), (Y, \bar{\psi}_n \circ d_Y))) = \\ &\quad = 2\hat{d}((X, d_X), (Y, d_Y)). \end{aligned} \quad (56)$$

Therefore,

$$\tilde{d}(H_k \text{VR} \bullet (X, d_X), H_k \text{VR} \bullet (Y, d_Y)) \leq 2\hat{d}((X, d_X), (Y, d_Y)). \quad (57)$$

□

6 Conclusions and future work

We have constructed suitable representative elements for the equivalence classes of the relation of weak isometry for finite metric spaces. Thanks to these representatives we can check in a simple way whether two spaces are weakly isometric

or not. We have given a definition of a notion of dissimilarity between finite metric spaces which measures how far they are from being weakly isometric. We have shown that curvature sets and Vietoris-Rips filtrations can help us in characterizing the classes of weak isometry. We have seen that we can use persistent homology to try to discriminate non-weakly isometric finite metric spaces, and we have defined a dissimilarity between persistence modules and shown that it satisfies a stability theorem for weak isometry. We would like to point out that in this work we never exploited the property of triangle inequality, therefore all the results could be easily extended to finite semi-metric spaces (see [7]). In the future, we would like to try to find other and simpler invariants for weak isometry and make a comparison between them. We have also seen the usefulness of persistent homology, and in future work we would like to study the problem of finding distances between persistence modules that are meaningful from the point of view of weak isometry.

Acknowledgement

AD, UF and FV have been supported by the SmartData@PoliTO center on Big Data and Data Science and by the Italian MIUR Award “Dipartimento di Eccellenza 2018-2022” - CUP: E11G18000350001.

References

- [1] Mireille Boutin and Gregor Kemper. On reconstructing n -point configurations from the distribution of distances or areas. *Adv. in Appl. Math.*, 32(4):709–735, 2004. doi:10.1016/S0196-8858(03)00101-5.
- [2] Peter Bubenik and Jonathan A. Scott. Categorification of persistent homology. *Discrete Comput. Geom.*, 51(3):600–627, 2014. doi:10.1007/s00454-014-9573-x.
- [3] Dmitri Burago, Yuri Burago, and Sergei Ivanov. *A course in metric geometry*, volume 33 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2001. doi:10.1090/gsm/033.
- [4] Gunnar Carlsson. Topology and data. *Bull. Amer. Math. Soc. (N.S.)*, 46(2):255–308, 2009. doi:10.1090/S0273-0979-09-01249-X.
- [5] Wojciech Chacholski, Alessandro De Gregorio, Nicola Quercioli, and Francesca Tombari. Landscapes of data sets and functoriality of persistent homology, 2020. arXiv:2002.05972.
- [6] Frédéric Chazal, Vin de Silva, and Steve Oudot. Persistence stability for geometric complexes. *Geom. Dedicata*, 173(1):193–214, 2014. doi:10.1007/s10711-013-9937-z.

- [7] O. Dovgoshey and E. Petrov. Weak similarities of metric and semimetric spaces. *Acta Math. Hungar.*, 141(4):301–319, 2013.
- [8] Herbert Edelsbrunner and John Harer. Persistent homology-a survey. In *Surveys on discrete and computational geometry*, volume 453 of *Contemp. Math.*, pages 257–282. Amer. Math. Soc., Providence, RI, 2008. doi:10.1090/conm/453/08802.
- [9] Patrizio Frosini. Measuring shapes by size functions. In David P. Casasent, editor, *Intelligent Robots and Computer Vision X: Algorithms and Techniques*, volume 1607, pages 122–133. International Society for Optics and Photonics, SPIE, 1992. doi:10.1117/12.57059.
- [10] A. G. Ganyushkin and V. V. Tsvirkunov. On the classification of finite metric spaces. *Mat. Zametki*, 56(4):48–58, 157, 1994. doi:10.1007/BF02362370.
- [11] Chad Giusti, Eva Pastalkova, Carina Curto, and Vladimir Itskov. Clique topology reveals intrinsic geometric structure in neural correlations. *Proc. Natl. Acad. Sci. USA*, 112(44):13455–13460, 2015. doi:10.1073/pnas.1506407112.
- [12] Misha Gromov. *Metric structures for Riemannian and non-Riemannian spaces*. Birkhäuser Boston, Inc., Boston, MA, english edition, 2007.
- [13] Mitsugu Hirasaka and Masashi Shinohara. Characterization of finite metric space by their isometric sequences, 2018. arXiv:1802.06097.
- [14] Facundo Mémoli. Some properties of gromov–hausdorff distances. *Discrete Comput. Geom.*, 48(2):416–440, Sep 2012. doi:10.1007/s00454-012-9406-8.
- [15] James R. Munkres. *Elements of algebraic topology*. Addison-Wesley Publishing Company, Menlo Park, CA, 1984.