



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Homogeneous and domain-wall topological Haldane conductors with dressed Rydberg atoms

Original

Homogeneous and domain-wall topological Haldane conductors with dressed Rydberg atoms / Montorsi, Arianna; Fazzini, Serena; Barbiero, Luca. - In: PHYSICAL REVIEW A. - ISSN 2469-9926. - 101:4(2020).

Availability:

This version is available at: 11583/2831585 since: 2020-06-01T16:30:47Z

Publisher:

American Physics Society

Published

DOI:10.1103/PhysRevA.101.043618

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Homogeneous and domain-wall topological Haldane conductors with dressed Rydberg atomsArianna Montorsi ¹, Serena Fazzini,² and Luca Barbiero ³¹*Institute for Condensed Matter Physics and Complex Systems, DISAT, Politecnico di Torino I-10129, Italy*²*Physics Department and Research Center OPTIMAS, University of Kaiserslautern, D-67663 Kaiserslautern, Germany*³*Center for Nonlinear Phenomena and Complex Systems, Université Libre de Bruxelles, CP 231, Campus Plaine, B-1050 Brussels, Belgium*

(Received 31 October 2019; revised manuscript received 23 March 2020; accepted 30 March 2020; published 24 April 2020)

The interplay between antiferromagnetic interaction and hole motion is capable of inducing conducting Haldane phases with topological features described by a finite nonlocal string order parameter. Here we show that these states of matter are captured by the one-dimensional t - J_z model, which can be experimentally realized with dressed Rydberg atoms trapped onto a one-dimensional optical lattice. In the sector with vanishing total magnetization, exact calculations associated with the bosonization technique allow us to predict that both metallic and superconducting topological Haldane states can be achieved. With the addition of an appropriate magnetic field, the system enters a domain-wall structure with finite total magnetization. In this regime, the conducting Haldane states are confined in domains separated by regions where a fully polarized Luttinger liquid occurs. A procedure to dynamically stabilize such topological phases starting from a confined Ising state is also described.

DOI: [10.1103/PhysRevA.101.043618](https://doi.org/10.1103/PhysRevA.101.043618)**I. INTRODUCTION**

Interacting quantum phases with topological features [1–6] associated with a class of nonlocal excitations represent one of the most intriguing topics in different areas of quantum physics. Besides providing a first case of quantum regimes that go beyond Landau’s theory of phase transitions, it has been further demonstrated that these peculiar excitations are deeply connected with symmetry-protected topological order [7,8]. A well-known example is the Haldane phase occurring in the spin-1 XXZ chain [9], where the presence of fractionalized edge states is captured by a nonlocal string operator [10]. Relevantly, a similar behavior is observed in other paradigmatic spin Hamiltonians like AKLT [11] and colorless Motzkin chains [12].

Haldane states with topological features can be induced by interaction also in itinerant bosonic [13–16] and fermionic [17–21] systems. Due to both the high level of parameter control and the fact that string order parameters can be measured by *in situ* imaging [22,23], many-body ultracold atomic systems [24] represent a very promising platform where these models and their topological states can be achieved. Indeed, many proposals involving particles with long-range dipolar interaction [25] trapped in optical lattices have been presented [26–30]. Crucially in all these possible setups, Haldane orders are associated with a finite value of the charge gap thus describing symmetry-protected topological insulators. On the other hand, it has been recently shown that a challenging implementation involving polar molecules can support the appearance of Haldane orders with a gapless charge sector [31]. In this special configuration, metallic states as well as regimes with dominant superconducting correlations can be achieved by manipulating the interaction strength between molecules in different dressed rotational states. Beside the

fact that a high density sample of polar molecules has been recently achieved [32] and spin exchange processes observed [33], the experimental implementation of the model supporting Haldane superconductivity remains very challenging.

At the same time, Rydberg atoms represent a very promising and flexible platform where long-range interaction can allow the study of intriguing properties of matter. Among many of them, it is worthwhile to mention recent results where crystal structures [34], Z_N density waves [35], interacting topological insulators [36], and Schrödinger cat states [37] have been experimentally observed. Moreover, recent impressive advances in manipulating such atoms allowed the production of “dressed” Rydberg excitations [38,39]. In this setup, the lifetime of the Rydberg excitations is able to exceed the time required by a particle to tunnel between two nearest-neighbor sites of an optical lattice. In the case of Ising-like spin-spin interactions, already observed in the laboratory, such an aspect allows in principle for the experimental realization of the usually called t - J_z model [40], whose topological properties have only recently been studied in the context of a generalized t - J model [31].

Motivated by this intriguing experimental platform, we address in detail here the topological properties of the aforementioned t - J_z model. In the case of nearest-neighbor couplings, the model is integrable and some ground-state properties can be extracted exactly. In particular, as a function of the filling and the Ising coupling, both a spin-gapped liquid and a phase-separated regime are present. Moreover, in the liquid phase it is known that, for sufficiently large J_z , dominant superconducting correlations can take place [41]. On the other hand, the bosonization analysis reported in [31] predicts topological features for such liquid in both the conducting and superconducting regimes: both should be hosted in a topological Haldane ground state, with a nonvanishing spin

string order correlation function and fractionalized spins at the edges. Thus the integrability of the model allows us to exactly derive the values of these topological quantities. In addition to this, we investigate by density-matrix renormalization-group (DMRG) [42] numerical analysis further quantities to enforce our findings related to the topological aspects of the model. As a first countercheck, we capture the constant finite value of the string order parameter as well as a doubly degenerate entanglement spectrum [3,7,43,44], which are both characteristic signatures of symmetry-protected topological states. Then we concentrate on the effect of a finite magnetic field that enriches the complexity of the treated system. Our numerical simulations show that for a weak magnetic field, the system remains in the sector of total vanishing magnetization. On the other hand, stronger fields produce a finite magnetization, namely a higher number of particles in one spin state than in the other. Although the system remains conducting, it forms a domain-wall structure composed by distinct regions with different features: a fully polarized Luttinger liquid (FPLL) of particles in just one internal state, and a Haldane topological region with vanishing total magnetization. For larger couplings, the latter phase is replaced by an insulating antiferromagnetic Ising domain with effective unit density. Finally, by means of time-dependent DMRG [45,46] we show how such topological states can be dynamically generated and stabilized by performing an expansion procedure, which is a well-established technique in cold atomic experiments; see, for instance, [47,48].

II. MODEL

As shown in [38,39], dressed Rydberg atoms interact via an Ising antiferromagnetic coupling. Due to the large lifetime of these excitations, tunneling processes at nonunit fillings have to be taken into account. In the case of nearest-neighbor coupling, which has been shown to be an accurate approximation in very similar models [31,49], these features are described by the so called t - J_z model,

$$H = -t \sum_{i,\sigma} (c_{i,\sigma}^\dagger c_{i+1,\sigma} + \text{H.c.}) + \sum_i (J_z S_i^z S_{i+1}^z + \delta S_i^z) \quad (1)$$

describing a system of $N = N_\uparrow + N_\downarrow$ particles [50] loaded onto L sites, with total density $n = N/L$. In particular, $c_{i,\sigma}^\dagger$ creates a particle in the internal or, analogously, spin state σ at the i th site and $S_i^z \doteq (n_{i,\uparrow} - n_{i,\downarrow})$. Besides $t = 1$, which fixes our energy and time scale and characterizes the hopping processes of a particle tunneling in a nearest-neighbor (NN) site, the other coupling constant J_z describes antiferromagnetic exchange in the z plane, and δ has the role of a magnetic field which fixes the total magnetization $S_{\text{tot}}^z = \sum_i S_i^z$. In addition to the fact that the aforementioned parameters are independently tunable, we also impose that double occupancies are strictly forbidden, which is a regime easily achievable by using Feshbach resonances.

Interestingly, the t - J_z Hamiltonian at $\delta = 0$ was introduced decades ago to model the behavior first of heavy-fermion compounds and later of high- T_c superconducting materials




FIG. 1. Phase diagram of the t - J_z model Eq. (1) with $\delta = 0$ as a function of the filling n and J_z . Here the topological conducting phase ($J_z < 2$) is denoted as a Haldane liquid (HL), hosting in particular a Haldane triplet superconducting regime (HTS; see text). The state with $J_z > 2$ is characterized by phase separation into an Ising antiferromagnet and empty sites, and is denoted as IPS. The illustrations reproduce a generic state in the HL and IPS phases; the three possible values of S_i^z , namely $+1$, 0 , and -1 are depicted as \uparrow , 0 , and \downarrow , respectively. The shorter arrows at the edges emphasize the fractional edge spins, in this case $+n$ at left and $-n$ at right. All the interaction parameters are in units of the hopping strength t .

in the regime of strong interaction. In particular, it allows us to emphasize how restricted charge fluctuations induce further spin correlations in these systems (see [40] and references therein) away from half-filling, where the system is conducting. While spin and charge structure factors in the ground state, as well as correlation functions of other local quantities, have been studied thoroughly, so far very little attention has been given to the topological aspects of the model. Recently, these have been unveiled in the context of the numerical investigation of a more general model, namely the t - J_z - J_\perp model [31]. There the results, also supported by a bosonization and renormalization-group analysis, identified in the weak J_\perp regime the presence of Haldane ordering of the spin degrees of freedom. This is captured by a nonvanishing value in the thermodynamic limit of solely the spin string order parameter, namely

$$O_S^z(r) \doteq -\langle S_i^z e^{i\pi \sum_{j=1}^{r-1} S_{i+j}^z S_{i+r}^z} \rangle. \quad (2)$$

A nonzero value of the above quantity can be recognized to describe the alternation along the chain of the spins of the particles, diluted in a background of empty sites, as depicted in Fig. 1.

The ground-state energy of the t - J_z Hamiltonian Eq. (1) at $\delta = 0$, and some of its properties, have been derived exactly in [41]. There it was noticed that, since doubly occupied sites are forbidden, singly occupied sites are strictly antiferromagnetically ordered at any $J_z > 0$. In this way, the problem can be mapped into that of finding the ground state of an equivalent attractive spinless fermions model. More precisely, the mapping amounts to replacing in the kinetic term $c_{i\sigma} \rightarrow f_i$, f_i being a spinless fermion annihilation operator. As for the Ising-like interaction term, one realizes that it is equivalent

to an attractive density-density term: $S_i^z S_{i+1}^z \rightarrow -n_i^f n_{i+1}^f$, with $n_j^f = f_j^\dagger f_j$. The resulting attractive t - V model ($V = -J_z$) reads

$$H = -t \sum_i (f_i^\dagger f_{i+1} + \text{H.c.}) + V \sum_i n_i^f n_{i+1}^f, \quad (3)$$

whose $T = 0$ phase diagram is well known. For $V < 0$ ($J_z > 0$) it is characterized by two critical values: when $J_z > 2$, a phase-separated state occurs, where the singly occupied sites separate from the empty ones, whereas for $J_z < 2$ the ground state is a liquid, with dominant superconducting correlations at large distance when the Luttinger exponent $K_c > 1$, i.e., for $J_z^c < J_z < 2$. In other words, this regime is characterized by the fact that the triplet superconducting correlation function

$$O_{\text{TS}}(r) = \langle C_{\text{TS}}^\dagger(i) C_{\text{TS}}(i+r) \rangle, \quad (4)$$

where $C_{\text{TS}}^\dagger(i) = \frac{1}{\sqrt{2}}(c_{i,\uparrow}^\dagger c_{i+1,\downarrow}^\dagger + c_{i,\downarrow}^\dagger c_{i+1,\uparrow}^\dagger)$ goes to zero with r slower with respect to the other correlation functions having a power-law decay. Relevantly, both K_c and J_z^c can be computed exactly by the Bethe ansatz [51,52]. In particular, at filling $n = 1/2$, one gets

$$K_c = \frac{\pi}{4(\pi - \mu)} \quad (5)$$

with $\cos \mu = -\frac{J_z}{2t}$.

III. GROUND-STATE TOPOLOGICAL PROPERTIES AT $\delta = 0$

Exploiting the rigorous results of the previous section on the phase diagram of the t - J_z model at $\delta = 0$, we can derive exactly the nonvanishing value of the spin string parameter (2) and other topological properties. In fact, restoring in the ground state of the above t - V model [Eq. (3)] the alternation of spin orientation of the fermions, one realizes that the phases must also be spin-gapped. In agreement with the general bosonization analysis reported in a previous work [31], the spin gap should be characterized by a nonvanishing value of the Haldane spin string order Eq. (2).

Since in the ground state the spins of the occupied sites are strictly alternating, the string at the exponent in (2) is either 0, in the case in which the two spins at the edges are opposite, or ± 1 if they are parallel. Thus we can rewrite the Haldane string order parameter as

$$O_S^s(r) = \langle (-)^{\frac{1}{2}(S_i^z - S_{i+r}^z)} S_i^z S_{i+r}^z \rangle. \quad (6)$$

The right-hand side here is such that contributions to the expectation value coming from opposite spins at the edges are identical to those coming from parallel spins. Thus $O_S^s(r) \equiv \langle |S_i^z| |S_{i+r}^z| \rangle$. The quantity can now be expressed in terms of the density-density correlation function [53] of the spinless t - V model (3). Explicitly, since the average density n of the spinless fermions is the same as that of the spinful liquid, in the thermodynamic limit we have

$$O_S^s(r) \doteq \lim_{r \rightarrow \infty} \langle n_i^f n_{i+r}^f \rangle = n^2. \quad (7)$$

On general grounds, in the case of Haldane spin string order, for an open chain of length L one expects two degenerate distinct ground states, $|\psi\rangle_\pm$. They are characterized

by opposite fractional spins accumulated at the two edges [21], as schematically shown in the illustration of Fig. 1. Its value for the Haldane ground state of the t - J_z model again can be calculated using the correspondence with the t - V model. Indeed at zero total magnetization, for each of the two degenerate topological states, the probability of having a particle with spin ± 1 in one of the system edges ($i = 1$ or L) is equal to the probability of having a spinless fermion, so that

$$\langle S_{1/\pm}^z \rangle = \pm \langle n_1^f \rangle = \pm n = -\langle S_L^z \rangle_\pm, \quad (8)$$

which is in fact nonvanishing and smaller than 1 at any filling $n \neq 1$ and thus fractionalized. Also, manifestly, the value of each of the two edge spins is different from the bulk magnetization and correlated to the other.

The aforementioned results are summarized in Fig. 1. In particular, the latter shows a more complete description of the phase diagram reported in [41] where the topological properties of Eq. (1), as already specified, were unknown.

More precisely, away from unit density we have three possible different phases. For $J_z < 2$ we observe a phase with no charge gap and a finite value in the thermodynamic limit of the spin string parameter given by (7), thus being a liquid phase with topological features that we denote as the Haldane liquid (HL) phase. In fact, the HL phase hosts two regimes: for $K_c < 1$ (i.e., $J_z < J_z^c$) it has dominant spin-spin correlations

$$C_s(r) = \langle S_i^z S_{i+r}^z \rangle - \langle S_i^z \rangle \langle S_{i+r}^z \rangle, \quad (9)$$

whereas for $K_c > 1$ (i.e., $J_z^c < J_z < 2$) it has dominant triplet pair-pair correlations (4). To emphasize its superconducting properties, we denote this latter regime of the HL phase as a Haldane triplet superconductor (HTS). Finally, for $J_z > 2$, where the spinless t - V model Eq. (3) enters the phase-separated state, correspondingly in the t - J_z model particles get separated from empty sites and a phase-separated antiferromagnetic Ising phase (IPS) takes place.

It is worth mentioning that the above HL phase is an example of a larger class of one-dimensional conducting phases of fermions characterized by nonlocal order in the spin channel. The most renowned is the Luther Emery phase, entered when spin-parity order becomes finite [20,21]. It was recently observed that parity orders can be generalized to two-dimensional arrays [54,55], in particular capturing the presence of a phase qualitatively similar to the Luther Emery liquid also in this case. On the contrary, it is expected that the topological nature the HL phase severely limits the possible generalization of string orders to higher dimension [56].

IV. DMRG ANALYSIS

To support the analytical results of the previous paragraph, here we employ the DMRG technique. Except when explicitly mentioned, in all the performed simulations we use up to 768 DMRG states and perform up to six finite-size sweeps, which ensure convergence with a truncation error smaller than 10^{-9} . As a first step, we evaluate the decay of the string correlation Eq. (2) when moving from the topological phase to the phase-separated regime. As is visible in Fig. 2, we find that this quantity is constant in the phase where we predicted the presence of fractionalized edge states, and its value is already




FIG. 2. Upper panel: decay of $O_S^s(r)$ for different J_z and $\delta = 0$ with $L = 81$, $N = (L + 1)/2$ particles and keeping only the central sites of the system. To pin the Ising domain in the right part of the lattice, we also include weak antiparallel magnetic fields at the systems edges. Lower panel: values of the entanglement spectrum for different J_z and $\delta = 0$ with $L = 81$ and $N = (L + 1)/2$ particles. All the interaction parameters are in units of the hopping strength t .

quite close to the one expected in the thermodynamic limit (7). On the other hand, at the transition point $J_z = 2$ this peculiar behavior is not present anymore. Moreover, once the system enters the phase-separated state the string approaches the value $O_S^s = 1$ in the region where true Ising antiferromagnetic order occurs while it goes to zero in the region where only empty sites are present.

A further probe signaling Haldane topological orders is the entanglement spectrum, which corresponds to the eigenvalues λ_j^N of the reduced density matrix $\rho_A = \sum_{N_j} \lambda_j^N \rho_j^N$ with respect to some system bipartition A , where ρ^N describes a pure state of N particles. In particular, it is known that in a topological phase the entanglement spectrum has to show even degeneracy in the lower λ_j^N 's [3,7,43,44]. Figure 2 indeed confirms that in the phase where edge states are predicted, we get a perfect twofold degeneracy. At the transition point the degeneracy is not accurate anymore and it totally disappears in IPS.

V. EFFECT OF FINITE MAGNETIZATION

Finite values of δ strongly enlarge the richness of the aforementioned phase diagram (Fig. 1). Intuitively, the role of a positive magnetic field is to make it energetically more favorable for the system to have more particles in one internal state than in the other, say $N_\downarrow > N_\uparrow$ with $N_\sigma = \sum_i n_{i\sigma}$. Of course, this contrasts with the effect of J_z , which for $N_\uparrow = N_\downarrow$ can maximize the number of antiferromagnetic surfaces. As shown in Fig. 3, our results indicate that for weak δ 's the system still prefers the solution with vanishing total magnetization, thus supporting the presence of the same phases as in Fig. 1, where for $J_z < 2$ the Haldane topological orders extend all over the lattice sites and for $J_z > 2$ IPS takes




FIG. 3. Density distribution for different values of δ at fixed $J_z = 1.5$, $L = 80$, and $N = L/2$. In all cases, a weak positive local magnetic field $\mu = 0.01$ is applied in the system center to lift the GS degeneracy. All the interaction parameters are in units of the hopping strength t .

place. An unbalanced phase with $S_{\text{tot}}^z \neq 0$ occurs only above a critical value of the magnetic field, almost independent of the strength of J_z . In this regime, as the example reported in the central panel of Fig. 3 shows, the system prefers to arrange into a domain-wall structure composed of two regions: one with $S_{\text{tot}}^z = 0$ and the other being a fully polarized Luttinger liquid. More precisely, we get that, for $J_z < 2$, the whole system remains conducting hosting a uniform finite amount of holes, though spatially distinguished regions with different magnetization can be noticed: one with $N_\uparrow = N_\downarrow$ and the other fully polarized ($N_\uparrow = 0$).

Both coexisting states are the ground state of an effective Hamiltonian of the form of the spinless t - V Hamiltonian Eq. (3) with one crucial difference. For the fully polarized state, besides an additional constant term, the effective interaction is now repulsive ($V = J_z > 0$). Thus the region amounts to a fully polarized Luttinger liquid with $K_c < 1$. The unpolarized region instead preserves its topological features since the effective model capturing such a domain is exactly Eq. (1). At $n = 0.5$, both states have the same filling of the uniform phase: the spatial extent of each of the two is determined by the strength of δ . The central panel in Fig. 3 indeed shows the presence of a central region with filling lower than 1 and diluted antiferromagnetic order; for the parameter value reported in the figure, it turns out to be an HTS region (HL in the case of lower J_z), whereas in the left and right parts two FPLs are seen. A further increase in δ makes the presence of \uparrow particles too energetically costly, and a fully polarized liquid with $N_\downarrow = N$ and $N_\uparrow = 0$ occurs in the whole lattice. To prove that the lattice part with vanishing magnetization is described by Eq. (1), we show in Fig. 4 the behavior of the correlation functions evaluated in such a region. In particular, on the one hand it is possible to notice that $O_S^s(r)$ is always




FIG. 4. Decay of $O_S^z(r)$, $O_{TS}(r)$, and $C_S(r)$ for $\delta = 0.4$, $L = 80$, and $N = L/2$ for two different values of J_z . The correlations are evaluated in the central region of the system where $N_\uparrow = N_\downarrow$. In both cases, a weak positive local magnetic field $\mu = 0.01$ is applied in the system center to lift the GS degeneracy. All the interaction parameters are in units of the hopping strength t .

constant, thus confirming the topological nature of the domain with $N_\uparrow = N_\downarrow$. On the other hand, we find that the triplet superconducting correlation function becomes the leading order only when $J_z^c < J_z < 2$, thus confirming the validity of our analysis.

The situation becomes different when a Ising coupling $J_z > 2$ is considered. Indeed, as known from the study of the $\delta = 0$ case, such a strong interaction destroys the Haldane order in favor of a true Ising antiferromagnet where particles and holes are totally demixed. In this context, the magnetic field plays a similar role as in the previous case. In particular, for weak δ 's the system remains balanced, whereas above a critical value δ_c a domain-wall structure composed of an Ising antiferromagnet and a FPLL is obtained. Moreover, as in the already discussed scenario, large magnetic fields drive the system in a regime where only \downarrow particles are present, and due to the specific choice of the considered density $n = 0.5$ a gapped phase with density wave (DW) order takes place. It is relevant to underline that for any other filling, the DW is substituted by a LL as in the previous case.

The above-discussed behavior is summarized in the phase diagram shown in Fig. 5. Here it is possible to notice that a weak magnetic field does not alter the phases of Fig. 1 until a critical value of δ where a regime of phase separation with finite total magnetization takes place. Here, in addition to a fully polarized region of particles behaving as a LL, as a function of the strength of J_z three possible phases can appear: a Haldane liquid (HL), a Haldane phase with dominant superconducting order (HTS), and an Ising antiferromagnet (Ising). On the other hand, larger values of J_z and δ give rise, due to the specific considered density, to an insulating DW state. We observe that the slope of the transition line to either FPLL or DW (green line) increases with J_z , and in fact it is expected to approach the value 2 in the strong-coupling limit. Moreover, we underline that for higher total fillings and $J_z < 2$, domain-wall structures composed of both Haldane orders and DW domains can occur.




FIG. 5. Phase diagram of the t - J_z model equation (1) as a function of δ and J_z with density $n = 0.5$ and $L = 80$. Here also regions characterized by phase-separated states are reported: the + symbol is used to emphasize this aspect. All the interaction parameters are in units of the hopping strength t .

VI. DYNAMICAL STABILIZATION OF THE HALDANE STATES

The above Haldane conductors can be dynamically stabilized in experiments involving dressed Rydberg states. Our starting point is precisely the state that has already been reached in such experiments, namely an Ising antiferromagnet with exactly one particle per lattice site. Due to the infinite on-site repulsion between particles in such an Ising state, tunneling processes are not energetically possible. To make Hamiltonian Eq. (1) a correct description of such a setup, holes, having the role of restoring the hopping processes, have to be injected in the system. A way to achieve that is to use optical tweezers [57], which can selectively remove atoms one by one in the lattice and hence generate empty sites. Another way to make the Rydberg excitations move along the lattice is to simply let the Ising state expand. In particular, our employed strategy is to prepare an initial state where all the atoms are compressed in the central part of the lattice by a strong harmonic confinement, which, in this region, excludes the presence of empty sites such as, for instance, $|00 \dots \uparrow \downarrow \uparrow \downarrow \dots 00\rangle$.

Due to the finite value of J_z , a perfect Ising antiferromagnet can be reached in the central part of the lattice. Once this state has been prepared, we remove the harmonic confinement. This has the effect of letting the atomic cloud expand and thus changing the effective particle density in the lattice. In this way, the system will equilibrate in a state whose physics will be captured by Eq. (1). To understand the validity of such an approach, we monitor the evolution in time t of the string order parameter Eq. (2) and of the spin-spin correlation function C_S . The latter should remain finite only when true Ising antiferromagnetism is present. Figure 6, where for simplicity we kept $\delta = 0$, shows indeed that such a procedure is able to stabilize Haldane conductors. In particular, in the upper panel the value $J_z = 0.5$ supports the presence of a topological state and clearly our results demonstrate that the string remains finite during the whole evolution and stabilizes to a




FIG. 6. Upper panel: Time evolution of C_s and O_s^s in a system with $L = 40$ and $N = L/2 + 1$ particles and starting with an initial state with fixed J_z and a strong harmonic potential to confine the particles in the central part of the system. The evolution is performed by keeping the same value of J_z and removing the harmonic potential. Both quantities are evaluated in the central 10 sites of the systems. The numerical simulations are performed by keeping up to 512 DMRG states in the evolution and a time step $\delta t = 0.01$. All the interaction parameters and the time are in units of the hopping strength t .

nonvanishing value. On the other hand, C_s displays oscillations and after a certain time reaches a constant vanishing value. This last feature changes when larger J_z are considered. Indeed, the lower panel of Fig. 6 shows the time-dependent expectation values of both the considered quantities as before but for an Ising coupling $J_z = 3$, which supports the presence of a true antiferromagnet. Here clearly both C_s and O_s^s remain finite during the evolution, confirming the fact that the system will thermalize in a phase-separated state with an Ising domain in its central part (IPS).

VII. CONCLUSIONS

We have shown that Rydberg dressed atoms trapped onto a one-dimensional optical lattice represent ideal candidates to probe Haldane topological orders with conducting features. In particular, for a vanishing magnetic field, the phase diagram can be derived analytically and the topological properties are extracted in an exact way. At the same time, our numerical results allow us to both confirm the analytically predicted topological nature of the studied model, and to investigate other peculiar cases. When a finite magnetic field is applied, we observed that the system is arranged in a domain-wall structure. For not too large J_z , the domains turn out to be composed of a fully polarized Luttinger liquid and by a Haldane topological state whose conducting or superconducting properties are determined by the strength of the antiferromagnetic coupling, whereas at larger J_z the Haldane state is replaced by an Ising domain with true antiferromagnetic order. Moreover, strong enough magnetic fields destroy the domain structure, and, depending on the antiferromagnetic coupling, either homogeneous Luttinger liquids or, in specific cases, a density wave state occurs. We also demonstrate that a protocol based on atomic expansion can be experimentally employed to dynamically stabilize such Haldane conductors. The present results enrich our understanding of interaction-induced symmetry-protected topological states of matter, giving a reliable procedure to experimentally reach and study such features with the currently available experimental platforms.

Note added. Recently, we became aware of recent unpublished work in which topological aspects of the t - J_z model are analyzed [58].

ACKNOWLEDGMENTS

The authors thank I. Bloch, C. Gross, F. Grusdt, and T. Macrì for interesting discussions. L.B. acknowledges ERC Starting Grant TopoCold for financial support.

- [1] Z.-C. Gu and X. G. Wen, *Phys. Rev. B* **80**, 155131 (2009).
- [2] X. Chen, Z.-C. Gu, Z.-X. Liu, and X.-G. Wen, *Science* **338**, 1604 (2012).
- [3] F. Pollmann, E. Berg, A. M. Turner, and M. Oshikawa, *Phys. Rev. B* **85**, 075125 (2012).
- [4] Z.-X. Liu and X.-G. Wen, *Phys. Rev. Lett.* **110**, 067205 (2013).
- [5] X. G. Wen, *Phys. Rev. D* **88**, 045013 (2013).
- [6] A. Vishwanath and T. Senthil, *Phys. Rev. X* **3**, 011016 (2013).
- [7] L. Fidkowski and A. Kitaev, *Phys. Rev. B* **83**, 075103 (2011).
- [8] F. Pollmann, A. M. Turner, E. Berg, and M. Oshikawa, *Phys. Rev. B* **81**, 064439 (2010).
- [9] F. D. M. Haldane, *Phys. Lett. A* **93**, 464 (1983); *Phys. Rev. Lett.* **50**, 1153 (1983).
- [10] M. den Nijs and K. Rommelse, *Phys. Rev. B* **40**, 4709 (1989).
- [11] I. Affleck, T. Kennedy, E. H. Lieb, and H. Tasaki, *Phys. Rev. Lett.* **59**, 799 (1987); *Commun. Math. Phys.* **115**, 477 (1988).
- [12] L. Barbiero, L. Dell'Anna, A. Trombettoni, and V. E. Korepin, *Phys. Rev. B* **96**, 180404(R) (2017).
- [13] E. G. Dalla Torre, E. Berg, and E. Altman, *Phys. Rev. Lett.* **97**, 260401 (2006).
- [14] E. Berg, E. G. Dalla Torre, T. Giamarchi, and E. Altman, *Phys. Rev. B* **77**, 245119 (2008).
- [15] D. Rossini and R. Fazio, *New J. Phys.* **14**, 065012 (2012).
- [16] G. G. Batrouni, R. T. Scalettar, V. G. Rousseau, and B. Grémaud, *Phys. Rev. Lett.* **110**, 265303 (2013).
- [17] H. Nonne, P. Lecheminant, S. Capponi, G. Roux, and E. Boulat, *Phys. Rev. B* **81**, 020408(R) (2010).
- [18] A. Montorsi and M. Roncaglia, *Phys. Rev. Lett.* **109**, 236404 (2012).
- [19] K. Kobayashi, M. Okumura, Y. Ota, S. Yamada, and M. Machida, *Phys. Rev. Lett.* **109**, 235302 (2012).
- [20] L. Barbiero, A. Montorsi, and M. Roncaglia, *Phys. Rev. B* **88**, 035109 (2013).
- [21] A. Montorsi, F. Dolcini, R. C. Iotti, and F. Rossi, *Phys. Rev. B* **95**, 245108 (2017).

- [22] M. Endres, M. Cheneau, T. Fukuhara, C. Weitenberg, P. Schauß, C. Gross, L. Mazza, M. C. Banuls, L. Pollet, I. Bloch, and S. Kuhr, *Science* **334**, 200 (2011).
- [23] T. A. Hilker, G. Salomon, F. Grusdt, A. Omran, M. Boll, E. Demler, I. Bloch, and C. Gross, *Science* **357**, 484 (2017).
- [24] I. Bloch, J. Dalibard, and W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
- [25] T. Lahaye, C. Menotti, L. Santos, M. Lewenstein, and T. Pfau, *Rep. Prog. Phys.* **72**, 126401 (2009).
- [26] M. Dalmonte, M. Di Dio, L. Barbiero, and F. Ortolani, *Phys. Rev. B* **83**, 155110 (2011).
- [27] X. Deng and L. Santos, *Phys. Rev. B* **84**, 085138 (2011).
- [28] S. Fazzini, A. Montorsi, M. Roncaglia, and L. Barbiero, *New J. Phys.* **19**, 123008 (2017).
- [29] S. Fazzini, L. Barbiero, and A. Montorsi, *J. Phys.: Conf. Ser.* **841**, 012016 (2017).
- [30] L. Barbiero, S. Fazzini, and A. Montorsi, *Eur. Phys. J. Spec. Top.* **226**, 2697 (2017).
- [31] S. Fazzini, L. Barbiero, and A. Montorsi, *Phys. Rev. Lett.* **122**, 106402 (2019).
- [32] L. De Marco, G. Valtolina, K. Matsuda, W. G. Tobias, J. P. Covey, and J. Ye, *Science* **363**, 853 (2019).
- [33] B. Yan, S. A. Moses, B. Gadway, J. P. Covey, K. R. A. Hazzard, A. M. Rey, D. S. Jin, and J. Ye, *Nature (London)* **501**, 521 (2013).
- [34] P. Schauß, M. Cheneau, M. Endres, T. Fukuhara, S. Hild, A. Omran, T. Pohl, C. Gross, S. Kuhr, and I. Bloch, *Nature (London)* **491**, 87 (2012).
- [35] H. Bernien *et al.*, *Nature (London)* **551**, 579 (2017).
- [36] S. de Léséleuc, V. Lienhard, P. Scholl, D. Barredo, S. Weber, N. Lang, H. P. Büchler, T. Lahaye, and A. Browaeys, *Science* **365**, 775 (2019).
- [37] A. Omran *et al.*, *Science* **365**, 570 (2019).
- [38] J. Zeiher, R. van Bijnen, P. Schauß, S. Hild, J.-Y. Choi, T. Pohl, I. Bloch, and C. Gross, *Nat. Phys.* **12**, 1095 (2016).
- [39] J. Zeiher, J.-Y. Choi, A. Rubio-Abadal, T. Pohl, R. van Bijnen, I. Bloch, and C. Gross, *Phys. Rev. X* **7**, 041063 (2017).
- [40] S. Zhang, M. Karbach, G. Müller, and J. Stolze, *Phys. Rev. B* **55**, 6491 (1997).
- [41] C. D. Batista and G. Ortiz, *Phys. Rev. Lett.* **85**, 4755 (2000).
- [42] S. R. White, *Phys. Rev. Lett.* **69**, 2863 (1992).
- [43] A. M. Turner, F. Pollmann, and E. Berg, *Phys. Rev. B* **83**, 075102 (2011).
- [44] L. Barbiero, L. Santos, and N. Goldman, *Phys. Rev. B* **97**, 201115(R) (2018).
- [45] S. R. White and A. E. Feiguin, *Phys. Rev. Lett.* **93**, 076401 (2004).
- [46] A. E. Feiguin and S. R. White, *Phys. Rev. B* **72**, 020404(R) (2005).
- [47] S. Trotzky, Y.-A. Chen, A. Flesch, I. P. McCulloch, U. Schollwöck, J. Eisert, and I. Bloch, *Nat. Phys.* **8**, 325 (2012).
- [48] J. P. Ronzheimer, M. Schreiber, S. Braun, S. S. Hodgman, S. Langer, I. P. McCulloch, F. Heidrich-Meisner, I. Bloch, and U. Schneider, *Phys. Rev. Lett.* **110**, 205301 (2013).
- [49] A. V. Gorshkov, S. R. Manmana, G. Chen, J. Ye, E. Demler, M. D. Lukin, and A. M. Rey, *Phys. Rev. Lett.* **107**, 115301 (2011).
- [50] Here we consider particles with fermionic statistics, but exactly the same results are expected in the case of a two-component Bose gas with hard-core on-site repulsion.
- [51] F. D. M. Haldane, *Phys. Lett. A* **81**, 153 (1980).
- [52] S. Qin, M. Fabrizio, L. Yu, M. Oshikawa, and I. Affleck, *Phys. Rev. B* **56**, 9766 (1997).
- [53] A. Luther and I. Peschel, *Phys. Rev. B* **12**, 3908 (1975).
- [54] S. Fazzini, F. Becca, and A. Montorsi, *Phys. Rev. Lett.* **118**, 157602 (2017).
- [55] L. Tocchio, F. Becca, and A. Montorsi, *SciPost Phys.* **6**, 018 (2019).
- [56] F. Anfuso and A. Rosch, *Phys. Rev. B* **75**, 144420 (2007).
- [57] N. Schlosser, G. Reymond, I. Protsenko, and P. Grangier, *Nature (London)* **411**, 1024 (2001).
- [58] T. Rakovszky, P. Sala, R. Verresen, M. Knap, and F. Pollmann, *Phys. Rev. B* **101**, 125126 (2020).