

Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: Effects on tomato plants growth and on soil properties

Original

Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: Effects on tomato plants growth and on soil properties / Cristina, G., Camelin, E., Tommasi, T., Fino, D., Pugliese, M.. - In: JOURNAL OF ENVIRONMENTAL MANAGEMENT. - ISSN 0301-4797. - ELETTRONICO. - 269:(2020), p. 110767. [10.1016/j.jenvman.2020.110767]

Availability:

This version is available at: 11583/2833534 since: 2020-06-08T10:26:20Z

Publisher:

Elsevier

Published

DOI:10.1016/j.jenvman.2020.110767

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.jenvman.2020.110767>

(Article begins on next page)

Manuscript Number: JEMA-D-19-07269R2

Title: Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: effects on tomato plants growth and on soil properties

Article Type: Research Article

Keywords: Waste management; nitrogen; soil organic matter; nutrient recycling; sewage sludge; tomato plants.

Corresponding Author: Dr. Tonia Tommasi, Ph.D

Corresponding Author's Institution: Politecnico di Torino

First Author: Giulio Cristina

Order of Authors: Giulio Cristina; Enrico Camelin; Tonia Tommasi, Ph.D; Debora Fino; Massimo Pugliese

Abstract: Anaerobic digestates from sewage sludge (SSADs) are a by-product of the wastewater treatment process that still preserves a certain agronomic interest for its richness in plant nutrients and organic matter. Fertilizing properties of two liquid and two dewatered SSADs were tested on tomato plants (*Solanum lycopersicum* L.). Pot experiments were performed on sandy soil and peat substrate under greenhouse conditions with a SSADs application rate of 170 kg N/ha over a period of three months. Beneficial effects of SSADs were reported on different growth parameters, revealing an increase in biomass and height up to 37.5 and 6-folds over untreated control. No phytotoxic effect occurred on SSAD-exposed plants. Chemical analysis of soils treated with SSADs showed enrichment of macro- and micro-nutrients as well as organic matter. In some cases, the chemical characterization of leaves revealed an enhancement of uptaken macronutrients. This study contributed in general to deepen the knowledge on the short-term growing season fertilizing effects of SSAD. Despite the treatment dosage was calculated only on nitrogen requirements, the study highlighted the importance of the other nutrients and organic matter on plant growth.

1 Title

2 **Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy**
3 **soil and on a peat substrate: effects on tomato plants growth and on soil properties**

4 **Authors**

5 Giulio Cristina^{a†}, Enrico Camelin^{a†}, Tonia Tommasi^{a*}, Debora Fino^a and Massimo Pugliese^{b,c*}.

6 ^a Department of Applied Science and Technology (DISAT), Politecnico di Torino, Corso Duca degli
7 Abruzzi 24, 10129 Torino (TO) – Italy

8 ^b Agricultural, Forestry and Food Science Department (DISAFA), University of Torino, Largo Paolo
9 Braccini 2, 10095 Grugliasco (TO) – Italy

10 ^c AGROINNOVA – Centre of Competence for the Innovation in the Agro-Environmental Sector,
11 University of Torino, Largo Paolo Braccini 2, 10095 Grugliasco (TO) – Italy

12

13 [†] These authors contributed equally to the work

14

15 * Corresponding authors:

16 Tonia Tommasi

17 Phone: +39 011 090 4774; e-mail: tonia.tommasi@polito.it

18 Department of Applied Science and Technology (DISAT), Politecnico di Torino

19 Corso Duca degli Abruzzi 24, 10129 Torino (TO) – Italy

20

21 Massimo Pugliese

22 Phone: + 39 011 670 8545; email: massimo.pugliese@unito.it

23 Agricultural, Forestry and Food Science Department (DISAFA), University of Torino

24 Largo Paolo Braccini 2, 10095 Grugliasco (TO) – Italy

25

26 **Declaration of Interest:**

27 Declaration of Interest: None

28 **Abstract**

29 Anaerobic digestates from sewage sludge (SSADs) are a by-product of the wastewater treatment
30 process that still preserves a certain agronomic interest for its richness in plant nutrients and organic
31 matter. Fertilizing properties of two liquid and two dewatered SSADs were tested on tomato plants
32 (*Solanum lycopersicum* L.). Pot experiments were performed on sandy soil and peat substrate under
33 greenhouse conditions with a SSADs application rate of 170 kg N/ha over a period of three months.
34 Beneficial effects of SSADs were reported on different growth parameters, revealing an increase in
35 biomass and height up to 37.5 and 6-folds over untreated control. No phytotoxic effect occurred on
36 SSAD-exposed plants. Chemical analysis of soils treated with SSADs showed enrichment of macro-
37 and micro-nutrients as well as organic matter. In some cases, the chemical characterization of leaves
38 revealed an enhancement of uptaken macronutrients. This study contributed in general to deepen the
39 knowledge on the short-term growing season fertilizing effects of SSAD. Despite the treatment
40 dosage was calculated only on nitrogen requirements, the study highlighted the importance of the
41 other nutrients and organic matter on plant growth.

42
43 **Keywords:** waste management; nitrogen; soil organic matter; nutrient recycling; sewage sludge;
44 tomato plants.

45

1. Introduction

Globally, the demand of the three primary plant nutrients used for soil fertilization (N, P_2O_5 and K_2O) is increasing (Vanotti et al., 2019). In 2015, the total fertilizer nutrient demand was around 184 Mt and, by the end of 2020, it is expected to overcome 200 Mt (FAO, 2017). The production processes of these fertilizers are very expensive in terms of energy (ammonia) and non-renewable resources (phosphorus and potassium), with heavy environmental costs (Li et al., 2009). Ammonia production is mainly performed via the Haber-Bosch process which requires a large amount of fossil fuel (Basosi et al., 2014). Phosphate rock is the principal raw material exploited in the production of nearly all phosphate fertilizers (Fixen and Johnston, 2012; Reijnders, 2014). This non-renewable resource may contain many toxic heavy metals such as As, Hg, Ni, V (Mortvedt, 1995), Cd, Cr, Cu, Pb, Zn (Sabiha-Javied et al., 2009), fluorine (Mirlean and Roisenberg, 2007) and uranium (Schnug and Lottermoser, 2013). The P_2O_5 extraction can cause environmental pollution by contaminants accumulating in air, soil, and water bodies around the manufacturing place (Mirlean et al., 2008; Sabiha-Javied et al., 2009). It has been observed that these impurities can persist into phosphate fertilizers, provoking a subsequent accumulation in agricultural soils (De López Camelo et al., 1997). Potassium derives from non-renewable resources like minerals such as sylvite, sylvinit, hartsalz and langbeinite (Fixen and Johnston, 2012). Furthermore, world distribution of phosphorous and potassium mines is not uniform: 45% of global phosphate rock is concentrated in Morocco and the Western Sahara (Fixen and Johnston, 2012).

Within a circular economy perspective, the reuse of sewage sludge (SS) as fertilizer is an interesting scenario. SS can be defined as “the residue generated from the treatment of wastewater” (Smith et al., 2009). This matrix is a valuable source in terms of plant nutrients: a study conducted on 240 dried samples from Pennsylvania revealed an average N, P and K content of 4.74%, 2.27%, and 0.31%, respectively (Stehouwer et al., 2000). Furthermore, SS can contain many micronutrients (e.g. Ca, Mg,

Abbreviations: A_N : assimilation; ANRE: Apparent Nitrogen Recovery Efficiency; ANUE: Agronomic Nitrogen Use Efficiency; C: centrifuged SSAD; CEC: cation exchange capacity; CCI: Chlorophyll Content Index; C_i : CO_2 concentration in substomatal cavity; CRF: controlled release fertilizer; D: dried SSAD; D.M.: dry matter; EC: electrical conductivity; EDC: endocrine disrupting compounds; EmC: Emerging Contaminants; EU: European Union; g_s : stomatal conductance; IRGA: infra-red gas analyzer; M: mineral fertilizer; OM: organic matter; P: primary SSAD; QL: quantification limits; S: secondary SSAD; SS: sewage sludge; SSAD: anaerobic digestate from sewage sludge; T: non-treated, control thesis; WWTP: wastewater treatment plant.

70 S, Fe, Mn, Cu, Zn and B) which are important for plant growth, but usually not included in most
71 commercial fertilizers (Warman and Termeer, 2005). The percentage of the nutrients appears low, but
72 it is important to underline that every year a huge amount of wastewater is produced. An empirical
73 study revealed that approximately 330 km³ of municipal wastewater are produced worldwide yearly
74 (Mateo-Sagasta et al., 2015). Therefore, also the SS production has dramatically risen thanks to
75 policies dealing with the improvement of wastewater treatment and of standard quality of effluents,
76 such as the E.U. directive 91/271/EEC (Council of the European Communities, 1991a).

77 The considerable presence of organic carbon and organic matter in SS is another strength of its reuse
78 (Alvarenga et al., 2015; Mateo-Sagasta et al., 2015). In fact, land application of organic matter (OM)
79 improves soil physical properties such as cation exchange capacity (CEC), soil structure, soil
80 moisture content and retention (Epstein, 2002). Furthermore, the addition of SS can enhance the
81 amount of organic carbon in soils (Kladivko and Nelson, 1979; Perez-Espinosa et al., 1999) and thus
82 reverse the current reduction of organic matter in soils (known as *SOM decline*) (Schulze and
83 Freibauer, 2005).

84 Today, SS is classified as waste and its safe disposal represents a very important issue in waste
85 management (Epstein, 2002; Singh and Agrawal, 2008). The four main destinations of SS are
86 incineration, landfilling, composting and agricultural use. In Italy, according to Eurostat data (Eurostat,
87 2019), the majority of SS is sent to landfill (50.8%), while 34.7% is reused in agriculture, 4% is
88 incinerated and 10.4% is sent to other destinations. The Council Directive 86/278 (Council of the
89 European Communities, 1986) regulates the agricultural SS reuse in Europe to prevent soil
90 contamination. In fact, this practice has three principal problems that limit its unconditioned use:
91 biological risk, heavy metal contamination and contamination by organic pollutants. The biological risk
92 is principally represented by pathogens such as *Salmonella* spp., *Escherichia coli* (enterotoxigenic
93 and enteropathogenic variants), *Campylobacter* spp., *Clostridium* spp., and *Yersinia* spp. (Arthurson,
94 2008); stabilization treatments can reduce significantly their presence in SS and are mandatory before
95 subsequent SS applications (Dumontet et al., 1999). For instance, one of the most diffused
96 stabilization techniques is anaerobic digestion (Liu et al., 2012), in which the reduction of pathogens,
97 putrescence and odor is coupled with biogas production, allowing energy recovery (Epstein, 2002).

98 Heavy metal content (normally represented by Cd, Cr, Cu, Hg, Ni, Pb, Zn) can be abated by means of
99 chemical (e.g. chelating addition), physical (e.g. electroremediation) or biological (e.g.
100 vermicomposting) treatments (Camargo et al., 2016). Finally, some organic pollutants (e.g. pesticides,
101 antibiotics and hormones) can be volatilized or degraded through biotic or abiotic processes (Harrison
102 et al., 2006). Concerning organic pollutants, their abatement is trickier. Indeed, class of emerging
103 contaminants (EmC) in wastewater is increasingly gaining more interest within the organic
104 compounds. EmC include molecules such as endocrine disrupting compounds (EDC, e.g. hormones),
105 pharmaceutically active compounds (e.g. antibiotics), illicit drugs and pesticides (Fijalkowski, 2019).
106 EmC abatement is becoming even more required both on the effluent of WWTPs with advanced
107 treatments (e.g. activated carbon absorption, advanced oxidation processes, reverse osmosis) and on
108 sewage sludge (Gadupudi et al., 2019). Some studies affirmed that anaerobic digestion is the
109 stabilization strategy ensuring the best EmC removal, especially when the sludge is pretreated (e.g.
110 via ozonation) (Neumann et al., 2016). However, further studies are still required to improve the
111 performances and to reduce the costs of these techniques, which nowadays are rarely applied at
112 WWTP level since they are money and/or time consuming (Camargo et al., 2016). The
113 abovementioned EU directive regulates the SS soil application in the EU and establishes threshold
114 values of some of these pathogens and pollutants in SS.

115 On the basis of these opportunities and threats related to SS, this work aims to deepen the
116 knowledge about SS fertilizing effects over time in terms of nutrients and OM on a poor alkaline sandy
117 soil. This kind of soil was selected because: i) nutrient depletion constrains plant growth to depend on
118 treatment application; ii) a high pH both hinders the nutrient adsorption and reduce the metal
119 bioavailability (Alvarenga et al., 2016); iii) sandy-textured soil lacks nutrients and has low water-
120 holding capacity. These results were compared to the one obtained with a richer peat substrate. Pot
121 experiments were performed on tomato plants (*Solanum lycopersicum* L.) in a greenhouse to
122 evaluate nutrient provision of anaerobic digestates from sewage sludge (SSADs). Tomato plant was
123 chosen because: i) it is one of the most exploited vegetables crop (Jones Jr, 2008); ii) there is an
124 increasing interest on alternative nutrient sources for this crop (Zucco et al., 2015); iii) it has a high
125 fertilizer requirements (Zucco et al., 2015); iv) plenty of scientific literature is available for this crop

(Jones Jr, 2008). In this work, no analysis on pathogens was carried out since anaerobic digestion is considered one of the safest technologies for pathogen reduction in SS (Epstein, 2002). Nevertheless, this aspect may be taken into consideration in future researches. Concerning the use of SS in agronomic experiments, a lack in details about SS typology is provided. Indeed, in many work no detail on stabilization strategy is provided (Bakshi et al., 2019), or the kind of SS digestion is not specified (Hossain et al., 2015). In the present work, the digestates used derived from the same WWTP and were obtained with consequent treatments (Cristina et al., 2019). As far as we know, this is the first example of use of four different and consequent SSADs to fertilize tomato plants. The paper examines agronomic parameters of tomato plants, the nutrient distribution in soil and nutrient absorption by plants after the application of SSADs. Furthermore, numerous plants physiological parameters were evaluated over a span of three months in order to better understand the effects in a time course approach.

2. Materials and methods

2.1. Characterizations

2.1.1. Anaerobic digestates from sewage sludge

Four SSADs were used in the experiment: two liquid (primary (P) and secondary (S)) and two dewatered (centrifuged (C) and dried (D)). Physical and chemical characterization of the SSADs is described in a previous work (Cristina et al., 2019); characterization of the four SSADs is reported in **Supplementary Material -Section I (Table S1)**.

2.1.2. Cultivation substrates

Two types of substrates were used: a sandy soil and a commercial peat substrate (**Table 1**). The sandy soil was sampled within 20 and 100 cm depth in Grugliasco (TO), Italy (45°03'58.4"N, 7°35'32.9"E). Analytical methods used for characterization of the sandy soil and the peat substrate are specified in **Supplementary material - Section II**. Based on the distribution of the particle size (sand: 94% ± 2; silt: 3% ± 1; clay: 3% ± 1), the selected soil was classified as sandy (Buol et al., 2011). Based on ARPAV soil analysis (Arpa Veneto, 2007), the soil was considered alkaline (8.2 ±

0.16), very poor in OM ($0.38 \pm 0.12\% < 0.8\%$) and very poor in macronutrients such as nitrogen ($0.29 \pm 0.09 \text{ g/kg} < 0.5 \text{ g/kg}$), phosphorous ($1.8 \pm 1.3 \text{ mg/kg} < 7 \text{ mg/kg}$), potassium ($18 \pm 1 \text{ mg/kg} < 40 \text{ mg/kg}$) and magnesium ($15 \pm 5 \text{ mg/kg} < 50 \text{ mg/kg}$). On the other hand, content of calcium ($675 \pm 27 \text{ mg/kg} < 1000 \text{ mg/kg}$) and some microelements such as iron ($2.5 \text{ mg/kg} < 6.7 \pm 1.1 \text{ mg/kg} < \text{mg/kg } 20$) and manganese ($2 \text{ mg/kg} < 6.5 \pm 3.0 \text{ mg/kg} < \text{mg/kg } 10$) resulted normal.

The peat substrate used consisted of a commercial blend of blond and black peat (15:85, Turco Silvestro, Italy), mixed with perlite (80:20 v/v). The substrate was steamed at 90°C for 30 minutes before use. The substrate had the following characteristics as indicated by the manufacturer: pH 6.1; E.C.: 0.56 dS/m; bulk density 250 kg/m³; C total 175 g kg⁻¹; N total 7 g kg⁻¹; organic matter 32% d.m.; P₂O₅ 10 g kg⁻¹; K₂O 11 g kg⁻¹. Hence, the peat substrate could be reasonably considered a good cultivation substrate, satisfying the requirements as a benchmark to be compared with the poor sandy soil.

2.2. Experimental set-up

A greenhouse experiment was performed over three months during the summer season in a greenhouse of the Centre of Competence AGROINNOVA – University of Torino, located in Grugliasco (TO), Italy. The experimental campaign was carried out with commercial plastic pots of 2.5 L (Ø 17 cm, height 20 cm, surface area 0.227 m²). Four types of SSADs (P, S, C, D) were applied as treatments, and compared to a commercial fertilizer (M) (NPK 22-5-6 + 2MgO, “Osmocote Topdress”, ICL, Israel) and an untreated control (T). The experiment was designed in a completely randomized block, with 15 replications per each thesis. The same experimental set-up was adopted on the two cultivation substrates (sandy soil and peat substrate). Each treatment was applied at the dosage of 170 kg N/ha, in line with the European Nitrates Directive (Council of the European Communities, 1991b). Moreover, this application rate was chosen as it showed the best results in a preliminary study (Cristina et al., 2019). Three untreated seeds of tomato (*Solanum lycopersicum* L. cv. Beefsteak, “Furia sementi”, Parma, Italy) were sown in each pot. Automatic sprinkler irrigation was set three times a day for 2-3 minutes in order to keep 40-50% WHC. Ten days after sowing a thinning was

178 conducted and the best plant from each pot was kept. At the end of each month, five pre-selected
179 replicates of each treatment were removed to carry out all the measurements.

180 2.3. Measurement of plant parameters

181 At the end of every month, the five removed replicates were examined. Firstly, height was measured,
182 then, leaves, inflorescences and fruits were counted, if present. After that, the Chlorophyll Content
183 Index (CCI) was evaluated with a SPAD 502 chlorophyll meter (CCM-200, Opti Sciences, Inc.,
184 Hudson, NH, USA) using the method described in the previous work (Cristina et al., 2019). One
185 month after sowing, it was not possible to measure CCI on the sand specimen because the minimum
186 leaves size was not satisfied. At the end of the second month, assimilation (A_N), stomatal
187 conductance (g_s) and CO_2 concentration in substomatal cavity (C_i) were measured by the means of
188 an Infrared Gas Analyzer (IRGA, ADC, Hoddesdon, UK). These measurements were performed on
189 three fully formed leaves in each replicate. The selected leaves had to be non-senescing, at the same
190 physiological age (in the middle part of the plant, considering the third to fourth leaf from the shoot
191 apex) and directly exposed to sunlight. After all the measurements were taken, each plant was
192 subsequently cut and immediately weighed to record the fresh biomass value. In order to evaluate the
193 mean dry biomass, each plant was dried at 105°C for at least 72 hours. Subsequently, agronomic
194 nitrogen use efficiency (ANUE) was calculated as:

$$ANUE = \frac{(Dry\ biomass\ treated\ samples - Dry\ biomass\ control\ samples)}{Amount\ of\ nitrogen\ applied\ in\ treated\ samples}$$

195 2.4. Chemical analysis

196 Substrates were chemically characterized at the end of the second month, once the aerial plant part
197 had been cut. Chemical analyses were performed on samples from the treatment with SSADs (P, S,
198 C, D) as well as on minerally fertilized ones (M) and untreated control (T). The samples were
199 collected excluding the upper 3 cm of topsoil and the rhizosphere area. The analyses were performed
200 with the same methods exploited for the chemical characterization of substrates prior to the
201 experiment (see **Supplementary material - Section II**). Chemical analyses of the leaves were

conducted at the end of the second month, after the biomass measurement, in order to assess the content of nitrogen, phosphorus and potassium in the leaves. In the case of the samples from the sandy soil, the measurements were performed on samples treated with one liquid digestate (P), one solid digestate (D) and the mineral fertilizer (M). It was not possible to analyze samples from the negative control (T) due to the low biomass production. On the peat substrate, it was possible to evaluate N-P-K content not only in the P, D, and M samples, but also in the negative control ones (T). The plant samples were firstly processed with a humid digestion protocol (Mills and Jones Jr, 1996). Then, nitrogen was measured through the Kjeldahl method, phosphorus was evaluated through colorimetry (molybdovanadate method) and potassium was quantified by Atomic Absorption Spectroscopy (AAS). Finally, the N, P and K percentages were used to calculate the mean total element present in the epigeal part of the plant using the following formula:

$$\frac{\text{dry sample biomass (g)}}{\text{N, P, K in sample (\%)}} * \frac{100}{1000} = \text{total N, P, K presence in single sample (mg)}$$

Using data of soil and leaves chemical analyses, nitrogen apparent balance was calculated as reported in **Supplementary material – Section III**.

2.5. Statistical analysis

The experimental data were subjected to statistical analyses. Two-way ANOVA was used to compare the average results of different treatments on plant measurements. Differently, one-way ANOVA was used to compare the mean results of different treatments on the chemical analyses of soils and leaf nutrient content. After the ANOVA, Tukey's post-hoc test ($P < 0.05$) was performed. The statistical software R (version 3.5.1 - Feather Spray - 2018) was used for all statistical analysis.

3. Results

3.1. Plant measurements

3.1.1. Dry biomass

On the sandy soil at the end of the first month, the dry biomass of the tomato plants grown with digestates did not show any difference between each other. Despite the absence of significant differences, it must be pointed out that biomass of S was 28.7 and 12.7-folds higher than control (T) and mineral fertilizer (M), respectively. At the end of second month, all digestates (P, S, C, D) showed a dry biomass production significantly higher (26.7, 33, 35.3 and 37.5-folds, respectively) than control. At the same time, S, C and D showed a higher biomass than mineral fertilizer (2.9, 3.1 and 3.3-folds, respectively). At the end of third month, dewatered SSADs proved to be the most productive treatments, with C and D displaying the highest yields (10.23 g and 10.97 g). Their biomasses doubled mineral fertilizer one (5.13 g), which was only comparable to the biomass produced by plants treated with SSADs after two months. Furthermore, C and D yields were 16 and 17-folds higher than T (0.64 g), respectively (**Figure 1.A**).

On the peat substrate, no significant differences between treatments were appreciable within the same month. The only significant differences emerged between biomass values between three different months (**Figure 1.B**).

Results of ANUE showed significant differences only on sandy soil (**Table S3**). Moreover, it must be pointed out that ANUE values of SSADs in sandy soil were up to 23, 3.5 and 2.4-folds higher than mineral ones after one, two and three months after sowing, respectively.

3.1.2.Height

On the sandy soil, no differences in plant height were present at the end of the first month. Nevertheless, S treatment revealed the tallest tomato plants, up to 2.6 and 2.5-folds higher than T and M. In the second month, all SSADs-treated plants were significantly taller than control and mineral fertilizer, with D treatment displaying a height 6 and 2.1-folds higher than T and M, respectively. After three months, the mean height of T was still the lowest. The mean height of the plants grown on P and D was comparable to plants grown on mineral fertilizer. Plants grown with S and C treatments had a statistically higher height than mineral fertilizer (M). It is worth highlighting that the mean heights of the plants grown on all digestates was at least 3.5-folds higher than the control ones (**Figure 2.A**).

On the peat substrate, no significant differences were observed between the different treatments within the same month. The only significant differences emerged between the height of the samples between three different months (data not shown).

3.1.3. Leaves and inflorescences

After the first month, the plants grown on sandy soil in presence of D and C treatments showed a number of leaves comparable to control and minerally fertilized plants. On the other hand, samples from liquid SSADs (P and S) revealed a higher mean leaf number than control. After two months, the leaves number on plants grown with digestates was significantly higher only than negative control plants. At the end of the experiment, samples from S and D treatments showed the highest number of leaves, which were not statistically different from samples from C treatment. Plants grown with P had similar number of leaves than C and mineral fertilizer, while leaves number in negative control was still the lowest one (**Figure 2.B**).

With regards to the number of inflorescences, no plant on sandy soil showed flowers one month after sowing. At the end of the second month, plants in T and M were still not revealing any flower. Differently, P, S, C and D had some inflorescences, but no significant difference between treatments was present. At the end of the experiment, negative control plants still did not show any flower. Plants treated with P and S had a number of inflorescences statistically similar to mineral fertilizer. The highest number of inflorescences was found on C and D treatments (**Figure 2.C**).

As regards the number of leaves and inflorescences of plants grown on peat substrate, no differences between treatments at the same month were highlighted by statistical analysis (data not shown).

3.1.4. Chlorophyll Content Index (CCI)

On sandy soil, leaves dimension after one month was too small to measure CCI. At the end of second month leaves of plants treated with P, S and C showed a CCI higher than control and comparable to mineral fertilizer. The mean CCI value of plants grown with D digestate was statistically higher than mineral fertilizer (M) but comparable to the others SSADs. CCI measures performed at the end of third month revealed a substantial decrease in CCI values registered in all SSADs and in mineral

277 fertilizer, whose values were not significantly different from the control. The only significant difference
278 emerging at the endpoint was between P and mineral fertilizer measure. **(Figure 2.D)**.
279 On peat substrate, the only differences were recorded between the CCI measure of D and P at the
280 end of second month, and S and control at the end of the third month (data not shown).

281 3.1.5. Infra-red gas analyzer (IRGA)

282 As regards IRGA measurements, on sandy soil the lowest A_N value was found in control, where
283 significantly higher values were recorded on C and S. Detailed results are reported in **Supplementary**
284 **material - Section V (Table S4)**.

285 3.2. Chemical analysis

286 3.2.1. Substrates analyses

287 Results of chemical analyses performed on the sandy soil after two months from treatments
288 application are summarized in **Table 2A**. SSADs showed all an intermediate mean pH included
289 between control (8.3) and mineral fertilizer samples (8.0). OM was significantly higher in P, C and D
290 treatments than in S, mineral fertilizer and control. As expected, values of organic carbon showed a
291 trend similar to OM. Total nitrogen (Kjeldahl) was lower in control, mineral fertilizer and S than P, C
292 and D treatments. All results of nitrite analysis were below quantification limits (QL). Nitrates were
293 detectable only in S, C and D treatments, showing very low concentrations (between 1 and 4 mg/kg)
294 with respect to M sample (60 mg/kg). Organic nitrogen values were roughly similar to total Kjeldahl
295 nitrogen ones. Regarding C/N ratio, the lowest value was calculated in control and mineral fertilizer,
296 while all SSADs revealed higher values. Olsen phosphorus was below QL in T and M samples;
297 differently, phosphorous content in samples treated with SSADs was higher. The lowest value of
298 exchangeable calcium was observed in S samples followed by negative control, D, P, C and mineral
299 fertilizer. A great difference in exchangeable sodium content was found between negative control
300 samples and all the treatments. Available zinc ranged between 0.21 mg/kg in control samples, and
301 1.00 mg/kg in D ones, with samples treated with liquid SSADs and mineral fertilizer showing an
302 intermediate behavior. Digestates showed intermediate values of CEC, included between control

(2.81 cmol/kg) and mineral fertilizer samples (3.58 cmol/kg). Values of electrical conductivity, ammonia nitrogen (NH_4^{4+}), exchangeable K, exchangeable Mg, available Mn and available Cu did not show any significant difference between treatments on sandy soil.

Results of chemical analyses performed on peat substrate two months after treatments application are summarized in **Table 2B**. pH values ranged from a minimum of 6.6 (M) to a maximum of 7.4 (D). Total Kjeldahl nitrogen was lower in control samples and gradually increased along with the dry matter of SSADs; the highest value was displayed by mineral fertilizer samples. Organic nitrogen values were analogous to total Kjeldahl nitrogen in terms of values, trend and differences between treatments. Nitrites, nitrates, extractable Mn, Cu and Zn were below detection limits. All other parameters did not show any significant difference.

3.2.2. Leaf analysis

On sandy soil, chemical characterization of leaves showed a concentration of nitrogen and potassium in P and D significantly lower than mineral fertilizer samples. As regards phosphorous, no significant difference emerged. The total nitrogen accumulated in leaves in D plants was significantly higher than in P ones. The mean phosphorous uptake by plants was significantly different across D, P and M samples. Finally, the potassium uptaken in leaves did not show significant differences between thesis (**Table 3A**).

On plants grown on peat substrate, concentrations and total uptake of both nitrogen and potassium on control, P and D were statistically similar to each other, but they resulted lower in comparison with mineral fertilizer ones. Concentration and total uptake of phosphorous in leaves, control showed the lowest values while D samples the highest ones (**Table 3B**).

4. Discussion

4.1. Agronomic and physiological evaluations

For many years extensive studies and reviews have shown that soil and plant benefit from SS. Indeed, SS is a good source of macro and micro nutrients as well as of OM; this enhances soil fertility and, as a consequence, crop production even in a more effective way than commercial fertilizers

(Singh and Agrawal, 2008). The results of the present work were in agreement with literature and the better performances of SS compared to inorganic fertilizers have been confirmed. **Table 4** shows technical details and results of other works dealing with SS treatment of tomato plant with pot experiments. It is important to notice that not only SSAD application rate was considerably lower in the present work, but also that the results obtained were remarkably higher. For instance, biomass and height of treated tomato plants at two months after sowing were up to 37.5 and 6-folds higher, respectively, than control plants (corresponding to an increase of 3652% and 500%), results never reached before in other works on tomato plants. Interestingly, fertilizing performances of SSAD also overcame the ones of mineral fertilizer, especially one month after sowing, when S treatment revealed biomass and height of tomato plants up to 12.7 and 2.5-folds higher than M. From here on out, differences between SSAD treatments and M samples were less accentuated, probably because nutrients release of the mineral fertilizer was faster after an initial “lag” phase. As a corollary, biomass values were reflected by ANUE ones, which were higher than the ones reported in literature for tomato plants grown in pot under greenhouse conditions treated with a 10-folds higher nitrogen application (Wang et al., 2013). Improvement in terms of leaves number and chlorophyll content were less intense, but still higher than the examples reported in literature (Bakshi et al., 2019; Elloumi et al., 2016; He et al., 2016; Hossain et al., 2015).

To a broader extent, results of the present study in terms of biomass and plant height can be compared to other works conducted with a similar experimental setup but exploiting different model species. In order to biomass, the general trend was an increase in dry matter ranging usually between 4 (*Capsicum annuum* L.; Pascual et al., 2008) and 16-folds (*Triticum aestivum* L.; Eid et al., 2019) more than untreated control. The findings of the present work confirmed and went beyond these results, considering also that the most used SS application rates ranged between the dosage used in this work and a 35-folds higher one (Eid et al., 2019). On the other hand, the improvements in plant height were in line with the results obtained by Eid and colleagues on cucumber (*Cucumis sativus* L.) (2017) and wheat (*Triticum aestivum* L.) (2019), reporting a stem length improvement up to 3 and 6-folds, respectively, over untreated control. The only case with a striking higher biomass production was described for the sunflower (*Heliantus annuus* L.), whose production increased up to 125-folds

357 more than the untreated control. However, the SS dosage was up to 35-folds higher than the present
358 study. Moreover improvement in terms of height was comparable to the present work (Bouriou et al.,
359 2018). Taking into account the works using SS dosages comparable to 170 kg N/ha, the majority
360 were open field experiments. For instance, triticale (*X Triticosecale Wittmack*) (Kchaou et al., 2018)
361 revealed a biomass increase of 2-folds. Furthermore, results of the present work corroborate positive
362 effects on biomass of SS application on soils poor in nutrients (Walter et al., 2000) and strongly
363 alkaline (Zuo et al., 2019).

364 SSAD application on tomato crops resulted also in an augmented number of leaves and
365 inflorescences with respect to control and mineral fertilizer. Moreover, inflorescences number of
366 SSAD-treated plants increased from 2 to 3-folds over the last month. These findings were in general
367 agreement with other results reported on tomato grown in presence of SS (Bakshi et al., 2019),
368 despite the higher treatment dosages.

369 Number of leaves and inflorescences are developmental parameters considered also with other plant
370 species when testing the fertilizing effects of SSAD. For instance, Eid and colleagues (2017)
371 registered on cucumber a boost in the number of leaves of more than 2-folds, which is in line with the
372 results of the present work. Similar outcomes have been reported in terms of number of flowers in
373 common bean (*Phaseolus vulgaris* L.)(Fernández-Luqueño et al., 2010) and marigold (*Tagetes erecta*
374 L.)(Solanki et al., 2017) grown in SS dosages lower and higher, respectively, than the present work.
375 In contrast with these results, Tariq and co-workers (2012) described a decrease up to 60% in flowers
376 number in *Dahlia x hortensis*, whose growth had probably been compromised by an excessive SS
377 dosage.

378 Results of the present work confirmed the positive effects of SS application on net photosynthesis
379 (Bouriou et al., 2018; Pascual et al., 2008) and chlorophyll content. Leaf chlorophyll content was
380 directly correlated with indirect chlorophyll measurements such as readings through SPAD and CCI-
381 meters (Xiong et al., 2015), whose value can be compared to each other with the equations proposed
382 by Parry and colleagues (2014). Application of SSAD improved chlorophyll content values of tomato
383 plants grown on sandy soil at the end of second month, as well as dry biomass and net
384 photosynthesis (A_N). This beneficial effect has been already observed also in sunflower (Bouriou et

al., 2018), sorghum (Alvarenga et al., 2016) and triticale (Kchaou et al., 2018). On the other hand, literature provides examples of reduction of leaf chlorophyll content in tomato after treatment with SS (Elloumi et al., 2016), which is probably due to the excessive heavy metals presence in the used SS (Singh and Agrawal, 2007). However, this aspect was likely not linked with the reduction of chlorophyll content over time observed in the present study. Indeed, this phenomenon has been already observed in other SS-treated plant species, such as common bean (Fernández-Luqueño et al., 2010). A possible explanation of this reduction in CCI at the end of the third month could be the deficiency of nutrients in soil. A second hypothesis for CCI decrease has been proposed by de Oliveira and co-workers (2017): after the initial blooming of the plant, gradual degradation of chlorophyll occurs due to the beginning of the fruit development phase, which induces a metabolic change in the plant, with a more sustained nutrients accumulation in the fruit. Taking into account the relationship between leaf nitrogen and chlorophyll content (Xiong et al., 2015), a third justification for CCI decrease can be provided by the so-called nitrogen dilution curve. In fact, biomass increase in tomato plant was accompanied by a reduction in nitrogen concentration (and, consequently in chlorophyll content) because the structural compartment (lower in N%) becomes proportionally more massive than metabolic active one (higher in N%) (Tei et al., 2002).

4.2. Chemical analysis

4.2.1. Substrates analyses

The application of SS on soil can affect different physical and chemical soil characteristics (Epstein, 2002). Likewise, many changes were documented in this experiment (both on sandy soil and on peat substrate) two months after treatments application. Although peat substrate was low in nutrient content, it showed a consistently higher amount of microelement than sandy soil. Moreover, peat substrate has many other advantages such as lightweight, high water holding capacity and high air space (Gruda et al., 2016). All these peculiarities most probably contributed to the minor differences registered on peat substrate.

Soil analysis results revealed a change in soil pH after the treatments application. Many works reported an increase (Bayoumi Hamuda et al., 2009; Ferreiro-Domínguez et al., 2011) or a decrease

(Mosquera-Losada et al., 2016; Singh and Agrawal, 2007) in soil pH. In the present work, acidification occurred in treated sandy soil samples, probably due to both the lower pH of SSADs and the nitrogen mineralization (Rasouli-Sadaghiani and Moradi, 2014). In particular, the nitrification process ($\text{NH}_4^+ \rightarrow \text{NO}_3^-$) (Stamatiadis et al., 1999) induces the release of H^+ in soil solution media and the leaching of NO_3^- by water (Whitehead, 1995). Another conceivable theory for soil acidification in SSAD-treated samples could be the generation of organic acids during sewage sludge mineralization (Angin et al., 2012; Bouriou et al., 2018). Additionally, the low buffering capacity might be yet another plausible effect occurring in the sandy soil case.

Electrical conductivity values (both on sandy soil and on peat substrate) did not statistically change after treatments application unlike many other works (Bouriou et al., 2018; Singh and Agrawal, 2007), likely due to the consistently lower SSAD application rates. Nevertheless, it must be pointed out that, concerning sandy soil, EC values in M were approximatively doubled compared to SSAD ones, which in turns were somewhat higher than control. High EC of M might be due to the particularly higher concentration of nitrates, likely released as bioavailable form nitrogen by the commercial fertilizer. However, these relatively elevate nitrate amounts were likely not necessary, as confirmed by the better growth parameters and ANUE values of tomato plants growing on SSAD amended soil. On the contrary, excess of nitrates may result in undesired drawbacks such as leaching and hyperaccumulation in plant tissues, feature in agreement with the foliar analyses. Moreover, sodium might have affected EC values both in mineral fertilizer and in SSAD treatments (probably influenced by sodium presence in the digestates). However, Na did not affect the physiological parameters of tomato plants as confirmed by IRGA measurements.

The thesis of a possible increasing of soil OM in soils treated with SSADs (Kladivko and Nelson, 1979; Perez-Espinosa et al., 1999) was confirmed by the present work. Despite the OM percentage was very low in all samples, the value in SSADs treated theses was higher than control and mineral fertilizer. This may partially justify the better performances of treated samples in term of biomass and height, according to the well-known soil OM benefits on plants growth (Bot and Benites, 2005).

CEC significantly increased in SSADs-treated soil, which was probably caused by the OM increment. This effect is even more pronounced on alkaline soils (Bohn et al., 2001) and similar results were found in other works (Angin et al., 2012; Ferreiro-Domínguez et al., 2011).

Total N, available P, exchangeable Ca and Na and available Fe and Zn concentrations increased in the sandy soil amended with SSADs due to their higher concentration in SS (Singh and Agrawal, 2007).

Two months after treatments application, N_{Tot} (Kjeldahl) was higher in C and D than liquid SSADs (P and S), probably due to their solid form that plausibly induced a slower release, both on sandy soil and peat substrate. Other studies revealed that total soil nitrogen can persist in higher concentrations also for longer periods after SSAD treatment application (Bourrioug et al., 2015). Anyway, all samples showed a total N content lower than before digestates application. It meant that a remarkable part of nitrogen both already present in sandy soil and added with digestates was absorbed, transformed or lost after two months, as suggested by the apparent nitrogen balance (**Table S2**). Concerning this balance, it worth specifying that no significant difference was found between P, D and M treatments.

The significant variation in N and OM content in treated sandy soils changed C/N ratio. The results obtained with SSADs were still low (< 9 ; Arpa Veneto, 2007), but higher than in control and mineral fertilizer. The small changes in C/N and the relatively low values across treatments likely indicated that nitrogen mineralization could have prevailed over microbial immobilization. Therefore, nitrogen in SSAD treated samples was surely bioavailable and used efficiently by plants, as also confirmed by ANUE values. However, it should be also noticed that mineralisation was likely a slow nitrogen release process, as evidenced by soil nitrate and leaf nitrogen analyses. Indeed, these evidenced that nitrogen was much more bioavailable in M treatments, but less efficiently utilizable, according to ANUE values.

In all SSADs treated soils, the available P was higher than control and mineral fertilizer. Considering that the different dosages were normalized on N dosage per each thesis, the difference in P content between the samples treated with SSADs can be explained by the different percentages of P in the four SSADs. This diversity could also explain the differences among different treatments on physiological parameters of tomato. Moreover, the addition of OM probably enhanced the availability

of P in soil treated with SSADs (Fekri et al., 2011). In fact, this can increase the abundance and the activity of microorganisms, favoring P capture (Nobile et al., 2019). Similar results in increase of soil P were obtained by Singh and Agrawal (2007) and Walter and colleagues (2000). For what it concerns K, no differences were registered in soil after digestates application, probably due to their low concentration in this macronutrient. These results agree with other works (Bouriou et al., 2015; Walter et al., 2000). Many SS are rich of Ca due to the stabilization by means of liming (Epstein, 2002). Although the SSADs exploited in this work did not undergo Ca addition at WWTP level, its content was pretty high (> 4.64% D.M.). Considering the medium content in the initial soil, exchangeable Ca increased in some cases in treated soils, confirming the results of Ferreiro Domínguez and Singh (Ferreiro-Domínguez et al., 2011; Singh and Agrawal, 2007). A significant increase of exchangeable Na was measured in all treated soils due to the sodium percentage in SSAD and confirmed by two abovementioned works (Ferreiro-Domínguez et al., 2011; Singh and Agrawal, 2007). The excess of Na is a well-known limiting factor for plants growing (Jones Jr., 2012) but Na has been recently defined as a “new beneficial element” (Morgan, 2000) that, in small quantities, can increase tomato yields (Jones Jr., 2012). The consistent presence of Fe and Zn in SSADs likely provoked the increase in their concentration in sandy soil, confirming the results of Angin and colleagues (2012).

4.2.2. Leaf analysis

In some cases, in literature the use of SS enhanced the percentage of macronutrients in leaves (Angin et al., 2012; Zuo et al., 2019), in other ones no change took place (Kotecki et al., 2014; Pinna et al., 2009) and still in other ones concentration increased only for some nutrients (Bakshi et al., 2019; De Andres et al., 2010). This work belongs to the third category, since only foliar P% and total uptaken P of control plants grown on peat substrate were significantly lower than SSADs ones. On sandy soil, content of uptaken P was significantly higher in D and P treatments, which was likely influenced by the phosphorous amount in the initial application. Nevertheless, no significant differences emerged in foliar P% despite the difference in uptaken P content between SSADs and

mineral fertilizer: probably, the controlled nutrient release of the mineral fertilizer compensated the higher quantity of P in the SSADs. Moreover, it could be inferred that differences in foliar macronutrient content could have been appreciated between control and treated samples on sandy soil. However, the too low biomass of untreated samples made impossible this investigation.

The total amount of N and K uptaken in leaves had varied results. On sandy soil, D samples revealed a significantly higher N content than P ones due to the different biomass production. Concerning plants grown on peat substrate, P and D showed a nitrogen plant uptake similar to negative control, but lower than mineral fertilizer, likely due to the characteristics of the fertilizer, such as the controlled nutrient release and the presence of readily bioavailable nitrogen forms. As regards K, despite its higher amount in mineral fertilizer, total K uptaken in leaves did not result significantly different between the treatments applied on sandy soil, due to the different aboveground biomass production. On the other hand, on peat substrate, the $K_{\text{Extractable}}$ content of plant with mineral fertilizer was the highest considering the similar biomass production.

5. Conclusions

In the present work, pot experiments under greenhouse conditions on two different substrates were performed to evaluate fertilizing effects of four different SSADs over a time span of three months. The application of these digestates clearly highlighted beneficial effects on different growth parameters of tomato plants, especially when cultivated on a sandy, alkaline and poor (in nutrient and OM) soil. For instance, it is important to point out that plant biomass and height reached values up to 37.5 and 6-folds, respectively, higher than untreated control; additionally, SSAD-treated plants showed values of biomass and height up to 12.7 and 2.5-folds, respectively, higher than mineral treatment, indicating that SSAD could be a valuable alternative to mineral fertilizers to boost fertility in poor and sandy soils. Moreover, the present work confirmed the thesis of the enhancement of soil OM with the use of SSAD. Furthermore, it is important to notice the increments of some macro- (nitrogen, phosphorous and calcium) and micro-nutrients (iron and zinc) in sandy soil, showing significant differences with respect to untreated control. Nevertheless, some of the registered values were low and it can be reasonably assumed that most of nutrients had already been assimilated to let the plant grow. This

520 aspect was confirmed by leaves analysis, which showed a remarkable uptake in N, P and K by
521 tomato plants. With respect to these macronutrients, it is worth emphasizing that the experiment was
522 designed to administer plants, across the different treatments, the same nitrogen dosage as sludge
523 application rate is usually based on plants nitrogen requirements. However, the differences in SSADs
524 composition implied a remarkable imbalance in terms of other nutrients and OM. Hence, we can
525 assume that these differences likely influenced plant growth, providing consistent differences between
526 different theses.

527 Future work should include on one side a deeper analysis of the issues tackled in the present paper,
528 and on the other hand it should consider also related aspects. Concerning the formers, chemical
529 characterization of the treated substrates and plants should be carried out in a time-course fashion,
530 allowing to properly describe the mass balance of the elements (including the study of leaching
531 effects) and their release dynamics over time. Consequently, it should allow a more detailed
532 evaluation of the fertilizing indexes (e.g. ANRE, ANUE). As regards the new related aspects to be
533 addressed, soil application of SSAD should be explored both analyzing the presence of organic
534 pollutants (e.g. antibiotics, EDC) as well as considering microbiological aspects, such as the effects
535 on microbial communities and the study of metagenomics and metatranscriptomics traits (e.g.
536 antibiotics resistance genes).

537 Despite reserves and resources for N, P and K appear adequate for the near future, it is necessary to
538 find less impactful solutions to produce fertilizers in the short term. In this way, the reuse of SS can
539 reduce the negative effects connected by the extraction, manufacturing and the use of mineral
540 fertilizers derived from non-renewable resources. Furthermore, this experiment showed how the
541 positive effects of SSADs are emphasized if applied on a poor alkaline soil.

542 Acknowledgements

543 Authors wish to thank Mr. Umberto Rossi for supporting this research with the admirable and ethical
544 aim to help people in either desertification areas and developing countries, in order to overcome
545 issues related to poverty and climate change.

546 Credits for icons in Graphical Abstract: Icon made by Freepik from www.flaticon.com.

547 **Funding**

548 This research was partially supported by the EU Horizon2020 NUTRIMAN project (GA 818470) to
549 University of Torino – Agroinnova.

550 **References**

551 Alvarenga, P., Farto, M., Mourinha, C., Palma, P., 2016. Beneficial Use of Dewatered and Composted
552 Sewage Sludge as Soil Amendments: Behaviour of Metals in Soils and Their Uptake by Plants.
553 Waste and Biomass Valorization 7, 1189–1201. <https://doi.org/10.1007/s12649-016-9519-z>

554 Alvarenga, P., Mourinha, C., Farto, M., Santos, T., Palma, P., Sengo, J., Morais, M.C., Cunha-Queda,
555 C., 2015. Sewage sludge, compost and other representative organic wastes as agricultural soil
556 amendments: Benefits versus limiting factors. Waste Manag. 40, 44–52.
557 <https://doi.org/10.1016/j.wasman.2015.01.027>

558 Angin, I., Aslantas, R., Kose, M., Karakurt, H., Ozkan, G., 2012. Changes in chemical properties of
559 soil and sour cherry as a result of sewage sludge application. Hortic. Sci. 39, 61–66.
560 <https://doi.org/10.17221/4/2011-HORTSCI>

561 Arpa Veneto, 2007. Capitolo 13 - Agrelan: un sistema di interpretazione dell’analisi del terreno, in:
562 L’interpretazione Delle Analisi Del Terreno - Strumento per La Sostenibilità Ambientale. Padova,
563 pp. 62–67.

564 Arthurson, V., 2008. Proper sanitization of sewage sludge: A critical issue for a sustainable society.
565 Appl. Environ. Microbiol. 74, 5267–5275. <https://doi.org/10.1128/AEM.00438-08>

566 Bakshi, M., Liné, C., Bedolla, D.E., Stein, R.J., Kaegi, R., Sarret, G., Pradas del Real, A.E., Castillo-
567 Michel, H., Abhilash, P.C., Larue, C., 2019. Assessing the impacts of sewage sludge
568 amendment containing nano-TiO2 on tomato plants: A life cycle study. J. Hazard. Mater. 369,
569 191–198. <https://doi.org/10.1016/j.jhazmat.2019.02.036>

570 Basosi, R., Spinelli, D., Fierro, A., Jez, S., 2014. Mineral nitrogen fertilizers: Environmental impact of

571 production and use. Fertil. Components, Uses Agric. Environ. Impacts 3–43.

572 Bayoumi Hamuda, H.E.A.F., Orosz, E., Horváth, M., Palágyi, A., Szederné Baranyai, B., Patkó, I.,
 573 Kecskés, M., 2009. Effect of sewage sludge on soil properties, the growth of *Lycopersicon*
 574 *esculentum* L. and the properties of the rhizosphere in a model experiment. *Agrokem. es Talajt.*
 575 58, 325–342. <https://doi.org/10.1556/Agrokem.58.2009.2.12>

576 Bohn, H.L., O'Connor, G.A., McNeal, B.L., 2001. *Soil Chemistry*, 3rd ed.

577 Bot, A., Benites, J., 2005. The importance of soil organic matter, *FAO Soils Bulletin*.
 578 <https://doi.org/10.1080/03650340214162>

579 Bouriou, M., Alaoui-Sehmer, L., Laffray, X., Benbrahim, M., Aleya, L., Alaoui-Sossé, B., 2015.
 580 Sewage sludge fertilization in larch seedlings: Effects on trace metal accumulation and growth
 581 performance. *Ecol. Eng.* <https://doi.org/10.1016/j.ecoleng.2015.01.031>

582 Bouriou, M., Krouna, M., Abouabdillah, A., Harraq, A., Bouabid, R., Aleya, L., 2018. Sewage sludge
 583 used as organic manure in Moroccan sunflower culture: Effects on certain soil properties, growth
 584 and yield components. *Sci. Total Environ.* 627, 681–688.
 585 <https://doi.org/10.1016/j.scitotenv.2018.01.258>

586 Buol, S.W., Southard, R.J., Graham, R.C., McDaniel, P.A., 2011. *Soil genesis and classification*, sixth
 587 edit. ed, John Wiley & Sons, Ltd., Publication. John Wiley & Sons, Inc.

588 Camargo, F.P., Sérgio Tonello, P., Dos Santos, A.C.A., Duarte, I.C.S., 2016. Removal of Toxic Metals
 589 from Sewage Sludge Through Chemical, Physical, and Biological Treatments—a Review. *Water.*
 590 *Air. Soil Pollut.* 227, 433. <https://doi.org/10.1007/s11270-016-3141-3>

591 Council of the European Communities, 1991a. Council Directive 91/271/EEC of 21 May 1991
 592 concerning urban waste-water treatment. *Off. J. Eur. Communities L* 135, 40–52.

593 Council of the European Communities, 1991b. Council Directive 91/676/EEC of 12 December 1991
 594 concerning the protection of waters against pollution caused by nitrates from agricultural
 595 sources. *Off. J. Eur. Communities L* 375, 1–8.

Council of the European Communities, 1986. Council Directive 86/278/EEC of 12 June 1986 on the
 protection of the environment, and in particular of the soil, when sewage sludge is used in
 agriculture. Off. J. Eur. Communities L 181, 6–12.

Cristina, G., Camelin, E., Pugliese, M., Tommasi, T., Fino, D., 2019. Evaluation of anaerobic
 digestates from sewage sludge as a potential solution for improvement of soil fertility. Waste
 Manag. 99, 122–134. <https://doi.org/10.1016/j.wasman.2019.08.018>

De Andres, E.F., Tenorio, J.L., Walter, I., 2010. Biomass production and nutrient concentration of
 kenaf grown on sewage sludge-amended soil. Spanish J. Agric. Res. 8, 472.
<https://doi.org/10.5424/sjar/2010082-1202>

De López Camelo, L.G., De Miguez, S.R., Marbán, L., 1997. Heavy metals input with phosphate
 fertilizers used in Argentina. Sci. Total Environ. 204, 245–250. [https://doi.org/10.1016/S0048-9697\(97\)00187-3](https://doi.org/10.1016/S0048-9697(97)00187-3)

de Oliveira, M.O.A., Alves, S.M. de F., Freitas, E. de F.M., de Faria, H.F.L., Lisboa, F.C., 2017.
 Relative chlorophyll index on doses of nitrogen fertilization for cherry tomato culture. African J.
 Agric. Res. 12, 2946–2953. <https://doi.org/10.5897/ajar2016.12051>

Dumontet, S., Dinel, H., Baloda, S.B., 1999. Pathogen Reduction in Sewage Sludge by Composting
 and Other Biological Treatments: A Review. Biol. Agric. Hortic. 16, 409–430.
<https://doi.org/10.1080/01448765.1999.9755243>

Eid, E.M., Alrumman, S.A., El-Bebany, A.F., Fawy, K.F., Taher, M.A., Hesham, A.E.-L., El-Shaboury,
 G.A., Ahmed, M.T., 2019. Evaluation of the potential of sewage sludge as a valuable fertilizer for
 wheat (*Triticum aestivum* L.) crops. Environ. Sci. Pollut. Res. 26, 392–401.
<https://doi.org/10.1007/s11356-018-3617-3>

Eid, E.M., Alrumman, S.A., El-Bebany, A.F., Hesham, A.E.L., Taher, M.A., Fawy, K.F., 2017. The
 effects of different sewage sludge amendment rates on the heavy metal bioaccumulation, growth
 and biomass of cucumbers (*Cucumis sativus* L.). Environ. Sci. Pollut. Res. 24, 16371–16382.
<https://doi.org/10.1007/s11356-017-9289-6>

622 Elloumi, N., Belhaj, D., Jerbi, B., Zouari, M., Kallel, M., 2016. Effects of sewage sludge on bio-
 623 accumulation of heavy metals in tomato seedlings. Spanish J. Agric. Res. 14, e0807.
 624 <https://doi.org/10.5424/sjar/2016144-9210>

625 Epstein, E., 2002. Land application of sewage sludge and biosolids. CRC Press.

626 Eurostat, 2019. Sewage sludge production and disposal.

627 FAO, 2017. World fertilizer trends and outlook to 2020 - Summary report. Food Agric. Organ. United
 628 Nations.

629 Fekri, M., Gorgin, N., Sadegh, L., 2011. Phosphorus desorption kinetics in two calcareous soils
 630 amended with P fertilizer and organic matter. Environ. Earth Sci. 64, 721–729.
 631 <https://doi.org/10.1007/s12665-010-0892-9>

632 Fernández-Luqueño, F., Reyes-Varela, V., Martínez-Suárez, C., Salomón-Hernández, G., Yáñez-
 633 Meneses, J., Ceballos-Ramírez, J.M., Dendooven, L., 2010. Effect of different nitrogen sources
 634 on plant characteristics and yield of common bean (*Phaseolus vulgaris* L.). Bioresour. Technol.
 635 101, 396–403. <https://doi.org/10.1016/j.biortech.2009.07.058>

636 Ferreiro-Domínguez, N., Rigueiro-Rodríguez, A., Mosquera-Losada, M.R., 2011. Response to
 637 sewage sludge fertilisation in a *Quercus rubra* L. silvopastoral system: Soil, plant biodiversity and
 638 tree and pasture production. Agric. Ecosyst. Environ. <https://doi.org/10.1016/j.agee.2011.02.009>

639 Fijalkowski, K., 2019. Emerging contaminants in sludge (endocrine disruptors, pesticides, and
 640 pharmaceutical residues, including illicit drugs/controlled substances, etc.), in: Industrial and
 641 Municipal Sludge. Elsevier, pp. 455–473. <https://doi.org/10.1016/B978-0-12-815907-1.00020-9>

642 Fixen, P.E., Johnston, A.M., 2012. World fertilizer nutrient reserves: A view to the future. J. Sci. Food
 643 Agric. 92, 1001–1005. <https://doi.org/10.1002/jsfa.4532>

644 Gadupudi, C.K., Rice, L., Xiao, L., Kantamaneni, K., 2019. Endocrine Disrupting Compounds
 645 Removal Methods from Wastewater in the United Kingdom: A Review. Sci 1, 15.
 646 <https://doi.org/10.3390/sci1010015.v1>

647 Gruda, N., Caron, J., Prasad, M., Maher, M., 2016. Growing Media, in: Lal, R. (Ed.), Encyclopedia of
648 Soil Science, Third Edition. CRC Press, pp. 1053–1058. [https://doi.org/10.1081/E-ESS3-](https://doi.org/10.1081/E-ESS3-120053784)
649 120053784

650 Harrison, E.Z., Oakes, S.R., Hysell, M., Hay, A., 2006. Organic chemicals in sewage sludges. Sci.
651 Total Environ. <https://doi.org/10.1016/j.scitotenv.2006.04.002>

652 He, S., Li, A., Lei, W., 2016. Effect of Sewage Sludge and its Biomass Composting Product on the
653 Soil Characteristics and N₂O Emission from the Tomato Planting Soil. Int. J. Agric. Biol. 18, 501–
654 508. <https://doi.org/10.17957/IJAB/15.0113>

655 Hossain, M.K., Stretzov, V., Nelson, P.F., 2015. Comparative Assessment of the Effect of Wastewater
656 Sludge Biochar on Growth, Yield and Metal Bioaccumulation of Cherry Tomato. Pedosphere 25,
657 680–685. [https://doi.org/10.1016/S1002-0160\(15\)30048-5](https://doi.org/10.1016/S1002-0160(15)30048-5)

658 Jones Jr., J.B., 2012. Plant Nutrition and Soil Fertility Manual, Plant Nutrition and Soil Fertility Manual.
659 <https://doi.org/10.1201/b11577>

660 Jones Jr, J.B., 2008. Tomato plant culture, Second edi. ed. CRC Press.

661 Kchaou, R., Baccar, R., Bouzid, J., Rejeb, S., 2018. Agricultural use of sewage sludge under sub-
662 humid Mediterranean conditions: effect on growth, yield, and metal content of a forage plant.
663 Arab. J. Geosci. 11, 746. <https://doi.org/10.1007/s12517-018-4103-4>

664 Kladvko, E.J., Nelson, D.W., 1979. Changes in soil properties from application of anaerobic sludge. J.
665 Water Pollut. Control Fed. 51, 325–332.

666 Kotecki, A., Kozak, M., Malarz, W., Kotecki, A., 2014. Wpływ nawożenia osadem ściekowym na
667 rozwój, plonowanie i skład chemiczny spartiny preriowej (*spartina pectinata* link.). J. Elem. 19,
668 1021–1036. <https://doi.org/10.5601/jelem.2014.19.3.725>

669 Li, L., XianJin, H., Hui, W., YuanHua, D., Sisi, X., 2009. Estimation of environmental costs of chemical
670 fertilizer utilization in China. Acta Pedol. Sin. 46, 63–69.

671 Liu, S., Zhu, N., Li, L.Y., 2012. The one-stage autothermal thermophilic aerobic digestion for sewage

672 sludge treatment: Stabilization process and mechanism. *Bioresour. Technol.* 104, 266–273.
673 <https://doi.org/10.1016/j.biortech.2011.11.041>

674 Mateo-Sagasta, J., Raschid-Sally, L., Thebo, A., 2015. Global Wastewater and Sludge Production,
675 Treatment and Use, in: *Wastewater*. Springer Netherlands, Dordrecht, pp. 15–38.
676 https://doi.org/10.1007/978-94-017-9545-6_2

677 Mills, H.A., Jones Jr, J.B., 1996. *Plant analysis handbook II: A practical sampling, preparation,*
678 *analysis, and interpretation guide.*

679 Mirlean, N., Baisch, P., Machado, I., Shumilin, E., 2008. Mercury contamination of soil as the result of
680 long-term phosphate fertilizer production. *Bull. Environ. Contam. Toxicol.* 81, 305–308.
681 <https://doi.org/10.1007/s00128-008-9480-z>

682 Mirlean, N., Roisenberg, A., 2007. Fluoride distribution in the environment along the gradient of a
683 phosphate-fertilizer production emission (southern Brazil). *Environ. Geochem. Health* 29, 179–
684 187. <https://doi.org/10.1007/s10653-006-9061-1>

685 Morgan, L., 2000. Beneficial elements for hydroponics: A new look at plant nutrition. *Growing Edge*.

686 Mortvedt, J.J., 1995. Heavy metal contaminants in inorganic and organic fertilizers. *Fertil. Res.* 43,
687 55–61. <https://doi.org/10.1007/BF00747683>

688 Mosquera-Losada, M.R., Ferreiro-Domínguez, N., Daboussi, S., Rigueiro-Rodríguez, A., 2016.
689 Sewage sludge stabilisation and fertiliser value in a silvopastoral system developed with
690 *Eucalyptus nitens* Maiden in Lugo (Spain). *Sci. Total Environ.* 566–567, 806–815.
691 <https://doi.org/10.1016/j.scitotenv.2016.05.003>

692 Neumann, P., Pesante, S., Venegas, M., Vidal, G., 2016. Developments in pre-treatment methods to
693 improve anaerobic digestion of sewage sludge. *Rev. Environ. Sci. Bio/Technology* 15, 173–211.
694 <https://doi.org/10.1007/s11157-016-9396-8>

695 Nobile, C., Houben, D., Michel, E., Firmin, S., Lambers, H., Kandeler, E., Faucon, M.P., 2019.
696 Phosphorus-acquisition strategies of canola, wheat and barley in soil amended with sewage

697 sludges. *Sci. Rep.* 9, 1–11. <https://doi.org/10.1038/s41598-019-51204-x>

698 Parry, C., Blonquist, J.M., Bugbee, B., 2014. In situ measurement of leaf chlorophyll concentration:
699 Analysis of the optical/absolute relationship. *Plant, Cell Environ.* 37, 2508–2520.
700 <https://doi.org/10.1111/pce.12324>

701 Pascual, I., Avilés, M., Aguirreolea, J., Sánchez-Díaz, M., 2008. Effect of sanitized and non-sanitized
702 sewage sludge on soil microbial community and the physiology of pepper plants. *Plant Soil* 310,
703 41–53. <https://doi.org/10.1007/s11104-008-9626-0>

704 Perez-Espinosa, A., Moreno-Caselles, J., Moral, R., Perez-Murcia, M.D., Gomez, I., 1999. Effect of
705 sewage sludge and cobalt treatments on tomato fruit yield, weight, and quality. *J. Plant Nutr.* 22,
706 379–385. <https://doi.org/10.1080/01904169909365635>

707 Pinna, M.V., Giunta, F., Nardi, S., Pusino, A., 2009. Effects of a municipal sewage sludge amendment
708 on triasulfuron soil sorption and wheat growth. *J. Agric. Food Chem.* 57, 11249–11253.
709 <https://doi.org/10.1021/jf902659f>

710 Rasouli-Sadaghiani, M.H., Moradi, N., 2014. Effect of poultry, cattle, sheep manures and sewage
711 sludge on N mineralisation. *Chem. Ecol.* 30, 666–675.
712 <https://doi.org/10.1080/02757540.2014.889122>

713 Reijnders, L., 2014. Phosphorus resources, their depletion and conservation, a review. *Resour.*
714 *Conserv. Recycl.* 93, 32–49. <https://doi.org/10.1016/j.resconrec.2014.09.006>

715 Sabiha-Javied, Mehmood, T., Chaudhry, M.M., Tufail, M., Irfan, N., 2009. Heavy metal pollution from
716 phosphate rock used for the production of fertilizer in Pakistan. *Microchem. J.* 91, 94–99.
717 <https://doi.org/10.1016/j.microc.2008.08.009>

718 Schnug, E., Lottermoser, B.G., 2013. Fertilizer-derived uranium and its threat to human health.
719 *Environ. Sci. Technol.* 47, 2433–2434. <https://doi.org/10.1021/es4002357>

720 Schulze, E.D., Freibauer, A., 2005. Environmental science: Carbon unlocked from soils. *Nature* 437,
721 205–206. <https://doi.org/10.1038/437205a>

722 Singh, R.P., Agrawal, M., 2008. Potential benefits and risks of land application of sewage sludge.
 723 Waste Manag. <https://doi.org/10.1016/j.wasman.2006.12.010>

724 Singh, R.P., Agrawal, M., 2007. Effects of sewage sludge amendment on heavy metal accumulation
 725 and consequent responses of Beta vulgaris plants. Chemosphere 67, 2229–2240.
 726 <https://doi.org/10.1016/j.chemosphere.2006.12.019>

727 Smith, K.M., Fowler, G.D., Pullket, S., Graham, N.J.D., 2009. Sewage sludge-based adsorbents: A
 728 review of their production, properties and use in water treatment applications. Water Res. 43,
 729 2569–2594. <https://doi.org/10.1016/j.watres.2009.02.038>

730 Solanki, P., Kalavagadda, B., Akula, B., Sharma, S.H.K., Reddy, D.J., 2017. Effect of Sewage Sludge
 731 on Marigold (*Tagetes erecta*). Int. J. Curr. Microbiol. Appl. Sci. 6, 825–831.
 732 <https://doi.org/10.20546/ijcmas.2017.606.097>

733 Stamatiadis, S., Doran, J.W., Kettler, T., 1999. Field and laboratory evaluation of soil quality changes
 734 resulting from injection of liquid sewage sludge. Appl. Soil Ecol. 12, 263–272.
 735 [https://doi.org/10.1016/S0929-1393\(99\)00007-4](https://doi.org/10.1016/S0929-1393(99)00007-4)

736 Stehouwer, R.C., Wolf, A.M., Doty, W.T., 2000. Chemical monitoring of sewage sludge in
 737 Pennsylvania: Variability and application uncertainty. J. Environ. Qual. 29, 1686–1695.
 738 <https://doi.org/10.2134/jeq2000.00472425002900050041x>

739 Tariq, U., ur Rehman, S., Khan, M.A., Younis, A., Yaseen, M., Ahsan, M., 2012. Agricultural and
 740 municipal waste as potting media components for the growth and flowering of *Dahlia hortensis*
 741 “Figaro.” Turk. J. Botany 36, 378–385. <https://doi.org/10.3906/bot-1109-16>

742 Tei, F., Benincasa, P., Guiducci, M., 2002. Critical nitrogen concentration in processing tomato, in:
 743 European Journal of Agronomy. [https://doi.org/10.1016/S1161-0301\(02\)00096-5](https://doi.org/10.1016/S1161-0301(02)00096-5)

744 Vanotti, M.B., García-González, M.C., Molinuevo-Salces, B., Riaño, B., 2019. Editorial: New
 745 Processes for Nutrient Recovery From Wastes. Front. Sustain. Food Syst. 3, 1–2.
 746 <https://doi.org/10.3389/fsufs.2019.00081>

747 Walter, I., Cuevas, G., García, S., Martínez, F., 2000. Biosolid effects on soil and native plant
748 production in a degraded semiarid ecosystem in central Spain. *Waste Manag. Res.*
749 <https://doi.org/10.1034/j.1399-3070.2000.00111.x>

750 Wang, Y., Liu, F., Jensen, L.S., de Neergaard, A., Jensen, C.R., 2013. Alternate partial root-zone
751 irrigation improves fertilizer-N use efficiency in tomatoes. *Irrig. Sci.* 31, 589–598.
752 <https://doi.org/10.1007/s00271-012-0335-3>

753 Warman, P.R., Termeer, W.C., 2005. Evaluation of sewage sludge, septic waste and sludge compost
754 applications to corn and forage: Ca, Mg, S, Fe, Mn, Cu, Zn and B content of crops and soils.
755 *Bioresour. Technol.* 96, 1029–1038. <https://doi.org/10.1016/j.biortech.2004.09.014>

756 Whitehead, D.C., 1995. Grassland nitrogen. CAB INTERNATIONAL, Wallingford.

757 Xiong, D., Chen, J., Yu, T., Gao, W., Ling, X., Li, Y., Peng, S., Huang, J., 2015. SPAD-based leaf
758 nitrogen estimation is impacted by environmental factors and crop leaf characteristics. *Sci. Rep.*
759 5, 13389. <https://doi.org/10.1038/srep13389>

760 Zucco, M.A., Walters, S.A., Chong, S.K., Klubek, B.P., Masabni, J.G., 2015. Effect of soil type and
761 vermicompost applications on tomato growth. *Int. J. Recycl. Org. Waste Agric.* 4, 135–141.
762 <https://doi.org/10.1007/s40093-015-0093-3>

763 Zuo, W., Gu, C., Zhang, W., Xu, K., Wang, Y., Bai, Y., Shan, Y., Dai, Q., 2019. Sewage sludge
764 amendment improved soil properties and sweet sorghum yield and quality in a newly reclaimed
765 mudflat land. *Sci. Total Environ.* 654, 541–549. <https://doi.org/10.1016/j.scitotenv.2018.11.127>

766

767

768 **Table 1.** Physiscal and chemical anlalysis of soil and peat used in the present work. CEC: Cation-Exchange Capacity; AAS: Atomic Absorption
769 Spectroscopy.

Sandy soil			Peat substrate		
Parameter	Unit	Value	Parameter	Unit	Value
Stones	-	absent	Stones	-	-
Sand (2.0 - 0.020 mm)	%	94 ± 2	Sand (2.0 - 0.020 mm)	-	-
Silt (0.020 - 0.002 mm)	%	3 ± 1	Silt (0.020 - 0.002 mm)	-	-
Clay (< 0.002 mm)	%	3 ± 1	Clay (< 0.002 mm)	-	-
Texture	-	sandy	Texture	-	-
pH	-	8.2 ± 0.16	pH	-	6.2 ± 0.1
Electrical conductivity	dS/m	0.131 ± 0.018	Electrical conductivity	dS/m	0.722 ± 0.146
Organic matter	%	0.38 ± 0.12	Organic matter	-	-
Organic carbon	%	0.22 ± 0.07	Organic carbon	-	-
N - Tot (Kjeldahl)	g/kg	0.29 ± 0.09	N - Tot (Kjeldahl)	%	0.42 ± 0.06
N - NO₂⁻	mg/kg	< 0,2	N - NO₂⁻	mg/l	< QL
N - NO₃⁻	mg/kg	6.33 ± 1.53	N - NO₃⁻	mg/l	30.4 ± 7.2
N - NH₄⁺	mg/kg	3 ± 1	N - NH₄⁺	mg/l	1.3 ± 0.3
N - Org	g/kg	0.29 ± 0.09	N - Org	%	0.4 ± 0.40
C/N		7.6 ± 0.2	C/N	-	-
P_{Olsen}	mg/kg	1.8 ± 1.3	P_{extractable}	mg/l	8.1 ± 2.3
K_{exchangeable}	mg/kg	18 ± 1	K_{extractable}	mg/l	41.1 ± 6.8
Mg_{exchangeable}	mg/kg	15 ± 5	Mg_{extractable}	mg/l	28 ± 7
Ca_{exchangeable}	mg/kg	675 ± 27	Ca_{extractable}	mg/l	36 ± 8
Na_{exchangeable}	mg/kg	6 ± 3	Na_{extractable}	mg/l	16 ± 11
Fe_{available}	mg/kg	6.7 ± 1.1	Fe_{extractable}	mg/l	0.79 ± 0.21
Mn_{available}	mg/kg	6.5 ± 3.0	Mn_{extractable}	mg/l	0.15 ± 0.04
Cu_{available}	mg/kg	0.69 ± 0.29	Cu_{extractable}	mg/l	< QL
Zn_{available}	mg/kg	0.47 ± 0.29	Zn_{extractable}	mg/l	0.02 ± 0.00
CEC	cmol/kg	3.65 ± 0.35	CEC	-	-

770 **Table 2.** Chemical characterization performed two months after treatments application on sandy soil (A) and on peat substrate (B). Data are
771 expressed as mean $\pm$ standard deviation. Asterisks mean significant differences according to ANOVA test (*, **, *** differences between means
772 significant at $P \leq 0.05$, 0.01 and 0.001, respectively). CEC, cation exchange capacity; QL, quantification limit.

773 **A**

Parameter	Unit of measure	Control (T)				Primary (P)				Secondary (S)				Centrifuged (C)				Dried (D)				Mineral fertiliser (M)			
pH	-	8.3	$\pm$	0.1	***	8.2	$\pm$	0.1	***	8.2	$\pm$	0.1	***	8.1	$\pm$	0.1	***	8.1	$\pm$	0.1	***	8.0	$\pm$	0.1	***
Electrical conductivity	dS/m	0.155	$\pm$	0.020		0.219	$\pm$	0.032		0.201	$\pm$	0.010		0.197	$\pm$	0.023		0.198	$\pm$	0.025		0.399	$\pm$	0.146	
Organic matter	%	0.16	$\pm$	0.01	***	0.24	$\pm$	0.01	***	0.18	$\pm$	0.02	***	0.25	$\pm$	0.02	***	0.26	$\pm$	0.01	***	0.16	$\pm$	0.02	***
Organic carbon	%	0.09	$\pm$	0.00	***	0.14	$\pm$	0.00	***	0.11	$\pm$	0.01	***	0.14	$\pm$	0.01	***	0.15	$\pm$	0.00	***	0.10	$\pm$	0.01	***
N - Tot (Kjeldahl)	g/kg	0.17	$\pm$	0.01	***	0.19	$\pm$	0.01	***	0.15	$\pm$	0.01	***	0.20	$\pm$	0.00	***	0.22	$\pm$	0.01	***	0.17	$\pm$	0.01	***
N - NO ₂ ⁻	mg/kg	< QL				< QL				< QL				< QL				< QL				< QL			
N - NO ₃ ⁻	mg/kg	< QL				< QL				1	$\pm$	1	*	4	$\pm$	4	*	2	$\pm$	1	*	60	$\pm$	40	*
N - NH ₄ ⁺	mg/kg	< QL				< QL				< QL				1	$\pm$	1		2	$\pm$	0		1	$\pm$	0	
N - Org	g/kg	0.17	$\pm$	0.01	***	0.19	$\pm$	0.01	***	0.15	$\pm$	0.01	***	0.20	$\pm$	0.00	***	0.22	$\pm$	0.01	***	0.17	$\pm$	0.01	***
C/N	-	5.5	$\pm$	0.2	**	7.3	$\pm$	0.6	**	6.9	$\pm$	0.5	**	7.3	$\pm$	0.5	**	7.1	$\pm$	0.3	**	5.8	$\pm$	1.0	**
P _{Olsen}	mg/kg	< QL				4.2	$\pm$	0.5	**	10.3	$\pm$	0.8	**	15.4	$\pm$	4.6	**	36.2	$\pm$	11.4	**	< QL			
K _{exchangeable}	mg/kg	14	$\pm$	3		11	$\pm$	3		12	$\pm$	1		12	$\pm$	2		9	$\pm$	1		13	$\pm$	2	
Mg _{exchangeable}	mg/kg	11	$\pm$	2		21	$\pm$	7		26	$\pm$	3		25	$\pm$	4		22	$\pm$	4		25	$\pm$	2	
Ca _{exchangeable}	mg/kg	524	$\pm$	26	*	594	$\pm$	25	*	491	$\pm$	62	*	626	$\pm$	94	*	579	$\pm$	48	*	646	$\pm$	62	*
Na _{exchangeable}	mg/kg	16	$\pm$	1	***	35	$\pm$	2	***	33	$\pm$	3	***	26	$\pm$	3	***	32	$\pm$	3	***	27	$\pm$	5	***
Fe _{available}	mg/kg	5.7	$\pm$	0.1	***	7.6	$\pm$	0.5	***	9.3	$\pm$	0.2	***	11.2	$\pm$	0.7	***	12.5	$\pm$	0.4	***	6.1	$\pm$	0.21	***
Mn _{available}	mg/kg	5.1	$\pm$	0.4		14.8	$\pm$	16.6		33.7	$\pm$	2.5		35.8	$\pm$	27.1		20.5	$\pm$	28.1		5.2	$\pm$	0.21	
Cu _{available}	mg/kg	0.40	$\pm$	0.08		0.47	$\pm$	0.13		0.60	$\pm$	0.06		0.85	$\pm$	0.12		0.96	$\pm$	0.29		0.40	$\pm$	0.01	
Zn _{available}	mg/kg	0.21	$\pm$	0.03	***	0.36	$\pm$	0.04	***	0.37	$\pm$	0.02	***	0.73	$\pm$	0.08	***	1.00	$\pm$	0.07	***	0.43	$\pm$	0.05	***
CEC	cmol/kg	2.81	$\pm$	0.13	*	3.32	$\pm$	0.17	*	2.83	$\pm$	0.33	*	3.47	$\pm$	0.43	*	3.24	$\pm$	0.20	*	3.58	$\pm$	0.32	*

QL: N - NO₂⁻ = 0.2 mg/kg; N - NO₃⁻ = 1 mg/kg; N - NH₄⁺ = 1 mg/kg; P = 1 mg/kg.

774

775 **B**

Parameter	Unit of measure	Control (T)				Primary SSAD (P)				Secondary SSAD (S)				Centrifuged SSAD (C)				Dried SSAD (D)				Mineral (M)			
pH	-	7.0	±	0.4	*	6.7	±	0.3	*	7.2	±	0.2	*	6.9	±	0.2	*	7.4	±	0.3	*	6.6	±	0.2	*
Electrical conductivity	dS/m	0.235	±	0.040		0.436	±	0.220		0.183	±	0.038		0.495	±	0.134		0.225	±	0.074		0.523	±	0.202	
N - Tot (Kjeldahl)	% D.M.	0.23	±	0.03	*	0.24	±	0.03	*	0.25	±	0.03	*	0.28	±	0.04	*	0.31	±	0.04	*	0.32	±	0.04	*
N - NO ₂ ⁻	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
N - NO ₃ ⁻	mg/l	1.0	±	0.9		1.5	±	0.5		1.0	±	0.2		2.6	±	0.4		1.2	±	0.3		2.6	±	1.7	
N - NH ₄ ⁺	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
N - Org	% D.M.	0.22	±	0.03	*	0.24	±	0.03	*	0.25	±	0.03	*	0.28	±	0.04	*	0.31	±	0.04	*	0.32	±	0.04	*
P _{extractable}	mg/l	0.4	±	0.1		1.2	±	0.4		1.2	±	0.5		0.6	±	0.3		1.1	±	0.1		< QL			
K _{extractable}	mg/l	2.9	±	0.3		2.0	±	0.3		2.4	±	0.8		2.4	±	0.4		5.0	±	2.2		2.6	±	0.2	
Mg _{extractable}	mg/l	8	±	2		22	±	16		5	±	2		24	±	12		6	±	3		26	±	17	
Ca _{extractable}	mg/l	13	±	4		26	±	16		12	±	3		33	±	14		14	±	3		32	±	17	
Na _{extractable}	mg/l	24	±	2		29	±	7		17	±	3		32	±	4		23	±	7		31	±	4	
Fe _{extractable}	mg/l	1.17	±	0.26		0.52	±	0.46		0.80	±	0.13		0.52	±	0.30		0.73	±	0.06		0.28	±	0.11	
Mn _{extractable}	mg/l	< QL				< QL				< QL				< QL				< QL				0.03	±	0.01	
Cu _{extractable}	mg/l	< QL				< QL				< QL				< QL				< QL				< QL			
Zn _{extractable}	mg/l	< QL				< QL				< QL				< QL				< QL				0.02	±	0	

QL: N - NO₂⁻ = 0.05 mg/l; N - NH₄⁺ = 0.06 mg/l; P = 0.3 mg/l; Mn = 0.03 mg/l; Cu = 0.03 mg/l; Zn = 0.02 mg/l.

776

777

778

779

780

781 **Table 3.** Results of leaves analyses performed after two months after treatments application on sandy soil (A) and on peat substrate (B). Different
782 letters indicate differences between treatments that are significant at $P < 0.05$ (Tukey HSD). Data are expressed as mean $\pm$ standard deviation.

783 **A**

Parameter	Unit of measure	Primary SSAD (P)			Dried SSAD (D)			Mineral fertilizer (M)		
N	%	1.10	$\pm$ 0.05	b	1.35	$\pm$ 0.28	b	2.95	$\pm$ 0.36	a
	Total (mg)	46.64	$\pm$ 7.45	b	81.08	$\pm$ 15.63	a	60.83	$\pm$ 9.26	ab
P	%	0.14	$\pm$ 0.01	a	0.16	$\pm$ 0.01	a	0.13	$\pm$ 0.02	a
	Total (mg)	5.97	$\pm$ 1.25	b	9.61	$\pm$ 1.26	a	2.56	$\pm$ 0.33	c
K	%	1.46	$\pm$ 0.38	b	1.40	$\pm$ 0.18	b	3.63	$\pm$ 0.57	a
	Total (mg)	61.61	$\pm$ 16.41		84.54	$\pm$ 10.95		74.24	$\pm$ 7.52	

784

785 **B**

Parameter	Unit of measure	Control (T)			Primary SSAD (P)			Dried SSAD (D)			Mineral fertilizer (M)		
N	%	1.26	$\pm$ 0.08	b	1.32	$\pm$ 0.04	b	1.29	$\pm$ 0.19	b	2.05	$\pm$ 0.36	a
	Total (mg)	319.01	$\pm$ 11.24	b	323.86	$\pm$ 20.48	b	360.41	$\pm$ 53.17	b	550.22	$\pm$ 102.49	a
P	%	0.23	$\pm$ 0.01	b	0.29	$\pm$ 0.02	ab	0.31	$\pm$ 0.04	a	0.27	$\pm$ 0.02	ab
	Total (mg)	58.02	$\pm$ 7.29	b	69.82	$\pm$ 2.92	ab	85.44	$\pm$ 9.77	a	72.95	$\pm$ 8.78	ab
K	%	1.83	$\pm$ 0.04	b	1.92	$\pm$ 0.04	b	1.74	$\pm$ 0.21	b	2.53	$\pm$ 0.27	a
	Total (mg)	465.11	$\pm$ 37.08	b	470.69	$\pm$ 24.49	b	486.97	$\pm$ 57.85	b	677.03	$\pm$ 53.15	a

786

787

788 **Table 4.** Comparison of the results from other works in literature on the effects of treatment with sewage sludge on tomato plants. Application
789 dosages are shown as reported in the original works; values in brackets indicate how many folds more is the SS application rate with respect to the
790 present study. n.a., not available.

Cultivar	SS typology	Dosage	Kind of experiment	Cultivation substrate	Differences with respect to untreated control				Reference
					Biomass increase	Plant height	Leaves and inflorescences	Chlorophyll content	
Cherry	Digested ^a	10 t/ha (2X)	Pot experiment Greenhouse 16 weeks	Chromosol	<i>Dry biomass</i> : + 20%	10 weeks: + 50% 13 weeks: + 20% 15 weeks: + 7%	n.a.	n.a.	Hossain et al., 2015
Red Robin	n.a.	SS:soil 1:10 (65X)	Pot experiment Growth chamber 120 days	Loamy soil	<i>Fresh biomass</i> Stem: + 70% Leaves: + 142%	+ 43%	<i>Leaves</i> : + 33% <i>Flowers</i> : +130%	<i>Chlorophyll</i> ^b : <i>a</i> : + 18.3% <i>b</i> : + 34.8%	Bakshi et al., 2019
Rio Grande	Aerobically digested	2.5%; 5.0%; 7.5% (11X; 22X; 33X)	Pot experiment Greenhouse 30 days	Sandy soil	<i>Dry biomass</i> : + 180%; + 280%; +140%	n.a	n.a.	<i>Chlorophyll a+b</i> ^b : + 17.5%; - 40%; - 68.5%	Elloumi et al., 2016
n.a.	Aerobically digested	400 - 800 kg N/ha (2.35X; 4.7X)	Pot experiment Greenhouse 90 days	Clay soil	<i>Dry biomass</i> : + 18.6% + 29.6%	+ 19.2%; + 24.5%	n.a.	n.a.	He et al. 2016
Beefsteak	Anaerobically digested (4 typologies: P, S, C, D)	170 kg N/ha	Pot experiment Greenhouse 120 days	Sandy soil	<i>Dry biomass</i> up to + 3652% (D treatment, II month)	up to + 500% (D treatment, II month)	<i>Leaves</i> : up to + 180% (S treatments, I month) <i>Flowers</i> : not observed in untreated control	<i>CCI</i> : up to + 172% (D treatment, II month)	This work
				Peat substrate	<i>Dry biomass</i> : up to + 70% (C treatment, I month)	up to + 24% (P treatment, I month)	n.a.	<i>CCI</i> : up to + 64% (D treatment, III month)	


^a In this work, no details about the typology of digestion are provided.

^b In these works, leaf chlorophyll content was evaluated with methods based on extraction with organic solvents followed by spectrophotometrical quantification.


791

792

Figure 1. Mean dry biomasses of *Solanum lycopersicum* L. grown on sandy soil (A) and peat substrate (B) with different treatments among three months. Different letters indicate differences between treatments that are significant at $P < 0.05$ (Tukey HSD). Each error bar represents one standard deviation. T: non-treated, control thesis; P: primary digestate; S: secondary digestate; C: centrifuged digestate; D: dried digestate; M: mineral fertilizer.



1 **Figure 2.** (A) Mean height of *Solanum lycopersicum* L. grown on sandy soil with different treatments
2 among three months; (B) Mean number of leaves of *Solanum lycopersicum* L. grown on sandy soil
3 with different treatments among three months; (C) Mean number of flower of *Solanum lycopersicum* L.
4 grown on sandy soil with different treatments among three months. (D) Mean Chlorophyll Content Index
5 (CCI) of leaves of *Solanum lycopersicum* L. grown on sandy soil with different treatments among 3
6 months. Different letters indicate differences between treatments that are significant at $P < 0.05$ (Tukey
7 HSD). Each error bar represents one standard deviation. T: non-treated, control thesis; P: primary
8 digestate; S: secondary digestate; C: centrifuged digestate; D: dried digestate; M: mineral fertilizer.



Supplementary material

of

“Anaerobic digestates from sewage sludge used as fertilizer on a poor alkaline sandy soil and on a peat substrate: effects on tomato plants growth and on soil properties”

Giulio Cristina, Enrico Camelin, Tonia Tommasi, Debora Fino and Massimo Pugliese.

Index

- Section I: Characterization of anaerobic digestates from sewage sludge (SSADs) used
- Section II: Methods of analysis of sandy soil and peat substrate
- Section III: Apparent balance of nitrogen in sandy soil
- Section IV: Agronomic nitrogen use efficiency (ANUE) calculation
- Section V: Infra-red gas analyzer (IRGA)
- References

Section I: Characterization of anaerobic digestates from sewage sludge (SSADs) used

Primary, secondary, centrifuged and dried SSADs were sampled directly at WWTP and stored at 4°C until chemical characterisation and further uses. The chemical analyses were performed according to “Analytical Methods for Fertilizers” by the Italian Ministry of Agriculture and Forestry (M.P.A.A.F., 2006) and “Methods for Analysis of Sewage Sludge by Water Research Institute of National Council of Researches (IRSA-CNR, 1985), unless specified differently. Results are reported in **Table S1**. pH and electrical conductivity were measured on distilled water extracts (1:10 m/v) by potentiometry and conductometry, respectively. Dry matter content and humidity were measured by gravimetry, drying the samples at 105°C until constant weight. Ashes were determined with calcination at 550°C for 5 hours.

Total organic carbon was evaluated as reported on “Official methods of soil analysis” by the Italian Ministry of Agriculture and Forestry (Italian Ministerial Decree, 1999), exploiting the Walkley-Black method: sample digestion with potassium dichromate and sulphuric acid is followed by titration with iron(II) sulphate heptahydrate. Organic matter content was calculated with the Van Bemmelen conversion factor (1.724) (Pribyl, 2010).

Total nitrogen (N_{Tot}) was measured with the Kjeldahl method, which allows to titrate both organic and inorganic forms of nitrogen. Ammonium nitrogen (NH_4^+) was evaluated through distillation with magnesium oxide followed by titration with sulphuric acid, while nitrates ($N-NO_3^-$) were determined by the means of ionic chromatography. Organic nitrogen (N_{Org}) was then calculated by subtraction: $N_{Org} = N_{Tot} - (N-NH_4^+)$. Other macronutrients (K and P), micronutrients (Ca, Mg, Na, Fe, Mn, B, Zn) and heavy metals (Pb, Cr, Ni, Cu) were extracted with mineral acid digestion and then analysed by the means of inductively coupled plasma optical emission spectrometry (ICP-OES). Other contaminants such as Cd and As were extracted with the same digestion protocol, but analysed with graphite furnace atomic absorption spectroscopy (GF-AAS). Hg was evaluated with hydride generation atomic absorption spectroscopy (HGAAS) after microwave mineralisation, while Cr^{6+} was determined by colorimetry after complexation with diphenylcarbazide.

Table S1. Physicochemical properties of the four anaerobic digestates from sewage sludge used. Last three columns on right specify analysis methods for sewage sludge, Italian law limits for Land application of sewage sludges (Italian Decree Law 99/1992, n.d.), and law limits for heavy metals in fertilizers (Italian Decree Law 75/2010). d.m.b., Dry matter basis.

Parameter	Unit of measure	Anaerobic digestates				Method of analysis	Technique	Italian Law Land application of sewage sludge (D. Lgs 99/92)	Italian Law Discipline on fertilizers (D.Lgs 75/2010)
		Primary (P)	Secondary (S)	Centrifuged (C)	Dried (D)				
Dry matter	%	4.4	4.8	25.8	88.8	Calculation	Calculation		
Humidity	%	95.6	95.2	74.2	11.2	M.P.A.A.F., 2006 Method III.1	Gravimetry		
Ashes	% d.m.b.	35.3	31.5	36.1	35.6	Calculation	Calculation		
pH (1:10)		7.7	7.5	7.3	6.8	M.P.A.A.F., 2006 Method III.3	Potentiometry		
E.C.	mS/cm	0.378	0.36	1.069	1.575	M.P.A.A.F., 2006 Method III.4	Conductometry		
Organic matter	% d.m.b.	64.7	68.5	63.9	64.4	Calculation	Calculation		
TOC	% d.m.b.	37.5	39.7	37.1	37.3	D.M. 13/09/99 GU 248 21/10/199 met. VII.3	Walkley & Black method	>20	
N - Tot	% d.m.b.	7.4	7.5	6.3	5	CNR IRSA 6 Q64 vol.3, 1985	Kjeldahl method	>1.5	
N - Org	% d.m.b.	5.84	6.16	5.33	4.75	M.P.A.A.F., 2006 Method IV.12	Calculation		
N - NO ₃ ⁻	% d.m.b.	<0.01	<0.01	<0.01	<0.01	M.P.A.A.F., 2006 Method IV.12	Ionic chromatography		
N - NH ₄ ⁺	% d.m.b.	1.56	1.34	0.97	0.25	M.P.A.A.F., 2006 Method IV.12	Distillation and titration		
N - org / N - Tot	%	79	82	84	94	Calculation	Calculation		
C/N		5.1	5.3	5.9	7.4	Calculation	Calculation		
P	% d.m.b.	4.16	5.75	6.74	6.26	M.P.A.A.F., 2006 Method VIII	Acid digestion + ICP-OES	>0.4	
K	% d.m.b.	0.55	0.69	0.39	0.18	M.P.A.A.F., 2006 Method VIII	Acid digestion + ICP-OES		
Ca	% d.m.b.	6.46	4.69	5.02	4.64	M.P.A.A.F., 2006 Method VIII	Acid digestion + ICP-OES		
Mg	% d.m.b.	1.78	1.53	1.45	1.16	M.P.A.A.F., 2006 Method VIII	Acid digestion + ICP-OES		
Na	% d.m.b.	1.05	1.03	0.34	0.19	M.P.A.A.F., 2006 Method VIII	Acid digestion + ICP-OES		
B	mg/kg d.m.b.	51	60	52	41	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES		
Zn	mg/kg d.m.b.	918	650	849	719	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES	2500	500
Fe	% d.m.b.	2.43	3.32	3.99	3.48	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES		
Mn	mg/kg d.m.b.	255	190	268	228	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES		
Cu	mg/kg d.m.b.	357	340	406	396	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES	1000	230
Pb	mg/kg d.m.b.	92	70	92	79	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES	750	140
Cr	mg/kg d.m.b.	245	210	245	217	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES	<200*	
Cd	mg/kg d.m.b.	1	0.6	0.8	<0.1	M.P.A.A.F., 2006 Method IX	Acid digestion + GF-AAS	20	1.5
Ni	mg/kg d.m.b.	163	120	155	137	M.P.A.A.F., 2006 Method IX	Acid digestion + ICP-OES	300	100
As	mg/kg d.m.b.	2.8	2.1	0.9	<0.1	M.P.A.A.F., 2006 Method IX	Acid digestion + GF-AAS	<20*	
Hg	mg/kg d.m.b.	<0.1	<0.1	<0.1	<0.1	Internal method	HGAAS	10	1.5
Cr ⁶⁺	mg/kg d.m.b.	<0.1	<0.1	<0.1	<0.1	CNR IRSA 16 Q64 vol.3, 1986	Colorimetry	<2*	0.5

* Values introduced with Italian Law 130/2018

Section II: Method of analysis of sandy soil and peat substrate

The soil used in this study was sampled in Grugliasco (TO), Italy (45°03'58.4"N, 7°35'32.9"E). It was collected within 20 and 100 cm depth, sieved at 2 mm and not previously sterilized.

Physical and chemical analysis were performed according to the official methods of soil analysis of Italian Ministry of Agriculture and Forestry (Italian Ministerial Decree, 1999), except for available Fe, Mn, Cu and Zn. Stones were evaluated by sieving (2 mm) (Method II.1) while soil texture was determined by granulometry (wet sieve analysis; Method II.6). Measure of pH and electrical conductivity, organic matter, nitrogen forms and phosphorous was conducted on an aqueous extract obtained following the Sonneveld method (Sonneveld and Voogt, 2009). pH and electrical conductivity were measured by potentiometry (Method III.1) and conductometry (Method IV.1). Organic carbon was measured with the Walkley-Black method (Method VII.3). Organic matter content was calculated with the Van Bemmelen conversion factor (1.724) (Pribyl, 2010). Total nitrogen was measured with the Kjeldahl method (Method XIV.3). Mineral forms of nitrogen were extracted with an aqueous solution of KCl 2M (Method XIV.4); ammonium was measured through distillation (Method XIV.6), while nitrate and nitrite were quantified through continuous flux colorimetry (Griess-Ilosvay reaction; Method XIV.12 and XIV.13). Organic nitrogen and C/N ratio were obtained by calculation. Available phosphorous was determined by Olsen method (Method XV.3).

Measure of cation exchange capacity (C.E.C.) and exchangeable bases (Na, K, Mg, Ca) were performed on an extract obtained with an aqueous solution of BaCl₂ – triethanolamine (pH 8.2); C.E.C. was determined through complexometric titration (Method XIII.2) while exchangeable bases were measured by the means of flame atomic absorption spectroscopy (FAAS) (Method XIII.5).

Analysis of available Fe, Mn, Cu and Zn was performed according to Italian Ministerial Decree (Italian Ministerial Decree, 1992) Method 37, which exploits the Lindsay-Norwell method, that is an extraction through an aqueous solution of DTPA, CaCl₂ and triethanolamine (pH 7.3) followed by quantification with flame atomic absorption spectroscopy (FAAS).

Peat substrate was mixed with perlite and then sterilised before each application. Chemical characterization of peat substrate was performed on an aqueous extract 1:2 (v/v water/peat

substrate) according to Sonneveld method (Sonneveld and van den Ende, 1971). The analytical methods for peat analysis were all internal methods. pH and electrical conductivity were measured by potentiometry and conductometry. Total Nitrogen was evaluated with Kjeldahl method while organic nitrogen was calculated. Inorganic forms of nitrogen (ammonium, nitrite, nitrate) were measured by colorimetry (indophenol-blue method, diazotization method and dimethylphenol method respectively). Phosphorous were measured by colorimetry (molybdovanadate method). K, Mg, Ca, Na, Fe, Mn, Cu and Zn were measured through flame atomic absorption spectroscopy (FAAS).

Section III: Apparent balance of nitrogen in sandy soil

Apparent N balance in sandy soil (**Table S2**) was calculated according to Yang and co-workers (2020), with some modifications. Total nitrogen was considered as the sum of Kjeldahl nitrogen, nitrites and nitrates per pot. Total initial nitrogen was calculated as the sum of N present in nude sandy soil and N added with treatments. Stored nitrogen in soil was the N still present after two months; stored nitrogen in plants was the N measured in plants (epigeal part) two months after sowing. N loss was calculated as difference between initial and stored nitrogen. Statistical analysis of N loss highlighted no significant differences between treatments. With regards to peat substrate, nitrogen storage was not calculated since no significant differences were observed in biomass and ANUE results.

Table S2. Results of apparent balance of nitrogen in sandy soil. Nitrogen amounts are reported as mean values $\pm$ standard deviation.

Treatment	Initial nitrogen			Nitrogen storage		Loss of nitrogen	
	Soil nitrogen storage (g)	Nitrogen added (g)	Total nitrogen (g)	Soil (g)	Plant (mg)	(g)	%
Primary SSAD	1.16 $\pm$ 0.32	0.39	1.55 $\pm$ 0.32	0.76 $\pm$ 0.03	46.64 $\pm$ 7.45	0.75 $\pm$ 0.04	48.14 $\pm$ 2.51
Dried SSAD	1.16 $\pm$ 0.32	0.39	1.55 $\pm$ 0.32	0.87 $\pm$ 0.04	81.08 $\pm$ 15.63	0.60 $\pm$ 0.03	38.46 $\pm$ 2.24
Mineral fertilizer	1.16 $\pm$ 0.32	0.39	1.55 $\pm$ 0.32	0.91 $\pm$ 0.20	60.83 $\pm$ 9.26	0.58 $\pm$ 0.17	37.45 $\pm$ 11.17

Section IV: Agronomic nitrogen use efficiency (ANUE) calculation

Table S3. Agronomic nitrogen use efficiency (ANUE) on sandy soil samples. Data are reported as mean value $\pm$ standard deviation and are expressed in g gN⁻¹. Different letters indicate differences between treatments that are significant at P < 0.05 (Tukey HSD).

Month	SSAD												Mineral fertilizer							
	Primary				Secondary				Centrifuged								Dried			
1	1.26	±	0.39	d	1.66	±	0.50	d	0.28	±	0.16	d	0.43	±	0.35	d	0.07	±	0.05	d
2	9.32	±	2.09	c	11.60	±	2.99	c	12.43	±	2.51	c	13.25	±	2.93	c	3.74	±	2.17	d
3	18.89	±	1.99	b	25.09	±	1.76	a	24.85	±	3.70	a	27.99	±	3.45	a	11.65	±	2.26	c

Section V: Infra-red gas analyzer (IRGA)

Net photosynthesis (A_N) of tomato plants leaves grown on sandy soil showed significant differences between treatments (**Table S4**). The lowest A_N value was found in control ($4.08 \mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$), while the significantly highest values were recorded on C ($10.56 \mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$) and S ($10.21 \mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$). P, D and mineral fertilizer (M) displayed values comparable with both the lowest and highest ones. Moving to the stomatal conductance (g_s) and CO_2 concentration in substomatal cavity (C_i), no statistically significant difference was registered. The overall mean g_s value was $0.25 \text{ mmol H}_2\text{O m}^{-2}\text{s}^{-1}$ while the overall mean C_i value was 276.22 ppm.

As regards IRGA measurements on peat substrate, A_N did not show any significative difference between the treatments and overall mean calculated value was $8.64 \mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$. The stomatal conductance of the digestates and of the control was higher than on mineral fertilizer. Concerning C_i , no significant difference between the treatments was found; the overall mean of C_i value was 280.89 ppm.

Table S4. Results of infra-red gas analyzer (IRGA) on plants grown on sandy soil. Different letters indicate differences between treatments that are significant at $P < 0.05$ (Tukey HSD). Data are expressed as mean $\pm$ standard deviation. A_N : net assimilation ($\mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$); g_s : stomatal conductance ($\text{mmol H}_2\text{O m}^{-2} \text{ s}^{-1}$); C_i : CO_2 concentration in substomatal cavity (ppm).

Parameter Treatment	A_N ($\mu\text{mol CO}_2 \text{ m}^{-2}\text{s}^{-1}$)			g_s ($\text{mmol H}_2\text{O m}^{-2}\text{s}^{-1}$)			C_i (ppm)		
Control (T)	4.08	$\pm$ 1.87	b	0.17	$\pm$ 0.09		300.11	$\pm$ 10.19	
Primary digestate (P)	9.96	$\pm$ 1.12	ab	0.27	$\pm$ 0.06		264.00	$\pm$ 11.35	
Secondary digestate (S)	10.21	$\pm$ 1.48	a	0.28	$\pm$ 0.02		269.56	$\pm$ 8.18	
Centrifuged digestate (C)	10.56	$\pm$ 2.37	a	0.28	$\pm$ 0.04		260.22	$\pm$ 20.72	
Dry digestate (D)	8.29	$\pm$ 3.37	ab	0.26	$\pm$ 0.07		281.56	$\pm$ 27.84	
Mineral fertilizer (M)	7.30	$\pm$ 1.97	ab	0.24	$\pm$ 0.07		281.89	$\pm$ 12.36	

References

- IRSA-CNR, 1985. Metodi analitici per i fanghi - Parametri chimico-fisici (Methods for Analysis of Sewage Sludge - Chemical and physical parameters). Quaderno 64 vol. 3. Istituto di ricerca sulle acque, Consiglio Nazionale delle Ricerche (in Italian).
- Italian Decree Law 75/2010, n.d. Decreto Legislativo 29 aprile 2010, n. 75. “Riordino e revisione della disciplina in materia di fertilizzanti, a norma dell’articolo 13 della legge 7 luglio 2009, n. 88”.
- Italian Decree Law 99/1992, n.d. Decreto Legislativo 27 gennaio 1992, n. 99. “Attuazione della direttiva n. 86/278/CEE concernente la protezione dell’ambiente, in particolare del suolo, nell’utilizzazione dei fanghi di depurazione in agricoltura.”
- Italian Law 130/2018, n.d. Legge 16 novembre 2018, n. 130. “Disposizioni urgenti per la città di Genova, la sicurezza della rete nazionale delle infrastrutture e dei trasporti, gli eventi sismici del 2016 e 2017, il lavoro e le altre emergenze.”
- Italian Ministerial Decree, 1999. Decreto Ministeriale 13 settembre 1999. Approvazione dei “Metodi ufficiali di analisi chimica del suolo.”
- Italian Ministerial Decree, 1992. Decreto Ministeriale 11 maggio 1992. Approvazione dei “Metodi ufficiali di analisi chimica del suolo.”
- M.P.A.A.F., 2006. Metodi di analisi per i fertilizzanti (Analytical methods for fertilizers). Ministero delle Politiche Agricole Alimentari e Forestali. Osservatorio Nazionale Pedologico e per la Qualità del Suolo Agricolo e Forestale (coord. Trinchera A., Leita L., Sequi P.). CRA (in Italian).
- Pribyl, D.W., 2010. A critical review of the conventional SOC to SOM conversion factor. *Geoderma* 156, 75–83. <https://doi.org/10.1016/j.geoderma.2010.02.003>
- Sonneveld, C., van den Ende, J., 1971. Soil analysis by means of a 1:2 volume extract. *Plant Soil* 35, 505–516. <https://doi.org/10.1007/BF01372683>

Sonneveld, C., Voogt, W., 2009. Substrates: Chemical Characteristics and Preparation, in: Plant Nutrition of Greenhouse Crops. Springer Netherlands, Dordrecht, pp. 227–256.
https://doi.org/10.1007/978-90-481-2532-6_11

Yang, J., Liang, X., Li, H., Chen, Y., Tian, G., 2020. Effects of no-till and rice varieties on nitrogen and phosphorus balance in rice fields. *Environ. Sci. Eur.* 32. <https://doi.org/10.1186/s12302-020-00302-z>

Declaration of interests

☒ The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

☐ The authors declare the following financial interests/personal relationships which may be considered as potential competing interests:

Author contributions

Giulio Cristina: Conceptualization, Methodology, Writing - Original Draft

Enrico Camelin: Writing - Review & Editing, Visualization, Formal analysis, Software

Tonia Tommasi: Data Curation, Project administration

Massimo Pugliese: Conceptualization, Methodology, Validation, Resources, Supervision

Debora Fino: Funding acquisition, Investigation