

The ice-water interface and protein stability: a review

*Original*

The ice-water interface and protein stability: a review / Arsiccio, A.; Pisano, R.. - In: JOURNAL OF PHARMACEUTICAL SCIENCES. - ISSN 0022-3549. - STAMPA. - 109:7(2020), pp. 2116-2130. [10.1016/j.xphs.2020.03.022]

*Availability:*

This version is available at: 11583/2862754 since: 2021-01-18T16:53:08Z

*Publisher:*

Elsevier B.V.

*Published*

DOI:10.1016/j.xphs.2020.03.022

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Elsevier postprint/Author's Accepted Manuscript

© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:  
<http://dx.doi.org/10.1016/j.xphs.2020.03.022>

(Article begins on next page)

1 Authors' post-prints

2 Andrea Arsiccio, Roberto Pisano (2020). The Ice-Water Interface and Protein Stability: A Review. Journal of  
3 Pharmaceutical Sciences **109**(7):2116-2130.

4 \* Corresponding author: [roberto.pisano@polito.it](mailto:roberto.pisano@polito.it)

5

6 **The Ice-Water Interface and Protein Stability: A Review**

7 Andrea Arsiccio, Roberto Pisano\*

8

9 Department of Applied Science and Technology, Politecnico di Torino, 24 corso Duca degli Abruzzi, Torino,  
10 10129 Italy

11

12 \*E-mail of the corresponding author: roberto.pisano@polito.it

13 **ABSTRACT**

14 The ice-water interface is commonly encountered in our life, and comes into play in a wide number of natural  
15 phenomena. Here, attention will be focused on its effects on protein stability, with specific reference to the  
16 case of pharmaceutical proteins. This field represents a fascinating, and not yet fully understood, subject of  
17 investigation. Some background information on the ice-water phase diagram, as well as to the mechanisms  
18 of nucleation and crystal growth, will be provided. We will eventually discuss the effect of ice on protein  
19 activity, reviewing the mechanisms of ice-induced denaturation that have been proposed so far and  
20 discussing the strategies that may help prevent, or minimize, undesired loss of therapeutic activity.

21

22 **Keywords:** Crystallization, Proteins, Freeze-drying, Stabilization

23

24 **1. INTRODUCTION**

25 The ice surface is part of our everyday life and represents one of the most routinely encountered solid  
26 interfaces. Despite being a very common and abundant material on Earth, ice shows unique and intriguing  
27 properties that make it a fascinating subject of investigation. For this reason it plays a crucial role in a variety  
28 of fields; it is central in glaciology and snow physics, but is also relevant in astrophysics and atmospheric  
29 science, oceanography, cryo-biology and nucleation studies. Thanks to its availability, accessibility and  
30 inherent safety (i.e., absence of toxicity, flammability, explosivity), it represents a superb model system for  
31 phase transitions and deformation phenomena. The range of different scales involved in ice research, from  
32 molecular phenomena to atmospheric events, is impressive, and clearly shows the transversality of this  
33 topic.

34 It is therefore not surprising that the number of studies and publications in this area is already massive,  
35 and growing fast. Several books have been written on the physics and chemistry of ice [1, 2, 3, 4, 5, 6]. More  
36 than 3 million papers published to date deal with ice chemistry and physics, and an international symposium  
37 (the International Conference on the Physics and Chemistry of Ice, PCI [7]) has been focusing on all aspects  
38 of ice research for more than 55 years.

39 Considering the wide range of topics related to ice research discussed above, it would be impossible to  
40 satisfactorily cover all of them. This review will therefore focus on the effect of the ice-water interface on  
41 protein stability, a specific aspect that is more closely related with the authors' expertise.

42 Pharmaceutical proteins are becoming increasingly important in the treatment of a variety of human  
43 diseases, and are commonly stored in the frozen or freeze dried state to reduce risks of degradation before

44 administration. Lyophilized formulations are generally preferred over frozen ones, because do not require to  
45 maintain a controlled cold chain. About 30% of currently marketed biopharmaceuticals are introduced as  
46 lyophilized products. Liquid formulations are also commonly used, and comprise about 50-60% of all biologic  
47 dosage forms. Liquid formulations are preferred because of the low cost of manufacturing and ease of  
48 administration, but cannot be used for very unstable molecules [8, 9].

49 Both freezing and freeze-drying involve the interaction of the protein with the ice-water surface, and this  
50 may affect the final therapeutic potency, as will be discussed in the following. Understanding the  
51 mechanisms and effects of protein-ice interaction would be essential for numerous applications, especially in  
52 the pharmaceutical industry where this knowledge may help to minimize undesired loss of activity.

53 However, many questions remain open in this field, and a clear picture of the problem is not available yet.  
54 We will show how experimental evidence suggests the central role of the ice surface on protein stability  
55 during freezing, even though no definitive explanation is available about the underlying mechanisms.

56 We aim here to provide an as comprehensive as possible review on this subject, with the hope that it  
57 could serve as a basis to further advance knowledge in this area.

58 In order to do this, some background information will first be provided. The crystal structure of ice will be  
59 discussed, focusing in particular on hexagonal ice, which represents the most common form in everyday  
60 application. The concept of the quasi-liquid layer (QLL, also called liquid-like layer), i.e., the thin film of liquid  
61 water that is formed on the surface of ice crystals and which shows a peculiar behavior, will then be  
62 introduced. Some hints to the mechanisms of growth and nucleation will also be provided, introducing the  
63 experimental techniques that have been proposed in the literature to monitor these processes.

64 Finally, the effect of ice on protein stability will be addressed. Experimental evidence of ice-induced  
65 destabilization will be presented, covering the extensive literature on this topic. Possible mechanisms at the  
66 basis of these observations will be reviewed, also trying to clarify whether direct protein adsorption is  
67 necessary or not. In this context, some space will be given to two very peculiar classes of molecules, i.e.,  
68 antifreeze and ice nucleating proteins, whose unique behavior at the ice surface may help shed some light  
69 on the freezing of biopharmaceuticals, as well. Finally, the relevant variables that influence ice-induced  
70 denaturation of proteins, and may help to prevent or modulate it, will be described. Among them, the effect of  
71 surfactants, cryoprotectants, protein concentration, cooling/thawing rate and nucleation temperature will be  
72 covered.

73

74

## 75        **2. CRYSTAL STRUCTURE OF ICE**

### 76        **2.1 Different Ice Polymorphs**

77        Liquid water can freeze into several ice forms, depending on the temperature ( $T$ ) and pressure ( $P$ ) at which  
78        crystallization is occurring. As a result, at least 17 crystalline ice phases have been experimentally observed  
79        to date [10, 11, 12, 13, 14, 15]. The molecular structure of these phases is known, mostly thanks to neutron-  
80        diffraction crystallographic studies. The large number of possible structures is mostly due to the open  
81        tetrahedral arrangement of water, which can adapt to very different environments.

82        In these structures, each molecule is hydrogen-bonded to four neighboring water molecules. While in low-  
83        pressure forms, such as the common hexagonal ice Ih, the O-O-O angles are close to the ideal value of  
84        109.47°, these angles are distorted in higher pressure structures. For example, they range between 80° and  
85        129° in ice II. When the pressure becomes very high, interpenetrating networks are eventually formed, as is  
86        the case, for instance, of ice VII [12].

87        While the oxygen atoms show topological order (i.e., a tetrahedral coordination) in each phase, all the  
88        polymorphs that can be directly crystallized from the liquid are hydrogen-disordered. These include the so-  
89        called liquidus phases, namely ices Ih, III, V, VI, VII, and the two metastable phases IV and XII (see Figure  
90        1).

91        When temperature is reduced, these disordered phases tend to order and transform into fully hydrogen-  
92        ordered structures. However, hydrogen ordering needs molecular reorientations processes, which are  
93        hindered at low temperature. Therefore, only ices III and VII undergo this transition spontaneously, ordering  
94        into ices IX and VIII, respectively. By contrast, the formation of ices XI, XIII, XIV and XV needs to be  
95        promoted by doping with bases or acids [16, 14, 17]. In this context, ice II is peculiar, because it is the only  
96        hydrogen-ordered phase that shows no hydrogen-disordered counterpart [18]. At the same time, no  
97        hydrogen-ordered analogue has been identified so far for ices XVI and XVII.

98        The different structures of ice mentioned so far are also shown in Figure 2. It is evident that the ice crystal  
99        phases differ widely. For instance, the crystals structure may be hexagonal (ices Ih and XVII), cubic (ices VII,  
100        X, XVI), rhombohedral (ices II and IV), tetragonal (ices III, VI, VIII, IX, XII), monoclinic (ices V, XIII), or  
101        orthorhombic (ices XI, XIV, XV). Moreover, while ices Ih, XI, II, and XVII show open structures and relatively  
102        low density, the ices VII, VIII and X network is extremely dense.

103        Amorphous structures of ice were also identified, including amorphous solid water (ASW) [20],  
104        hyperquenched glassy water (HGW) [21], very high-density (VHDA) [22], high-density (HDA) [23] and low-  
105        density (LDA) amorphous ice [24]. In the past years, neutron-diffraction techniques showed that the three

106 lower density forms, namely ASW, HGW and LDA, share a very similar structure, which resembles a  
107 disordered, but tetrahedrally coordinated, fully hydrogen bonded network [25]. In contrast, the two denser  
108 forms, HDA and VHDA, are structurally different.

109 The landscape of possible ice structures is therefore extremely complex. However, the picture seems not  
110 complete yet, and many more ice polymorphs may be observed in the future. Several ice phases have been  
111 predicted by molecular dynamics (MD) simulations or density functional theory (DFT) computations [26, 27,  
112 28, 29, 30, 31, 32, 33, 34, 13]. For instance, 74963 potential ice structures have been recently identified by  
113 DFT [35], even though it is not clear if these structures may be prepared experimentally.

114

## 115 **2.2 A Focus on Hexagonal Ice Ih**

116 As previously mentioned, structure Ih is the ordinary form of ice, which is most easily observed in common  
117 experimental setup. Ice Ih displays a hexagonal lattice, relatively open and low-density. As a result, the large  
118 hexagonal rings characteristics of ice Ih leave almost enough room for an interstitial water molecule. In the  
119 ice Ih unit cell, the top and bottom faces are basal planes  $\{0001\}$ , while the six equivalent side faces are  
120 called primary prism faces  $\{10\bar{1}0\}$ . Secondary prism faces  $\{\bar{1}2\bar{1}0\}$  are also observed (see Figure 3).

121 Along with ice Ih, another metastable, hydrogen-disordered phase of ice exists at ambient pressure,  
122 namely, ice Ic, which is characterized by a cubic crystal [36, 2, 3, 37, 38], and shows a lower nucleation  
123 barrier compared to the Ih form [39]. Ice Ic is generally observed either in the Earth's atmosphere and in  
124 extra-terrestrial environments [40], or in small-size droplets at very low temperature [41]. It was further shown  
125 that ice Ic can be formed directly in the bulk water phase of a vitrified solution, for instance glucose at low  
126 temperature [42]. More recently, it was also hypothesized that ice Ic may be an intermediate phase in the  
127 pathway of heterogeneous ice nucleation [43]. However, it is not clear at present if fully cubic ice I (ice Ic)  
128 can be prepared. In fact, it was observed from diffraction data that ice Ic samples were not entirely cubic, but  
129 contained hexagonal stacking, as well [44]. This structure, where the hexagonal and the cubic stackings are  
130 mixed, is called stacking-disordered ice I (ice Isd), and is hydrogen-disordered. Interestingly, it was  
131 suggested that Isd may be even more stable than pure Ih ice [45]. No hydrogen-ordered counterpart has  
132 been identified so far for ice Isd.

133

134

135

136

137 **2.3 The Quasi-Liquid Layer (QLL)**

138 Ice interfaces may be disordered even below the equilibrium melting temperature, resulting in the so-called  
139 surface premelting [46, 3, 47, 12]. A quasi-liquid layer (QLL), that represents an intermediate state between  
140 the solid and bulk liquid, is therefore formed.

141 The extension of the QLL increases rapidly as temperature approaches the equilibrium melting value [46,  
142 48, 49, 50], because thermal fluctuations of the intermolecular distance become larger in these conditions.  
143 The actual thickness of the QLL as a function of temperature is not well known, but, as a rough estimate, it  
144 should be comparable to the lattice spacing at  $T \approx -10$  °C, and should rapidly increase approaching the  
145 melting temperature [46].

146

147 **3. ICE NUCLEATION AND GROWTH**

148 **3.1 Fundamentals of Nucleation**

149 During the freezing process, water needs to nucleate first, and the ice nuclei which are formed in this stage  
150 subsequently grow to form crystals. In this context, nucleation plays a central role [51].

151 According to the classic nucleation theory (CNT) [52], the free energy change  $\Delta G$  (see Figure 4)  
152 associated with the formation of a nucleus in a supercooled solution is the sum of a negative bulk  
153 contribution ( $-V\Delta g_v$ , where  $V$  is the volume of the nucleus), and a positive surface term ( $A\gamma$ , where  $A$  is the  
154 surface of the nucleus, and  $\gamma$  the interfacial tension),

155

$$\Delta G = -V\Delta g_v + A\gamma \quad (1)$$

156

157 For pure water, the ice/liquid interfacial tension is generally assumed to be on the order of 0.033 J/m<sup>2</sup>  
158 near the triple point, dropping to approximately 0.020 J/m<sup>2</sup> at -40°C [53].

159 Foreign bodies, e.g. the walls of the container, impurities or other particles present in the solution, can  
160 lower the interfacial free energy and reduce the energy barrier for nucleation. In this case, heterogeneous  
161 nucleation ensues, and a correlation factor  $f$  may be used to quantify the decrease in the energy barrier [54,  
162 55, 56],

$$f = \Delta G_{heter}^* / \Delta G_{homo}^* \quad (2)$$

163

164 where  $\Delta G_{heter}^*$  is the heterogeneous nucleation barrier (see Figure 4).



165 The factor  $f$  depends on the chemical compatibility between the nucleus and the substrate, which can be  
166 described in terms of the contact angle  $\theta$ . Moreover, it also depends on the radius of curvature  $r_s$  of the  
167 substrate. Defining two new variables  $m = (\gamma_{sf} - \gamma_{sc})/\gamma_{cf} \approx \cos \theta$  and  $r' = r_s/r_{cr}$ , it is therefore possible to  
168 write  $f = f(m, r')$ . In the previous equation for  $m$ ,  $\gamma_{sf}$ ,  $\gamma_{sc}$  and  $\gamma_{cf}$  are the interfacial free tension values  
169 between the substrate and the fluid, the substrate and the crystal or the crystal and the fluid, respectively.

170 While Ih ice is thermodynamically stable below 0 °C, homogeneous nucleation does not occur  
171 spontaneously until very low temperatures. It has been suggested that the onset of homogeneous nucleation  
172 may be -40 °C [57, 58, 59].

173 However, computational simulations predicted that no ice crystals would spontaneously form above -71  
174 °C [60], and also experimental results suggest that homogeneous nucleation scarcely occurs even far below  
175 -40 °C [61]. The possibility to reach this high degree of supercooling is due to the high energy barrier for  
176 homogeneous nucleation.

177 Because of this, ice crystallization is generally controlled by heterogeneous nucleation events. In this  
178 case, the presence of foreign particles lowers the energy barrier by decreasing the interfacial free energy.

179

### 180 **3.2 Understanding the Growth of Ice Crystals**

181 Ice crystals growth is governed by two physical phenomena, namely, surface attachment kinetics and the  
182 diffusion processes within the solution [47].

183 A key variable in this context is the degree of supercooling, which is often defined as  $\Delta T_{surf} = T_{eq} - T_{surf}$ , where  
184  $T_{eq}$  is the equilibrium freezing temperature, and  $T_{surf}$  the temperature at the ice-water surface during growth.

185 The growth rate  $v_g$  can then be written as,

186

$$v_g \approx K_g \Delta T_{surf} \quad (3)$$

187

188 where  $K_g$  is a kinetic coefficient. Ice crystal growth is influenced by the different growth rates on each facet,  
189 which in turn depends on their different step energies.

190 The step energy is defined as the energy required to create the edge of a one-molecule-high terrace on a  
191 faceted surface. For instance, the ice/water step energy on the basal face near the triple point is about  $5.6 \times$   
192  $10^{-13}$  J/m [62, 63] while is essentially zero for the prism faces. Experimental data suggest that the most  
193 stable ice-water interface at 0 °C is the secondary prismatic face (see Figure 3), followed by the primary

194 prismatic plane and, as a distant third, by the basal facet. This results in the typical hexagonal shape of ice Ih  
195 [64].

196 However, the morphology of the ice formed is extremely variable. For instance, it was reported to vary  
197 from circular disks when  $T_{surf} < 1$  °C, to dendritic plates when  $T_{surf} < 2$  °C, unbranched needle-like structures  
198 if  $T_{surf} \approx 4$  °C, branched needle-like structures if  $T_{surf} \approx 8$  °C and platelets if  $10$  °C  $< T_{surf} < 30$  °C [65]. Either  
199 the surface attachment kinetics, or the mass and heat transfer through the system may be the controlling  
200 mechanism, thus determining the final expression of the kinetic coefficient.

201 Figure 5 shows an example of a typical temperature profile during freezing of a pharmaceutical solution.  
202 The nucleation and growth phases have been highlighted. The liquid solution is first supercooled to a value  
203 below the equilibrium freezing point (segment A-\*). Supercooling represents a metastable state, during which  
204 water molecules tend to form clusters with long-living hydrogen bonds [66]. However, these clusters are still  
205 unstable, and break up quickly, as discussed in section 3.1. When the temperature is low enough to allow  
206 crossing of the energy barrier for the nucleation process, ice crystallization occurs rapidly in the whole  
207 product (point \* in Figure 5). In all pharmaceutical solutions heterogeneous nucleation is observed, and the  
208 degree of supercooling often lies in the range of 10-15 °C or more [67]. A sharp increase in product  
209 temperature to a value close to the equilibrium freezing point is observed at the onset of nucleation. The ice  
210 crystals growth (segment B-C) then proceeds through the addition of molecules to the interface. Here, the  
211 latent heat of crystallization is almost compensated for by the heat removed through the already frozen  
212 product, and the temperature remains nearly constant. Finally, to ensure complete solidification, the frozen  
213 product is typically cooled down to -40 / -50 °C (segment C-D).

214

#### 215 **4. MONITORING, PREDICTING AND CONTROLLING THE ICE CRYSTAL SIZE**

##### 216 **4.1 Monitoring the Freezing Process**

217 Larger ice crystals are generally preferred for protein-based pharmaceuticals [68], as will be discussed in  
218 detail in section 5, because this results in a smaller ice/freeze-concentrate surface area, and therefore  
219 reduced risk of adsorption.

220 The extension of the ice/freeze-concentrate surface is, therefore, a fundamental parameter for protein  
221 activity preservation during freezing, and should be strictly controlled. However, at present, this task cannot  
222 be easily achieved.

223 Generally, the parameters that mostly influence the ice/freeze-concentrate surface area are the cooling  
224 rate used during freezing, and the nucleation temperature. A high nucleation temperature, and/or a low

225 cooling rate, result in larger crystals, and hence smaller surface area [69, 70]. In this context, it should be  
226 considered that, in many fields, the range of cooling rates that can be explored is limited by equipment  
227 capabilities. For instance, common freeze dryers cannot achieve very fast cooling rates (e.g.,  $> 1 \div 2 \text{ K min}^{-1}$ ),  
228 and are therefore limited to slow cooling protocols.

229 The temperature profile during freezing of a pharmaceutical solution, which gives information about both  
230 nucleation temperature and cooling rate, may be monitored using several tools. For instance, thermocouples  
231 or resistance thermal detectors (RTDs) may be used. Thermocouples are typical in lab-scale equipment,  
232 while RTDs are common in production-scale ones because are more robust and can be sterilized [71, 72].  
233 Another option for temperature monitoring are the optical fiber sensors [73], where the sensing element  
234 shows a temperature-dependent refractive index. Alternatively, if wire connections should be avoided,  
235 passive transponders may be used, like the temperature remote interrogation system (TEMPRIS).

236 However, the presence of a sensor within the sample undergoing freezing could affect its ice nucleation  
237 behavior, therefore modifying the ice crystal size and ice-water surface area. For this reason, plasma  
238 sputtering was proposed to embed thermocouples within the containers walls, so as to avoid interaction with  
239 the product [74, 75]. Optical fiber sensors may be also used noninvasively, for instance fusing the fiber in the  
240 container bottom or embedding it in the controlled-temperature shelf where freezing is occurring [73].

241 Recently, the use of an infrared thermocamera was proposed to noninvasively monitor the freezing  
242 process [76], and was successfully combined with mathematical modelling to predict the final ice crystal size  
243 [77]. However, this method directly measures the temperature of the vial glass, and a model needs to be  
244 used to extract the actual product temperature [76]. Through vial impedance spectroscopy (TVIS) [78, 79] is  
245 another promising approach for the noninvasive monitoring of freezing, which exploits the variations in the  
246 bulk electrical properties during phase transformations.

247 These techniques make it possible to monitor the temperature profile, and some of them also the position  
248 of the freezing front and therefore the duration of the process. However, coupling with a modelling approach  
249 is needed to obtain information on the ice/freeze-concentrate surface area extension, as will be  
250 described in section 4.2.

251 During freezing, liquid water is separated as ice crystals, and the solutes that may be present in the  
252 system cryo-concentrate, forming a viscous matrix. If the ice crystals are then removed by sublimation, as it  
253 is done in the context of freeze drying, a porous structure is eventually formed in the freeze-concentrated  
254 matrix. The few techniques that are currently available to directly measure the extension of the ice/freeze-  
255 concentrate interface rely on the fact that the ice crystal size after freezing coincides with the pore size after

256 lyophilization, provided that neither collapse nor shrinkage is observed during drying. Therefore, the specific  
257 surface area of the dried cake provides a measure of the extension of the ice-water interface, and could be  
258 obtained by BET [80]. Scanning electron microscopy of lyophilized products (see Figure 6) may also be used  
259 to obtain information on the average pore size  $D_p$ , which could then be linked to the ice-water surface area  
260  $S_{iw}$ , for instance assuming that the crystals (and therefore also the pores) have cylindrical shapes,  
261

$$S_{iw} = \frac{4m}{D_p \rho} \quad (4)$$

262 where  $m$  is the mass of crystallizable water, and  $\rho$  the ice density. Image analysis could help in the  
263 automation of SEM image analysis, as well [81, 82].

264 Microscopy observation of frozen products was also suggested [83], and another promising experimental  
265 technique is the X-ray micro-computed tomography, which allows the reconstruction of the entire porous  
266 internal structure of lyophilized samples [84].

267 Monitoring the freezing process, and its impact on product morphology, is therefore possible. In the next  
268 section, we will briefly describe how these measurements may be integrated, or used to validate, modelling  
269 approaches for the prediction of the ice/freeze-concentrate surface area.

270

#### 271 **4.2 Modelling Approaches to Predict the Ice/Freeze-Concentrate Surface Area**

272

273 The prediction of ice crystal size generally requires knowledge of the temperature profiles within the solution  
274 being frozen. These could be measured, as described in the previous section, or extracted from simulations  
275 [85, 86], and may then be combined with empirical [87, 88, 89], or mechanistic [90, 77] models for crystal  
276 sizing. A simulation approach, that takes into account the variation in temperature during freezing, as well as  
277 cryoconcentration effects and the ice/freeze-concentrate interface formation, has been recently proposed to  
278 predict the risk of both cold- and ice-induced denaturation of proteins [91].

279 Numerical methods have also been developed to reproduce growth morphologies of ice crystals, for  
280 instance phase-field models [92, 93, 94], that should apply well to the growth of ice from liquid water [47].

281 Our understanding of the effect of the ice surface on protein stability may also substantially benefit from a  
282 deeper knowledge of the molecular-scale phenomena involved. For this purpose, molecular dynamics  
283 simulations represent a promising tool, which has been extensively used to simulate the ice-water interface

284 [95, 96, 97, 98, 99, 100, 101, 102, 103], and may provide some interesting information about protein/ice  
285 interaction, as will be discussed in detail in section 5.

286 The ability to predict the extension of the ice/freeze-concentrate surface is not enough if we cannot then  
287 control it, at least to some extent. This will therefore be the subject of the next section.

288

### 289 **4.3 Controlling the Ice/Freeze-Concentrate Surface Area**

290 If we consider that the ice/freeze-concentrate interface is a major source of protein instability during freezing,  
291 it is clear that the possibility to control its extension would be extremely beneficial. In principle, adsorption  
292 and denaturation phenomena may be minimized by promoting the formation of a small surface area. This  
293 could be achieved reducing the cooling rate, and inducing nucleation at a high temperature. However, while  
294 the cooling rate can be easily adjusted, the nucleation temperature is generally a stochastic variable, and its  
295 value is randomly distributed [104]. This results in huge heterogeneity, and limited possibility to control  
296 product morphology. This heterogeneity is hardly compatible with the stringent requirements of the  
297 pharmaceutical industry, and several techniques have been developed over the years to address this  
298 problem [67, 105, 106, 107].

299 An approach that may be used to increase the ice crystal size is annealing, which consists in holding the  
300 frozen product at a temperature above the glass transition for a given amount of time. During annealing,  
301 large ice crystals grow at the expense of smaller ones [108].

302 A number of techniques have also been developed, that allow control of the nucleation temperature. For  
303 instance, electrofreezing [109, 110] uses a high voltage pulse to trigger nucleation in supercooled water.  
304 However, individual electrodes need to be inserted in each sample, and in direct contact with the product,  
305 which is not compatible with GMP (Good Manufacturing Practice) conditions.

306 In the case of the ice-fog technique [111, 112, 113, 114, 115] nucleation is triggered by the introduction of  
307 small ice particles into the vials. These particles act as foreign bodies and promote heterogeneous  
308 nucleation, as discussed in section 3.1.

309 Another possible solution is ultrasound-induced ice nucleation [116, 117, 118, 119, 120, 121], where  
310 ultrasounds are used to start the nucleation process in a noninvasive way. However, it is not yet clear  
311 whether this technique may be applied to sensitive proteins, as the localized high temperatures generated by  
312 cavitation, and the bubble themselves, may potentially favor aggregation or other denaturation routes.

313 Two methods have also been developed, that are based on variations in pressure within the chamber. In  
314 the high-pressure-shift or depressurization method [122, 123, 124], the pressure is first increased to 1.5-4.5

315 bar, and then rapidly brought back to the atmospheric value. In the second approach, known as vacuum  
316 induced surface freezing, or vacuum induced nucleation [125, 126, 127, 128, 129, 130], pressure is reduced  
317 to induce nucleation by promoting evaporation from the surface of the product, and then quickly released to  
318 avoid undesired boiling or blow-up of the formulation.

319 Large ice crystals are obtained when using these controlled nucleation techniques, and a possible impact  
320 of that on protein stability will be addressed in section 5.4.

321

## 322 **5. ICE AND PROTEIN STABILITY**

### 323 **5.1 Evidence of Ice Effect on Protein Stability**

324 It is well known that the freezing process may have detrimental effects for a protein, leading to denaturation,  
325 aggregation and loss of biological activity. Numerous phenomena could contribute to the undesired  
326 conformational changes of the protein, including cold denaturation, cryo-concentration, or the formation  
327 of ice crystals [131, 132].

328 The physical environment of the protein is strongly affected by the cryo-concentration process, which  
329 induces variations in ionic strength and relative composition of solutes. pH shifts due to crystallization of  
330 buffer components [133], or phase separation phenomena [134, 135] may occur. On the one hand, if the  
331 protein being processed is stable in a narrow range of pHs, the precipitation of buffer components may lead  
332 to undesired denaturation. Actually, it was even shown that freeze-drying may result in remarkable changes  
333 in apparent acidity even without buffer crystallization. These changes in the apparent acidity may be related  
334 to changes in pK<sub>a</sub> upon water removal, or preferential inclusion of a basic component by ice crystals, and  
335 were observed to correlate with the degradation rates of acid-sensitive compounds [136]. On the other hand,  
336 the generation of a new interface during phase separation, as well as the possible partitioning of the protein  
337 into a phase with low concentration of stabilizers, may have detrimental consequences. The rate of some  
338 degradative reactions, such as hydrolysis and oxidation, may even increase during the freezing process

339 [137, 138, 139]. Recently it was proposed that the formation of water clusters at subzero temperature  
340 catalyzes proton transfer reactions, that are involved in many chemical degradation pathways like hydrolysis  
341 and deamidation [140].

342 At low temperature the hydration of nonpolar residues also becomes less unfavorable, and this leads to  
343 the possibility for water molecules to penetrate the protein structure, promoting the exposure of its  
344 hydrophobic core and the partial loss of secondary structure. This phenomenon is referred to as cold  
345 unfolding, and is driven by the reduced enthalpic barrier against water-hydrophobes interaction that occurs

346 when lowering the temperature. The result is the formation of compact, partially unfolded states of the  
347 protein, characterized by a high degree of solvent penetration [141, 142, 143, 144, 145, 146, 147]. A  
348 possible way to study the effects of supercooled water on protein stability is the confinement in nanoporous  
349 matrices, that inhibits crystallization. Exploiting this technique, it has been shown that the structural  
350 transitions of proteins are dictated by the thermodynamic and kinetic modifications of their hydration water  
351 [148, 149].

352 However, the formation of ice crystals was shown to represent the most critical destabilizing factor. A  
353 dramatic decrease in the average phosphorescence lifetime of the Trp-48 residue was detected in solutions  
354 of the azurin protein upon ice formation [150]. This suggested that protein destabilization occurred as soon  
355 as ice crystals were formed.

356 The denaturation of several proteins during freeze-thawing, including ciliary neurotropic factor (CNTF),  
357 phosphofructokinase (PFK), LDH, glutamate dehydrogenase (GDH), interleukin-1 receptor antagonist (IL-  
358 1ra) and tumor necrosis factor binding protein (TNFbp), was found to be mostly related to the increase in the  
359 area of the ice-water surface [151].

360 Lower recovery of catalase,  $\beta$ -galactosidase and LDH activity was observed during fast freezing [152],  
361 and the formation of insoluble human growth hormone (hGH) aggregates was also found to increase with  
362 increasing cooling rates [153]. This occurs because more rapid cooling leads to smaller ice crystals, which  
363 have a greater surface area to volume ratio than larger crystals.

364 Sarciaux et al. [154] proposed a mechanism of aggregate formation involving denaturation of bovine  
365 immunoglobulin (IgG) at the ice/freeze-concentrate interface. This denaturation was reversible upon freeze-  
366 thawing, but became irreversible after freeze-drying and reconstitution. Also in this case, quick cooling  
367 resulted in increased aggregation. Increasing the protein concentration improved the percentage of  
368 recovered protein [152, 154]. This occurs because the extension of the ice/freeze-concentrate interface is  
369 finite, and the number of protein molecules adsorbed at the surface cannot exceed a given value. When this  
370 value is reached, increasing the bulk concentration reduces the percentage of adsorbed

371 molecules [154].

372 Exploiting the change in Trp fluorescence between the native and denatured state, it was observed that  
373 the stability of the azurin mutant C112S from *Pseudomonas aeruginosa* was remarkably perturbed in ice  
374 [155]. Moreover, the extent of destabilization depended mainly on the size of the liquid water pool in  
375 equilibrium with ice. At the same time, it was observed that protein-ice interactions increased the solvent  
376 accessible surface area of the native fold, and/or decreased that of the denatured conformation [155].



377 In line with this observation, a remarkable loss of activity was observed in frozen lactate dehydrogenase  
378 (LDH) matrices. However, concentrated solutions at the same temperature and composition, but without ice,  
379 resulted in no degradation, suggesting that ice formation is the controlling factor for protein  
380 denaturation [68].

381 Similarly, an infrared spectroscopy investigation of LDH and human immune globulin (IgG) revealed an  
382 increase in intermolecular  $\beta$ -sheet structures close to the ice crystals, which is indicative of aggregation  
383 [156]. In contrast, the infrared spectra for these two protein molecules collected distant from the surface of  
384 ice crystals were very similar to spectra collected from the initial solution. This suggests that the vicinity of  
385 the ice surface is key to promote conformational changes. In a similar study, the aggregation of IgG2 at -10  
386 °C was related to the formation of the ice/freeze-concentrate interface [157].

387 Molecular dynamics simulations of hGH also suggested a change in protein conformation at the ice  
388 surface, with a significant increase in the non-polar surface area exposed, while no notable conformational  
389 change was detected in unfrozen solutions at the same temperature [158]. Partial unfolding of the GB1  
390 hairpin was similarly observed in metadynamics simulations at the ice-water interface [159].

391 Overall, these results suggest that prevention of ice-induced denaturation represents a key issue  
392 whenever a protein is subjected to a freezing process. In the next section, we will review the mechanisms  
393 that have been proposed so far in the literature to explain this phenomenon.

394

## 395 **5.2 How Does the Ice Interface Affect Protein Stability?**

396 In the previous section, evidence about the negative impact of ice formation on protein stability was  
397 presented. However, a question that spontaneously arises from the previous discussion is: what are the  
398 reasons at the basis of the ice-induced denaturation of proteins? What makes it difficult to answer is the  
399 absence of chemical differences between water and ice. Moreover, the structural differences are also  
400 minimal, as both water and ice are composed of a tetrahedral network of hydrogen bonds. Why two  
401 substances that are so similar would affect protein behavior in such a different way? This represents an  
402 intriguing and complex field of investigation, also because not many experimental techniques are currently  
403 available to fully address this problem.

404 Strambini and Gabellieri [150] suggested that the conformational changes of proteins may arise from the  
405 direct interaction between the protein and the ice surface, as schematized in Figure 7a. In this context,  
406 accumulation of milk proteins [160] and albumin [161, 162] near the ice surface was detected, using either  
407 light and transmission electron microscopy or confocal Raman microspectroscopy. Recombinant human



408 interferon- $\gamma$  (rhIFN- $\gamma$ ) was found to accumulate at the ice/liquid interface during lyophilization, but with  
409 significantly smaller intensity than at the air/liquid surface [163]. Also, accumulation of rhIFN- $\gamma$  to the  
410 ice/liquid surface alone was found to be not responsible for aggregation, and a subsequent drying step was  
411 necessary to induce particle formation [164].

412 However, unless the protein has an antifreeze behavior (see section 5.3), there is no real evidence of  
413 direct protein adsorption onto the ice crystals. By contrast, solid-state NMR studies seem to indicate that the  
414 hydration shell of non-antifreeze proteins does not freeze below the freezing temperature of the bulk solution  
415 [165, 166]. For instance, ubiquitin keeps its entire hydration shell even at -35 °C, and this prevents direct  
416 interaction with the ice surface [167]. It is generally believed that most proteins should behave like ubiquitin,  
417 and that their hydration shell should remain in the liquid form until a temperature which is much lower than  
418 the equilibrium freezing value [168]. No direct interaction with ice would hence be possible above this  
419 temperature.

420 High-resolution synchrotron X-ray diffraction results further suggest that bovine serum albumin tends to  
421 partition into the quasi-liquid layer above ice crystals (see section 2.3), but without being directly adsorbed  
422 onto the ice surface [169]. A similar behavior was observed for two other proteins, namely recombinant  
423 human albumin and a monoclonal antibody, in a more recent investigation [170].

424 A different behavior is generally observed only in presence of antifreeze proteins (AFPs) [170, 171, 172],  
425 whose peculiar behavior will be briefly addressed in section 5.3. Therefore, another explanation for the ice-  
426 induced denaturation of proteins, which is not related to direct adsorption at the ice-water surface, should  
427 exist.

428 In this context, it should be remembered that the physical microenvironment of a protein confined within  
429 the QLL may be substantially different compared to the bulk. More specifically, the stabilizer may remain in  
430 the bulk, and may not be able to exert anymore its cryoprotective effect. In this context, Bhatnagar et al.  
431 [170] proposed a picture of the protein/ice interaction where the concentration gradients in the liquid phase  
432 adjacent to the ice crystals depend on the mobility of the species. Larger species tend to concentrate closer  
433 to the ice surface, because of their lower mobility. At the same time, the pH in the QLL may decrease  
434 because of the negative charge on the surface of ice crystals [170, 173]. The electrical double layer which is  
435 eventually formed may result in an increased concentration of cations, including protons (see Figure 7b).

436 Another possible explanation involves the entrapment of air at the ice surface. Using fluorescence  
437 microscopy [156], it was observed that several air bubbles may remain trapped between ice crystals. Air  
438 bubbles have also been observed by optical microscopy [174] and small angle neutron scattering [175].

439 Therefore, proteins may denature at this hydrophobic air-water interface, as outlined in Figure 7c. A  
440 similar mechanism was also proposed in a recent work [132], where it was pointed out that the growth of ice  
441 crystals leads to an increase in the concentration of dissolved air gasses. The cryo-concentration of oxygen  
442 may accelerate oxidation reactions during freezing [176, 177]. It was also estimated that the extension of the  
443 air bubbles interface may be significantly larger than the already existing interface on the top of the container  
444 [132].

445 Recently, a disordered population of ice crystals was detected by high-resolution synchrotron X-ray  
446 diffraction in frozen solutions of recombinant human albumin, lysozyme, an insect AFP and a monoclonal  
447 antibody [170]. This disordered population of ice crystals was identified as a high pressure form of ice,  
448 tentatively Ice III or Ice IX (see section 2.1). This observation was explained by the volume expansion that  
449 may occur during ice formation, and which may result in mechanical stresses and high local pressure.  
450 Values of freeze-induced pressure exceeding 2 kBar have been theoretically estimated [132]. Pressure-  
451 induced unfolding of the protein may hence occur [178], and the combination of elevated pressure and low  
452 temperature may have a synergetic effect on protein unfolding [179] (see Figure 7d).

453 Finally, in a recent molecular dynamics investigation of protein L [180], the effect of the ice surface was  
454 explained as an enhancement of cold denaturation phenomena. The protein was found to be destabilized in  
455 presence of an ice slab, compared to the bulk solution at the same temperature but without ice. No direct  
456 interaction between the protein and the surface was evidenced, but the observed denaturing effect seemed  
457 to be mediated by the nearby layer of liquid (or, better, liquid-like) water molecules. These molecules were  
458 significantly slowed down by the presence of ice, and could form a significantly larger number of hydrogen  
459 bonds with the protein, especially with the nonpolar patches that are generally poorly hydrated. These  
460 hydrogen bonds were also remarkably strong, and promoted the solvent-penetration of protein L and  
461 consequent exposure of its hydrophobic core, which is a common feature of cold denaturation [147] (see  
462 Figure 7e).

463 In the next sections, we will first present two very peculiar classes of molecules, namely, antifreeze and  
464 ice-nucleating proteins, whose unique behavior may help us understand the phenomenology of ice-peptide  
465 interaction. Afterwards, possible approaches to counteract ice-induced denaturation of proteins will be  
466 discussed.

467  
468  
469

### 470 **5.3 The Peculiarity of Antifreeze (AFPs) and Ice-Nucleating (INPs) Proteins**

471 In this section, attention will be devoted to two specific classes of proteins, namely, antifreeze (AFPs) and  
472 ice-nucleating (INPs) proteins. These molecules have peculiar characteristics, and our general  
473 understanding of protein-ice interaction may substantially benefit from a deeper analysis of their behavior.

474 AFPs have the ability to bind to nascent ice nuclei, therefore inhibiting their growth [181, 182]. For  
475 instance, it was reported that AFPs can inhibit the growth of both basal and prism planes of ice I<sub>h</sub> [183].  
476 They represent the natural way many organisms protect themselves against freezing damage, by  
477 depressing, in a noncolligative way, the freezing point of water. In a broader perspective, AFPs represent a  
478 central element in the problem of ice recognition by biomolecules. What makes their role and action even  
479 more puzzling is the remarkable difference of AFP structure between different classes of organism. This  
480 indicates that the depression of the freezing point can be exerted by extremely different structural motifs.

481 A spontaneous question about AFPs is what mechanisms drive their interaction with ice. Since the ice  
482 surface is characterized by the presence of OH groups, it was first proposed that the hydrophilic groups on  
483 the ice-binding sites (IBS) of AFPs may hydrogen-bond with ice [182, 184]. However, many AFPs do not  
484 expose H-bonding groups at their IBS. On the contrary, it was observed that hydrophobic residues are  
485 largely present on the IBS [185]. It was therefore suggested that hydrophobic groups were crucial [186, 187],  
486 together with a flat-binding surface [188]. However, neither H-bonding nor hydrophobic groups can be the  
487 only explanation. H-bonding groups would provide affinity, but no specificity, as they tend to interact more  
488 with liquid than ice water [189]. At the same time, hydrophobic groups would make it possible to achieve  
489 specificity, but are not compatible enough with the hydrophilic ice surface. This implicates that both groups  
490 should be involved [190].

491 Exploiting a simulation approach, it was suggested that the arrangement of water molecules in proximity  
492 of the IBS may play a role in the action of AFPs [191, 192, 193]. In fact, both a slight increase in tetrahedral  
493 order [194] and slower relaxation dynamics were observed in the hydration water of AFPs [195, 196]. These  
494 water molecules seem to adopt a highly ordered structure, forming a clathrate-like structure, similar to ice,  
495 around the ice-binding site of AFPs [190, 189]. This structure surrounds the hydrophobic groups of the IBS,  
496 while being anchored at the edges by H-bonds to the polar groups. Therefore, the AFPs may bind to ice  
497 thank to this ice-like structure that is formed at their IBS. These ordered layer of water molecules may be  
498 released upon binding, leading to a net entropic gain [197, 198]. NMR suggests that only the portion of AFPs  
499 hydration shell at the IBS is in direct contact with the ice surface. The remaining part of the hydration shell  
500 does not undergo freezing, behaving similarly to the hydration shell of non-AFPs proteins [167].

501 A further question that may arise is whether this ice-like structure is always present, or, on the contrary,  
502 forms only close to the ice surface. Molecular simulations of TmAFP (a hyperactive insect AFP, which shows  
503 the best lattice matching with ice Ih, see Figure 8a) indicate that the second option is correct [199],  
504 suggesting that a preordering of hydration water is not necessary for ice recognition. The formation of the  
505 clathrate-like structure at the IBS is therefore probably induced by the close presence of an ice layer. This  
506 suggests that ice has an effect on the surrounding liquid water layers [172].

507 Another class of interesting molecules is represented by ice-nucleating proteins (INPs), that are used by  
508 several bacteria, such as *Pseudomonas syringae*, to trigger ice formation at temperatures close to the  
509 equilibrium value [200].

510 INPs are structurally and chemically similar to AFPs, but are considerably bigger and can result in the  
511 formation of molecular clusters. The larger surfaces that are eventually formed are capable of nucleating ice.  
512 A common example of INP is InaZ, produced by *Pseudomonas syringae*, which is approximately 1200  
513 residues-long [201]. Its structure includes degenerate octapeptide repeats, a subpopulation of which includes  
514 the sequence GSTXT(A/S), where X is an unconserved amino acid. Recently, *ab initio* structures of the  
515 GSTSTA segment were resolved by electron microdiffraction [202]. It was discovered that both homochiral  
516 and racemic GSTSTA (see Figure 8b-c) form amyloid-like protofibrils, that display antiparallel  $\beta$ -sheets.

517 Molecular dynamics simulations suggested that INPs may be anchored to the cell surface in such a way  
518 to expose ice-active sites to the surrounding water [204, 205]. It was proposed that threonine and serine  
519 amino acids, that are largely represented on these sites, should mimic the basal plane of ice. This is possible  
520 because of their OH groups and the presence of clathrate water molecules that effectively induce nucleation  
521 [190, 206].

522 Later on, sum frequency generation (SFG) spectroscopy was used to study the INP of *Pseudomonas*  
523 *syringae*, namely, the protein inaZ [207]. This study demonstrated that hydrogen bonding at the water-  
524 bacteria contact promotes structural ordering on the adjacent water network. The unique hydrophilic-  
525 hydrophobic motifs at the ice-nucleating site, together with the effective removal of latent heat due to  
526 nucleation, are highly effective in triggering the formation of stable ice crystals. Later on, it was also  
527 suggested that only large surfaces, like INPs clusters, may effectively order hydration water into ice-like  
528 structures [199].

529 *Pseudomonas syringae* may also be added to a solution being cooled down to trigger nucleation. This  
530 was done to promote the formation of nuclei at higher temperature [69, 208, 209, 210]. However, this

531 approach cannot provide a real control of the nucleation temperature, and cannot therefore substitute the  
532 techniques discussed in section 4.3.

533

#### 534 **5.4 How to Counteract Ice-Induced Denaturation of Proteins**

535 From the previous discussion, it is evident that the ice-induced denaturation of proteins should be adequately  
536 controlled to maximize the recovery of therapeutic activity after freezing. In this context, two main classes of  
537 strategies could be adopted; the first relies on the selection of suitable operating conditions, while the second  
538 involves optimization of the formulation.

539 From the point of view of the operating conditions, the two variables that mostly affect protein stability  
540 during freezing are cooling rate and nucleation temperature. As previously mentioned, a low cooling rate and  
541 a high nucleation temperature are associated with the formation of larger ice crystals, and therefore smaller  
542 ice/freeze-concentrate surface area. This, in turn, should minimize ice-induced denaturation of proteins. In  
543 line with these considerations, a correlation has often been observed between cooling rate and protein  
544 stability, with higher recovery at lower cooling rates [151, 152, 153]. However, this is not true for all proteins.  
545 In the case of particularly unstable molecules, such as myoglobin at low pH, a fast freezing protocol was  
546 found to maximize protein stability [91]. In this case, the formation of a cryoconcentrated matrix at high  
547 viscosity, where cold unfolding of the protein is kinetically inhibited, should be promoted as quickly as  
548 possible.

549 While the cooling rate can easily be controlled and adjusted, the nucleation temperature is a stochastic  
550 variable. However, as discussed in section 4.3, several strategies have been developed to address this  
551 problem. The impact of controlled nucleation techniques on protein stability has also been investigated in  
552 recent works. For instance, the controlled nucleation strategy based on the depressurization method was  
553 found to improve LDH stability after freeze-thawing [211]. The depressurization method applied to a highly  
554 concentrated monoclonal antibody also suppressed glass fogging, which is the undesired migration of  
555 protein solutions up on the inner walls of glass vials [212]. Another controlled nucleation technique, namely,  
556 the ice-fog method, was applied to monoclonal antibody formulations [213], and resulted in reduced particle  
557 formation in highly concentrated systems. However, the addition of polysorbates was more effective in  
558 decreasing particle level. Also, the authors observed no difference in particle formation between the  
559 controlled and spontaneously nucleated samples at low concentration [213]. The benefits of vacuum induced  
560 surface freezing (VISF) on the stability of the human growth hormone (hGH) were also investigated [214].

561 In this case, both HPLC-SEC and a cell-based potency assay indicated that there was no dramatic  
562 difference in the behaviour of hGH at low concentration when either VISF or spontaneous nucleation were  
563 used. In the case of very unstable molecules, such as myoglobin at low pH, the application of VISF proved to  
564 be detrimental. In this case, the benefits of a smaller ice/freeze-concentrate interface are offset by the  
565 necessity to introduce two holding steps at low temperature and low product viscosity during application of  
566 the VISF. During these equilibration stages, cold denaturation of myoglobin may quickly occur [91].

567 Finally, annealing is another technique that may affect protein stability. While it does not make it possible  
568 to control the nucleation temperature, it nevertheless promotes the formation of larger ice crystals. The  
569 resulting decrease in specific surface area of the ice/freeze-concentrate interface may improve protein  
570 stability. This was the case, for instance, of bovine IgG as model protein [154]. It should also be considered  
571 that the thawing process may also be dangerous for protein activity.

572 In this context, the thawing rate is a crucial parameter, as slow thawing promotes the recrystallization  
573 process, with small ice crystals growing into larger ones. While the growth of larger ice crystals during  
574 freezing, for instance by annealing, was found to be beneficial, recrystallization has been observed to be  
575 detrimental during thawing, as it promoted undesired interfacial or shear stress for proteins at the ice-water  
576 interface [215]. Moreover, during the process of ice melting, the amorphous cryo-concentrated phase will still  
577 not have the same composition as the initial fully-liquid solution. Problems due to unequal distribution of  
578 solutes may therefore occur during thawing, as well. For this reason, it is generally advisable to perform a  
579 fast thawing [132].

580 For what concerns the choice of the formulation, different possibilities exist to improve protein recovery  
581 after freezing. Typical biopharmaceutical formulations include cryo/lyo-protectants (e.g., sugars, polyols,  
582 amino acids, polymers), buffering species, to control the pH, and in many cases also surfactants, to  
583 counteract surface-induced denaturation, and bulking agents (e.g., mannitol, glycine), to improve cake  
584 resistance. A comprehensive description of the type/amount of excipients found in commercial products is  
585 out of the scope of this review. We will therefore focus on those case studies where formulation components  
586 have been shown to affect protein stability at the ice surface, starting from surfactants. For instance, it was  
587 observed that the addition of small amounts of Tween 80 or Tween 20 protected proteins from ice-induced  
588 denaturation [151, 163, 216]. Addition of the same surfactant to LDH and a human IgG formulation resulted  
589 in a decrease in intermolecular  $\beta$ -sheet structure formation at the ice surface, which is indicative of  
590 diminished aggregation [156]. Surfactants are amphiphilic molecules that tend to locate at interfaces, thereby  
591 sterically preventing protein adsorption [217, 218]. This may explain why they have often been reported to

592 prevent ice-induced denaturation of proteins. However, as discussed in section 5.2, there is increasing  
593 evidence that non-AFP proteins do not bind to the ice interface, but rather remain confined in the QLL above  
594 the surface. The steric repulsion exerted by surfactants that coat the ice interface may, therefore, be not  
595 enough to account for the observed stabilizing effect. Molecular dynamics simulations of the GB1 peptide  
596 suggest that surfactants tend to surround the protein at the ice surface, with their hydrophilic heads oriented  
597 towards the peptide [159]. On the one hand, this orientation prevents denaturation, because the exposure of  
598 the protein hydrophobic core is unfavorable in these conditions. On the other hand, the formation of a  
599 protein-surfactant complex hinders aggregation, as well. The stabilizing role of surfactants may hence be  
600 explained according to an orientation-dependent mechanism.

601 Apart from surfactants, cryoprotectants may be added, including sugars (for instance sucrose, trehalose,  
602 glucose, lactose), polyols (e.g., sorbitol and glycerol), polymers (albumin, dextran, polyvinylpyrrolidone and  
603 hydroxyethyl cellulose are common examples), and amino acids (such as glycine, proline, arginine etc.). For  
604 instance, Strambini and Gabellieri [150] observed that the addition of glycerol and sucrose reduced the  
605 protein conformational changes induced by ice-formation during freezing. They suggested that the stabilizing  
606 action of these excipients may be considered as a combined effect of decreasing the freezing temperature  
607 and reducing the adsorption affinity of the protein by coating the surface of ice. However, sterical exclusion  
608 from the ice interface can hardly be the only mechanism to explain the role of cryoprotectants. Actually, the  
609 opposite was often observed, with excipients being expelled in the bulk freeze-concentrated solution and  
610 promoting a closer approach between the protein and the ice surface [170, 162].

611 The crystallite size and microstrain for Ih crystals was also evaluated for different protein formulations  
612 [170]. A higher microstrain, compared to pure water, was observed in all protein-containing samples,  
613 indicating that proteins may promote the formation of defects during ice crystal growth. In contrast, the  
614 microstrain level was consistently lower in all non-protein samples (containing, for instance, sucrose or  
615 histidine), suggesting that the effect of excipients on the ice structure should be minimal. At the same time,  
616 SEM analysis of freeze-dried cakes shows that the choice of the formulation strongly affects the pore size,  
617 and, therefore, the extension of the ice/freeze-concentrate interface. It has been hypothesized that this may  
618 be related to a different energetic cost for forming the ice/freeze-concentrate interface (i.e., different  
619 interfacial tension) in presence of different cosolutes [90].

620 It was also suggested [150] that the cosolutes may exert their protective action by being preferentially  
621 excluded from the protein surface. This should make the unfolding process thermodynamically unfavorable  
622 [219]. In particular, molecular dynamics simulations suggested that exclusion from specific patches on the



623 protein surface should be crucial for protein stability at the ice surface [158, 180]. However, the mechanism  
624 of preferential exclusion should be prevailing only at the beginning of the freezing process, when  
625 cryoconcentration is not yet complete. During the last stages of freezing, the matrix that separates from ice is  
626 so concentrated that other mechanisms come into play [220, 221]. At high excipient concentration, the  
627 vitrification theory [222, 223], the water replacement scenario [224, 225] and the water entrapment  
628 hypothesis [226, 227, 228] apply. Vitrification invokes the formation of a viscous, glassy matrix [229] where  
629 protein movements are hindered, while water entrapment envisions the formation of a cage of excipient  
630 molecules around the protein, where water is entrapped and slowed down. While these two are kinetic  
631 mechanisms, the water replacement theory suggests that a thermodynamic stabilization applies, where the  
632 excipient substitutes water in satisfying the hydrogen-bonding requirements of the protein. All these  
633 mechanisms of stabilization are effective only if the stabilizer remains in the same amorphous phase where  
634 the protein is. Therefore, crystallization of the cryoprotectant may reduce protein stabilization, also because it  
635 results in another surface onto which the protein may adsorb and denature [216].

636 Another typical component of a protein formulation is the buffer, which should maintain the pH in a range  
637 of values where the active ingredient is stable. In this context, it should be remembered that some buffer  
638 species may undergo selective crystallization, and this could result in undesired pH shifts. This is the case,  
639 for instance, of sodium and potassium phosphate [230, 231].

640 Finally, as previously mentioned, a high protein concentration is beneficial to minimize the effect of the  
641 ice-water interface [152, 154]. On the one hand, the percentage of protein molecules that are in the region  
642 perturbed by the presence of ice decreases when the concentration is increased. On the other hand, volume  
643 exclusion effects may arise at higher concentration. For instance, the cold denaturation temperature of  $\beta$ -  
644 lactoglobulin was found to decrease significantly when increasing the protein concentration [232].

645

## 646 **6. CONCLUSIONS**

647 In this work, an overview of the chemical and physical properties of the ice-water interface, and of its effects  
648 on protein stability, was provided. Rather than giving answers, the objective of this work was to raise  
649 questions that, according to the authors opinion, are worth further investigation.

650 More specifically, the following points should be addressed:

- 651 i. Is the Ih form of ice the only crystal structure that is relevant for pharmaceutical applications? As  
652 evidenced using high-resolution synchrotron X-ray diffraction [170], other structures of the ice



653 phase diagram described in section 2.1 may be important, and this aspect should be further  
654 investigated.

655 ii. Do the advantages of controlled nucleation approaches make them worth application in the  
656 manufacturing of pharmaceuticals and biopharmaceuticals? At present, the controlled nucleation  
657 techniques discussed in section 4.3 are not applied by pharmaceutical industries in the  
658 production of commercially available drugs. Further investigations should be added to evaluate  
659 their effect on protein stability, and the first steps in this direction have been discussed in section  
660 5.4 .

661 iii. What are the mechanisms at the basis of ice-induced denaturation of proteins? Some hypotheses  
662 in this context have been described in section 5.2, and should further by addressed. The answer  
663 to this question may benefit from a deeper understanding of the peculiarity of antifreeze and ice-  
664 nucleating proteins, as discussed in section 8.

665 iv. What are the conditions that minimize the denaturing effect of ice on protein stability? Some  
666 possible strategies to prevent, or minimize, ice-induced denaturation of proteins have been  
667 described in section 5.4. However, a more robust approach to the selection of appropriate  
668 conditions for the freezing of protein-based therapeutics would be desirable. It is evident that  
669 many variables come into play, and it is therefore probably impossible to give a unique answer,  
670 valid for all classes of proteins. In this regard, the application of mathematical models, such as  
671 those mentioned in section 4.2, may help to reduce the number of experiments to be performed,  
672 providing at the same time a better understanding of the underlying phenomena. The modelling  
673 approaches should be multiscale, covering both the molecular mechanisms of protein  
674 conformational changes and the effect of process variables on the critical attributes of the  
675 product.

676

677 The hope of the authors is that this work may help to focus the attention of both industries and academies in  
678 this direction, indicating possible objectives and providing a basis for future investigations.

679

680 **REFERENCES**

- 681 [1] W. Kuhs, *Physics and Chemistry of Ice*, Special Publications, The Royal Society of Chemistry, 2007.
- 682 [2] C. A. Hobbs, P. V. Hobbs, *Ice Physics*, Clarendon Press, 1974.
- 683 [3] V. F. Petrenko, R. W. Whitworth, *Physics of Ice*, OUP Oxford, 1999.
- 684 [4] N. H. Fletcher, *The Chemical Physics of Ice*, Cambridge University Press, 2009.
- 685 [5] E. R. Pounder, J. A. Jacobs, J. T. Wilson, *The Physics of Ice*, Commonwealth and international library of  
686 science, technology, engineering, and liberal studies. Geophysics division, Elsevier Science, 2013.
- 687 [6] J. Wettlaufer, J. Dash, N. Untersteiner, *Ice Physics and the Natural Environment*, Nato ASI Subseries I.,  
688 Springer Berlin Heidelberg, 2013.
- 689 [7] T. Bartels-Rausch, M. Montagnat, *The physics and chemistry of ice*, *Philosophical Transactions of the*  
690 *Royal Society A: Mathematical, Physical and Engineering Sciences* 377 (2146) (2019) 20190138.
- 691 [8] V. Gervasi, R. D. Agnol, S. Cullen, T. McCoy, S. Vucen, A. Crean, *Parenteral protein formulations: An*  
692 *overview of approved products within the european union*, *Eur. J. Pharm. Biopharm.* 131 (2018) 8 – 24.
- 693 [9] B. K. Muralidhara, M. Wong, *Critical considerations in the formulation development of parenteral biologic*  
694 *drugs*, *Drug Discov. Today* 25 (2020) 574 – 581.
- 695 [10] C. G. Salzmann, *Advances in the experimental exploration of water's phase diagram*, *J. Chem. Phys.*  
696 150 (6) (2019) 060901.
- 697 [11] J. L. Finney, *Ice: Structures*, in: K. H. J. Buschow, R. W. Cahn, M. C. Flemings, B. Ilschner, E. J.  
698 Kramer, S. Mahajan (Eds.), *Encyclopedia of Materials: Science and Technology*, Vol. 5, Elsevier  
699 Science, Oxford, 2001, p. 4018.
- 700 [12] T. Bartels-Rausch, V. Bergeron, J. H. E. Cartwright, R. Escibano, J. L. Finney, H. Grothe, P. J.  
701 Gutierrez, J. Haapala, W. F. Kuhs, J. B. C. Pettersson, S. D. Price, C. I. Sainz-D'iaz, D. J. Stokes, G.  
702 Strazzulla, E. S. Thomson, H. Trinks, N. Uras-Aytemiz, *Ice structures, patterns, and processes: A view*  
703 *across the icefields*, *Rev. Mod. Phys.* 84 (2) (2012) 885–944.
- 704 [13] W. Zhu, Y. Huang, C. Zhu, H.-H. Wu, L. Wang, J. Bai, J. Yang, J. S. Francisco, J. Zhao, L.-F. Yuan, X.  
705 C. Zeng, *Room temperature electrofreezing of water yields a missing dense ice phase in the phase*  
706 *diagram*, *Nat. Commun.* 10 (2019) 1925.
- 707 [14] C. G. Salzmann, P. G. Radaelli, E. Mayer, J. L. Finney, *Ice XV: A new thermodynamically stable phase*  
708 *of ice*, *Phys. Rev. Lett.* 103 (10) (2009) 105701.
- 709 [15] A. Falenty, T. C. Hansen, W. F. Kuhs, *Formation and properties of ice XVI obtained by emptying a type*  
710 *sII clathrate hydrate*, *Nature* 516 (2014) 231–233.

- 711 [16] Y. Tajima, T. Matsuo, H. Suga, Phase transition in KOH-doped hexagonal ice, *Nature* 299 (1982) 810–  
712 812.
- 713 [17] C. G. Salzmann, P. G. Radaelli, A. Hallbrucker, E. Mayer, J. L. Finney, The preparation and structures  
714 of hydrogen ordered phases of ice, *Science* 311 (2006) 1758–1761.
- 715 [18] J. J. Shephard, B. Slater, P. Harvey, M. Hart, C. L. Bull, S. T. Bramwell, C. G. Salzmann, Doping-  
716 induced disappearance of ice II from water's phase diagram, *Nature Phys.* 14 (2018) 569–572.
- 717 [19] M. Masakazu, Y. Takuma, T. Hideki, Genice: Hydrogen-disordered ice generator, *J. Comput. Chem.* 39  
718 (1) (2018) 61–64.
- 719 [20] E. F. Burton, W. F. Oliver, J. C. McLennan, The crystal structure of ice at low temperatures, *Proc. R.*  
720 *Soc. London. A* 153 (1935) 166–172.
- 721 [21] E. Mayer, P. Bruggeller, Vitrification of pure liquid water by high pressure jet freezing, *Nature* 298 (1982)  
722 715–718.
- 723 [22] T. Loerting, C. Salzmann, I. Kohl, E. Mayer, A. Hallbrucker, A second distinct structural “state” of high-  
724 density amorphous ice at 77 K and 1 bar, *Phys. Chem. Chem. Phys.* 3 (2001) 5355–5357.
- 725 [23] O. Mishima, L. D. Calvert, E. Whalley, ‘Melting ice’ I at 77 K and 10 kbar: a new method of making  
726 amorphous solids, *Nature* 310 (1984) 393–395.
- 727 [24] O. Mishima, L. D. Calvert, E. Whalley, An apparently first-order transition between two amorphous  
728 phases of ice induced by pressure, *Nature* 314 (1985) 76–78.
- 729 [25] D. T. Bowron, J. L. Finney, A. Hallbrucker, I. Kohl, T. Loerting, E. Mayer, A. K. Soper, The local and  
730 intermediate range structures of the five amorphous ices at 80K and ambient pressure: A Faber-Ziman  
731 and Bhatia-Thornton analysis, *J. Chem. Phys.* 125 (2006) 194502.
- 732 [26] Y. Huang, C. Zhu, L. Wang, X. Cao, Y. Su, X. Jiang, S. Meng, J. Zhao, X. C. Zeng, A new phase  
733 diagram of water under negative pressure: The rise of the lowest-density clathrate s-III, *Sci. Adv.* 2  
734 (2016) e1501010.
- 735 [27] Y. Huang, C. Zhu, L. Wang, J. Zhao, X. C. Zeng, Prediction of a new ice clathrate with record low  
736 density: A potential candidate as ice XIX in guest-free form, *Chem. Phys. Lett.* 671 (2017) 186 – 191.
- 737 [28] T. Matsui, M. Hirata, T. Yagasaki, M. Matsumoto, H. Tanaka, Communication: Hypothetical ultralow-  
738 density ice polymorphs, *J. Chem. Phys.* 147 (2017) 091101.
- 739 [29] Y. Liu, L. Ojamae, Clathrate ice sL: a new crystalline phase of ice with ultralow density predicted by  
740 first-principles phase diagram computations, *Phys. Chem. Chem. Phys.* 20 (2018) 8333–8340.
- 741 [30] G. A. Tribello, B. Slater, M. A. Zwijnenburg, R. G. Bell, Isomorphism between ice and silica, *Phys.*

742 Chem. Chem. Phys. 12 (2010) 8597–8606.

743 [31] J. Russo, F. Romano, H. Tanaka, New metastable form of ice and its role in the homogeneous  
744 crystallization of water, *Nat. Mat.* 13 (2014) 733–739.

745 [32] C. J. Fennell, J. D. Gezelter, Computational free energy studies of a new ice polymorph which exhibits  
746 greater stability than ice Ih, *J. Chem. Theory Comput.* 1 (2005) 662–667.

747 [33] L. A. Baez, P. Clancy, Phase equilibria in extended simple point charge ice-water systems, *J. Chem.*  
748 *Phys.* 103 (1995) 9744–9755.

749 [34] M. Ji, K. Umemoto, C.-Z. Wang, K.-M. Ho, R. M. Wentzcovitch, Ultrahigh-pressure phases of H<sub>2</sub>O ice  
750 predicted using an adaptive genetic algorithm, *Phys. Rev. B* 84 (2011) 220105.

751 [35] E. A. Engel, A. Anelli, C. M. C. J. Pickard, R. J. Needs, Mapping uncharted territory in ice from zeolite  
752 networks to ice structures, *Nat. Commun.* 9 (2018) 2173.

753 [36] H. Konig, Eine kubische eismodifikation, *Z. Kristallogr.* 105 (1943) 279–286.

754 [37] I. Kohl, E. Mayer, A. Hallbrucker, The glassy water–cubic ice system: a comparative study by X-ray  
755 diffraction and differential scanning calorimetry, *Phys. Chem. Chem. Phys.* 2 (2000) 1579–1586.

756 [38] W. F. Kuhs, G. Genov, D. K. Staykova, T. Hansen, Ice perfection and onset of anomalous preservation  
757 of gas hydrates, *Phys. Chem. Chem. Phys.* 6 (2004) 4917–4920.

758 [39] T. Kobayashi, T. Kuroda, Snow crystals, in: I. Sunagawa (Ed.), *Morphology of Crystals*, Terra Scientific  
759 Publishing, Tokyo, 1987, p. 649.

760 [40] B. J. Murray, T. L. Malkin, C. G. Salzmann, The crystal structure of ice under mesospheric conditions, *J.*  
761 *Atmos. Sol.-Terr. Phys.* 127 (2015) 78–82.

762 [41] G. P. Johari, Water’s size-dependent freezing to cubic ice, *J. Chem. Phys.* 122 (2005) 194504.

763 [42] P. Thanatuksorn, K. Kajiwara, N. Murase, F. Franks, Freeze–thaw behaviour of aqueous glucose  
764 solutions—the crystallisation of cubic ice, *Phys. Chem. Chem. Phys.* 10 (2008) 5452–5458.

765 [43] C. Li, X. Gao, Z. Li, Surface energy-mediated multistep pathways for heterogeneous ice nucleation, *J.*  
766 *Phys. Chem. C* 122 (2018) 9474–9479.

767 [44] W. F. Kuhs, D. V. Bliss, J. L. Finney, High-resolution neutron powder diffraction study of ice Ic, *J. Phys.*  
768 *Colloques* 48 (1987) C1–631–C1–636.

769 [45] L. Lupi, A. Hudait, B. Peters, M. Gruunwald, R. G. Mullen, A. H. Nguyen, V. Molinero, Role of stacking  
770 disorder in ice nucleation, *Nature* 551 (2017) 218–222.

771 [46] J. G. Dash, A. W. Rempel, J. S. Wettlaufer, The physics of premelted ice and its geophysical  
772 consequences, *Rev. Mod. Phys.* 78 (2006) 695–741.

- 773 [47] K. G. Libbrecht, Physical dynamics of ice crystal growth, *Annu. Rev. Mater. Res.* 47 (1) (2017) 271–295.
- 774 [48] M. Elbaum, M. Schick, Application of the theory of dispersion forces to the surface melting of ice, *Phys.*  
775 *Rev. Lett.* 66 (1991) 1713–1716.
- 776 [49] M. Elbaum, S. Lipson, J. Dash, Optical study of surface melting on ice, *J. Cryst. Growth* 129 (1993) 491  
777 – 505.
- 778 [50] V. Sadtchenko, G. E. Ewing, Interfacial melting of thin ice films: An infrared study, *J. Chem. Phys.* 116  
779 (2002) 4686–4697.
- 780 [51] Z. Zhang, X.-Y. Liu, Control of ice nucleation: freezing and antifreeze strategies, *Chem. Soc. Rev.* 47  
781 (2018) 7116–7139.
- 782 [52] J. Frenkel, A general theory of heterophase fluctuations and pretransition phenomena, *J. Chem. Phys.*  
783 7 (1939) 538–547.
- 784 [53] C. A. Jeffery, P. H. Austin, Homogeneous nucleation of supercooled water: Results from a new equation  
785 of state, *J. Geophys. Res.* 102 (1997) 25269–25279.
- 786 [54] X. Y. Liu, Interfacial effect of molecules on nucleation kinetics, *J. Phys. Chem. B* 105 (47) (2001) 11550–  
787 11558.
- 788 [55] X. Y. Liu, - generic mechanism of heterogeneous nucleation and molecular interfacial effects, in: K.  
789 Sato, Y. Furukawa, K. Nakajima (Eds.), *Advances in Crystal Growth Research*, Elsevier Science B.V.,  
790 Amsterdam, 2001, pp. 42 – 61.
- 791 [56] X. Y. Liu, K. Maiwa, K. Tsukamoto, Heterogeneous two-dimensional nucleation and growth kinetics, *J.*  
792 *Chem. Phys.* 106 (5) (1997) 1870–1879.
- 793 [57] T. Koop, H. P. Ng, L. T. Molina, M. J. Molina, A new optical technique to study aerosol phase transitions:  
794 The nucleation of ice from H<sub>2</sub>SO<sub>4</sub> aerosols, *J. Phys. Chem. A* 102 (1998) 8924–8931.
- 795 [58] C. A. Knight, Adding to the antifreeze agenda, *Nature* 406 (2000) 249–251.
- 796 [59] P. L. Davies, B. D. Sykes, Antifreeze proteins, *Curr. Opin. Struct. Biol.* 7 (1997) 828 – 834.
- 797 [60] E. B. Moore, V. Molinero, Structural transformation in supercooled water controls the crystallization rate  
798 of ice, *Nature* 479 (2011) 506 – 508.
- 799 [61] X. Y. Liu, N. Du, Zero-sized effect of nano-particles and inverse homogeneous nucleation: Principles of  
800 freezing and antifreeze, *J. Biol. Chem.* 279 (2004) 6124–6131.
- 801 [62] A. S. Michaels, P. L. T. Brian, P. R. Sperry, Impurity effects on the basal plane solidification kinetics of  
802 supercooled water, *J. Appl. Phys.* 37 (1966) 4649–4661.
- 803 [63] K. G. Libbrecht, Toward a comprehensive model of snow crystal growth: 3. The correspondence

804 between ice growth from water vapor and ice growth from liquid water, arXiv:1407.0740 [cond-mat.mtrl-  
805 sci] (2014) electronic.

806 [64] M. J. Shultz, P. J. Bisson, A. Brumberg, Best face forward: Crystal-face competition at the ice–water  
807 interface, *J. Phys. Chem. B* 118 (2014) 7972–7980.

808 [65] A. Shibkov, Y. Golovin, M. Zheltov, A. Korolev, A. Leonov, Morphology diagram of nonequilibrium  
809 patterns of ice crystals growing in supercooled water, *Physica A: Statistical Mechanics and its*  
810 *Applications* 319 (2003) 65–79.

811 [66] M. Matsumoto, S. Saito, I. Ohmine, Molecular dynamics simulation of the ice nucleation and growth  
812 process leading to water freezing, *Nature* 416 (2002) 409–413.

813 [67] J. C. Kasper, W. F. Friess, The freezing step in lyophilization: Physico-chemical fundamentals, freezing  
814 methods and consequences on process performance and quality attributes of biopharmaceuticals, *Eur.*  
815 *J. Pharm. Biopharm.* 78 (2011) 248–263.

816 [68] B. S. Bhatnagar, M. J. Pikal, H. B. Robin, Study of the individual contributions of ice formation and  
817 freeze-concentration on isothermal stability of lactate dehydrogenase during freezing., *J. Pharm. Sci.* 97  
818 (2008) 798–814.

819 [69] J. A. Searles, J. F. Carpenter, T. W. Randolph, The ice nucleation temperature determines the primary  
820 drying rate of lyophilization for samples frozen on a temperature-controlled shelf, *J. Pharm. Sci.* 90  
821 (2001) 860–871.

822 [70] A. Arsiccio, R. Pisano, Application of the quality by design approach to the freezing step of freeze-  
823 drying: Building the design space, *J. Pharm.Sci.* 107 (2018) 1586–1596.

824 [71] D. Fissore, R. Pisano, A. A. Barresi, Process analytical technology for monitoring pharmaceuticals  
825 freeze-drying – a comprehensive review, *Dry. Technol.* 36 (15) (2018) 1839–1865.

826 [72] S. Nail, S. Tchessalov, E. Shalaev, A. Ganguly, E. Renzi, F. Dimarco, L. Wegiel, S. Ferris, W. Kessler,  
827 M. Pikal, G. Sacha, A. Alexeenko, T. N. Thompson, C. Reiter, J. Searles, P. Coiteux, Recommended  
828 best practices for process monitoring instrumentation in pharmaceutical freeze drying—2017, *AAPS*  
829 *PharmSciTech* 18 (2017) 2379–2393.

830 [73] J. C. Kasper, M. Wiggernhorn, M. Resch, W. Friess, Implementation and evaluation of an optical fiber  
831 system as novel process monitoring tool during lyophilization, *Eur. J. Pharm. Biopharm.* 83 (3) (2013)  
832 449 – 459.

833 [74] S. Grassini, M. Parvis, A. A. Barresi, Inert thermocouple with nanometric thickness for lyophilization  
834 monitoring, *IEEE Trans. Instrum. Meas.* 62 (5) (2013) 1276–1283.

- 835 [75] I. Oddone, D. Fulginiti, A. A. Barresi, S. Grassini, R. Pisano, Non-invasive temperature monitoring in  
836 freeze drying: Control of freezing as a case study, *Drying Technology* 33 (13) (2015) 1621–1630.
- 837 [76] D. Colucci, R. Maniaci, D. Fissore, Monitoring of the freezing stage in a freeze-drying process using IR  
838 thermography, *Int. J. Pharm.* 566 (2019) 488 – 499.
- 839 [77] D. Colucci, D. Fissore, A. A. Barresi, R. D. Braatz, A new mathematical model for monitoring the  
840 temporal evolution of the ice crystal size distribution during freezing in pharmaceutical solutions, *Eur. J.*  
841 *Pharm. Biopharm.* 148 (2020) 148–159.
- 842 [78] G. Smith, E. Polygalov, Through vial impedance spectroscopy (TVIS): A novel approach to process  
843 understanding for freeze-drying cycle development, in: K. R. Ward, P. Matejtschuk (Eds.), *Lyophilization*  
844 *of Pharmaceuticals and Biologicals: New Technologies and Approaches*, Springer New York, New York,  
845 NY, 2019, pp. 241–290.
- 846 [79] G. Smith, Y. Jeeraruangrattana, A new method for determining the ice nucleation temperature and the  
847 solidification end point, in: D. Fissore, R. Pisano, A. Barresi (Eds.), *Freeze Drying of Pharmaceutical*  
848 *Products*, CRC Press, 2019.
- 849 [80] S. Rambhatla, R. Ramot, C. Bhugra, M. J. Pikal, Heat and mass transfer scale-up issues during freeze  
850 drying: II. Control and characterization of the degree of supercooling, *AAPS PharmSciTech* 5 (2004) 54–  
851 62.
- 852 [81] S. Grassini, R. Pisano, A. Barresi, E. Angelini, M. Parvis, Frequency domain image analysis for the  
853 characterization of porous products, *Measurement* 94 (2016) 515 – 522.
- 854 [82] A. Arsiccio, A. C. Sparavigna, R. Pisano, A. A. Barresi, Measuring and predicting pore size distribution  
855 of freeze-dried solutions, *Drying Technol.* 37 (4) (2019) 435–447.
- 856 [83] H. Goshima, G. Do, K. Nakagawa, Impact of ice morphology on design space of pharmaceutical freeze-  
857 drying, *J. Pharm. Sci.* 105 (2016) 1920 –1933.
- 858 [84] R. Pisano, A. A. Barresi, L. C. Capozzi, G. Novajra, I. Oddone, C. Vitale-Brovarone, Characterization of  
859 the mass transfer of lyophilized products based on x-ray micro-computed tomography images, *Dry.*  
860 *Technol.* 35 (2017) 933–938.
- 861 [85] K. Nakagawa, A. Hottot, S. Vessot, J. Andrieu, Modeling of freezing step during freeze-drying of drugs in  
862 vials, *AIChE J.* 53 (2007) 1362–1372.
- 863 [86] R. Pisano, L. C. Capozzi, Prediction of product morphology of lyophilized drugs in the case of vacuum  
864 induced surface freezing, *Chem. Eng. Res. Des.* 125 (2017) 119 – 129.
- 865 [87] M. Kochs, C. Korber, B. Nunner, I. Heschel, The influence of the freezing process on vapour transport

866 during sublimation in vacuum-freeze-drying, *Int. J. Heat Mass Transf.* 34 (1991) 2395 – 2408.

867 [88] J. L. Bomben, C. J. King, Heat and mass transport in the freezing of apple tissue, *Int. J. Food Sci.*  
868 *Technol.* 17 (1982) 615–632.

869 [89] W. Kurz, D. Fisher, *Fundamentals of solidification*, Trans Tech Publications, 1986.

870 [90] A. Arsiccio, A. A. Barresi, R. Pisano, Prediction of ice crystal size distribution after freezing of  
871 pharmaceutical solutions, *Cryst. Growth Des.* 17 (2017) 4573–4581.

872 [91] A. Arsiccio, P. Giorcello, L. Marengo, R. Pisano, Considerations on protein stability during freezing and  
873 its impact on the freeze-drying cycle: A design space approach, *J. Pharm. Sci.* 109 (2020) 464 – 475.

874 [92] A. Karma, W.-J. Rappel, Phase-field method for computationally efficient modeling of solidification with  
875 arbitrary interface kinetics, *Phys. Rev. E* 53 (1996) R3017–R3020.

876 [93] T. Takaki, Phase-field modeling and simulations of dendrite growth, *ISIJ Int.* 54 (2014) 437–444.

877 [94] J. W. Barrett, H. Garcke, R. Nürnberg, Stable phase field approximations of anisotropic solidification,  
878 *IMA J. Numer. Anal.* 34 (2013) 1289–1327.

879 [95] O. A. Karim, A. Haymet, The ice/water interface, *Chem. Phys. Lett.* 138 (1987) 531 – 534.

880 [96] O. A. Karim, A. D. J. Haymet, The ice/water interface: A molecular dynamics simulation study, *J. Chem.*  
881 *Phys.* 89 (1988) 6889–6896.

882 [97] H. Nada, Y. Furukawa, Anisotropic growth kinetics of ice crystals from water studied by molecular  
883 dynamics simulation, *J. Cryst. Growth* 169 (1996) 587 – 597.

884 [98] D. Rozmanov, P. G. Kusalik, Anisotropy in the crystal growth of hexagonal ice, *Ih*, *J. Chem. Phys.* 137  
885 (2012) 094702.

886 [99] M. Seo, E. Jang, K. Kim, S. Choi, J. S. Kim, Understanding anisotropic growth behavior of hexagonal ice  
887 on a molecular scale: A molecular dynamics simulation study, *J. Chem. Phys.* 137 (2012) 154503.

888 [100] M. Carignano, E. Baskaran, P. Shepson, I. Szleifer, Molecular dynamics simulation of ice growth  
889 from supercooled pure water and from salt solution, *Ann. Glaciol.* 44 (2006) 113–117.

890 [101] J. A. Hayward, A. D. J. Haymet, The ice/water interface: Molecular dynamics simulations of the  
891 basal, prism,  $20\bar{2}1$ , and  $2\bar{1}\bar{1}0$  interfaces of ice *Ih*, *J. Chem. Phys.* 114 (2001) 3713–3726.

892 [102] T. Bryk, A. D. J. Haymet, Ice *Ih*/water interface of the SPC/E model: Molecular dynamics simulations  
893 of the equilibrium basal and prism interfaces, *J. Chem. Phys.* 117 (2002) 10258–10268.

894 [103] J. L. F. Abascal, E. Sanz, R. García Fernández, C. Vega, A potential model for the study of ices  
895 and amorphous water: TIP4P/Ice, *J. Chem. Phys.* 122 (23) (2005) 234511.

896 [104] L. C. Capozzi, R. Pisano, Looking inside the ‘black box’: Freezing engineering to ensure the quality



897 of freeze-dried biopharmaceuticals., *Eur. J. Pharm. Biopharm.* 129 (2018) 58–65.

898 [105] R. Pisano, Alternative methods of controlling nucleation in freeze drying, in: K. R. Ward, P.  
899 Matejtschuk (Eds.), *Lyophilization of Pharmaceuticals and Biologicals: New Technologies and*  
900 *Approaches*, Springer New York, New York, NY, 2019, pp. 79–111.

901 [106] R. Pisano, A. Arsiccio, K. Nakagawa, A. A. Barresi, Tuning, measurement and prediction of the  
902 impact of freezing on product morphology: A step toward improved design of freeze-drying cycles, *Dry.*  
903 *Technol.* 37 (2019) 579–599.

904 [107] R. Geidobler, G. Winter, Controlled ice nucleation in the field of freeze-drying: Fundamentals and  
905 technology review, *Eur. J. Pharm. Biopharm.* 85 (2013) 214–222.

906 [108] J. A. Searles, J. F. Carpenter, T. W. Randolph, Annealing to optimize the primary drying rate, reduce  
907 freezing-induced drying rate heterogeneity, and determine  $T_g$  in pharmaceutical lyophilization, *J.*  
908 *Pharm. Sci.* 90 (2001) 872–887.

909 [109] W. Rau, Eiskeimbildung durch Dielektrische Polarisaton, *Zeitschrift fur Naturforschung A* 6 (1951)  
910 649–657.

911 [110] A. Petersen, H. Schneider, G. Rau, B. Glasmacher, A new approach for freezing of aqueous  
912 solutions under active control of the nucleation temperature, *Cryobiology* 53 (2006) 248–257.

913 [111] S. M. Patel, C. Bhugra, M. J. Pikal, Reduced pressure ice fog technique for controlled ice nucleation  
914 during freeze-drying, *AAPS PharmSciTech* 10 (2009) 1406–1411.

915 [112] R. Geidobler, S. Mannschedel, G. Winter, A new approach to achieve controlled ice nucleation of  
916 supercooled solutions during the freezing step in freeze-drying, *J. Pharm. Sci.* 101 (2012) 4409–4413.

917 [113] M. Umbach, Freeze drying plant, EP Patent 3093597 B1 (December 2017).

918 [114] W. Ling, Controlled nucleation during freezing step of freeze drying cycle using pressure differential  
919 ice crystals distribution from condensed frost, US Patent 8875413 B2 (November 2014).

920 [115] I. Vollrath, W. Friess, A. Freitag, A. Hawe, G. Winter, Comparison of ice fog methods and monitoring  
921 of controlled nucleation success after freeze-drying, *Int. J. Pharm.* 558 (2019) 18 – 28.

922 [116] T. Inada, X. Zhang, A. Yabe, Y. Kozawa, Active control of phase change from supercooled water to  
923 ice by ultrasonic vibration, part 1: Control of freezing temperature, *Int. J. Heat Mass Transfer* 44 (2001)  
924 4523–4531.

925 [117] X. Zhang, T. Inada, A. Yabe, S. Lu, Y. Kozawa, Active control of phase change from supercooled  
926 water to ice by ultrasonic vibration, part 2: Generation of ice slurries and effect of bubble nuclei, *Int. J.*  
927 *Heat Mass Transfer* 44 (2001) 4533–4539.

- 928 [118] M. Saclier, R. Peczalski, J. Andrieu, A theoretical model for ice primary nucleation induced by  
929 acoustic cavitation, *Ultrason. Sonochem.* 17 (2010) 98–105.
- 930 [119] M. Saclier, R. Peczalski, J. Andrieu, Effect of ultrasonically induced nucleation on ice crystals size  
931 and shape during freezing in vials, *Chem. Eng. Sci.* 65 (2010) 3064–3071.
- 932 [120] K. Nakagawa, A. Hottot, S. Vessot, J. Andrieu, Influence of controlled nucleation by ultrasounds on  
933 ice morphology of frozen formulations for pharmaceutical proteins freeze-drying, *Chem. Eng. Process.*  
934 45 (2006) 783–791.
- 935 [121] S. Passot, I. C. Trelea, M. Marin, M. Galan, G. J. Morris, F. Fonseca, Effect of controlled ice  
936 nucleation on primary drying stage and protein recovery in vials cooled in a modified freeze-dryer, *J.*  
937 *Biochem. Eng.* 131 (2009) 074511.
- 938 [122] T. H. Gasteyer, R. R. Sever, B. Hunek, N. Grinter, M. L. Verdone, Lyophilization system and method,  
939 US Patent 9651305 B2 (May 2017).
- 940 [123] R. Bursac, R. Sever, B. Hunek, A practical method for resolving the nucleation problem in  
941 lyophilization, *Bioproc. Int.* 7 (2009) 6672.
- 942 [124] A. K. Konstantinidis, W. Kuu, L. Otten, S. L. Nail, R. R. Sever, V. Bons, D. Debo, M. J. Pikal,  
943 Controlled nucleation in freeze-drying: Effects on pore size in the dried product layer, mass transfer  
944 resistance, and primary drying rate, *J. Pharm. Sci.* 100 (2011) 3453–3470.
- 945 [125] M. Kramer, B. Sennhenn, G. Lee, Freeze-drying using vacuum-induced surface freezing, *J. Pharm.*  
946 *Sci.* 91 (2002) 433–443.
- 947 [126] J. Liu, T. Viverette, M. Virgin, M. Anderson, P. Dalal, A study of the impact of freezing on the  
948 lyophilization of a concentrated formulation with a high fill depth, *Pharm. Dev. Technol.* 10 (2005) 261–  
949 272.
- 950 [127] I. Oddone, R. Pisano, R. Bullich, P. Stewart, Vacuum-induced nucleation as a method for freeze-  
951 drying cycle optimization, *Ind. Eng. Chem. Res.* 53 (2014) 18236–18244.
- 952 [128] I. Oddone, P.-J. Van Bockstal, T. De Beer, R. Pisano, Impact of vacuum-induced surface freezing on  
953 inter- and intra-vial heterogeneity, *Eur. J. Pharm. Biopharm.* 103 (2016) 167–178.
- 954 [129] I. Oddone, A. A. Barresi, R. Pisano, Influence of controlled ice nucleation on the freeze-drying of  
955 pharmaceutical products: The secondary drying step, *Int. J. Pharm.* 524 (2017) 134–140.
- 956 [130] A. Arsiccio, A. A. Barresi, T. De Beer, I. Oddone, P.-J. Van Bockstal, R. Pisano, Vacuum induced  
957 surface freezing as an effective method for improved inter- and intra-vial product homogeneity, *Eur. J.*  
958 *Pharm. Biopharm.* 128 (2018) 210–219.

- 959 [131] B. S. Bhatnagar, R. H. Bogner, M. J. Pikal, Protein stability during freezing: Separation of stresses  
960 and mechanisms of protein stabilization, *Pharm. Dev. Technol.* 12 (5) (2007) 505–523.
- 961 [132] J.-R. Authelin, M. A. Rodrigues, S. Tchessalov, S. Singh, T. McCoy, S. Wang, E. Shalaev, Freezing  
962 of biologicals revisited: scale, stability, excipients, and degradation stresses, *J. Pharm. Sci.* 109 (2020)  
963 44–61.
- 964 [133] K. A. Pikal-Cleland, N. Rodriguez-Hornedo, G. L. Amidon, J. F. Carpenter, Protein denaturation  
965 during freezing and thawing in phosphate buffer systems: Monomeric and tetrameric  $\beta$ -galactosidase,  
966 *Arch. Biochem. Biophys.* 384 (2000) 398–406.
- 967 [134] M. C. Heller, J. F. Carpenter, T. W. Randolph, Effects of phase separating systems on lyophilized  
968 hemoglobin, *J. Pharm. Sci.* 85 (1996) 1358–1362.
- 969 [135] M. C. Heller, J. F. Carpenter, T. W. Randolph, Manipulation of lyophilization induced phase  
970 separation: Implications for pharmaceutical proteins, *Biotechnol. Prog.* 13 (1997) 590–596.
- 971 [136] C. Wu, S. Shamblin, D. Varshney, E. Shalaev, *Advance Understanding of Buffer Behavior during*  
972 *Lyophilization*, Springer New York, New York, NY, 2015, pp. 25–41.
- 973 [137] R. E. Pincock, T. E. Kiovsky, Kinetics of reactions in frozen solutions, *J. Chem. Educ.* 43 (1966) 358.
- 974 [138] N. Takenaka, H. Bandow, Chemical kinetics of reactions in the unfrozen solution of ice, *J. Phys.*  
975 *Chem. A* 111 (36) (2007) 8780–8786.
- 976 [139] K. Anzo, M. Harada, T. Okada, Enhanced kinetics of pseudo first-order hydrolysis in liquid phase  
977 coexistent with ice, *J. Phys. Chem. A* 117 (2013) 10619–10625.
- 978 [140] E. Shalaev, A. Soper, J. A. Zeitler, S. Ohtake, C. J. Roberts, M. J. Pikal, K. Wu, E. Boldyreva,  
979 Freezing of aqueous solutions and chemical stability of amorphous pharmaceuticals: Water clusters  
980 hypothesis, *J. Pharm. Sci.* 108 (2019) 36 – 49.
- 981 [141] M. Davidovic, C. Mattea, J. Qvist, B. Halle, Protein cold denaturation as seen from the solvent, *J.*  
982 *Am. Chem. Soc.* 131 (2009) 1025–1036.
- 983 [142] K. A. Dill, D. O. V. Alonso, K. Hutchinson, Thermal stabilities of globular proteins, *Biochemistry* 28  
984 (1989) 5439–5449.
- 985 [143] P. L. Privalov, Cold denaturation of proteins, *Crit. Rev. Biochem. Mol. Biol.* 25 (1990) 281–305.
- 986 [144] F. Franks, Protein destailization at low temperatures, *Adv. Protein Chem.* 46 (1995) 105–139.
- 987 [145] G. Graziano, F. Catanzano, A. Riccio, G. Barone, A reassessment of the molecular origin of cold  
988 denaturation, *J. Biochem.* 122 (1997) 395–401.
- 989 [146] C. F. Lopez, R. K. Darst, P. J. Rossky, Mechanistic elements of protein cold denaturation, *J. Phys.*

- 990 Chem. B 112 (2008) 5961–5967.
- 991 [147] S. Matysiak, P. G. Debenedetti, P. J. Rossky, Role of hydrophobic hydration in protein stability: A  
992 3D water-explicit protein model exhibiting cold and heat denaturation, *J. Phys. Chem. B* 116 (2012)  
993 8095–8104.
- 994 [148] E. Reategui, A. Aksan, Effects of the low-temperature transitions of confined water on the structures  
995 of isolated and cytoplasmic proteins, *J. Phys. Chem. B* 113 (2009) 13048–13060.
- 996 [149] E. Reategui, A. Aksan, Effects of water on the structure and low/high temperature stability of  
997 confined proteins, *Phys. Chem. Chem. Phys.* 12 (2010) 10161–10172.
- 998 [150] G. B. Strambini, E. Gabellieri, Proteins in frozen solutions: Evidence of ice-induced partial unfolding,  
999 *Biophys. J.* 70 (1996) 971–976.
- 1000 [151] B. S. Chang, B. S. Kendrick, J. F. Carpenter, Surface-induced denaturation of proteins during  
1001 freezing and its inhibition by surfactants, *J. Pharm. Sci.* 85 (1996) 1325–1330.
- 1002 [152] S. Jiang, S. L. Nail, Effect of process conditions on recovery of protein activity after freezing and  
1003 freeze-drying, *Eur. J. Pharm. Biopharm.* 45 (1998) 249 – 257.
- 1004 [153] B. M. Eckhardt, J. Q. Oeswein, T. A. Bewley, Effect of freezing on aggregation of human growth  
1005 hormone, *Pharm. Res.* 8 (1991) 1360–1364.
- 1006 [154] J. M. Sarciaux, S. Mansour, M. J. Hageman, S. L. Nail, Effects of buffer composition and processing  
1007 conditions on aggregation of bovine IgG during freeze-drying, *J. Pharm. Sci.* 88 (1999) 1354–1361.
- 1008 [155] G. B. Strambini, M. Gonnelli, Protein stability in ice, *Biophys. J.* 92 (2007) 2131–2138.
- 1009 [156] J. J. Schwegman, J. F. Carpenter, S. L. Nail, Evidence of partial unfolding of proteins at the  
1010 ice/freeze-concentrate interface by infrared microscopy, *J. Pharm. Sci.* 98 (9) (2009) 3239 – 3246.
- 1011 [157] J. H. Gu, A. Beekman, T. Wu, D. M. Piedmonte, P. Baker, M. Eschenberg, M. Hale, M. Goldenberg,  
1012 Beyond glass transitions: Studying the highly viscous and elastic behavior of frozen protein formulations  
1013 using low temperature rheology and its potential implications on protein stability, *Pharm. Res.* 30 (2013)  
1014 387–401.
- 1015 [158] A. Arsiccio, R. Pisano, Clarifying the role of cryo- and lyo-protectants in the biopreservation of  
1016 proteins, *Phys. Chem. Chem. Phys.* 20 (2018) 8267–8277.
- 1017 [159] A. Arsiccio, J. McCarty, R. Pisano, J.-E. Shea, Effect of surfactants on surface-induced denaturation  
1018 of proteins: Evidence of an orientation-dependent mechanism, *J. Phys. Chem. B* 122 (2018) 11390–  
1019 11399.
- 1020 [160] A. Regand, H. Goff, Effect of biopolymers on structure and ice recrystallization in dynamically frozen

1021 ice cream model systems, *J. Dairy Sci.* 85 (2002) 2722 – 2732.

1022 [161] A. Twomey, R. Less, K. Kurata, H. Takamatsu, A. Aksan, In situ spectroscopic quantification of  
1023 protein–ice interactions, *J. Phys. Chem. B* 117 (2013) 7889–7897.

1024 [162] A. Twomey, K. Kurata, Y. Nagare, H. Takamatsu, A. Aksan, Microheterogeneity in frozen protein  
1025 solutions, *Int. J. Pharm.* 487 (2015) 91 –100.

1026 [163] S. Webb, S. Golledge, J. Cleland, J. Carpenter, T. Randolph, Surface adsorption of recombinant  
1027 human interferon- $\gamma$  in lyophilized and spray-lyophilized formulations, *J. Pharm. Sci.* 91 (2002) 1474–  
1028 1487.

1029 [164] S. Webb, J. Cleland, J. Carpenter, T. Randolph, Effects of annealing lyophilized and spray-  
1030 lyophilized formulations of recombinant human interferon- $\gamma$ , *J. Pharm. Sci.* 92 (2003) 715–729.

1031 [165] I. D. Kuntz, T. S. Brassfield, G. D. Law, G. V. Purcell, Hydration of macromolecules, *Science* 163  
1032 (1969) 1329–1331.

1033 [166] M. G. Usha, R. J. Wittebort, Orientational ordering and dynamics of the hydrate and exchangeable  
1034 hydrogen atoms in crystalline crambin, *J. Mol. Biol.* 208 (1989) 669–678.

1035 [167] A. B. Siemer, K.-Y. Huang, A. E. McDermott, Protein–ice interaction of an antifreeze protein  
1036 observed with solid-state NMR, *Proc. Natl. Acad. Sci. U.S.A.* 107 (2010) 17580–17585.

1037 [168] K. Tompa, P. Banki, M. Bokor, P. Kamasa, G. Lasanda, P. Tompa, Interfacial water at protein  
1038 surfaces: Wide-line NMR and DSC characterization of hydration in ubiquitin solutions, *Biophys. J.* 96  
1039 (2009) 2789–2798.

1040 [169] B. Zakharov, A. Fisyuk, A. Fitch, Y. Watier, A. Kostyuchenko, D. Varshney, M. Sztucki, E. Boldyreva,  
1041 E. Shalaev, Ice recrystallization in a solution of a cryoprotector and its inhibition by a protein:  
1042 Synchrotron X-ray diffraction study, *J. Pharm. Sci.* 105 (2016) 2129 – 2138.

1043 [170] B. Bhatnagar, B. A. Zakharov, A. S. Fisyuk, X. Wen, F. Z. Karim, K. Lee, Y. V. Seryotkin, M. Mogodi,  
1044 A. Fitch, E. Boldyreva, A. Kostyuchenko, E. Shalaev, Protein/ice interaction: High-resolution synchrotron  
1045 X-ray diffraction differentiates pharmaceutical proteins from lysozyme, *J. Phys. Chem. B* 123 (2019)  
1046 5690–5699.

1047 [171] C. A. Knight, A. L. DeVries, Ice growth in supercooled solutions of a biological “antifreeze”, AFGP 1–  
1048 5: an explanation in terms of adsorption rate for the concentration dependence of the freezing point,  
1049 *Phys. Chem. Chem. Phys.* 11 (2009) 5749–5761.

1050 [172] S. M. Marks, A. J. Patel, Antifreeze protein hydration waters: Unstructured unless bound to ice, *Proc.*  
1051 *Natl. Acad. Sci.* 115 (2018) 8244–8246.

- 1052 [173] P. R. Edgington, P. McCabe, C. F. Macrae, E. Pidcock, G. P. Shields, R. Taylor, M. Towler, J. Van  
1053 De Streek, Mercury: visualization and analysis of crystal structures, *J. Appl. Crystallogr.* 39 (2006) 453–  
1054 457.
- 1055 [174] C. Korber, Phenomena at the advancing ice–liquid interface: solutes, particles and biological cells,  
1056 *Q. Rev. Biophys.* 21 (1988) 229–298.
- 1057 [175] J. E. Curtis, H. Nanda, S. Khodadadi, M. Cicerone, H. J. Lee, A. McAuley, S. Krueger, Small-angle  
1058 neutron scattering study of protein crowding in liquid and solid phases: Lysozyme in aqueous solution,  
1059 frozen solution, and carbohydrate powders, *J. Phys. Chem. B* 116 (2012) 9653–9667.
- 1060 [176] N. Takenaka, A. Ueda, T. Daimon, H. Bandow, T. Dohmaru, Y. Maeda, Acceleration mechanism of  
1061 chemical reaction by freezing: The reaction of nitrous acid with dissolved oxygen, *J. Phys. Chem.* 100  
1062 (1996) 13874–13884.
- 1063 [177] N. Takenaka, A. Ueda, Y. Maeda, Acceleration of the rate of nitrite oxidation by freezing in aqueous  
1064 solution, *Nature* 358 (1992) 736–738.
- 1065 [178] S. Kunugi, N. Tanaka, Cold denaturation of proteins under high pressure, *Biochim. Biophys. Acta*  
1066 1595 (2002) 329 – 344.
- 1067 [179] K. Heremans, The phase diagram and the pressure-temperature behavior of proteins, in: R. Winter,  
1068 J. Jonas (Eds.), *High Pressure Molecular Science*, Springer Netherlands, Dordrecht, 1999, pp. 437–472.
- 1069 [180] A. Arsiccio, J. McCarty, R. Pisano, J.-E. Shea, Heightened cold-denaturation of proteins at the ice-  
1070 water interface, *J. Am. Chem. Soc.* (2020) In press, DOI: 10.1021/jacs.9b13454.
- 1071 [181] J. A. Raymond, A. L. DeVries, Adsorption inhibition as a mechanism of freezing resistance in polar  
1072 fishes, *Proc. Natl. Acad. Sci.* 74 (1977) 2589–2593.
- 1073 [182] Z. Jia, P. L. Davies, Antifreeze proteins: an unusual receptor–ligand interaction, *Trends Biochem.*  
1074 *Sci.* 27 (2002) 101–106.
- 1075 [183] L. L. C. Olijve, K. Meister, A. L. DeVries, J. G. Duman, S. Guo, H. J. Bakker, I. K. Voets, Blocking  
1076 rapid ice crystal growth through nonbasal plane adsorption of antifreeze proteins, *Proc. Natl. Acad. Sci.*  
1077 113 (14) (2016) 3740–3745.
- 1078 [184] A. L. Devries, Y. Lin, Structure of a peptide antifreeze and mechanism of adsorption to ice, *Biochim.*  
1079 *Biophys. Acta* 495 (1977) 388 – 392.
- 1080 [185] P. L. Davies, Ice-binding proteins: A remarkable diversity of structures for stopping and starting ice  
1081 growth, *Trends Biochem. Sci.* 39 (2014) 548– 555.
- 1082 [186] F. D. Sonnichsen, C. I. DeLuca, P. L. Davies, B. D. Sykes, Refined solution structure of type III

- 1083 antifreeze protein: hydrophobic groups may be involved in the energetics of the protein–ice interaction,  
1084 *Structure* 4 (1996) 1325 – 1337.
- 1085 [187] A. Haymet, L. G. Ward, M. M. Harding, C. A. Knight, Valine substituted winter flounder ‘antifreeze’:  
1086 preservation of ice growth hysteresis, *FEBS Lett.* 430 (1998) 301–306.
- 1087 [188] D. S. C. Yang, W.-C. Hon, S. Bubanko, Y. Xue, J. Seetharaman, C. L. Hew, F. Sicheri, Identification  
1088 of the ice-binding surface on a type iii antifreeze protein with a “flatness function” algorithm, *Biophys. J.*  
1089 74 (1998) 2142–2151.
- 1090 [189] K. A. Sharp, A peek at ice binding by antifreeze proteins, *Proc. Natl. Acad. Sci.* 108 (2011) 7281–  
1091 7282.
- 1092 [190] C. P. Garnham, R. L. Campbell, P. L. Davies, Anchored clathrate waters bind antifreeze proteins to  
1093 ice, *Proc. Natl. Acad. Sci.* 108 (18) (2011) 7363–7367.
- 1094 [191] K. R. Gallagher, K. A. Sharp, Analysis of thermal hysteresis protein hydration using the random  
1095 network model, *Biophys. Chem.* 105 (2) (2003) 195 – 209.
- 1096 [192] C. Yang, K. A. Sharp, The mechanism of the type III antifreeze protein action: A computational  
1097 study, *Biophys. Chem.* 109 (1) (2004) 137 – 148.
- 1098 [193] C. Yang, K. A. Sharp, Hydrophobic tendency of polar group hydration as a major force in type I  
1099 antifreeze protein recognition, *Proteins* 59 (2) (2005) 266–274.
- 1100 [194] D. R. Nutt, J. C. Smith, Dual function of the hydration layer around an antifreeze protein revealed by  
1101 atomistic molecular dynamics simulations, *J. Am. Chem. Soc.* 130 (39) (2008) 13066–13073.
- 1102 [195] K. Meister, S. Ebbinghaus, Y. Xu, J. G. Duman, A. DeVries, M. Gruebele, D. M. Leitner, M. Havenith,  
1103 Long-range protein–water dynamics in hyperactive insect antifreeze proteins, *Proc. Natl. Acad. Sci.* 110  
1104 (5) (2013) 1617–1622.
- 1105 [196] K. Meister, S. Strazdaite, A. L. DeVries, S. Lotze, L. L. C. Olijve, I. K. Voets, H. J. Bakker,  
1106 Observation of ice-like water layers at an aqueous protein surface, *Proc. Natl. Acad. Sci.* 111 (50)  
1107 (2014) 17732–17736.
- 1108 [197] H. Chao, M. E. Houston, R. S. Hodges, C. M. Kay, B. D. Sykes, M. C. Loewen, P. L. Davies, F. D.  
1109 Sonnichsen, A diminished role for hydrogen bonds in antifreeze protein binding to ice, *Biochemistry* 36  
1110 (48) (1997) 14652–14660.
- 1111 [198] J. Baardsnes, L. H. Kondejewski, R. S. Hodges, H. Chao, C. Kay, P. L. Davies, New ice-binding face  
1112 for type I antifreeze protein, *FEBS Letters* 463 (1) (1999) 87 – 91.
- 1113 [199] A. Hudait, D. R. Moberg, Y. Qiu, N. Odendahl, F. Paesani, V. Molinero, Preordering of water is not

1114 needed for ice recognition by hyperactive antifreeze proteins, Proc. Natl. Acad. Sci. 115 (33) (2018)  
1115 8266–8271.

1116 [200] S. E. Lindow, D. C. Army, C. D. Upper, Bacterial ice nucleation: A factor in frost injury to plants, Plant  
1117 Physiol. 70 (1982) 1084–1089.

1118 [201] R. L. Green, G. J. Warren, Physical and functional repetition in a bacterial ice nucleation gene,  
1119 Nature 317 (1985) 645–648.

1120 [202] C. T. Zee, C. Glynn, M. Gallagher-Jones, J. Miao, C. G. Santiago, D. Cascio, T. Gonen, M. R.  
1121 Sawaya, J. A. Rodriguez, Homochiral and racemic microed structures of a peptide repeat from the ice-  
1122 nucleation protein inaZ, IUCrJ 6 (2019) 197–205.

1123 [203] Y.-C. Liou, A. Tocilj, P. L. Davies, Z. Jia, Mimicry of ice structure by surface hydroxyls and water of a  
1124  $\beta$ -helix antifreeze protein, Nature 406 (2000) 322–324.

1125 [204] A. V. Kajava, S. E. Lindow, A model of the three-dimensional structure of ice nucleation proteins, J.  
1126 Mol. Biol. 232 (1993) 709 – 717.

1127 [205] C. P. Garnham, R. L. Campbell, V. K. Walker, P. L. Davies, Novel dimeric  $\beta$ -helical model of an ice  
1128 nucleation protein with bridged active sites, BMC Struct. Biol. 11 (2011) 36.

1129 [206] C. I. DeLuca, P. L. Davies, Q. Ye, Z. Jia, The effects of steric mutation on the structure of type iii  
1130 antifreeze protein and its interaction with ice, J. Mol. Biol. 275 (1998) 515 – 525.

1131 [207] R. Pandey, K. Usui, R. A. Livingstone, S. A. Fischer, J. Pfaendtner, E. H. G. Backus, Y. Nagata, J.  
1132 Frohlich-Nowoisky, L. Schmuser, S. Mauri, J. F. Scheel, D. A. Knopf, U. Poschl, M. Bonn, T. Weidner,  
1133 Ice-nucleating bacteria control the order and dynamics of interfacial water, Sci. Adv. 2 (2016) e1501630.

1134 [208] A. Margaritis, A. Bassi, Principles and biotechnological applications of bacterial ice nucleation, Crit.  
1135 Rev. Biotechnol. 11 (1991) 277–295.

1136 [209] W. Lindong, N. Shannon, S. Anisa, L. Shannon, T. Mehmet, Controlled ice nucleation using freeze-  
1137 dried *Pseudomonas syringae* encapsulated in alginate beads, Cryobiology 75 (2017) 1–6.

1138 [210] N. Cochet, P. Widehem, Ice crystallization by *Pseudomonas syringae*, Appl. Microbiol. Biotechnol.  
1139 54 (2000) 153–161.

1140 [211] R. Fang, K. Tanaka, V. Mudhivarthi, R. H. Bogner, M. J. Pikal, Effect of controlled ice nucleation on  
1141 stability of lactate dehydrogenase during freeze-drying, J. Pharm. Sci. 107 (2018) 824–830.

1142 [212] S. N. Singh, S. Kumar, V. Bondar, N. Wang, R. Forcino, J. Colandene, D. Nesta, Unexplored  
1143 benefits of controlled ice nucleation: Lyophilization of a highly concentrated monoclonal antibody  
1144 solution, Int. J. Pharm. 552 (2018) 171 – 179.



- 1145 [213] I. Vollrath, W. Friess, A. Freitag, A. Hawe, G. Winter, Does controlled nucleation impact the  
1146 properties and stability of lyophilized monoclonal antibody formulations?, *Eur. J. Pharm. Biopharm.* 129  
1147 (2018) 134–144.
- 1148 [214] I. Oddone, A. Arsiccio, C. Duru, K. Malik, J. Ferguson, R. Pisano, P. Matejtschuk, Vacuum induced  
1149 surface freezing for the freeze-drying of the human growth hormone: How does nucleation control affect  
1150 protein stability?, *J. Pharm. Sci.* 109 (2020) 254–263.
- 1151 [215] E. Cao, Y. Chen, Z. Cui, P. R. Foster, Effect of freezing and thawing rates on denaturation of  
1152 proteins in aqueous solutions, *Biotechnol. Bioeng.* 82 (2003) 684–690.
- 1153 [216] J. Horn, S. Jena, A. Aksan, W. Friess, Freeze/thaw of IGG solutions, *Eur. J. Pharm. Biopharm.* 134  
1154 (2019) 185 – 189.
- 1155 [217] H. J. Lee, A. McAuley, K. F. Schilke, J. McGuire, Molecular origins of surfactant-mediated  
1156 stabilization of protein drugs, *Adv. Drug. Deliv. Rev.* 63 (2011) 1160–1171.
- 1157 [218] T. W. Randolph, L. S. Jones, Surfactant-Protein Interactions, In: "Rational Design of Stable Protein  
1158 Formulations: Theory and Practice", Kluwer Academic, New York, 2002, pp. 159–175.
- 1159 [219] S. Timasheff, The control of protein stability and association by weak interactions with water: How do  
1160 solvents affect these processes?, *Ann. Rev. Biophys. Biomol. Struct.* 22 (1993) 67–97.
- 1161 [220] A. Arsiccio, R. Pisano, Stability of proteins in carbohydrates and other additives during freezing: The  
1162 human growth hormone as a case study, *J. Phys. Chem. B* 121 (2017) 8652–8660.
- 1163 [221] S. Ohtake, Y. Kita, T. Arakawa, Interactions of formulation excipients with proteins in solution and in  
1164 the dried state, *Adv. Drug Deliv. Rev.* 63 (2011) 1053–1073.
- 1165 [222] F. Franks, Long-term stabilization of biologicals, *Biotechnology* 12 (1994) 253–256.
- 1166 [223] S. J. Hagen, J. Hofrichter, W. A. Eaton, Protein reaction kinetics in a room-temperature glass,  
1167 *Science* 269 (1995) 959–962.
- 1168 [224] J. F. Carpenter, J. H. Crowe, An infrared spectroscopic study of the interactions of carbohydrates  
1169 with dried proteins, *Biochemistry* 28 (1989) 3916–3922.
- 1170 [225] J. H. Crowe, L. M. Crowe, D. Chapman, Preservation of membranes in anhydrobiotic organisms:  
1171 The role of trehalose, *Science* 223 (1984) 701–703.
- 1172 [226] D. Corradini, E. G. Strekalova, H. E. Stanley, P. Gallo, Microscopic mechanism of protein  
1173 cryopreservation in an aqueous solution with trehalose, *Sci. Rep.* 3 (2013) 1218.
- 1174 [227] A. Arsiccio, R. Pisano, Water entrapment and structure ordering as protection mechanisms for  
1175 protein structural preservation, *J. Chem. Phys.* 148 (2018) 055102.

- 1176 [228] P. S. Belton, A. M. Gil, IR and Raman spectroscopic studies of the interaction of trehalose with hen  
1177 egg white lysozyme, *Biopolymers* 34 (1994) 957–961.
- 1178 [229] X. C. Tang, M. J. Pikal, Measurement of the kinetics of protein unfolding in viscous systems and  
1179 implications for protein stability in freeze-drying, *Pharm. Res.* 22 (2005) 1176–1185.
- 1180 [230] G. Gomez, M. J. Pikal, N. Rodriguez-Hornedo, Effect of initial buffer composition on pH changes  
1181 during far-from-equilibrium freezing of sodium phosphate buffer solutions, *Pharm. Res.* 18 (2001) 90–97.
- 1182 [231] N. Murase, F. Franks, Salt precipitation during the freeze-concentration of phosphate buffer  
1183 solutions, *Biophys. Chem.* 34 (1989) 293–300.
- 1184 [232] X. C. Tang, M. J. Pikal, The effect of stabilizers and denaturants on the cold denaturation  
1185 temperatures of proteins and implications for freeze-drying, *Pharm. Res.* 22 (2005) 1167–1175.
- 1186
- 1187

1188 **LIST OF FIGURES**

1189

1190 Figure 1: Phase diagram of ice, where the hydrogen-disordered forms are shown in orange, the ordered  
1191 ones in green and the polymeric ice X in blue. The stable phases are written in red and bold, while  
1192 metastable ones are displayed in smaller size. An exclamation point is used to indicate transitions that are  
1193 not yet known. Reproduced from reference [10], with modifications.

1194

1195 Figure 2: Structures of the various polymorphs of ice (generated using the GenIce algorithm [19]). The  
1196 tetrahedral arrangement of hydrogen bonds in ice is also shown in the right panel on the last row of the  
1197 figure. In all crystal structures, this same motif is repeated, where each molecule accepts two hydrogen  
1198 bonds, and donates two to its neighbors.

1199

1200 Figure 3: Hexagonal ice Ih, where the main crystal faces (basal, primary prismatic and secondary prismatic)  
1201 are shown.

1202

1203 Figure 4: Free energy variation upon homogeneous (black) and heterogeneous (red) nucleation at different  
1204 nuclei size  $r$ . These free energy changes are the sum of a negative bulk (dotted) and a positive surface  
1205 (dashed) contributions, and reach a maximum at  $r = r_{cr}$ .

1206

1207 Figure 5: Evolution of temperature during freezing of an aqueous solution.

1208

1209 Figure 6: Sample SEM images of freeze-dried products. Unpublished data from the authors.

1210

1211 Figure 7: Possible mechanisms of ice-induced denaturation of proteins. (a) Adsorption at the ice interface  
1212 [150] (b) Partitioning of the protein in the QLL, where the concentration of stabilizer is decreased, and the  
1213 local pH is more acid than in the freeze-concentrated solution (FCS) [170] (c) Accumulation of air bubbles at  
1214 the ice surface [156, 132] (d) Pressure-induced unfolding due to mechanical stresses associated with the ice  
1215 growth [170] (e) Enhancement of cold denaturation phenomena, mediated by the liquid molecules in the QLL  
1216 [180].

1217

1218

1219 Figure 8: (a) Cartoon representation of TmAFP (PDB structure 1EZG [203]), where  $\beta$ -sheets are in yellow,  
1220 turns in cyan and coils in white. The N-terminus and C-terminus are highlighted in red and blue respectively.  
1221 (b-c) L-GSTSTA (PDB code 6M9I [202]) (b) and racemic-GSTSTA (PDB code 6M9J [202]) (c) from inaZ (INP  
1222 protein from *Pseudomonas syringae*). Glycine amino acids are shown in white, serine in yellow, threonine in  
1223 red and alanine in blue.