

The fractional porous medium equation on the hyperbolic space

Original

The fractional porous medium equation on the hyperbolic space / Berchio, E.; Bonforte, M.; Ganguly, D.; Grillo, G.. - In: CALCULUS OF VARIATIONS AND PARTIAL DIFFERENTIAL EQUATIONS. - ISSN 0944-2669. - STAMPA. - 59:5(2020). [10.1007/s00526-020-01817-2]

Availability:

This version is available at: 11583/2873254 since: 2021-03-05T12:04:44Z

Publisher:

Springer Science and Business Media Deutschland GmbH

Published

DOI:10.1007/s00526-020-01817-2

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

THE FRACTIONAL POROUS MEDIUM EQUATION ON THE HYPERBOLIC SPACE

ELVISE BERCHIO, MATTEO BONFORTE, DEBDIP GANGULY, AND GABRIELE GRILLO

ABSTRACT. We consider a nonlinear degenerate parabolic equation of porous medium type, whose diffusion is driven by the (spectral) fractional Laplacian on the hyperbolic space. We provide existence results for solutions, in an appropriate weak sense, for data belonging either to the usual L^p spaces or to larger (weighted) spaces determined either in terms of a ground state of $\Delta_{\mathbb{H}^N}$, or of the (fractional) Green's function. For such solutions, we also prove different kind of smoothing effects, in the form of quantitative $L^1 - L^\infty$ estimates. To the best of our knowledge, this seems the first time in which the fractional porous medium equation has been treated on non-compact, geometrically non-trivial examples.

CONTENTS

1. Introduction	2
2. Preliminaries and main results	3
2.1. Definition of weak dual solutions	3
2.2. Main results	5
3. Fundamental estimates for weak dual solutions (WDS)	7
4. Boundedness of WDS. Proof of the Smoothing Effects	13
4.1. Proof of Theorem 2.2	13
4.2. Proof of Theorem 2.3	16
4.3. Proof of Theorem 2.4	17
5. Semigroup theory in Banach and Hilbert spaces: existence, uniqueness, contractivity and comparison.	18
5.1. Nonlinear Semigroup in $L^1(\mathbb{H}^N)$. Mild VS Weak Dual Solutions.	19
5.2. Nonlinear Semigroup in $H^{-s}(\mathbb{H}^N)$ and Gradient Flows.	23
5.3. Proof of Theorem 2.1	25
Appendix A. Green function estimates in \mathbb{H}^N	28
References	33

2010 *Mathematics Subject Classification*. Primary: 35R01. Secondary: 35K65, 35A01, 35R11, 58J35.

Key words and phrases. Fractional porous medium equation; Hyperbolic space; a priori estimates; smoothing effects.

1. INTRODUCTION

This article is devoted to the study of *nonnegative* solutions to the Fractional Porous Medium Equation (FPME) on the hyperbolic space \mathbb{H}^N . More precisely, we consider the Cauchy problem:

$$\begin{cases} \partial_t u + (-\Delta_{\mathbb{H}^N})^s u^m = 0, & (t, x) \in (0, \infty) \times \mathbb{H}^N; \\ u(0, x) = u_0(x) \geq 0, & x \in \mathbb{H}^N, \end{cases} \quad (1.1)$$

where $(-\Delta_{\mathbb{H}^N})^s$ denotes the spectral fractional Laplacian on the hyperbolic space, $0 < s < 1$, $m > 1$, and we use the standard convention $u^m = |u|^{m-1}u$. The operator $(-\Delta_{\mathbb{H}^N})^s$ is in fact defined by functional calculus, and well-posedness of (1.1) in an appropriate sense will be one of the main issues in this paper, the other one being the validity of suitable *smoothing effects* for such evolution, this meaning quantitative bounds on the L^∞ norm of the solution at time $t > 0$ in terms of a (possibly weighted) L^p norm ($p \geq 1$) of the initial datum. In fact, we shall prove three different estimates of that type, each dealing with a larger class of initial data and different time behaviour as $t \rightarrow 0$ and $t \rightarrow \infty$ accordingly.

The study of the FPME in the Euclidean setting has been initiated in [22] for the special case $s = 1/2$, and continued in the case of general exponents in [23]. In such papers well-posedness of the evolutions for L^p data is proved even when $m < 1$ (the *fractional fast diffusion case*), and appropriate smoothing effects are proved for solutions. By a *smoothing effect* we mean a bound of the form

$$\|u(t)\|_\infty \leq C \frac{\|u_0\|_p^\alpha}{t^\beta} \quad \forall t > 0 \quad (1.2)$$

for suitable exponents α, β , and possible generalizations of (1.2) in which in the r.h.s. a *weighted L^p norm* appears. Such kind of instantaneous regularization (L^p data are smoothed out instantaneously into bounded solutions) is typical of heat-like evolution equations, see e.g. [21] for the linear case and [41, 42] for porous medium-type evolutions. A number of subsequent results concerning e.g. pointwise bounds on solutions, propagation of positivity, existence, uniqueness and properties of fundamental (or Barenblatt) solution i.e. solutions corresponding to a Dirac delta as initial datum, regularity of solutions, were proved later in [3, 13, 32, 43, 45].

The FPME has been later studied in bounded domains, with appropriate boundary conditions, mainly of homogeneous Dirichlet type. In fact, there are several different versions of what one might call a Dirichlet fractional Laplacian on domains and these different versions have been being actively investigated recently, first in [14, 15] and then e.g. in [8, 9, 12]. It is particularly important in regard to the contents of the present paper that some of these papers deal with the FPME by using properties of the *Green's function* associated to the version of the fractional Laplacian considered, starting from the very definition of solution, a strategy we shall use in this paper too. This is a particularly useful approach also in view of the technical difficulties in dealing with extension methods *à la* Caffarelli-Silvestre [18] on manifolds, as developed in [5], in [25] for the higher rank case, see also [4] for the case of nonsingular kernels.

The analysis of the heat equation in the setting of Riemannian manifolds is a widely studied topic, see e.g. [26, 27, 28] and references therein. The analysis of *nonlinear* diffusions of porous medium type on manifolds started instead just recently. The first results in this connection are to our knowledge given in [11], in which basic properties of the porous medium equation on Cartan-Hadamard manifolds, namely simply connected manifolds of nonpositive sectional curvature, are considered. A number of subsequent recent contributions deal with the porous and fast diffusion

equations (in the nonfractional case) on manifolds, mainly in the case of negative curvature, starting with [30, 44] and continuing e.g. in [29, 31, 33, 34, 35, 36].

It should be noted that the long-time behaviour of solutions even to the *linear* heat equation posed on negatively curved manifolds is completely different from the Euclidean one: in fact, one has

$$K_{\mathbb{R}^N}(t, x, x) \asymp t^{-N/2}, \quad K_{\mathbb{H}^N}(t, x, x) \asymp t^{-3/2} e^{-(N-1)^2 t} \quad \text{as } t \rightarrow +\infty, \quad (1.3)$$

where $K_{\mathbb{R}^N}$ is the heat kernel on the N -dimensional Euclidean space, and $K_{\mathbb{H}^N}$ is the heat kernel on the most important example of negatively curved, noncompact manifold, namely on the *hyperbolic space*, i.e. the simply connected manifold whose sectional curvatures are constant and equal to -1 . In the series of papers just quoted above, significant differences are shown to appear when dealing with the nonlinear setting as well: in particular, the form of the smoothing effects may be quite different.

Little is known, however, in the fractional case. Some recent results on the FPME on compact manifolds with conical singularities are given in [38], see also [39, 40] for the non-fractional case. It seems that no results are available in geometrically nontrivial, noncompact cases. Our goal here will be indeed to deal with the FPME on the hyperbolic space \mathbb{H}^N .

The first aim of the paper will be to introduce a concept of solution which will be based on the existence and properties of the *fractional Green function* $\mathbb{G}_{\mathbb{H}^N}^s$, namely the kernel of the operator $(-\Delta_{\mathbb{H}^N})^{-s}$. The latter operator and the corresponding kernel can be defined by functional calculus, see (2.1) below. Such definition is given precisely in Definition 2.1, and involves data that are integrable w.r.t. the Riemannian measure or, at least, that are integrable *with respect to suitable weights*. Such weights can be chosen to be either the *ground state eigenfunction* of $-\Delta_{\mathbb{H}^N}$, or any smooth, strictly positive function whose tail has the same decay of $\mathbb{G}_{\mathbb{H}^N}^s$ at infinity. This latter class is shown to be larger than the former, as it follows by comparing the behaviour of the ground state at infinity, see (2.4), with the one of the fractional Green function, see (3.2). Existence results for weak dual solutions are then given in Theorem 2.1, whereas the main smoothing effects are stated in the other main results of this paper, namely Theorems 2.2, 2.3, 2.4. Of course, the specific smoothing effect depends on the class of data chosen.

The paper is organized as follows: in Section 2 we introduce some of the notations, the definition of Weak Dual Solution to (1.1) and its relation with appropriate weighted spaces. We state our main existence and uniqueness result in Theorem 2.1 and then we state the smoothing estimates in Theorems 2.2, 2.3 and 2.4. In Section 3, we derive fundamental estimates for Weak Dual Solutions in Proposition 3.3 and we derive monotonicity in time estimates for solutions in weighted L^1 -spaces in Propositions 3.4, 3.5. Section 4 is devoted to the proof of the smoothing estimates stated in Section 2. Section 5 contains existence of mild solutions using Crandall-Liggett Theory or Brezis-Komura theory and we prove that mild solutions are also Weak Dual Solutions, see Theorem 5.2. Finally, we prove the existence Theorem 2.1, by approximating Weak Dual Solutions (from below) in terms of mild solutions constructed in Section 5. Appendix A contains a detailed proof of all crucial Green function estimates on the hyperbolic space which play a pivotal role in our study

2. PRELIMINARIES AND MAIN RESULTS

2.1. Definition of weak dual solutions. An important aspect is the right notion of solution to (1.1). There are many notions starting from weak solution to strong solutions. Here we shall be dealing with the notion of *weak dual solutions* introduced in [14]. This notion has its own advantage

in obtaining a priori estimates of solutions. Before formally defining weak dual solutions, we recall some crucial facts.

The fractional power of the Laplacian on \mathbb{H}^N can be written in terms of the heat semigroup as follows:

$$(-\Delta_{\mathbb{H}^N})^s u = \frac{1}{\Gamma(-s)} \int_0^\infty (e^{t\Delta_{\mathbb{H}^N}} u - u) \frac{dt}{t^{1+s}},$$

for an appropriate set of functions u and where Γ denotes the Euler Gamma function. On the other hand, we denote by $\mathbb{G}_{\mathbb{H}^N}^s(x, x_0)$ the fractional Green function on \mathbb{H}^N with fixed pole $x_0 \in \mathbb{H}^N$, and by $\mu_{\mathbb{H}^N}$ its Riemannian measure. Functional calculus allows to define the inverse of $(-\Delta_{\mathbb{H}^N})^s$, on an appropriate set of functions u , as

$$(-\Delta_{\mathbb{H}^N})^{-s} u = \int_0^{+\infty} \frac{e^{t\Delta_{\mathbb{H}^N}} u}{t^{1-s}} dt. \quad (2.1)$$

Since the kernel of $(-\Delta_{\mathbb{H}^N})^{-s}$ is $\mathbb{G}_{\mathbb{H}^N}^s$, there holds:

$$(-\Delta_{\mathbb{H}^N})^{-s} u = \int_{\mathbb{H}^N} \mathbb{G}_{\mathbb{H}^N}^s(\cdot, y) u(y) d\mu_{\mathbb{H}^N}(y).$$

In the following we will deal with solutions with initial data in $L^1(\mathbb{H}^N)$ or in suitable weighted spaces containing $L^1(\mathbb{H}^N)$. More precisely, we will deal with two kinds of weights, described below. The smoothing effects proved in the two classes will of course be different.

The first class of weights we will deal with is defined as follows:

$$\mathcal{W} = \{ \Phi = (-\Delta_{\mathbb{H}^N})^{-s} \psi \text{ for some nonnegative } \psi \in C_c^\infty(\mathbb{H}^N) \}. \quad (2.2)$$

It's worth noting that $\mathcal{W} \subset L^\infty(\mathbb{H}^N)$ and weights in \mathcal{W} behave asymptotically like the Green function at infinity. More precisely, in Lemma A.3 below, we prove that for all $\Phi \in \mathcal{W}$ there exist $R, C_1, C_2 > 0$ such that

$$C_1 r^{s-1} e^{-(N-1)r} \leq \Phi(x) \leq C_2 r^{s-1} e^{-(N-1)r} \quad \text{for } r \geq R \quad (2.3)$$

where $r := r(x_0, x)$ denotes the geodesic distance from a fixed $x_0 \in \mathbb{H}^N$.

To introduce the second weight, we recall the definition of *generalised ground state* for $(-\Delta_{\mathbb{H}^N})^s$. Let $\Lambda = \frac{(N-1)^2}{4}$ be the bottom of the spectrum of the Laplace-Beltrami operator on \mathbb{H}^N . Then the equation

$$(-\Delta_{\mathbb{H}^N})^s u = \Lambda^s u \quad \text{in } \mathbb{H}^N$$

admits a positive radial solution $\Phi_1(x) = \Phi_1(r(x_0, x))$, r denoting the geodesic distance. Φ_1 is then called a generalised ground state or the eigenfunction, and is defined up to multiplicative constants and up to the choice of the pole x_0 . It is known e.g. from [10, Section 3.3], that Φ_1 satisfies

$$\alpha(1+r)e^{-\frac{(N-1)}{2}r} \leq \Phi_1(x) \leq \beta(1+r)e^{-\frac{(N-1)}{2}r} \quad \text{for all } r > 0 \quad (2.4)$$

and for some $\alpha, \beta > 0$. Note that (2.3) and (2.4) readily imply that $\Phi \leq C\Phi_1$ on \mathbb{H}^N , for a suitable $C > 0$, so that $L_{\Phi_1}^1(\mathbb{H}^N) \subset L_\Phi^1(\mathbb{H}^N)$ for all $\Phi \in \mathcal{W}$. In particular, since $\Phi = o(\Phi_1)$ at infinity, the inclusion is strict.

We are finally ready to state

Definition 2.1. Let $\Phi \in \mathcal{W}$, where \mathcal{W} is given in (2.2), or $\Phi = \Phi_1$, a ground state of $-\Delta_{\mathbb{H}^N}$, or $\Phi \equiv 1$. We say that u is a Weak Dual Solution (WDS) to Problem (1.1) in $[0, T) \times \mathbb{H}^N$ if

- $u \in C([0, T) : L_\Phi^1(\mathbb{H}^N))$, and $u^m \in L^1((0, T) : L_{loc}^1(\mathbb{H}^N))$;

- u satisfies the following identity

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u) \partial_t \psi \, d\mu_{\mathbb{H}^N} \, dt - \int_0^T \int_{\mathbb{H}^N} u^m \psi \, d\mu_{\mathbb{H}^N} \, dt = 0 \quad (2.5)$$

for every test function $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$;

- $u(0, x) = u_0(x)$ a.e. in \mathbb{H}^N .

Remark 2.1. We note that the first term of (2.5) is finite. Indeed, using Fubini Theorem, we have

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u) \partial_t \psi \, d\mu_{\mathbb{H}^N} \, dt = \int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(\partial_t \psi) u \, d\mu_{\mathbb{H}^N} \, dt$$

for all $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$. Hence, if $u \in C([0, T] : L_\Phi^1(\mathbb{H}^N))$ with $\Phi \in \mathcal{W}$, then $\Phi = (-\Delta_{\mathbb{H}^N})^{-s} \bar{\psi} \in \mathcal{W}$ for some positive $\bar{\psi} \in C_c^\infty(\mathbb{H}^N)$ and, since $\frac{(-\Delta_{\mathbb{H}^N})^{-s}(\partial_t \bar{\psi})}{(-\Delta_{\mathbb{H}^N})^{-s} \bar{\psi}} \in L^\infty(\mathbb{H}^N)$ (see Lemma A.3 below) we conclude that the above integrals are finite.

Since $L^1(\mathbb{H}^N) \subset L_{\Phi_1}^1(\mathbb{H}^N) \subset L_\Phi^1(\mathbb{H}^N)$ for all $\Phi \in \mathcal{W}$, the same conclusion holds if either $\Phi = \Phi_1$ or $\Phi \equiv 1$.

This class of solutions is quite big, indeed it contains mild (semigroup) L^1 -solutions, see Section 5. On the other hand, in Remark 5.1 we see that WDS are contained in the class of very weak (distributional) solutions.

2.2. Main results. The construction of a WDS will be done by means of approximating with semigroup (mild) solutions constructed either in $L^1(\mathbb{H}^N)$ or in $H^{-s}(\mathbb{H}^N)$, by means of appropriate nonlinear semigroup techniques both in Banach and Hilbert spaces. In Section 5 we show that mild solutions are weak dual solutions.

Theorem 2.1 (Existence and uniqueness of Nonnegative Minimal Weak Dual Solutions in L_Φ^1). *Let $\Phi \in \mathcal{W}$, where \mathcal{W} is given in (2.2), or $\Phi = \Phi_1$, a ground state of $-\Delta_{\mathbb{H}^N}$, or $\Phi \equiv 1$. For every nonnegative $u_0 \in L_\Phi^1(\Omega)$ there exists a weak dual solution to Problem (1.1), continuous in the weighted space $u \in C([0, T] : L_\Phi^1(\Omega))$ for all $T > 0$. Such a solution is obtained as the unique monotone limit of nonnegative L^1 -semigroup (mild) solutions which exist and are unique. We call such solution the minimal WDS. In this class of solutions the standard comparison principle holds.*

Remark 2.2. *i) Existence and uniqueness of signed (semigroup) solutions.* When dealing with signed solutions, existence and uniqueness of mild (hence WDS) solutions follows from the nowadays standard theory of nonlinear contractive semigroups on Banach and Hilbert spaces. In the first case, the $L^1(\mathbb{H}^N)$ -theory of m-Accretive operators, developed by Benilan, Crandall, Pazy and Pierre provides existence, uniqueness and comparison, see [6, 7, 20] and Theorem 5.1. In the Hilbertian setting, the (nowadays called Gradient Flow) theory by Brezis and Komura, gives the same results in $H^{-s}(\mathbb{H}^N)$, see [16, 17, 37] and Theorem 5.4. More details will be provided in Section 5.2.

ii) Boundedness of solutions. On one hand, showing that mild solutions are bounded is in general a difficult task, unless the initial datum is already bounded, since the a-priori regularity is too low. On the other hand, WDS turn out to be bounded, as we will see below. We show in Section 5 that mild solutions are indeed WDS, hence bounded. Informally speaking, using WDS means to find appropriate weak solutions to the dual equation $(-\Delta_{\mathbb{H}^N})^{-s} u_t = -u^m$: this is useful since it allows to extend the approach via Green functions estimates of [15, 14] to the present setting, and to prove quantitative boundedness of WDS, avoiding DeGiorgi-Nash-Moser iterations, which rely on integration by parts, not available in this non-local framework.

It is convenient to define the exponent

$$\vartheta_1 := \frac{1}{2s + N(m-1)}$$

which appears in our $L^1 - L^\infty$ smoothing estimates.

Theorem 2.2. *Let u be a WDS corresponding to the nonnegative initial datum $u_0 \in L^1(\mathbb{H}^N)$. There exists $\bar{\kappa}_1 = \bar{\kappa}_1(s, m, N) > 0$ such that*

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \bar{\kappa}_1 \frac{\|u(t)\|_{L^1(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}} \leq \bar{\kappa}_1 \frac{\|u_0\|_{L^1(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}} \quad \text{for all } t > 0. \quad (2.6)$$

Moreover, for $t \geq e^{2(N-1)(m-1)} \|u_0\|_{L^1(\mathbb{H}^N)}^{-(m-1)}$ there exists $\bar{\kappa}_2 = \bar{\kappa}_2(s, m, N) > 0$ such that

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \frac{\bar{\kappa}_2}{t^{\frac{1}{m-1}}} \left[\log \left(t \|u_0\|_{L^1(\mathbb{H}^N)}^{m-1} \right) \right]^{\frac{s}{m-1}}. \quad (2.7)$$

A second smoothing effect can be shown in the larger class, namely for data belonging to $L^1_{\Phi_1}(\mathbb{H}^N)$.

Theorem 2.3. *Let Φ_1 be a ground state of $-\Delta_{\mathbb{H}^N}$ and let u be a WDS corresponding to the nonnegative initial datum $u_0 \in L^1_{\Phi_1}(\mathbb{H}^N)$. There exists $\bar{\kappa}_3 = \bar{\kappa}_3(s, m, N) > 0$ such that*

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \bar{\kappa}_3 \frac{\|u(t)\|_{L^1_{\Phi_1}(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}} \leq \bar{\kappa}_3 \frac{\|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}} \quad \text{for all } t > 0. \quad (2.8)$$

Moreover, for $t \geq e^{(m-1)(N-1)} \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}^{-(m-1)}$ there exists $\bar{\kappa}_4 = \bar{\kappa}_4(s, m, N) > 0$ such that

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \frac{\bar{\kappa}_4}{t^{\frac{1}{m-1}}} \left[\log(t \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}^{m-1}) \right]^{\frac{s}{m-1}}. \quad (2.9)$$

Remark 2.3. It should be commented that, in the above Theorems, the bounds (2.6), (2.8) hold for all $t > 0$ but are mainly significant for t small, as for long time faster decay estimates are provided by (2.7), (2.9). This is reminiscent of the situation valid in Euclidean *bounded* domains, in which short and long time asymptotics of solutions to the porous medium equation (or even to the heat equation), fractional or not, are different, mainly due to the validity of the Poincaré inequality, a fact that holds on \mathbb{H}^N as well. Note that (2.6) is identical to the corresponding Euclidean bound and is consistent with the Euclidean scaling. The bound (2.8) has the same time dependence but, being written in terms of a weighted norm, has no Euclidean analogue. That for large time a faster time decay overtakes the one valid in the Euclidean case might also informally be seen as an effect of negative curvature, that somehow increases the speed of propagation and produces a better decay for large time (recall the bounds (1.3) in the linear case).

We conclude by enlarging further the class of allowed initial data, i.e. by allowing them to be integrable w.r.t. weight in \mathcal{W} , namely having the tail of the fractional Green function.

Theorem 2.4. *Let u be a WDS corresponding to the nonnegative initial datum $u_0 \in L^1_{\Phi}(\mathbb{H}^N)$, with $\Phi \in \mathcal{W}$ defined in (2.2). There exists $\bar{\kappa}_5 = \bar{\kappa}_5(s, m, N, \Phi) > 0$ such that*

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \frac{\bar{\kappa}_5}{t^{1/m}} \|u_0\|_{L^1_{\Phi}(\mathbb{H}^N)}^{1/m} \quad \text{for all } t > 0.$$

Remark 2.4. *About the space $L^1_{\Phi}(\mathbb{H}^N)$ versus $H^{-s}(\mathbb{H}^N)$.* The above smoothing effect shows that nonnegative WDS corresponding to data in $L^1_{\Phi}(\mathbb{H}^N)$ are bounded. As we shall see in Remark 5.2, the gradient flow solutions with data in $H^{-s}(\mathbb{H}^N)$, which exist and are unique, are also WDS, hence the above Theorem applies also to nonnegative gradient flow solutions and gives for all $t > 0$:

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \frac{\bar{\kappa}_5}{t^{1/m}} \|u_0\|_{L^1_{\Phi}(\mathbb{H}^N)}^{1/m} \leq \frac{\bar{\kappa}_5 c_\psi^{1/m}}{t^{1/m}} \|u_0\|_{H^{-s}(\mathbb{H}^N)}^{1/m},$$

where we have used inequality (5.12), i.e. $\|u_0\|_{L^1_{\Phi}(\mathbb{H}^N)} \leq c_\psi \|u_0\|_{H^{-s}(\mathbb{H}^N)}$ with $c_\psi = \|\psi\|_{L^1_{\Phi}} < +\infty$. The latter inequality also shows that for nonnegative functions, H^s_+ is contained in $L^1_{\Phi,+}(\mathbb{H}^N)$.

3. FUNDAMENTAL ESTIMATES FOR WEAK DUAL SOLUTIONS (WDS)

In this section we derive a priori estimates of WDS in the spirit of [15, Sections 5 and 6] and [14]. To this end, we will use repeatedly the next lemma.

Lemma 3.1. *There exists $\kappa_1 = \kappa_1(N, s) > 0$ such that for a.e. $x, y \in \mathbb{H}^N$ we have*

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \kappa_1 \frac{1}{r(x, y)^{N-2s}} \quad (3.1)$$

where $r(\cdot, \cdot)$ is the geodesic distance of \mathbb{H}^N . When $r(x, y) \geq 1$ (eventually by taking a bigger κ_1) we also have

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \kappa_1 \frac{e^{-(N-1)r(x,y)}}{r(x, y)^{1-s}}. \quad (3.2)$$

As a consequence, for all $R \geq 0$ and $y \in \mathbb{H}^N$ we have

$$\int_{B_R(y)} \mathbb{G}_{\mathbb{H}^N}^s(x, y) d\mu_{\mathbb{H}^N}(x) \leq \frac{\kappa_1}{2s} \cosh(R)^{N-1} R^{2s}, \quad (3.3)$$

and for all $R \geq 2$ and $y \in \mathbb{H}^N$ we have

$$\int_{B_R(y)} \mathbb{G}_{\mathbb{H}^N}^s(x, y) d\mu_{\mathbb{H}^N}(x) \leq \tilde{\kappa}_1 R^s, \quad (3.4)$$

for some $\tilde{\kappa}_1 = \tilde{\kappa}_1(N, s, \kappa_1) > 0$.

Proof. In order not to break the flow of the section we will postpone the (technical) proofs of (3.1) and (3.2) to Appendix A, see Corollary A.2. As concerns the proof of (3.3) it follows from (3.1) by passing to radial coordinates and estimating the volume element by means of the elementary inequality: $\sinh r \leq r \cosh R$ for all $0 < r < R$. Indeed

$$\begin{aligned} \int_{B_R(y)} \mathbb{G}_{\mathbb{H}^N}^s(x, y) d\mu_{\mathbb{H}^N}(x) &\leq \kappa_1 \int_0^R \frac{(\sinh r)^{N-1}}{r^{N-2s}} dr = \kappa_1 \int_0^R \frac{r^{N-1}}{r^{N-2s}} \left(\frac{\sinh r}{r}\right)^{N-1} dr \\ &\leq \frac{\kappa_1}{2s} (\cosh R)^{N-1} R^{2s}. \end{aligned}$$

As for (3.4), for $R \geq 2$ we estimate using the fact that $e^{-(N-1)r} (\sinh r)^{N-1} \leq 1$,

$$\int_{B_R(y)} \mathbb{G}_{\mathbb{H}^N}^s(x, y) d\mu_{\mathbb{H}^N}(x) d\mu_{\mathbb{H}^N}(x) \leq \frac{\kappa_1}{2s} \cosh(2)^{N-1} 2^{2s} + \kappa_1 \int_2^R \frac{e^{-(N-1)r}}{r^{1-s}} (\sinh r)^{N-1} dr$$

$$\leq \frac{\kappa_1}{s} [\cosh(2)^{N-1} 2^{2s-1} - 2^s + R^s] \leq \tilde{\kappa}_1 R^s$$

for some $\tilde{\kappa}_1 = \tilde{\kappa}_1(N, s, \kappa_1) > 0$. \square

We are now ready to state and prove what we call the fundamental pointwise formulae, that in the nonlinear case under study, play the role of the representation formula in the linear case.

Following the strategy of [14], it is convenient at this point to introduce a special class of WDS, useful to ensure that all the integral quantities in the proofs are finite. Indeed, in most of the proofs we will work with bounded integral solutions and then extend the result to general data by a standard limiting process. To this end, we shall proceed initially by considering WDS having initial data in $L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$.

Lemma 3.2 (Comparison and time monotonicity for WDS). *Let u, v be WDS corresponding to $u_0, v_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Then comparison holds, more precisely the T -contraction property (5.10) holds, namely*

$$\left\| (u(t) - v(t))_+ \right\|_{H^{-s}(\mathbb{H}^N)} \leq \left\| (u_0 - v_0)_+ \right\|_{H^{-s}(\mathbb{H}^N)} \quad \text{for all } t \geq 0. \quad (3.5)$$

Let u be a nonnegative WDS corresponding to $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Then u enjoys the following time monotonicity property:

$$\text{the map } t \mapsto t^{\frac{1}{m-1}} u(t, x) \quad \text{is nondecreasing in } t > 0 \text{ for a.e. } x \in \mathbb{H}^N, \quad (3.6)$$

and the following L^p -stability property

$$\|u(t)\|_{L^p(\mathbb{H}^N)} \leq \|u_0\|_{L^p(\mathbb{H}^N)} \quad \text{for all } t \geq 0 \text{ and all } 1 \leq p \leq \infty. \quad (3.7)$$

Proof. This proof follows by the T -contraction in H^{-s} -spaces, that we adapt from an original proof of Brezis, we just emphasize the main key points, the rest being completely analogous to the H^{-s} -case, see more details in Section 5.2. By the Poincaré inequality $\lambda_1 \|f\|_{H^{-s}(\mathbb{H}^N)} \leq \|f\|_{L^2(\mathbb{H}^N)}$. Then, since $u_0, v_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, we have that $(u_0 - v_0)_+ \in H^{-s}(\mathbb{H}^N)$ and the following holds

$$\begin{aligned} & \frac{d}{dt} \frac{1}{2} \left\| (u(t) - v(t))_+ \right\|_{H^{-s}(\mathbb{H}^N)}^2 \\ &= \int_{\mathbb{H}^N} \text{sign}^+(u(t) - v(t)) (\partial_t u(t) - \partial_t v(t)) (-\Delta_{\mathbb{H}^N})^{-s} (u(t) - v(t))_+ d\mu_{\mathbb{H}^N} \\ &= - \int_{\mathbb{H}^N} \text{sign}^+(u(t) - v(t)) [(-\Delta_{\mathbb{H}^N})^s (u^m - v^m)] (-\Delta_{\mathbb{H}^N})^{-s} (u(t) - v(t))_+ d\mu_{\mathbb{H}^N} \\ &= - \int_{\mathbb{H}^N} \text{sign}^+(u^m(t) - v^m(t)) [(-\Delta_{\mathbb{H}^N})^s (u^m - v^m)] (-\Delta_{\mathbb{H}^N})^{-s} (u(t) - v(t))_+ d\mu_{\mathbb{H}^N} \\ &\leq - \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^s (u^m - v^m)_+ (-\Delta_{\mathbb{H}^N})^{-s} (u(t) - v(t))_+ d\mu_{\mathbb{H}^N} \\ &\leq - \int_{\mathbb{H}^N} (u^m - v^m)_+ (u(t) - v(t))_+ d\mu_{\mathbb{H}^N} \leq 0 \end{aligned}$$

which clearly implies (3.5). The inequality of line 4 follows by Kato's inequality $\text{sign}^+(f) (-\Delta_{\mathbb{H}^N})^s f \leq (-\Delta_{\mathbb{H}^N})^s (f)_+$, which holds in the sense of distributions.

The above computations can be rigorously justified by the results of Theorem 5.4, in which strong H^{-s} -solutions are constructed. When u_0 is sufficiently integrable (and $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$) is

more than enough in view of the above discussion) then the strong H^{-s} solutions are the same as Weak Dual Solutions in the sense of Definition 2.1, see Remark 5.2.

Time Monotonicity Estimates. We show here for the reader's convenience a proof based on scaling and comparison that we have learned from Vázquez [41]. Consider the rescaled solution $u_\lambda(t, x) = \lambda^{\frac{1}{m-1}} u(\lambda t, x)$, then $u_\lambda(0, \cdot) = \lambda^{\frac{1}{m-1}} u_0$. Then, letting $\lambda = (t+h)/t \geq 1$, we obtain for all $h \geq 0$

$$0 \leq u_\lambda(t, x) - u(t, x) = \lambda^{\frac{1}{m-1}} u(\lambda t, x) - u(t, x) = \left(\frac{t+h}{t} \right)^{\frac{1}{m-1}} u(t+h, x) - u(t, x)$$

which is equivalent to the desired monotonicity of the map $t \mapsto t^{\frac{1}{m-1}} u(t, x)$. The first inequality in the above formula, is true for all $t \geq 0$ and a.e. $x \in \mathbb{H}^N$ by comparison (which follows by the T-contraction inequality (3.5)), since $\lambda \geq 1$ implies that $u_\lambda(0, \cdot) = \lambda^{\frac{1}{m-1}} u_0 \geq u_0$.

L^p stability of nonnegative WDS with $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. A sketch of the proof of inequality (3.7) is given in Remark 5.1. \square

We are now ready to prove our main result of this section:

Proposition 3.3. *Let $u \geq 0$ be a WDS to problem (1.1) with $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Then,*

$$\int_{\mathbb{H}^N} u(t, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} u_0(x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \text{ for all } t > 0 \quad (3.8)$$

and

$$\begin{aligned} \left(\frac{t_0}{t_1} \right)^{\frac{m}{m-1}} (t_1 - t_0) u^m(t_0, x_0) &\leq \int_{\mathbb{H}^N} [u(t_0, x) - u(t_1, x)] \mathbb{G}_{\mathbb{H}^N}^s(x_0, x) d\mu_{\mathbb{H}^N}(x) \\ &\leq (m-1) \frac{t_0^{\frac{m}{m-1}}}{t_1^{\frac{m}{m-1}}} u^m(t, x_0), \end{aligned} \quad (3.9)$$

for a. e. $0 < t_0 \leq t_1 \leq t$ and a. e. $x_0 \in \mathbb{H}^N$.

Proof. We give the proof for $\Phi \in \mathcal{W}$. The case $\Phi = \Phi_1$ follows by noting that $L_{\Phi_1}^1(\mathbb{H}^N) \subset L_\Phi^1(\mathbb{H}^N)$ for all $\Phi \in \mathcal{W}$. The statement is proved in four steps.

Step 1 : For all non-negative ψ such that $\psi \in L_c^\infty(\mathbb{H}^N)$, there holds

$$\begin{aligned} &\int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi(x)) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N} u(t_1, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi(x)) d\mu_{\mathbb{H}^N}(x) \\ &= \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi(x) d\mu_{\mathbb{H}^N}(x) d\tau. \end{aligned} \quad (3.10)$$

The proof follows by adapting the ideas of [14, Proposition 4.2-Step 1]. We use the Definition 2.1 of weak dual solution, with test function $\psi(t, x) = \psi_1(t) \psi_2(x)$, where $\psi_1(t) \in C_c^1(0, +\infty)$ and $\psi_2 \in L_c^\infty(\mathbb{H}^N)$. Namely, we have that $u \in C([0, T] : L_\Phi^1(\mathbb{H}^N))$, $u^m \in L^1((0, T) : L_{loc}^1(\mathbb{H}^N))$, and the following identity holds

$$\begin{aligned} &\int_0^\infty \psi_1'(\tau) \int_{\mathbb{H}^N} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \\ &= \int_0^\infty \psi_1(\tau) \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau. \end{aligned} \quad (3.11)$$

Now, to prove (3.10), we exploit the following approximate procedure. For $0 \leq t_0 \leq t_1 < +\infty$ we consider the characteristic function $\chi_{[t_0, t_1]}(\tau)$. By standard approximation arguments, there exists $\psi_{1,n} \in C_c^1(0, +\infty)$ with $\text{supp}(\psi_{1,n}) \subset [t_0 - \frac{1}{n}, t_1 + \frac{1}{n}]$, $\chi_{[t_0, t_1]} \leq \psi_{1,n} \leq 1$ and such that $\psi_{1,n} \rightarrow \chi_{[t_0, t_1]}(\tau)$ a.e. in $(0, +\infty)$. Since, by Remark 2.1, $\tau \mapsto \int_{\mathbb{H}^N} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_2(x) d\mu_{\mathbb{H}^N}(x) \in C^0([0, T])$, as $n \rightarrow \infty$, it follows that

$$\begin{aligned} & \int_0^\infty \psi'_{1,n}(\tau) \int_{\mathbb{H}^N} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \\ & \xrightarrow{n \rightarrow \infty} \int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_2(x) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N} u(t_1, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_2(x) d\mu_{\mathbb{H}^N}(x). \end{aligned}$$

On the other hand, since

$$\int_0^\infty \psi_{1,n}(\tau) \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \leq \int_0^{t_1+1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau < +\infty$$

by Lebesgue dominated Theorem, as $n \rightarrow \infty$, we get:

$$\int_0^\infty \psi_{1,n}(\tau) \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \xrightarrow{n \rightarrow \infty} \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau.$$

Hence, (3.10) follows by writing (3.11) with $\psi_1 = \psi_{1,n}$ and passing to the limit.

Step 2 : From (3.10) we derive estimate (3.8). Fix $x_0 \in \mathbb{H}^N$ and consider the sequence of non-negative test functions $\psi_n^{(x_0)}(x) := \frac{1}{|B_{1/n}(x_0)|} \chi_{B_{1/n}(x_0)}(x)$ with $n > 1$. Clearly, $\psi_n^{(x_0)} \in L_c^\infty(\mathbb{H}^N)$. Furthermore, $\psi_n^{(x_0)} \xrightarrow{n \rightarrow \infty} \delta_{x_0}$ in the sense of Radon measure.

A direct application of Lebesgue differentiation theorem to the function $y \mapsto \mathbb{G}_{\mathbb{H}^N}^s(y, x)$ (without loss of generality we may assume that x_0 is a Lebesgue point) implies

$$(-\Delta_{\mathbb{H}^N})^{-s}(\psi_n^{(x_0)})(x) = \frac{1}{|B_{1/n}(x_0)|} \int_{B_{1/n}(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(y, x) d\mu_{\mathbb{H}^N}(y) \xrightarrow{n \rightarrow \infty} \mathbb{G}_{\mathbb{H}^N}^s(x, x_0)$$

as $n \rightarrow \infty$ and for a.e. $x \in \mathbb{H}^N$. Next we write

$$\begin{aligned} & \left| \int_{\mathbb{H}^N} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right| \tag{3.12} \\ & \leq \left| \int_{B_R(x_0)} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) - \int_{B_R(x_0)} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right| \\ & + \left| \int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right|. \end{aligned}$$

By Lemma 3.1, $\mathbb{G}_{\mathbb{H}^N}^s(x_0, \cdot) \in L_{loc}^q(\mathbb{H}^N)$ with $0 < q < \frac{N}{N-2s}$ and we readily derive that

$$(-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) \xrightarrow{n \rightarrow \infty} \mathbb{G}_{\mathbb{H}^N}^s(x_0, x) \quad \text{in } L_{loc}^q(\mathbb{H}^N)$$

for all $0 < q < \frac{N}{N-2s}$. Then, the first expression on the r.h.s. of (3.12) can be estimated as follows

$$\begin{aligned} & \left| \int_{B_R(x_0)} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) - \int_{B_R(x_0)} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right| \\ & \leq \|u(\tau)\|_{L^p(\mathbb{H}^N)} \|(-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)} - \mathbb{G}_{\mathbb{H}^N}^s(\cdot, x_0)\|_{L^p(B_R(x_0))} \xrightarrow{n \rightarrow \infty} 0. \end{aligned}$$

Recall that by assumption $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, hence $u(t) \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ for all $t > 0$. This is true since WDS are L^p -stable, see Lemma 3.2 and also Remark 5.1.

Now we tackle the second term on the r.h.s. of (3.12) and we show that

$$\left| \int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right| \xrightarrow{n \rightarrow \infty} 0.$$

To this aim, we estimate $(-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x)$ for $r(x_0, x) \geq R$. Without loss of generality, we may assume $R \geq 2$. By repeating the proof of Lemma A.3, it is readily deduced that there exists a positive constant \bar{C} (not depending on n) such that

$$(-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) \leq \bar{C} \mathbb{G}_{\mathbb{H}^N}^s(x_0, x) \quad \text{for all } r(x_0, x) \geq 2.$$

Therefore, invoking again Lemma A.3 and using the fact $u(\tau, x) \in L^1_{\Phi}(\mathbb{H}^N)$ for each fixed τ , we deduce that

$$u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) \leq C u(\tau, x) \Phi(x) \in L^1(\mathbb{H}^N \setminus B_R(x_0))$$

for a suitable $C > 0$. Hence, by Lebesgue Theorem, it follows that

$$\int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) (-\Delta_{\mathbb{H}^N})^{-s} \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) \xrightarrow{n \rightarrow \infty} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(\tau, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x),$$

Writing (3.10) with $\psi = \psi_n^{(x_0)}$ and recalling that the right-hand side of (3.10) is non-negative, we have established (3.8).

Step 3 : We make use of the time monotonicity property (3.6) to prove

$$\begin{aligned} \left(\frac{t_0}{t_1} \right)^{\frac{m}{m-1}} (t_1 - t_0) \int_{\mathbb{H}^N} u^m(t_0, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) &\leq \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \quad (3.13) \\ &\leq \frac{m-1}{t_0^{\frac{1}{m-1}}} t^{\frac{m}{m-1}} \int_{\mathbb{H}^N} u^m(t, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x), \end{aligned}$$

for a.e. $0 \leq t_0 \leq t_1 \leq t$ and all $\psi_2 \in L_c^\infty(\mathbb{H}^N)$.

Now let us start by proving the upper estimates in (3.13). To this aim, we consider the sequence $\psi_{1,n} \in C_c^\infty(0, +\infty)$ defined in Step 1. Let us choose n so large that $0 \leq t_0 - \frac{1}{n} \leq t_1 + \frac{1}{n} \leq t$, for all nonnegative $\psi_2 \in L_c^\infty(\mathbb{H}^N)$ we have

$$\begin{aligned} &\int_0^\infty \psi_{1,n}(\tau) \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \\ &\leq \int_{t_0 - \frac{1}{n}}^{t_1 + \frac{1}{n}} \left(\frac{t}{\tau} \right)^{\frac{m}{m-1}} \int_{\mathbb{H}^N} u^m(t, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \\ &\leq \left(\int_{t_0 - \frac{1}{n}}^{t_1 + \frac{1}{n}} \left(\frac{t}{\tau} \right)^{\frac{m}{m-1}} d\tau \right) \left(\int_{\mathbb{H}^N} u^m(t, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) \right) \\ &\leq (m-1) \frac{1}{(t_0 - \frac{1}{n})^{\frac{1}{m-1}}} t^{\frac{m}{m-1}} \int_{\mathbb{H}^N} u^m(t, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x). \end{aligned}$$

Now, letting $n \rightarrow \infty$, we get

$$\int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x) d\tau \leq \frac{(m-1)}{t_0^{\frac{1}{m-1}}} t^{\frac{m}{m-1}} \int_{\mathbb{H}^N} u^m(t, x) \psi_2(x) d\mu_{\mathbb{H}^N}(x),$$

for all $t_0 \leq t_1 \leq t$. Hence, we obtain the upper bound estimates of (3.13). Following a similar approach we obtain the left-hand side of (3.13), see [14, Proposition 4.2-Step 3] for more details.

Step 4 : The final conclusion, i.e. (3.9), can be obtained by an approximation procedure again. Indeed, by combining and (3.10) and (3.13), we know

$$\begin{aligned} & \left(\frac{t_0}{t_1}\right)^{\frac{m}{m-1}} (t_1 - t_0) \int_{\mathbb{H}^N} u^m(t_0, x) \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) \\ & \int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi_n^{(x_0)}(x)) d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N} u(t_1, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi_n^{(x_0)}(x)) d\mu_{\mathbb{H}^N}(x) \\ & \leq \frac{m-1}{t_0^{\frac{1}{m-1}}} t^{\frac{m}{m-1}} \int_{\mathbb{H}^N} u^m(t, x) \psi_n^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) \end{aligned} \quad (3.14)$$

where the sequence $\psi_n^{(x_0)}$ is as defined in Step 2. So, as before, as $n \rightarrow +\infty$, we have

$$(-\Delta_{\mathbb{H}^N})^{-s}(\psi_n^{(x_0)})(x) = \frac{1}{|B_{1/n}(x_0)|} \int_{B_{1/n}(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(y, x) d\mu_{\mathbb{H}^N}(x)(y) \longrightarrow \mathbb{G}_{\mathbb{H}^N}^s(x, x_0),$$

provided x_0 is the Lebesgue point of the function $x \mapsto \mathbb{G}_{\mathbb{H}^N}^s(y, x)$. Furthermore, for all $\tau \geq 0$, as $n \rightarrow +\infty$, we have

$$\int_{\mathbb{H}^N} u^m(\tau, x) \psi_{2,n}^{(x_0)}(x) d\mu_{\mathbb{H}^N}(x) = |B_{\frac{1}{n}}(x_0)|^{-1} \int_{B_{\frac{1}{n}}(x_0)} u^m(\tau, x) d\mu_{\mathbb{H}^N}(x) \longrightarrow u^m(\tau, x_0),$$

provided x_0 is the Lebesgue point of the function $x \mapsto u^m(\tau, x)$. With no loss of generality we may choose x_0 belonging to the Lebesgue point set for both functions. With this information in hand, one can let $n \rightarrow \infty$ in (3.14) to obtain (3.9). This completes the proof of Proposition 3.3. \square

From step 1 in the proof of the above proposition it readily follows the monotonicity result below which is of fundamental importance to get existence of solutions in $L^1_{\Phi}(\mathbb{H}^N)$.

Proposition 3.4. *Let $u \geq 0$ be a WDS to problem (1.1) with $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Then for all $\psi \in C_c^\infty(\mathbb{H}^N)$ with $\psi \geq 0$ there holds*

$$\int_{\mathbb{H}^N} u(t_1, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi)(x) d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s}(\psi)(x) d\mu_{\mathbb{H}^N}(x)$$

for all $t_1 > t_0 \geq 0$. As a consequence, if $\Phi \in \mathcal{W}$, the class \mathcal{W} being as in (2.2), the integral $\int_{\mathbb{H}^N} u(t, x) \Phi(x) d\mu_{\mathbb{H}^N}(x)$ is monotonically non-increasing in time, i.e.

$$\int_{\mathbb{H}^N} u(t_1, x) \Phi(x) d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} u(t_0, x) \Phi(x) d\mu_{\mathbb{H}^N}(x)$$

for all $t_1 > t_0 \geq 0$.

Next, we get the monotonicity for solutions in $L^1_{\Phi_1}(\mathbb{H}^N)$, Φ_1 being a ground state of $-\Delta_{\mathbb{H}^N}$.

Proposition 3.5. *Let $u \geq 0$ be a WDS to problem (1.1) with $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Let Φ_1 be a ground state of $-\Delta_{\mathbb{H}^N}$. Then the integral $\int_{\mathbb{H}^N} u(t, x) \Phi_1(x) d\mu_{\mathbb{H}^N}(x)$ is monotonically non-increasing in time, i.e.*

$$\int_{\mathbb{H}^N} u(t_1, x) \Phi_1(x) d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} u(t_0, x) \Phi_1(x) d\mu_{\mathbb{H}^N}(x)$$

for all $t_1 > t_0 \geq 0$.

Proof. The proof follows from Step 1 in the proof of Proposition 3.3. To this aim, let $\psi_n \in L_c^\infty(\mathbb{H}^N)$ be a sequence of nonnegative functions such that $\psi_n \rightarrow \Phi_1$ a.e. in \mathbb{H}^N and such that $\psi_n \leq \Phi_1$, we write (3.10) with $\psi = \psi_n$:

$$\begin{aligned} & \int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s} (\psi_n(x)) \, d\mu_{\mathbb{H}^N}(x) - \int_{\mathbb{H}^N} u(t_1, x) (-\Delta_{\mathbb{H}^N})^{-s} (\psi(x)) \, d\mu_{\mathbb{H}^N}(x) \\ &= \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_n(x) \, d\mu_{\mathbb{H}^N}(x) \, d\tau. \end{aligned}$$

Since

$$\int_{\mathbb{H}^N} u(t_0, x) (-\Delta_{\mathbb{H}^N})^{-s} (\psi_n(x)) \, d\mu_{\mathbb{H}^N}(x) \leq \Lambda^{-s} \int_{\mathbb{H}^N} u(t_0, x) \Phi_1(x) \, d\mu_{\mathbb{H}^N}(x) < +\infty$$

and

$$\int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \psi_n(x) \, d\mu_{\mathbb{H}^N}(x) \, d\tau \leq \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \Phi_1(x) \, d\mu_{\mathbb{H}^N}(x) \, d\tau < +\infty,$$

we may pass to the limit and get

$$\begin{aligned} & \Lambda^{-s} \int_{\mathbb{H}^N} u(t_0, x) \Phi_1(x) \, d\mu_{\mathbb{H}^N}(x) - \Lambda^{-s} \int_{\mathbb{H}^N} u(t_1, x) \Phi_1(x) \, d\mu_{\mathbb{H}^N}(x) \\ &= \int_{t_0}^{t_1} \int_{\mathbb{H}^N} u^m(\tau, x) \Phi_1(x) \, d\mu_{\mathbb{H}^N}(x) \, d\tau \geq 0. \end{aligned}$$

This completes the proof. \square

4. BOUNDEDNESS OF WDS. PROOF OF THE SMOOTHING EFFECTS

In this section we prove the three different smoothing effects for Weak Dual Solutions, namely the results of Theorems 2.2, 2.3 and 2.4.

4.1. Proof of Theorem 2.2. Without loss of generality, we can assume $0 \leq u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, we will explain how to remove these apparent restrictions at the end of the proof. First we recall that estimate (3.9) with $t_1 = 2t_0$, immediately implies

$$\begin{aligned} u^m(t_0, x_0) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) \, d\mu_{\mathbb{H}^N}(x) \\ &= \underbrace{\frac{2^{\frac{m}{m-1}}}{t_0} \int_{B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) \, d\mu_{\mathbb{H}^N}(x)}_{(I)} + \underbrace{\frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) \, d\mu_{\mathbb{H}^N}(x)}_{(II)} \end{aligned} \quad (4.1)$$

for some $R > 0$.

Proof of (2.6). We are going to use the Green function estimates (3.1) and (3.3) of Lemma 3.1 to estimate the two terms of inequality (4.1).

As for the first term:

$$\begin{aligned} (I) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)} \int_{B_R(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) \, d\mu_{\mathbb{H}^N}(x) \\ &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} \left[\int_{B_R(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) \, d\mu_{\mathbb{H}^N}(x) \right]^{\frac{m}{m-1}} \end{aligned} \quad (4.2)$$

$$\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} \left(\frac{\kappa_1}{2s}\right)^{\frac{m}{m-1}} \cosh(R)^{\frac{m(N-1)}{m-1}} R^{\frac{2sm}{m-1}},$$

where we have used Young's inequality $ab \leq \frac{a^m}{m} + \frac{m-1}{m} b^{\frac{m}{m-1}}$ and the estimate (3.3). The constant $c_m = \frac{m-1}{m} 2^{\frac{m^2}{(m-1)^2}}$ only depends on $m > 1$.

As for the second term, from the estimate (3.1), we derive

$$\begin{aligned} (II) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \\ &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(t_0, x) \frac{1}{(r(x, x_0))^{N-2s}} d\mu_{\mathbb{H}^N}(x) \leq \frac{2^{\frac{m}{m-1}} \kappa_1}{t_0} \frac{\|u(t_0, x)\|_{L^1(\mathbb{H}^N)}}{R^{N-2s}}. \end{aligned} \quad (4.3)$$

Combining inequalities (4.1), (4.2) and (4.3) we obtain

$$\begin{aligned} u^m(t_0, x_0) &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} \left(\frac{\kappa_1}{2s}\right)^{\frac{m}{m-1}} \cosh(R)^{\frac{m(N-1)}{m-1}} R^{\frac{2sm}{m-1}} + \frac{2^{\frac{m}{m-1}} \kappa_1}{t_0} \frac{\|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{N-2s}} \\ &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \kappa_4 \frac{R^{\frac{2sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[\cosh(R)^{\frac{m(N-1)}{m-1}} + \frac{t_0^{\frac{1}{m-1}} \|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{\frac{1}{(m-1)\vartheta_1}}} \right] \end{aligned}$$

where $\kappa_4 = \kappa_4(s, m, \kappa_1)$. By taking the supremum in $x_0 \in \mathbb{H}^N$ we obtain

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{m \kappa_4}{m-1} \frac{R^{\frac{2sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[\cosh(R)^{\frac{m(N-1)}{m-1}} + \frac{t_0^{\frac{1}{m-1}} \|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{\frac{1}{(m-1)\vartheta_1}}} \right]. \quad (4.4)$$

Choosing now $R = \left(t_0^{\frac{1}{m-1}} \|u(t_0)\|_{L^1(\mathbb{H}^N)}\right)^{(m-1)\vartheta_1}$ gives

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{m \kappa_4}{m-1} \frac{\|u(t_0)\|_{L^1(\mathbb{H}^N)}^{2sm\vartheta_1}}{t_0^{Nm\vartheta_1}} \left[1 + \cosh\left(t_0^{\vartheta_1} \|u(t_0)\|_{L^1(\mathbb{H}^N)}^{(m-1)\vartheta_1}\right)^{\frac{m(N-1)}{m-1}} \right],$$

which proves inequality (2.6) for all $t_0 \leq \|u_0\|_{L^1(\mathbb{H}^N)}^{-(m-1)}$, recalling that the L^1 -norm is decreasing in time, $\|u(t_0)\|_{L^1(\mathbb{H}^N)} \leq \|u_0\|_{L^1(\mathbb{H}^N)}$. As for the constant

$$\bar{\kappa}_1^m = \frac{m \kappa_4}{m-1} \left[1 + \cosh(1)^{\frac{m(N-1)}{m-1}} \right].$$

The inequality can be easily extended to $t_0 \geq \|u_0\|_{L^1(\mathbb{H}^N)}^{-(m-1)}$ by a standard time-scaling argument.

Once we have proven the estimate (2.6) for $0 \leq u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, it is easy to see that this can be extended to all $0 \leq u_0 \in L^1(\mathbb{H}^N)$. Indeed, given $0 \leq u_0 \in L^1(\mathbb{H}^N)$ consider a sequence of data $0 \leq u_{0,n} \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ converging pointwise monotonically from below (consider for instance $u_{0,n} = u_0 \wedge n$) to u_0 and also in the strong $L^1(\mathbb{H}^N)$ topology. Let $u_n(t) \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ be the corresponding solutions. Then it is clear that $u_n(t)$ converges to $u(t)$ for any fixed t , where $u(t)$ is the solution corresponding to u_0 . Moreover, by lower semicontinuity of the L^∞ norm and applying estimate (2.6) to u_n , we conclude

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \liminf_{n \rightarrow \infty} \|u_n(t)\|_{L^\infty(\mathbb{H}^N)} \leq \bar{\kappa}_0 \liminf_{n \rightarrow \infty} \frac{\|u_{0,n}\|_{L^1(\mathbb{H}^N)}^{2s\theta_1}}{t^{N\theta_1}} = \bar{\kappa}_0 \frac{\|u_0\|_{L^1(\mathbb{H}^N)}^{2s\theta_1}}{t^{N\theta_1}}.$$

Proof of (2.7). We are going to use the Green function estimates (3.2) and (3.4) of Lemma 3.1 to estimate the two terms of inequality (4.1), when $R \geq 2$. As for the first term:

$$\begin{aligned}
 (I) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)} \int_{B_R(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \\
 &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} \left[\int_{B_R(x_0)} \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \right]^{\frac{m}{m-1}} \\
 &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} (\tilde{\kappa}_1)^{\frac{m}{m-1}} R^{\frac{sm}{m-1}},
 \end{aligned} \tag{4.5}$$

where we have used Young's inequality $ab \leq \frac{a^m}{m} + \frac{m-1}{m} b^{\frac{m}{m-1}}$ and the estimate (3.4). The constant $c_m = \frac{m-1}{m} 2^{\frac{m^2}{(m-1)^2}}$ only depends on $m > 1$.

As for the second term:

$$\begin{aligned}
 (II) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \\
 &\leq \frac{\kappa_1 2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} \frac{u(t_0, x)}{r(x, x_0)^{1-s}} e^{-(N-1)r(x, x_0)} d\mu_{\mathbb{H}^N}(x) \\
 &\leq \frac{\kappa_1 2^{\frac{m}{m-1}}}{t_0} \frac{\|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{1-s} e^{(N-1)R}},
 \end{aligned} \tag{4.6}$$

where we have used the estimate (3.2). Combining inequalities (4.1), (4.5) and (4.6) we obtain

$$\begin{aligned}
 u^m(t_0, x_0) &\leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} (\tilde{\kappa}_1)^{\frac{m}{m-1}} R^{\frac{sm}{m-1}} + \frac{\kappa_1 2^{\frac{m}{m-1}}}{t_0} \frac{\|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{1-s} e^{(N-1)R}} \\
 &= \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \kappa_5 \frac{R^{\frac{sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[1 + \frac{t_0^{\frac{1}{m-1}} \|u(t_0)\|_{L^1(\mathbb{H}^N)}}{R^{1-s + \frac{sm}{m-1}} e^{(N-1)R}} \right]
 \end{aligned}$$

where $\kappa_5 = \kappa_5(s, m, \kappa_1)$. By taking the supremum in $x_0 \in \mathbb{H}^N$, recalling that we are assuming $R \geq 2$ and that $\|u(t_0)\|_{L^1(\mathbb{H}^N)} \leq \|u_0\|_{L^1(\mathbb{H}^N)}$, we obtain

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{m \kappa_5}{m-1} \frac{R^{\frac{sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[1 + \frac{t_0^{\frac{1}{m-1}} \|u_0\|_{L^1(\mathbb{H}^N)}}{2^{1-s + \frac{sm}{m-1}} e^{(N-1)R}} \right].$$

Choosing now $R = \frac{1}{N-1} \log \left(t_0^{\frac{1}{m-1}} \|u_0\|_{L^1(\mathbb{H}^N)} \right)$, the above inequality gives

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{\bar{\kappa}_2^m}{t_0^{\frac{m}{m-1}}} \left[\log \left(t_0 \|u_0\|_{L^1(\mathbb{H}^N)}^{m-1} \right) \right]^{\frac{sm}{m-1}}$$

with constant

$$\bar{\kappa}_2^m := \frac{m \kappa_5}{m-1} \left[1 + \frac{1}{2^{1 + \frac{s}{m-1}}} \right] \left(\frac{1}{(N-1)(m-1)} \right)^{\frac{sm}{m-1}}.$$

We finally remark that the assumption $R \geq 2$ and our choice of R , restrict the validity of the above inequality to large times, namely we have

$$R \geq 2 \quad \text{if} \quad t_0 \geq e^{2(N-1)(m-1)} \|u_0\|_{L^1(\mathbb{H}^N)}^{-(m-1)}.$$

We have thus obtained (2.7) and the proof is concluded.

4.2. Proof of Theorem 2.3. Let us start with the fundamental estimates :

$$\begin{aligned} u^m(t_0, x_0) &\leq \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x) \\ &= \underbrace{\frac{2^{\frac{m}{m-1}}}{t_0} \int_{B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x)}_{(I)} + \underbrace{\frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} u(t_0, x) \mathbb{G}_{\mathbb{H}^N}^s(x, x_0) d\mu_{\mathbb{H}^N}(x)}_{(II)}. \end{aligned} \quad (4.7)$$

The estimate for (I) works exactly in the proof of Theorem 2.2 i.e., for all $R > 0$ we have

$$(I) \leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} \left(\frac{\kappa_1}{2s}\right)^{\frac{m}{m-1}} \cosh(R)^{\frac{m(N-1)}{m-1}} R^{\frac{2sm}{m-1}} \quad (4.8)$$

where $c_m = \frac{m}{m-1} 2^{\frac{m^2}{(m-1)^2}}$ and κ_1 is as in Lemma 3.1. Instead, if $R \geq 2$, as in (4.5) we get

$$(I) \leq \frac{1}{m} \|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m + \frac{c_m}{t_0^{\frac{m}{m-1}}} (\tilde{\kappa}_1)^{\frac{m}{m-1}} R^{\frac{sm}{m-1}} \quad (4.9)$$

with $\tilde{\kappa}_1$ defined in (3.4).

Proof of (2.8). First, combining (2.4) with Corollary A.2, we observe that

$$\frac{\mathbb{G}_{\mathbb{H}^N}^s(x_0, x)}{\Phi_1(x)} \leq \begin{cases} \frac{\bar{C}_3 \alpha^{-1} e^{\frac{(N-1)r(x_0, x)}{2}}}{(r(x_0, x))^{N-2s} (1+r(x_0, x))} & \text{for } r(x_0, x) \leq 1, \\ \frac{\bar{C}_4 \alpha^{-1} e^{-\frac{(N-1)r(x_0, x)}{2}}}{r(x_0, x)^{1-s} (1+r(x_0, x))} & \text{for } r(x_0, x) \geq 1. \end{cases} \quad (4.10)$$

Hence, by the fact that $e^{-\frac{(N-1)r}{2}}/r^{-N+s+1}$ is bounded for $r > 1$, we infer that

$$\frac{\mathbb{G}_{\mathbb{H}^N}^s(x_0, x)}{\Phi_1(x)} \leq \frac{\kappa_6}{(r(x_0, x))^{N-2s} (1+r(x_0, x))} \quad \text{for all } r(x_0, x) > 0$$

for some $\kappa_6 = \kappa_6(s, m, \kappa_1)$. Using the above, (II) can be estimated as follows

$$(II) \leq \frac{2^{\frac{m}{m-1}} \kappa_6}{t_0 R^{N-2s} (1+R)} \|u(t_0)\|_{L_{\Phi_1}^1(\mathbb{H}^N)}.$$

Now, by (4.8) and Young's inequality we obtain

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{m \kappa_7}{m-1} \frac{R^{\frac{2sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[\cosh(R)^{\frac{m(N-1)}{m-1}} + \frac{t_0^{\frac{1}{m-1}} \|u_0\|_{L_{\Phi_1}^1(\mathbb{H}^N)}}{R^{\frac{1}{(m-1)\vartheta_1}} (1+R)} \right] \quad (4.11)$$

where $\kappa_7 = \kappa_7(s, m, \kappa_1)$. If we estimate the r.h.s. of (4.11) by using $1+R > 1$, then the same argument adopted for estimating (4.4) yields (2.8) with

$$\bar{\kappa}_3^m := \frac{m \kappa_7}{m-1} \left[1 + \cosh(1)^{\frac{m(N-1)}{m-1}} \right].$$

Note that here we use the fact that the $L^1_{\Phi_1}$ -norm is decreasing in time, i.e. $\|u(t_0)\|_{L^1_{\Phi_1}(\mathbb{H}^N)} \leq \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}$, which follows from Proposition 3.5. Note that the inequality can be extended to all $t_0 > 0$ by a standard time-scaling argument.

Proof of (2.9). Take $R \geq 2$ in (4.7), by (4.10) we have

$$\begin{aligned} (II) &= \frac{2^{\frac{m}{m-1}}}{t_0} \int_{\mathbb{H}^N \setminus B_R(x_0)} \frac{\mathbb{G}_{\mathbb{H}^N}^s(x, x_0)}{\Phi_1(x)} u(t_0, x) \Phi_1(x) d\mu_{\mathbb{H}^N}(x) \\ &\leq \frac{2^{\frac{m}{m-1}} \bar{C}_4}{\alpha t_0} \frac{e^{-\frac{(N-1)R}{2}}}{R^{2-s}} \|u(t_0)\|_{L^1_{\Phi_1}(\mathbb{H}^N)} \\ &\leq \frac{2^{\frac{m}{m-1}} \bar{C}_4}{\alpha t_0} \frac{e^{-\frac{(N-1)R}{2}}}{R^{2-s}} \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}, \end{aligned} \quad (4.12)$$

where for the last estimate we have exploited the fact that $\|u(t_0)\|_{L^1_{\Phi_1}(\mathbb{H}^N)} \leq \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}$. Therefore, by (4.9) and (4.12), we conclude that

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \frac{m \kappa_8}{m-1} \frac{R^{\frac{sm}{m-1}}}{t_0^{\frac{m}{m-1}}} \left[1 + \frac{e^{-\frac{(N-1)R}{2}}}{R^{2+\frac{s}{m-1}}} t_0^{\frac{1}{m-1}} \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)} \right], \quad (4.13)$$

where $\kappa_8 = \kappa_8(s, m, \kappa_1)$. Finally, we choose

$$R = \frac{2}{(N-1)} \log \left(t_0^{\frac{1}{m-1}} \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)} \right)$$

and substituting in (4.13) we obtain

$$\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}^m \leq \bar{\kappa}_4^m \frac{\left[\log(t_0 \|u_0\|_{L^1_{\Phi_1}(\mathbb{H}^N)}) \right]^{\frac{sm}{m-1}}}{t_0^{\frac{m}{m-1}}},$$

where

$$\bar{\kappa}_4^m := \frac{m \kappa_8}{m-1} \left[1 + \frac{1}{2^{2+\frac{s}{m-1}}} \right] \left(\frac{2}{(N-1)(m-1)} \right)^{\frac{sm}{m-1}}.$$

The assumption $R \geq 2$ and our choice of R , restrict the validity of the above inequality to large times. More precisely, we infer that

$$R \geq 2 \quad \text{if} \quad t_0 \geq e^{(N-1)(m-1)} \|u_0\|_{L^1(\mathbb{H}^N)}^{-(m-1)}.$$

We have thus obtained (2.9) and the proof is concluded.

4.3. Proof of Theorem 2.4. We begin again by exploiting the fundamental estimate (4.7). I_1 can be estimated as in (4.8), with the only difference that we take the supremum over the ball instead of all the space, indeed we have

$$I_1 \leq \frac{c(\cosh(R_0))^{N-1}}{t_0} \|u(t_0, x)\|_{L^\infty(B_{R_0}(x_0))} R_0^{2s} \quad \text{for } 0 < R_0 \leq 2.$$

Let

$$\tilde{g}(R_0) := (\cosh(R_0))^{N-1} R_0^{2s}$$

so that

$$I_1 \leq \frac{c}{t_0} \tilde{g}(R_0) \|u(t_0, x)\|_{L^\infty(B_{R_0}(x_0))} \quad \text{for } 0 \leq R_0 \leq 2.$$

For I_2 , recalling that $\Phi \in \mathcal{W}$ yields $\Phi = (-\Delta_{\mathbb{H}^N})^{-s} \bar{\psi} \in \mathcal{W}$ for some positive $\bar{\psi} \in C_c^\infty(\mathbb{H}^N)$, we fix $z_0 \in \mathbb{H}^N$ and we set

$$\bar{g}(R_0, z_0) := \sup_{x_0 \in B_{R_0}(z_0)} \sup_{x \in \mathbb{H}^N \setminus B_{R_0}(x_0)} \frac{\mathbb{G}_{\mathbb{H}^N}^s(x_0, x)}{(-\Delta_{\mathbb{H}^N})^{-s} \bar{\psi}(x)},$$

where, by Lemma A.3, $\bar{g}(R_0, z_0)$ is bounded. Then, exploiting the monotonicity formula of Proposition 3.4, we get

$$I_2 \leq \frac{2^{\frac{m}{m-1}}}{t_0} \bar{g}(R_0, z_0) \int_{\mathbb{H}^N \setminus B_{R_0}(x_0)} u(t_0, x) \Phi(x) dv_{\mathbb{H}^N} \leq \frac{c}{t_0} \bar{g}(R_0, z_0) \|u_0(x)\|_{L_{\Phi}^1(\mathbb{H}^N)},$$

for some $c > 0$ depending only on m, s, N .

Joining the above estimates we obtain

$$u^m(t_0, x_0) \leq \frac{c}{t_0} \tilde{g}(R_0) \|u(t_0)\|_{L^\infty(B_{R_0}(x_0))} + \frac{c}{t_0} \bar{g}(R_0, z_0) \|u_0\|_{L_{\Phi}^1(\mathbb{H}^N)}.$$

Taking the supremum over $x_0 \in B_{R_0}(z_0)$ we obtain

$$\begin{aligned} \|u(t_0)\|_{L^\infty(B_{R_0}(z_0))}^m &\leq \frac{c}{t_0} \tilde{g}(R_0) \|u(t_0)\|_{L^\infty(B_{2R_0}(z_0))} + \frac{c}{t_0} \bar{g}(R_0) \|u(t_0)\|_{L_{\Phi}^1(\mathbb{H}^N)} \\ &\leq \frac{1}{2} \|u(t_0)\|_{L^\infty(B_{2R_0}(z_0))}^m + \frac{c}{t_0^{\frac{m}{m-1}}} \tilde{g}(R_0)^{\frac{m}{m-1}} + \frac{c}{t_0} \bar{g}(R_0, z_0) \|u(t_0)\|_{L_{\Phi}^1(\mathbb{H}^N)}. \end{aligned}$$

Eliminate the first term on the right-hand side by a De Giorgi lemma [24, Lemma 6.1] we obtain:

$$\|u(t_0)\|_{L^\infty(B_{R_0}(z_0))}^m \leq \frac{c}{t_0^{\frac{m}{m-1}}} \tilde{g}(R_0)^{\frac{m}{m-1}} + \frac{c}{t_0} \bar{g}(R_0, z_0) \|u(t_0)\|_{L_{\Phi}^1(\mathbb{H}^N)} \quad (4.14)$$

for all $R_0 \leq 2$. Since $\tilde{g}(R_0) \leq (\cosh(2))^{N-1} R_0^{2s}$ and $\bar{g}(R_0, z_0) \leq c$, by taking

$$R_0 = \left(t_0^{\frac{1}{m-1}} \|u_0\|_{L_{\Phi}^1(\mathbb{H}^N)} \right)^{2s},$$

(4.14) gives:

$$\|u(t_0)\|_{L^\infty(B_{R_0}(z_0))}^m \leq \frac{\bar{\kappa}_5^m}{t_0} \|u_0\|_{L_{\Phi}^1(\mathbb{H}^N)}$$

for some $\bar{\kappa}_5^m$. Note that the right-hand side does not depend on z_0 , hence the same estimate holds for $\|u(t_0)\|_{L^\infty(\mathbb{H}^N)}$ and this concludes the proof for small times; the inequality can be extended to all $t_0 > 0$ by a standard time-scaling argument.

5. SEMIGROUP THEORY IN BANACH AND HILBERT SPACES: EXISTENCE, UNIQUENESS, CONTRACTIVITY AND COMPARISON.

There are two parallel theories based on nonlinear semigroup and gradient flow techniques, respectively posed in the $L^1(\mathbb{H}^N)$ or $H^{-s}(\mathbb{H}^N)$ framework. The nowadays classical theories of Benilan-Crandall-Pazy-Pierre [6, 7, 20], in the spirit of the celebrated Crandall-Liggett theorem in Banach spaces [19], or the Brezis-Komura theory [16, 17, 37] of maximal monotone operators in Hilbert spaces, both apply to our problem. Therefore, it is possible to prove existence, uniqueness and comparison results of mild solutions (solutions obtained as limit of an Implicit Time Discretization -ITD- scheme, see also [1]). Such mild (or semigroup) solutions, turn out to be unique as a consequence of contractivity of the solution map S_t in the respective spaces, and actually the so-called T-contractivity (contractivity with absolute value replaced by the positive part) holds and implies

comparison. However, in general mild solutions have low regularity, not enough to guarantee boundedness (the energy may not be finite). Once constructed, we have to show that mild solutions are indeed Weak Dual Solutions and hence bounded, as a consequence of the smoothing effects proven in Section 4.

We devote the first part of this section to briefly explain the relevant ideas of the two nonlinear semigroup approaches, omitting tedious and standard details: we need these solutions in order to approximate the WDS that we want to construct in suitable weighted spaces, which is one of our main results. The last part of the section is devoted to the proof of the existence and uniqueness theorems for WDS, Theorem 2.1.

5.1. Nonlinear Semigroup in $L^1(\mathbb{H}^N)$. Mild VS Weak Dual Solutions. The basic theory in $L^1(\mathbb{H}^N)$ developed by Benilan-Crandall-Pazy-Pierre [6, 7, 20] applies to our setting. Indeed, our aim is to solve an equation of the form $\partial_t u = -\mathcal{L}\varphi(u)$, where the operator $\mathcal{L} : \text{dom}(\mathcal{L}) \subset L^1(\mathbb{H}^N) \rightarrow L^1(\mathbb{H}^N)$ is a densely defined linear m -accretive (see also Remark 5.1) operator of sub-Markovian type, and the nonlinearity $\varphi(r) = r^m$ satisfies suitable ‘‘monotonicity conditions’’ that we omit here, since the nonlinearity $\varphi(r) = r^m$ with $m > 1$ is the prototype example and obviously fulfills the conditions. Indeed, this theory allows to prove existence and uniqueness for a larger class of nonlinearities and operators, but we prefer to stick to the simplest nontrivial example, in order to focus on the main ideas. In the present case, the operator is $\mathcal{L} = (-\Delta_{\mathbb{H}^N})^s$ and the nonlinearity $\varphi(r) = r^m$, and both clearly satisfy the assumptions required for Theorem 3 and 4 of [20] to hold. The same assumptions imply that the semigroup is indeed T -contractive, or order preserving, cf. Chapter 19.4 of [7]. We summarize the above mentioned results in the following

Theorem 5.1 (Nonlinear Semigroup in $L^1(\mathbb{H}^N)$, [6, 7, 20]). *Let $u_0, v_0 \in L^1(\mathbb{H}^N)$. Then there exist unique mild solutions $u, v \in C^0([0, \infty) : L^1(\mathbb{H}^N))$ to Problem 1.1, such that*

$$\int_{\mathbb{H}^N} (u(t, x) - v(t, x))_+ d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} (u(0, x) - v(0, x))_+ d\mu_{\mathbb{H}^N}(x) \quad \text{for all } t \geq 0,$$

and the same holds replacing $(\cdot)_+$ with $(\cdot)_-$. As a consequence,

$$\|u(t) - v(t)\|_{L^1(\mathbb{H}^N)} \leq \|u_0 - v_0\|_{L^1(\mathbb{H}^N)} \quad \text{for all } t \geq 0.$$

Moreover, nonnegative mild solutions enjoy the following time monotonicity property:

$$\text{the map } t \mapsto t^{\frac{1}{m-1}} u(t, x) \quad \text{is nondecreasing in } t > 0 \text{ for a.e. } x \in \mathbb{H}^N.$$

As we have already explained above, now that we have built unique mild solutions in $L^1(\mathbb{H}^N)$, we need to show that these are also WDS, hence bounded, as a consequence of Theorem 2.2. We devote the rest of this subsection to the proof of this fact, borrowing ideas from Section 7 of [15].

Theorem 5.2. *Let u be the mild solution to (1.1) corresponding to the non-negative initial datum $u_0 \in L^1(\mathbb{H}^N)$. Then, u is a WDS in the sense of Definition 2.1 and it is bounded, in particular it satisfies the smoothing effects of Theorem 2.2.*

To prove the above theorem we need the following auxiliary result:

Proposition 5.3 (L^p -stability of Mild Solutions). *Let $0 \leq u_0 \in L^1(\mathbb{H}^N) \cap L^p(\mathbb{H}^N)$. Then, the unique mild solution to (1.1) satisfies*

$$\|u(t)\|_{L^p(\mathbb{H}^N)} \leq \|u_0\|_{L^p(\mathbb{H}^N)} \quad \text{for all } t \geq 0 \text{ and all } 1 \leq p \leq \infty. \quad (5.1)$$

Remark 5.1. *L^p stability of very weak solutions.* Let us remark that the L^p stability is easily enjoyed by WDS. Indeed, first we have to notice that WDS are indeed very weak (or distributional) solutions, namely they satisfy the following identity for every test function $\varphi \in C_c^\infty((0, T) \times \mathbb{H}^N)$:

$$\int_0^T \int_{\mathbb{H}^N} u(t, x) \partial_t \varphi(t, x) d\mu_{\mathbb{H}^N} dt - \int_0^T \int_{\mathbb{H}^N} u^m(t, x) (-\Delta_{\mathbb{H}^N})^s \varphi(t, x) d\mu_{\mathbb{H}^N} dt = 0 \quad (5.2)$$

The idea to show that WDS are very weak solutions is to put $\psi = (-\Delta_{\mathbb{H}^N})^s \varphi$ in the integral identity of Definition 2.1 of WDS: however, to make it rigorous some approximations are needed. Note that even if $0 \leq \varphi \in C_c^\infty$, we always have that $(-\Delta_{\mathbb{H}^N})^s \varphi \asymp \Phi \asymp e^{(N-1)r} r^{s-1}$, see Lemma A.3. This is a peculiarity of the nonlocal case, in contrast with the local case $s = 1$, for which $\text{supp}(\mathcal{L}u) \subseteq \text{supp}(u)$. For this reason, we need to require further integrability conditions also for very weak solutions, for instance $u \in L_{loc}^1((0, \infty) : L_{loc}^1(\mathbb{H}^N))$ and that $u^m \in L_{loc}^1((0, T) : L_\Phi^1(\mathbb{H}^N))$. It is not clear whether or not WDS and very weak solutions are indeed the same class.

For very weak solutions, the L^p stability is just a consequence of the m -accretivity of the operator $\mathcal{L} = (-\Delta_{\mathbb{H}^N})^s$, which reads: if β is a maximal monotone graph in $\mathbb{R} \times \mathbb{R}$, with $0 \in \beta(0)$, $u \in \text{dom}(\mathcal{L})$, $\mathcal{L}u \in L^p(\mathbb{H}^N)$, with $1 \leq p \leq \infty$, and $v \in L^{\frac{p}{p-1}}(\mathbb{H}^N)$, $v(x) \in \beta(u(x))$ a.e., then

$$\int_{\mathbb{H}^N} v(x) \mathcal{L}u(x) d\mu_{\mathbb{H}^N}(x) \geq 0.$$

In particular, it is not difficult to see that if $0 \leq u \in \text{dom}(\mathcal{L}) \cap L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, we always have

$$\int_{\mathbb{H}^N} u^q(x) \mathcal{L}u^m(x) d\mu_{\mathbb{H}^N}(x) \geq 0 \quad \text{for all } q, m > 0. \quad (5.3)$$

Let now u be a very weak solution of $u_t = -\mathcal{L}u^m$, corresponding to $0 \leq u_0 \in C_c^\infty(\mathbb{H}^N) \subset \text{dom}(\mathcal{L}) \cap L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Inequality (5.1) then follows formally by integrating the following differential inequality for any $p \in [1, \infty]$,

$$\frac{d}{dt} \int_{\mathbb{H}^N} u^p(t, x) d\mu_{\mathbb{H}^N}(x) = -p \int_{\mathbb{H}^N} u^{p-1}(t, x) \mathcal{L}u^m(t, x) d\mu_{\mathbb{H}^N}(x) \leq 0,$$

where in the last step we have used (5.3). The above proof can be made rigorous through a long but standard approximation argument: the above differential inequality (integrated on $[0, T]$), is nothing but the definition of very weak solution (5.2) with the choice of test function $\psi(t, x) = u^{p-1}$. Of course, that choice is not admissible, but it can be approximated by a sequence of admissible test functions $\psi_n \in C_c^\infty((0, \infty) \times \mathbb{H}^N)$ through a careful but standard limiting process.

We now sketch the proof -which follows the ideas of Proposition 7.1 of [15]- of the L^p stability for mild solutions, which only relies on the definition of mild solutions (without passing through very weak solutions, hence avoiding tedious and long approximations).

Proof. The case $p = 1$ follows directly by Theorem 5.1. Fix $1 < p < \infty$ and for $n \geq 1$ set $u_{0,n}(x) := T_n u_0(x)$ where $T_n : \mathbb{R} \rightarrow \mathbb{R}$ is defined as follows

$$T_n(x) = \begin{cases} x & \text{if } |x| \leq n, \\ \frac{x}{|x|} n & \text{if } |x| > n. \end{cases} \quad (5.4)$$

Clearly, $u_{0,n} \in L^p(\mathbb{H}^N) \subset L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ and $u_{0,n} \rightarrow u_0$ a.e. in \mathbb{H}^N , hence by dominated convergence $u_{0,n} \rightarrow u_0$ strongly in $L^p(\mathbb{H}^N)$.

We claim that it is sufficient to show that for any fixed $t > 0$, the semigroup solution $u_n(t)$ corresponding to the initial datum $u_{0,n}$ satisfies

$$\|u_n(t)\|_{L^p(\mathbb{H}^N)} \leq \|u_{0,n}\|_{L^p(\mathbb{H}^N)} \leq \|u_0\|_{L^p(\mathbb{H}^N)} \quad (5.5)$$

to conclude the proof of the Proposition, since the above inequality easily implies (5.1). Indeed, since $u_n(t)$ is bounded in $L^p(\mathbb{H}^N)$, then, up to subsequences, $u_n \rightharpoonup u$ in $L^p(\mathbb{H}^N)$ and, using lower semicontinuity of weak convergence, we obtain

$$\|u(t)\|_{L^p(\mathbb{H}^N)} \leq \liminf_n \|u_n(t)\|_{L^p(\mathbb{H}^N)} \leq \|u_0\|_{L^p(\mathbb{H}^N)}.$$

Hence, (5.1) holds true for all $p < \infty$. For $p = \infty$ we just let $p \rightarrow \infty$ in the above inequality.

It just remains to prove (5.5), and without loss of generality we can assume $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$. Mild solutions are obtained via an ITD scheme: consider the following partition of $[0, T]$ (with $t_0 = 0$ and $t_n = T$)

$$t_k = \frac{k}{n}T, \quad \text{for any } 0 \leq k \leq n,$$

with $h = t_{k+1} - t_k = \frac{T}{n}$. For any $t \in (0, T)$, the unique semigroup solution $u(t, \cdot)$ is obtained as the strong $L^1(\mathbb{H}^N)$ -limit of solutions $u_{k+1}(\cdot) = u(t_{k+1}, \cdot)$ to the following elliptic equation:

$$h(-\Delta_{\mathbb{H}^N})^s(u_{k+1})^m + u_{k+1} = u_k \quad \text{in } \mathbb{H}^N \quad (5.6)$$

whose solvability is guaranteed by the running assumption on the operator and nonlinearity, cf. Theorem 5.2, in particular

$$\int_{\mathbb{H}^N} (u_{k+1}^p - u_k u_{k+1}^{p-1}) d\mu_{\mathbb{H}^N} = -h \int_{\mathbb{H}^N} u_{k+1}^m (-\Delta_{\mathbb{H}^N})^s(u_{k+1})^{p-1} d\mu_{\mathbb{H}^N} \geq 0.$$

(This is equivalent to the sub-Markovianity of \mathcal{L}). Therefore, we obtain

$$\begin{aligned} \int_{\mathbb{H}^N} u_{k+1}^p d\mu_{\mathbb{H}^N} &\leq \int_{\mathbb{H}^N} u_k u_{k+1}^{p-1} d\mu_{\mathbb{H}^N} \\ &\leq \|u_k\|_{L^p(\mathbb{H}^N)} \left(\int_{\mathbb{H}^N} u_{k+1}^p d\mu_{\mathbb{H}^N} \right)^{\frac{p-1}{p}}. \end{aligned}$$

A simple inductive argument shows that

$$\|u_{k+1}\|_{L^p(\mathbb{H}^N)} \leq \|u_k\|_{L^p(\mathbb{H}^N)} \leq \dots \leq \|u_0\|_{L^p(\mathbb{H}^N)}.$$

We conclude the proof by letting $k \rightarrow \infty$ to obtain

$$\|u(t)\|_{L^p(\mathbb{H}^N)} \leq \liminf_{k \rightarrow \infty} \|u_{k+1}(t)\|_{L^p(\mathbb{H}^N)} \leq \|u_0\|_{L^p(\mathbb{H}^N)}.$$

□

Proof of Theorem 5.2. We sketch the proof which follows the ideas of Proposition 7.2 of [15] which deals with FPME on bounded Euclidean domains. We will just emphasize the main points for convenience of the reader. We first pass to a weak dual formulation of the approximate problem elliptic problem (5.6), i.e. we apply $(-\Delta_{\mathbb{H}^N})^{-s}$ to both members of (5.6) to get:

$$(-\Delta_{\mathbb{H}^N})^{-s}u_{k+1} - (-\Delta_{\mathbb{H}^N})^{-s}u_k = -h(u_{k+1})^m \quad \text{in } \mathbb{H}^N.$$

By multiplying the above equation by $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$, we obtain (upon using the notation $\psi_k = \psi(\cdot, t_k)$)

$$\sum_{k=0}^{n-1} \int_{\mathbb{H}^N} [(-\Delta_{\mathbb{H}^N})^{-s}u_{k+1} - (-\Delta_{\mathbb{H}^N})^{-s}u_k] \psi_k d\mu_{\mathbb{H}^N} = -h \sum_{k=0}^{n-1} \int_{\mathbb{H}^N} (u_{k+1})^m \psi_k d\mu_{\mathbb{H}^N}.$$

Since ψ is compactly supported in time, then for sufficiently large n , we have

$$\psi_1 = \psi(\cdot, t_1) = \psi(\cdot, \frac{T}{n}) = 0 \quad \text{and} \quad \psi_{n-1} \rightarrow \psi(\cdot, T) = 0.$$

By this we infer that

$$\begin{aligned} & \int_{\mathbb{H}^N} [\psi_{n-1} (-\Delta_{\mathbb{H}^N})^{-s} u_n - \psi_0 (-\Delta_{\mathbb{H}^N})^{-s} u_0] d\mu_{\mathbb{H}^N} \\ & - \underbrace{\sum_{k=1}^{n-1} \int_{\mathbb{H}^N} (\psi_k - \psi_{k-1}) (-\Delta_{\mathbb{H}^N})^{-s} u_k d\mu_{\mathbb{H}^N}}_{=I_1} = - \underbrace{h \sum_{k=0}^{n-1} \int_{\mathbb{H}^N} (u_{k+1})^m \psi_k(x) d\mu_{\mathbb{H}^N}}_{=I_2}. \end{aligned}$$

Let us carefully look at I_1 :

$$\begin{aligned} I_1 &= \sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} \frac{(\psi_k - \psi_{k-1})}{h} (-\Delta_{\mathbb{H}^N})^{-s} u_k d\mu_{\mathbb{H}^N} \\ &= \sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} (\psi_k)_t(x, \hat{t}_k) (-\Delta_{\mathbb{H}^N})^{-s} (u_k - u) d\mu_{\mathbb{H}^N} + \sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} (\psi_k)_t(x, \hat{t}_k) (-\Delta_{\mathbb{H}^N})^{-s} u d\mu_{\mathbb{H}^N}. \end{aligned}$$

Now consider the above two integrals separately. Since ψ is compactly supported, we obtain

$$\begin{aligned} & \sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} (\psi_k)_t(\cdot, \hat{t}_k) (-\Delta_{\mathbb{H}^N})^{-s} (u_k - u) d\mu_{\mathbb{H}^N} \\ &= \sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s} ((\psi_k)_t(\cdot, \hat{t}_k)) (u_k - u) d\mu_{\mathbb{H}^N} \\ &\leq \|u_k - u\|_{L^1(\mathbb{H}^N)} \sum_{k=1}^{n-1} h \|(-\Delta_{\mathbb{H}^N})^{-s} ((\psi_k)_t(\cdot, \hat{t}_k))\|_{L^\infty(K)} \xrightarrow{n \rightarrow \infty} 0, \end{aligned}$$

where we have exploited the fact that, $\|u_k - u\|_{L^1(\mathbb{H}^N)} \xrightarrow{n \rightarrow \infty} 0$ and that,

$$\sum_{k=1}^{n-1} h \|(-\Delta_{\mathbb{H}^N})^{-s} ((\psi_k)_t(\cdot, \hat{t}_k))\|_{L^\infty(K)} \xrightarrow{n \rightarrow \infty} \int_0^T \|(-\Delta_{\mathbb{H}^N})^{-s} \psi_t(\cdot, t)\|_{L^\infty(\mathbb{H}^N)} dt < \infty,$$

the latter integral being bounded by arguing as in Lemma A.3.

On the other hand, we recognize that the second integral is nothing but a Riemann sum, so that

$$\sum_{k=1}^{n-1} h \int_{\mathbb{H}^N} (\psi_k)_t(x, \hat{t}_k) (-\Delta_{\mathbb{H}^N})^{-s} u d\mu_{\mathbb{H}^N} \longrightarrow \int_0^T \int_{\mathbb{H}^N} (\psi_t) (-\Delta_{\mathbb{H}^N})^{-s} (u) d\mu_{\mathbb{H}^N} dt,$$

as $n \rightarrow \infty$. Hence,

$$I_1 \xrightarrow{n \rightarrow \infty} 0 \int_0^T \int_{\mathbb{H}^N} (\psi_t) (-\Delta_{\mathbb{H}^N})^{-s} (u) d\mu_{\mathbb{H}^N} dt. \quad (5.7)$$

A similar computation yields

$$I_2 \xrightarrow{n \rightarrow \infty} 0 \int_0^T \int_{\mathbb{H}^N} u^m \psi d\mu_{\mathbb{H}^N} dt. \quad (5.8)$$

Moreover, letting $n \rightarrow \infty$, we obtain

$$\int_{\mathbb{H}^N} [\psi_{n-1} (-\Delta_{\mathbb{H}^N})^{-s} u_n - \psi_0 (-\Delta_{\mathbb{H}^N})^{-s} u_0] d\mu_{\mathbb{H}^N} \xrightarrow{n \rightarrow \infty} 0. \quad (5.9)$$

Finally, as a consequence of (5.7), (5.8) and (5.9) we obtain

$$\int_0^T \int_{\mathbb{H}^N} (\psi_t) (-\Delta_{\mathbb{H}^N})^{-s} (u) d\mu_{\mathbb{H}^N} dt - \int_0^T \int_{\mathbb{H}^N} u^m \psi d\mu_{\mathbb{H}^N} dt = 0,$$

for all $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$. The above expression shows that u is a WDS, recalling that $L^1(\mathbb{H}^N) \subset L_{\Phi}^1(\mathbb{H}^N)$ and $L^1(\mathbb{H}^N) \subset L_{\Phi_1}^1(\mathbb{H}^N)$. Furthermore, the boundedness of solutions follows by Theorem 2.2, which holds for WDS with $0 \leq u_0 \in L^1(\mathbb{H}^N)$. \square

5.2. Nonlinear Semigroup in $H^{-s}(\mathbb{H}^N)$ and Gradient Flows. The celebrated theory of maximal monotone operators in Hilbert spaces of Brezis and Komura, [16, 17, 37], applies in this framework, see for instance Brezis' paper [16]. The link between such theory and the theory of Gradient Flows has been very well explained by Ambrosio et al. in [1], see also [2]. We will briefly explain how the previously mentioned theory applies to the present case.

We choose as Hilbert space $H^{-s}(\mathbb{H}^N)$, the dual of $H^s(\mathbb{H}^N)$, with the Hilbertian norm given by

$$\|f\|_{H^{-s}(\mathbb{H}^N)}^2 = \int_{\mathbb{H}^N} f (-\Delta_{\mathbb{H}^N})^{-s} f d\mu_{\mathbb{H}^N} = \int_{\mathbb{H}^N} |(-\Delta_{\mathbb{H}^N})^{-s/2} f|^2 d\mu_{\mathbb{H}^N}$$

and the corresponding scalar product. Define the (convex, nonlinear) energy functional

$$E_m[f] = \frac{1}{m+1} \int_{\mathbb{H}^N} |f|^{m+1} d\mu_{\mathbb{H}^N}$$

whenever finite, $+\infty$ elsewhere. The equation $u_t = -(-\Delta)^s u^m$ can be interpreted as the subdifferential inclusion $u_t \in \partial[E_m](u)$, where $\partial[E_m]$ is the sub-differential in $H^{-s}(\mathbb{H}^N)$ of the convex energy functional E_m .

We summarize in the following theorem the results of the Brezis-Komura theory.

Theorem 5.4 (Brezis-Komura, [16, 17, 37]). *For every $u_0 \in \overline{\text{dom} E_m} = H^{-s}(\mathbb{H}^N)$, there exists a unique gradient flow solution starting from u_0 , that we denote by $u(t) = S_t u_0$. This defines a strongly continuous semigroup $S_t : H^{-s}(\mathbb{H}^N) \rightarrow H^{-s}(\mathbb{H}^N)$ for $t > 0$, i.e. $S_{t+\tau} = S_t S_\tau$ for any $t, \tau > 0$ with the contraction property*

$$\|u(t) - v(t)\|_{H^{-s}(\mathbb{H}^N)} \leq \|u_0 - v_0\|_{H^{-s}(\mathbb{H}^N)} \quad \text{for all } u_0, v_0 \in H^{-s}(\mathbb{H}^N).$$

Moreover, the T -contraction property holds true,

$$\|(u(t) - v(t))_+\|_{H^{-s}(\mathbb{H}^N)} \leq \|(u_0 - v_0)_+\|_{H^{-s}(\mathbb{H}^N)} \quad \text{for all } u_0, v_0 \in H^{-s}(\mathbb{H}^N). \quad (5.10)$$

Hence standard comparison holds in $H^{-s}(\mathbb{H}^N)$. Moreover, for a.e. $t > 0$ we have

$$\frac{1}{2} \frac{d}{dt} \|u(t)\|_{H^{-s}(\mathbb{H}^N)}^2 = -\frac{1}{m+1} \int_{\mathbb{H}^N} |u(t)|^{m+1} d\mu_{\mathbb{H}^N} = -E_m[u(t)] \leq 0$$

and

$$\frac{d}{dt} E_m[u(t)] = - \int_{\mathbb{H}^N} u^m (-\Delta_{\mathbb{H}^N})^s u^m d\mu_{\mathbb{H}^N} \leq 0.$$

Finally, the equation $u_t = (-\Delta_{\mathbb{H}^N})^s u^m$ holds true for almost every $t > 0$ as equality among functions in $H^{-s}(\mathbb{H}^N)$. We call this solutions H^{-s} -strong solutions.

The concept of gradient flow solution corresponds in this case to H^{-s} -strong solutions, which are a priori, a smaller class than mild solutions, hence more regular. More details can be found in [1, 17, 37]. The proof of inequality (5.10) is due to Brezis and it has been sketched in Lemma 3.5.

The proof of this theorem, considered nowadays the nonlinear analogous of the Hille-Yosida Theorem, follows by a careful analysis of the convergence of the ITD, or the Implicit Euler Scheme. An excellent exposition is given in the Lecture Notes of Ambrosio et al. [1], where a characterization in terms of EVI, Evolution Variational Inequalities, is given. Recall that an absolutely continuous curve in $u(t) \in H^{-s}(\mathbb{H}^N)$ is called an EVI solution if for any $y \in \text{dom}E_m \subset H^{-s}(\mathbb{H}^N)$ we have

$$\frac{1}{2} \frac{d}{dt} \|u(t) - y\|_{H^{-s}(\mathbb{H}^N)}^2 + E_m[u(t)] \leq E_m[y] \quad \text{for a.e. } t > 0. \quad (5.11)$$

In Section 11 of [1] it is carefully explained that u is an EVI solution if and only if it is a gradient flow solution as in Theorem 5.4. Hence we just have to check inequality (5.11), namely

$$\begin{aligned} \frac{1}{2} \frac{d}{dt} \|u(t) - y\|_{H^{-s}(\mathbb{H}^N)}^2 &= \int_{\mathbb{H}^N} u_t(t) (-\Delta_{\mathbb{H}^N})^{-s} (u(t) - y) d\mu_{\mathbb{H}^N} = \int_{\mathbb{H}^N} (u - y) (-\Delta_{\mathbb{H}^N})^{-s} u_t d\mu_{\mathbb{H}^N} \\ &= - \int_{\mathbb{H}^N} (u - y) u^m d\mu_{\mathbb{H}^N} = - \int_{\mathbb{H}^N} u^{m+1} d\mu_{\mathbb{H}^N} + \int_{\mathbb{H}^N} y u^m d\mu_{\mathbb{H}^N} \\ &\leq - \left(1 - \frac{m}{m+1}\right) \int_{\mathbb{H}^N} u^{m+1} d\mu_{\mathbb{H}^N} + \frac{1}{m+1} \int_{\mathbb{H}^N} y^{m+1} d\mu_{\mathbb{H}^N} = -E_m[u(t)] + E_m[y], \end{aligned}$$

where we have used Fubini's theorem, the dual equation $(-\Delta_{\mathbb{H}^N})^{-s} u_t = -u^m$, and Young's inequality $ab \leq \frac{m}{m+1} a^{\frac{m+1}{m}} + \frac{1}{m+1} b^{m+1}$.

Remark 5.2. Strong H^{-s} solutions are WDS. When u_0 is sufficiently integrable, for instance $u_0 \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ (weaker assumptions are possible), then the gradient flow solutions (or strong H^{-s} solutions) constructed in Theorem 5.4, first introduced by Brezis and Komura [16, 17, 37], turn out to be also WDS. Let us recall the definition of strong H^{-s} solution: let $u \in C^0([0, T] : H^{-s}(\mathbb{H}^N))$, $u^m \in L^1([0, T] : H^s(\mathbb{H}^N))$, be such that

$$\int_0^T \int_{\mathbb{H}^N} u \partial_t \psi d\mu_{\mathbb{H}^N} dt = \int_0^T \int_{\mathbb{H}^N} u^m (-\Delta_{\mathbb{H}^N})^s \psi d\mu_{\mathbb{H}^N} dt \quad \forall \psi \in C_c^1([0, T] : H^s(\mathbb{H}^N))$$

or, equivalently, since $(-\Delta_{\mathbb{H}^N})^s : H^s(\mathbb{H}^N) \rightarrow H^{-s}(\mathbb{H}^N)$ is an isomorphism, let $(-\Delta_{\mathbb{H}^N})^s \psi = \eta$,

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s} u \partial_t \eta d\mu_{\mathbb{H}^N} dt = \int_0^T \int_{\mathbb{H}^N} u^m \eta d\mu_{\mathbb{H}^N} dt \quad \forall \eta \in C_c^1([0, T] : H^{-s}(\mathbb{H}^N)).$$

The above solutions turn out to be equivalent to Weak Dual Solutions in the sense of Definition 2.1, when the initial datum is sufficiently integrable. This can be seen using the Poincaré inequality $\lambda_1 \|f\|_{H^{-s}(\mathbb{H}^N)} \leq \|f\|_{L^2(\mathbb{H}^N)}$, together with the L^p stability of very weak solutions, see Lemma 3.7 and Remark 5.1. As a consequence, we have that H^{-s} -strong solutions are also WDS: this is clear since the class of test functions of WDS is strictly contained in the above class of admissible test functions, more precisely $C_c^1([0, T] : H^{-s}(\mathbb{H}^N)) \subset C_c^1([0, T] : L_c^\infty(\mathbb{H}^N))$.

Notice that the concept of WDS is somehow more general than the one of H^{-s} strong solutions, indeed nonnegative H^{-s} functions are in L_Φ^1 , namely for any $0 \leq f \in H^{-s}(\mathbb{H}^N)$ we have

$$\begin{aligned} \|f\|_{L_\Phi^1(\mathbb{H}^N)} &= \int_{\mathbb{H}^N} f (-\Delta_{\mathbb{H}^N})^{-s} \psi d\mu_{\mathbb{H}^N} = \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s/2} f (-\Delta_{\mathbb{H}^N})^{-s/2} \psi d\mu_{\mathbb{H}^N} \\ &\leq \left(\int_{\mathbb{H}^N} |(-\Delta_{\mathbb{H}^N})^{-s/2} \psi|^2 d\mu_{\mathbb{H}^N} \right)^{\frac{1}{2}} \left(\int_{\mathbb{H}^N} |(-\Delta_{\mathbb{H}^N})^{-s/2} f|^2 d\mu_{\mathbb{H}^N} \right)^{\frac{1}{2}} = c_\psi \|f\|_{H^{-s}(\mathbb{H}^N)}. \end{aligned} \quad (5.12)$$

Hence $H_+^{-s} \subseteq L_{\Phi,+}^1(\mathbb{H}^N)$. Note that $c_\psi = \int_{\mathbb{H}^N} \psi(-\Delta_{\mathbb{H}^N})^{-s} \psi d\mu_{\mathbb{H}^N} = \int_{\mathbb{H}^N} \psi \Phi d\mu_{\mathbb{H}^N} = \|\psi\|_{L_{\Phi}^1} < +\infty$ since $0 \leq \psi \in C_c^\infty(\mathbb{H}^N)$ and $\Phi = (-\Delta_{\mathbb{H}^N})^{-s} \psi$.

We would like to stress that as a consequence of our $L_{\Phi}^1 - L^\infty$ smoothing effects, Theorem 2.4, we have shown that H^{-s} -strong solutions are bounded, see Remark 2.4.

5.3. Proof of Theorem 2.1. The proof of existence of (minimal) weak dual solutions will be obtained by approximations in terms of mild solutions, more precisely the minimal weak dual solution will be the limit of a pointwise monotone sequence of (unique) mild solutions: we have decided to take, as approximating sequences, L^1 -mild solutions, whose existence and uniqueness is guaranteed by Theorem 5.1. However, a similar proof would hold by replacing L^1 -mild solutions with H^{-s} -gradient-flow solutions, whose existence and uniqueness is guaranteed by Theorem 5.4.

Approximation: Let $u_0 \geq 0$ be the initial datum (the same construction will work in $L^1(\mathbb{H}^N)$, $L_{\Phi_1}^1(\mathbb{H}^N)$ or $L_{\Phi}^1(\mathbb{H}^N)$). Define, for $n \geq 1$, $u_{0,n}(x) := \chi_{B_n(x_0)}(x) T_n u_0(x)$, where T_n is defined as in (5.4). Hence, $u_{0,n} \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$ and $u_{0,n}(x) \leq u_{0,n+1}(x) \leq u_0(x) \in L^1(\mathbb{H}^N)$ for a.e. $x \in \mathbb{H}^N$. By comparison, we know that the unique L^1 -mild solutions satisfy: $u_n(t, x) \leq u_{n+1}(t, x)$ for a.e. $x \in \mathbb{H}^N$ and a.e. $t > 0$, hence the limit

$$u(t, x) := \liminf_{n \rightarrow \infty} u_{n+1}(t, x) \quad (5.13)$$

exists for a.e. $x \in \mathbb{H}^N$ and a.e. $t > 0$ as limit of monotone sequences of real numbers. This is our candidate to be the minimal weak dual solution. We will analyze different setups below. As a consequence of the Monotone Convergence Theorem, we have that $u_n(t) \rightarrow u(t)$ as $n \rightarrow \infty$ in the strong $L^1(\mathbb{H}^N)$, $L_{\Phi_1}^1(\mathbb{H}^N)$ or $L_{\Phi}^1(\mathbb{H}^N)$ topology, depending on where u_0 belongs.

Proof of Theorem 2.1. The case $u_0 \in L^1(\mathbb{H}^N)$. First we show that the WDS defined in (5.13) is bounded, more precisely, by semicontinuity of the L^∞ -norm, we have that for all $t > 0$

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \liminf_{n \rightarrow +\infty} \|u_n(t)\|_{L^\infty(\mathbb{H}^N)} \leq \bar{\kappa}_1 \liminf_{n \rightarrow +\infty} \frac{\|u_{0,n}\|_{L^1(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}} = \bar{\kappa}_1 \frac{\|u_0\|_{L^1(\mathbb{H}^N)}^{2s\vartheta_1}}{t^{N\vartheta_1}},$$

and we have used the $L^1 - L^\infty$ smoothing estimates (2.6), which hold true for the mild solutions of the approximating sequence u_n , as shown in Theorem 5.2.

As a consequence, for all $t > 0$, $u(t, x) \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N) \subset L^p(\mathbb{H}^N)$ for all $p \geq 1$. Now, for $1 < p < +\infty$, the elementary inequality $a^p - b^p \leq pa^{p-1}(a - b)$ gives

$$0 \leq \int_{\mathbb{H}^N} (u^p(t, x) - u_n^p(t, x)) d\mu_{\mathbb{H}^N}(x) \leq p \|u(t, x)\|_{L^\infty(\mathbb{H}^N)}^{p-1} \int_{\mathbb{H}^N} (u(t, x) - u_n(t, x)) d\mu_{\mathbb{H}^N}(x). \quad (5.14)$$

Hence, $u_n(t) \rightarrow u(t)$ strongly in L^p , for all $p \in (1, \infty)$ and $t > 0$, as a consequence of the L^1 -strong convergence.

We are finally ready to verify that u is a weak dual solution in the sense of Definition 2.1:

- $u \in C([0, T] : L^1(\mathbb{H}^N))$. This follows by noting that for all $0 \leq \tau < t$ or $0 \leq t < \tau$, one has

$$\begin{aligned} \int_{\mathbb{H}^N} |u(t, x) - u(\tau, x)| d\mu_{\mathbb{H}^N}(x) &\leq \int_{\mathbb{H}^N} |u(t, x) - u_n(t, x)| d\mu_{\mathbb{H}^N}(x) \\ &\quad + \int_{\mathbb{H}^N} |u_n(t, x) - u_n(\tau, x)| d\mu_{\mathbb{H}^N}(x) + \int_{\mathbb{H}^N} |u_n(\tau, x) - u(\tau, x)| d\mu_{\mathbb{H}^N}(x). \end{aligned}$$

Then, recalling that $u_n \in C([0, \infty) : L^1(\mathbb{H}^N))$, the continuity for $t > 0$ readily follows from above; similarly, for $t = 0$ we also obtain that $u(0, x) = u_0(x)$ a.e. in \mathbb{H}^N , by

$$\begin{aligned} \int_{\mathbb{H}^N} |u(t, x) - u_0(x)| d\mu_{\mathbb{H}^N}(x) &\leq \int_{\mathbb{H}^N} |u(t, x) - u_n(t, x)| d\mu_{\mathbb{H}^N}(x) \\ &\quad + \int_{\mathbb{H}^N} |u_n(t, x) - u_{n,0}(x)| d\mu_{\mathbb{H}^N}(x) + \int_{\mathbb{H}^N} |u_{n,0}(x) - u_0(x)| d\mu_{\mathbb{H}^N}(x). \end{aligned}$$

- $u^m \in L^1((0, T) : L^1(\mathbb{H}^N))$ for any fixed $T > 0$. To this aim, we note that, from inequality (2.6) of Theorem 2.2, we have

$$\begin{aligned} \int_0^T \int_{\mathbb{H}^N} u^m(t, x) d\mu_{\mathbb{H}^N}(x) dt &\leq \int_0^T \|u(t, x)\|_{L^\infty(\mathbb{H}^N)}^{m-1} \int_{\mathbb{H}^N} u(t, x) d\mu_{\mathbb{H}^N}(x) dt \\ &\leq \bar{\kappa}_1^{m-1} \|u_0(x)\|_{L^1(\mathbb{H}^N)}^{2s\vartheta_1(m-1)+1} \int_0^T \frac{1}{t^{N\vartheta_1(m-1)}} dt < +\infty, \end{aligned}$$

where the latter integral is bounded since $N\vartheta_1(m-1) < 1$.

- u satisfies the following integral identity

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u) \frac{\partial \psi}{\partial t} d\mu_{\mathbb{H}^N} dt - \int_0^T \int_{\mathbb{H}^N} u^m \psi d\mu_{\mathbb{H}^N} dt = 0$$

for every test function $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$. This follows by recalling that we have proven in Theorem 5.2 that being u_n an L^1 -mild solution, it is also a weak dual solution, hence we just have to take limits in the following equality:

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u_n) \frac{\partial \psi}{\partial t} d\mu_{\mathbb{H}^N} dt - \int_0^T \int_{\mathbb{H}^N} u_n^m \psi d\mu_{\mathbb{H}^N} dt = 0$$

for every test function $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$. Having in mind this goal, we note that

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u_n) \frac{\partial \psi}{\partial t} d\mu_{\mathbb{H}^N} dt = \int_0^T \int_{\mathbb{H}^N} u_n (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) d\mu_{\mathbb{H}^N} dt$$

and that, as $n \rightarrow +\infty$, we have

$$\int_0^T \int_{\mathbb{H}^N} u_n (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) d\mu_{\mathbb{H}^N} dt \rightarrow \int_0^T \int_{\mathbb{H}^N} u (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) d\mu_{\mathbb{H}^N} dt,$$

since the sequence $\{u_n\}$ is monotone, $u \in L^1((0, T); L^1(\mathbb{H}^N))$ and, by Lemma A.3, $(-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) \in L^\infty((0, T) \times \mathbb{H}^N)$. Furthermore, as $n \rightarrow +\infty$, we have

$$\int_0^T \int_{\mathbb{H}^N} u_n^m \psi d\mu_{\mathbb{H}^N} dt \rightarrow \int_0^T \int_{\mathbb{H}^N} u^m \psi d\mu_{\mathbb{H}^N} dt$$

since, the sequence $\{u_n^m\}$ is monotone and $u^m \in L^1((0, T); L^1(\mathbb{H}^N))$.

The proof of Theorem 2.1 when $u_0 \in L^1(\mathbb{H}^N)$ is concluded.

Proof of Theorem 2.1. The cases $u_0 \in L_{\Phi}^1(\mathbb{H}^N)$ with $\Phi \in \mathcal{W}$ and $u_0 \in L_{\Phi_1}^1(\mathbb{H}^N)$. The proof is similar to the one for $u_0 \in L^1(\mathbb{H}^N)$, hence we just sketch it.

Again, the first step consist in showing that the WDS defined in (5.13) is bounded, more precisely, by semicontinuity of the L^∞ -norm, we have that for all $t > 0$

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \liminf_{n \rightarrow +\infty} \|u_n(t)\|_{L^\infty(\mathbb{H}^N)} \leq \bar{\kappa}_1 \liminf_{n \rightarrow +\infty} \frac{\|u_{0,n}\|_{L_{\Phi_1}^{1, 2s\vartheta_1}(\mathbb{H}^N)}}{t^{N\vartheta_1}} = \bar{\kappa}_1 \frac{\|u_0\|_{L_{\Phi_1}^{1, 2s\vartheta_1}(\mathbb{H}^N)}}{t^{N\vartheta_1}},$$

and we have used the $L^1_{\Phi_1} - L^\infty$ smoothing estimates (2.6), which hold true for the mild solutions of the approximating sequence u_n , as shown in Theorem 5.2. Similarly for $\Phi \in \mathcal{W}$, using the smoothing estimate (2.8), we get

$$\|u(t)\|_{L^\infty(\mathbb{H}^N)} \leq \liminf_{n \rightarrow +\infty} \|u_n(t)\|_{L^\infty(\mathbb{H}^N)} \leq \frac{\bar{\kappa}_5}{t^{1/m}} \liminf_{n \rightarrow +\infty} \|u_{0,n}\|_{L^1_{\Phi}(\mathbb{H}^N)}^{1/m} = \frac{\bar{\kappa}_5}{t^{1/m}} \|u_0\|_{L^1_{\Phi}(\mathbb{H}^N)}^{1/m}. \quad (5.15)$$

As a consequence, $u_n(t) \rightarrow u(t)$ strongly in $L^p_{\Phi_1}$ (resp. L^p_{Φ}), for all $p \in (1, \infty)$ and $t > 0$, as a consequence of the $L^1_{\Phi_1}$ -strong convergence (resp. L^1_{Φ} -strong convergence), using the weighted analogous of inequality (5.14) which has essentially the same proof.

We are finally ready to verify that u is a weak dual solution in the sense of Definition 2.1:

The case of $\Phi \in \mathcal{W}$.

- $u \in C([0, T] : L^1_{\Phi}(\mathbb{H}^N))$. This follows by noting that for all $0 \leq \tau < t$ or $0 \leq t < \tau$, one has

$$\begin{aligned} \int_{\mathbb{H}^N} |u(t, x) - u(\tau, x)| \Phi(x) \, d\mu_{\mathbb{H}^N}(x) &\leq \int_{\mathbb{H}^N} |u(t, x) - u_n(t, x)| \Phi(x) \, d\mu_{\mathbb{H}^N}(x) \\ &\quad + \int_{\mathbb{H}^N} |u_n(t, x) - u_n(\tau, x)| \Phi(x) \, d\mu_{\mathbb{H}^N}(x) + \int_{\mathbb{H}^N} |u_n(\tau, x) - u(\tau, x)| \Phi(x) \, d\mu_{\mathbb{H}^N}(x). \end{aligned}$$

Using that $u_n(t) \rightarrow u(t)$ strongly in L^1_{Φ} and that $u_n \in C([0, \infty) : L^1(\mathbb{H}^N)) \subset C([0, T] : L^1_{\Phi}(\mathbb{H}^N))$, yield the continuity for $t > 0$; when $t = 0$ we proceed analogously.

- $u^m \in L^1((0, T) : L^1_{\Phi}(\mathbb{H}^N))$. This follows by the smoothing estimate (5.15). Indeed, we have

$$\begin{aligned} \int_0^T \int_{\mathbb{H}^N} u(t, x)^m \Phi(x) \, d\mu_{\mathbb{H}^N}(x) \, dt &\leq \int_0^T \|u(t, x)\|_{L^\infty(\mathbb{H}^N)}^{m-1} \int_{\mathbb{H}^N} u(t, x) \Phi(x) \, d\mu_{\mathbb{H}^N}(x) \, dt \\ &\leq \|u_0(x)\|_{L^1_{\Phi}(\mathbb{H}^N)}^{\frac{2m-1}{m}} \int_0^T \frac{\bar{\kappa}_5^{m-1}}{t_0^{\frac{m-1}{m}}} \, dt < +\infty, \end{aligned}$$

since $\frac{m-1}{m} < 1$.

- u satisfies the following identity

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u) \frac{\partial \psi}{\partial t} \, d\mu_{\mathbb{H}^N}(x) \, dt - \int_0^T \int_{\mathbb{H}^N} u^m \psi \, d\mu_{\mathbb{H}^N}(x) \, dt = 0$$

for every test function $\psi \in C^1_c(0, T; L^\infty_c(\mathbb{H}^N))$. This follows by recalling that u_n is a weak dual solution and then passing to the limit in the equality:

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u_n) \frac{\partial \psi}{\partial t} \, d\mu_{\mathbb{H}^N}(x) \, dt - \int_0^T \int_{\mathbb{H}^N} u_n^m \psi \, d\mu_{\mathbb{H}^N}(x) \, dt = 0$$

for every test function $\psi \in C^1_c(0, T; L^\infty_c(\mathbb{H}^N))$. To this aim, we note that

$$\int_0^T \int_{\mathbb{H}^N} (-\Delta_{\mathbb{H}^N})^{-s}(u_n) \frac{\partial \psi}{\partial t} \, d\mu_{\mathbb{H}^N}(x) \, dt = \int_0^T \int_{\mathbb{H}^N} u_n (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) \, d\mu_{\mathbb{H}^N}(x) \, dt$$

and that, as $n \rightarrow +\infty$, we have

$$\int_0^T \int_{\mathbb{H}^N} u_n (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) \, d\mu_{\mathbb{H}^N}(x) \, dt \rightarrow \int_0^T \int_{\mathbb{H}^N} u (-\Delta_{\mathbb{H}^N})^{-s} \left(\frac{\partial \psi}{\partial t} \right) \, d\mu_{\mathbb{H}^N}(x) \, dt,$$

since the sequence $\{u_n\}$ is monotone, $u \in L^1((0, T); L^1_{\Phi}(\mathbb{H}^N))$ and, arguing as in the proof of by Lemma A.3, it readily follows that $|(-\Delta_{\mathbb{H}^N})^{-s}(\frac{\partial \psi}{\partial t})| \leq C(t)\Phi(x)$ for some $C \in C^0_c(0, T)$.

Furthermore, as $n \rightarrow +\infty$, we have

$$\int_0^T \int_{\mathbb{H}^N} u_n^m \psi \, d\mu_{\mathbb{H}^N}(x) \, dt \rightarrow \int_0^T \int_{\mathbb{H}^N} u^m \psi \, d\mu_{\mathbb{H}^N}(x) \, dt$$

since the sequence $\{u_n^m\}$ is monotone and $u^m \in L^1((0, T); L^1_{\Phi}(\mathbb{H}^N))$.

The case of Φ_1 . This case is completely analogous to the previous ones, hence we omit the proof.

Uniqueness of Minimal Weak Dual Solutions.

The proof is an adaptation to the present setting of the Proof of Theorem 4.5 of [15], we sketch it here for convenience of the reader. Here Φ will denote either an element of the class \mathcal{L} or the ground state Φ_1 . Essentially, this uniqueness for the minimal WDS follows by the T-contraction property. Let u be the WDS corresponding to the initial datum u_0 , obtained as limit of the monotone non-decreasing sequence u_k . Consider another monotone non-decreasing sequence $0 \leq v_{0,k} \leq v_{0,k+1} \leq u_0$, with $v_{0,k} \in L^1(\mathbb{H}^N) \cap L^\infty(\mathbb{H}^N)$, monotonically converging from below to $u_0 \in L^1_{\Phi}(\mathbb{H}^N)$ in the strong $L^1_{\Phi}(\mathbb{H}^N)$ topology. Repeating the construction, we obtain another weak dual solution $v(t, x) \in C^0([0, \infty) : L^1_{\Phi}(\mathbb{H}^N))$. We want to show that $u = v$ by showing that $v \leq u$ and then that $u \leq v$. To prove that $v \leq u$ we use the estimates

$$\int_{\mathbb{H}^N} [v_k(t, x) - u_n(t, x)]_+ \, d\mu_{\mathbb{H}^N}(x) \leq \int_{\mathbb{H}^N} [v_k(0, x) - u_n(0, x)]_+ \, d\mu_{\mathbb{H}^N}(x)$$

which hold for all $u_n(t, \cdot)$ and $v_k(t, \cdot)$, see Theorem 5.1. Letting $n \rightarrow \infty$ we get that

$$\begin{aligned} \lim_{n \rightarrow \infty} \int_{\mathbb{H}^N} [v_k(t, x) - u_n(t, x)]_+ \, d\mu_{\mathbb{H}^N}(x) &\leq \lim_{n \rightarrow \infty} \int_{\mathbb{H}^N} [v_k(0, x) - u_n(0, x)]_+ \, d\mu_{\mathbb{H}^N}(x) \\ &= \int_{\mathbb{H}^N} [v_k(0, x) - u_0(x)]_+ \, d\mu_{\mathbb{H}^N}(x) = 0 \end{aligned}$$

since $v_k(0, x) \leq u_0$ by construction. Therefore also $v_k(t, x) \leq u(t, x)$ for $t > 0$, so that in the limit $k \rightarrow \infty$ we obtain $v(t, x) \leq u(t, x)$. The inequality $u \leq v$ can be obtained simply by switching the roles of u_n and v_k . \square

APPENDIX A. GREEN FUNCTION ESTIMATES IN \mathbb{H}^N

This section is devoted to provide proper estimates of Green function $\mathbb{G}_{\mathbb{H}^N}^s(x, y)$ for the fractional laplacian on the hyperbolic space for $0 < s < 1$. It is well-known that the Green function is given by the following explicit formula

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) = \int_0^{+\infty} \frac{k_{\mathbb{H}^N}(t, x, y)}{t^{1-s}} \, dt,$$

where $k_{\mathbb{H}^N}(t, x, y)$ is the heat kernel for $-\Delta_{\mathbb{H}^N}$. Furthermore, $k_{\mathbb{H}^N}$ satisfies the following estimates (see [21]): for $N \geq 2$, there exist some positive constants A_N and B_N such that

$$A_N h_N(t, x, y) \leq k_{\mathbb{H}^N}(t, x, y) \leq B_N h_N(t, x, y) \quad \text{for all } t > 0 \text{ and } x, y \in \mathbb{H}^N, \quad (\text{A.1})$$

where $h_N(t, x, y)$ is given by

$$h_N(t, x, y) := h_N(t, r) = (4\pi t)^{-\frac{N}{2}} e^{-\frac{(N-1)^2 t}{4} - \frac{(N-1)r}{2} - \frac{r^2}{4t}} (1+r+t)^{\frac{(N-3)}{2}} (1+r), \quad (\text{A.2})$$

where $r := r(x, y) = \text{dist}(x, y)$.

Using (A.1) we shall obtain the following Green functions estimates:

Lemma A.1. *Let $N \geq 3$ and $0 < s < 1$. Then there exist $R_1, R_2, \underline{C}_1, \overline{C}_1, \underline{C}_2, \overline{C}_2 > 0$, only depending on N and s , such that*

$$\frac{\underline{C}_1}{(r(x, y))^{N-2s}} \leq \mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \frac{\overline{C}_1}{(r(x, y))^{N-2s}} \quad \text{for all } (x, y) \in \mathbb{H}^N : 0 < r(x, y) \leq R_1; \quad (\text{A.3})$$

and

$$\frac{\underline{C}_2 e^{-(N-1)r(x, y)}}{(r(x, y))^{1-s}} \leq \mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \frac{\overline{C}_2 e^{-(N-1)r(x, y)}}{(r(x, y))^{1-s}} \quad \text{for all } (x, y) \in \mathbb{H}^N : r(x, y) \geq R_2. \quad (\text{A.4})$$

Proof. Let $h_N(t, r)$ be as defined in (A.2), we compute

$$\begin{aligned} \int_0^{+\infty} \frac{h_N(t, r)}{t^{1-s}} dt &= \int_0^\infty \frac{(4\pi t)^{N/2} e^{-\frac{(N-1)^2}{4}t - \frac{(N-1)r}{2} - \frac{r^2}{4t}} (1+r+t)^{(N-3)/2} (1+r)}{t^{1-s}} dt \\ &= \frac{e^{-(N-1)r/2} (1+r)}{(4\pi)^{N/2}} \int_0^\infty t^{-N/2-1+s} (1+r+t)^{(N-3)/2} e^{-\left(\frac{(N-1)\sqrt{t}}{2} - \frac{r}{2\sqrt{t}}\right)^2} e^{-(N-1)r/2} dt \\ &= \frac{e^{-(N-1)r} (1+r)}{(4\pi)^{N/2}} \int_0^\infty t^{-\frac{N+2s-2}{2}} (1+r+t)^{(N-3)/2} e^{-\left(\frac{(N-1)\sqrt{t}}{2} - \frac{r}{2\sqrt{t}}\right)^2} dt. \end{aligned} \quad (\text{A.5})$$

Now making the following substitution one can write

$$z = \frac{r}{2\sqrt{t}} - \frac{(N-1)\sqrt{t}}{2} \Leftrightarrow \sqrt{t} = \frac{-z + \sqrt{z^2 + (N-1)r}}{N-1}.$$

Therefore, we have

$$dt = -\frac{2}{(N-1)^2} \frac{(\sqrt{z^2 + (N-1)r} - z)^2}{\sqrt{z^2 + (N-1)r}} dz.$$

Further substituting back in (A.5) yields

$$\begin{aligned} \int_0^{+\infty} \frac{h_N(t, r)}{t^{1-s}} dt &= \frac{e^{-(N-1)r} (1+r)}{(N-1)^{2s-3} (4\pi)^{N/2}} \times \\ &\left\{ \int_{-\infty}^\infty \frac{(1+r + (\sqrt{z^2 + (N-1)r} - z)^2)^{(N-3)/2} (\sqrt{z^2 + (N-1)r} - z)^2 e^{-z^2}}{(-z + \sqrt{z^2 + (N-1)r})^{N-2s+2} \sqrt{z^2 + (N-1)r}} dz \right\}. \end{aligned}$$

Now for $r \rightarrow \infty$, one can look for the leading term as follows

$$\begin{aligned} \int_0^{+\infty} \frac{h_N(t, r)}{t^{1-s}} dt &= C_N e^{-(N-1)r} (1+r) r^{-\frac{N+2s-2}{2} + \frac{N-3}{2} + \frac{1}{2}} \times \\ &\left\{ \int_{-\infty}^\infty \frac{((1/r + 1)(N-1)^2 + (\sqrt{z^2/r + (N-1)} - z/\sqrt{r})^2)^{(N-3)/2} (\sqrt{z^2/r + (N-1)} - z/\sqrt{r})^2 e^{-z^2}}{(-z/r + \sqrt{z^2 + (N-1)})^{N-2s+2} \sqrt{z^2/r + (N-1)}} dz \right\}. \end{aligned}$$

$= I_N$

The finiteness of I_N follows easily from the fact that e^{-z^2} is the leading term at infinity. Therefore, recalling (A.1), we obtain

$$\int_0^{+\infty} \frac{h_N(t, r)}{t^{1-s}} dt = C_N I_N e^{-(N-1)r} r^{s-1} + o(e^{-(N-1)r} r^{s-1}) \quad \text{as } r \rightarrow \infty.$$

Hence, we obtain (A.4). Now we turn to prove (A.3). To see this we can make the change of variable as $\alpha = \frac{r^2}{4t}$ and therefore we can write

$$\int_0^{+\infty} \frac{h_N(t, r)}{t^{1-s}} dt = C_N \frac{e^{-(N-1)r/2}(1+r)}{r^{(N-2s)}} \underbrace{\int_0^\infty \alpha^{\frac{N-2s-2}{2}} e^{-\frac{(N-1)^2 r^2}{16\alpha}} - \alpha \left(1+r+\frac{r^2}{4\alpha}\right)^{\frac{(N-3)}{2}} d\alpha}_{= J_N}.$$

Again, as before, the integral is finite for $0 < s < 1$ and hence

$$G_s(r, t) = C_N J_N r^{-(N-2s)} + o(r^{-(N-2s)}) \quad \text{as } r \rightarrow 0^+.$$

This proves the lemma. □

From Lemma A.1 we derive

Corollary A.2. *Let $N \geq 3$ and $0 < s < 1$. Then there exist positive constants $\underline{C}_3, \overline{C}_3, \underline{C}_4, \overline{C}_4$, only depending on N and s , such that*

$$\underline{C}_3 \frac{1}{(r(x, y))^{N-2s}} \leq \mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \overline{C}_3 \frac{1}{(r(x, y))^{N-2s}} \quad \text{for all } (x, y) \in \mathbb{H}^N : 0 < r(x, y) \leq 1, \quad (\text{A.6})$$

$$\underline{C}_4 \frac{e^{-(N-1)r(x, y)}}{(r(x, y))^{1-s}} \leq \mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \overline{C}_4 \frac{e^{-(N-1)r(x, y)}}{(r(x, y))^{1-s}} \quad \text{for all } (x, y) \in \mathbb{H}^N : r(x, y) \geq 1. \quad (\text{A.7})$$

Furthermore, the following global estimate holds:

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \frac{\overline{C}_5}{(r(x, y))^{N-2s}} \quad \text{for all } (x, y) \in \mathbb{H}^N : r(x, y) > 0, \quad (\text{A.8})$$

where $\overline{C}_5 = \min\{\overline{C}_3, \overline{C}_4\}$.

Proof. The proof of this corollary is a straightforward consequence of Lemma A.1. We give a detailed proof of (A.7), the proof of (A.6) can be achieved with similar arguments. On the other hand, the proof of (A.8) follows by noting that $e^{-(N-1)r} \leq r^{-N+s+1}$ for all $r > 1$ and then combining (A.6) with (A.7).

Proof of (A.7). We may assume $R_2 > 1$, where R_2 is as given in Lemma A.1 (otherwise, there is nothing to prove). Next we set

$$g(r) := \frac{e^{-(N-1)r}}{r^{1-s}} \quad \text{with } r > 0 \quad (\text{A.9})$$

and we note that

$$g(r(x, y)) \geq g(R_2) \quad \text{for all } 1 \leq r(x, y) \leq R_2.$$

Set $M := \max_{1 \leq r(x, y) \leq R_2} \mathbb{G}_{\mathbb{H}^N}^s(x, y)$, then

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq M = \frac{M}{g(R_2)} g(R_2) \leq \frac{M}{g(R_2)} g(r(x, y)) \quad \text{for all } 1 \leq r(x, y) \leq R_2.$$

By taking $\overline{C}_4 := \max\{\frac{M}{g(R_2)}, \overline{C}_2\}$ we get the upper bound in (A.7).

Now, to obtain the lower bound, we note that

$$g(r(x, y)) \leq g(1) \quad \text{for all } 1 \leq r(x, y) \leq R_2.$$

Then, we set $m := \min_{1 \leq r(x, y) \leq R_2} \mathbb{G}_{\mathbb{H}^N}^s(x, y)$ and we conclude

$$\mathbb{G}_{\mathbb{H}^N}^s(x, y) \geq m = \frac{m}{g(1)}g(1) \geq \frac{m}{g(1)}g(r(x, y)) \quad \text{for all } 1 \leq r(x, y) \leq R_2.$$

Taking $\underline{C}_4 := \min\{\frac{m}{g(1)}, \underline{C}_2\}$, we obtain the lower bound in (A.7). □

It can be useful to check that, if $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$ with $\psi \geq 0$, then $(-\Delta_{\mathbb{H}^N})^{-s}(\psi)$ behaves like $G_s(x_0, x)$ near infinity:

Lemma A.3. *For all $\psi \in C_c^1(0, T; L_c^\infty(\mathbb{H}^N))$ with $\psi \geq 0$, there exist $\hat{R} > 0$ and two positive functions $\underline{C}, \overline{C} \in C_c^1(0, T)$ such that*

$$\underline{C}(t) \frac{e^{-(N-1)r(x_0, x)}}{(r(x_0, x))^{1-s}} \leq (-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) \leq \overline{C}(t) \frac{e^{-(N-1)r(x_0, x)}}{(r(x_0, x))^{1-s}} \quad \text{for } r(x_0, x) \geq \hat{R}. \quad (\text{A.10})$$

Furthermore, for all $R > 0$, there exists a positive function $\overline{D} \in C_c^1(0, T)$ such that

$$(-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) \leq \overline{D}(t) \quad \text{for } r(x_0, x) \leq R. \quad (\text{A.11})$$

Proof. Proof of (A.10). We first prove the upper bound. Let $g = g(r)$ be as in (A.9) and denote with $K \subset \mathbb{H}^N$ the compact support of ψ with respect to the space variable. Then we can write

$$\begin{aligned} (-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) &= \int_K \mathbb{G}_{\mathbb{H}^N}^s(x, y) \psi(y, t) d\mu_{\mathbb{H}^N}(y) \\ &= g(r(x_0, x)) \int_K \frac{\mathbb{G}_{\mathbb{H}^N}^s(x, y)}{g(r(x_0, x))} \psi(y, t) d\mu_{\mathbb{H}^N}(y). \end{aligned} \quad (\text{A.12})$$

Furthermore, by Corollary A.2 there exist constants \underline{C}_4 and \overline{C}_4 such that there holds

$$\underline{C}_4 g(r(x, y)) \leq \mathbb{G}_{\mathbb{H}^N}^s(x, y) \leq \overline{C}_4 g(r(x, y)) \quad \text{for } r(x, y) > 1.$$

Let $\gamma > 0$ be such that $r(x_0, y) \leq \gamma$ for all $y \in K$ and assume $r(x_0, x) \geq 1 + \gamma$, we have

$$r(x, y) \geq r(x_0, x) - r(x_0, y) > 1 + \gamma - \gamma = 1.$$

Hence,

$$\frac{\mathbb{G}_{\mathbb{H}^N}^s(x, y)}{g(r(x_0, x))} \leq \overline{C}_4 \frac{g(r(x, y))}{g(r(x_0, x))} \quad \text{for } r(x_0, x) \geq 1 + \gamma.$$

Now, using the fact that the function g is decreasing, for $r(x_0, x) \geq 1 + \gamma$ and for all $y \in K$ we have

$$\begin{aligned} \frac{\mathbb{G}_{\mathbb{H}^N}^s(x, y)}{g(r(x_0, x))} &\leq \overline{C}_4 \frac{g(r(x, y))}{g(r(x_0, x))} \leq \overline{C}_4 \frac{g(r(x_0, x) - r(x_0, y))}{g(r(x_0, x))} \\ &= \overline{C}_4 e^{(N-1)r(x_0, y)} \left(\frac{r(x_0, x)}{r(x_0, x) - r(x_0, y)} \right)^{1-s} \end{aligned}$$

$$= \bar{C}_4 e^{(N-1)r(x_0,y)} \frac{1}{\left(1 - \frac{r(x_0,y)}{r(x_0,x)}\right)^{1-s}} \leq \bar{C}_4 \frac{e^{(N-1)r(x_0,y)}}{\left(1 - \frac{r(x_0,y)}{1+\gamma}\right)^{1-s}},$$

where the latter term is well defined since $r(x_0, y) \leq \gamma$ for all $y \in K$. Now, plugging the above estimate into (A.12) we conclude that

$$(-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) \leq g(r(x_0, x)) \underbrace{\int_K \bar{C}_4 \frac{e^{(N-1)r(x_0,y)}}{\left(1 - \frac{r(x_0,y)}{1+\gamma}\right)^{1-s}} \psi(y, t) d\mu_{\mathbb{H}^N}(y)}_{=\bar{C}(t) \in C_c^1(0,T)} \quad \text{for all } r(x_0, x) \geq 1 + \gamma.$$

Next we turn to prove lower the bound in (A.10). For $r(x_0, x) \geq 1 + \gamma$, invoking again Corollary A.2, we have

$$\begin{aligned} \int_K \frac{\mathbb{G}_{\mathbb{H}^N}^s(x, y)}{g(r(x_0, x))} \psi(y, t) d\mu_{\mathbb{H}^N}(y) &\geq \underline{C}_4 \int_K \frac{g(r(x, y))}{g(r(x_0, x))} \psi(y, t) d\mu_{\mathbb{H}^N}(y) \\ &= \underline{C}_4 \int_K e^{(N-1)(r(x_0,x)-r(x,y))} \left(\frac{r(x_0, x)}{r(x, y)}\right)^{1-s} \psi(y, t) d\mu_{\mathbb{H}^N}(y) \\ &\geq \underline{C}_4 \int_K e^{-(N-1)r(x_0,y)} \left(\frac{r(x_0, x)}{r(x, y)}\right)^{1-s} \psi(y, t) d\mu_{\mathbb{H}^N}(y), \end{aligned}$$

where the last estimate follows from

$$r(x_0, x) - r(x, y) \geq -r(x_0, y).$$

Assume now that $r(x_0, x) > 3\gamma$, recalling that $r(x_0, y) \leq \gamma$, we infer

$$r(x, y) \geq r(x_0, x) - r(x_0, y) \geq 2\gamma$$

which yield

$$\left(1 - \frac{r(x_0, y)}{r(x, y)}\right)^{1-s} \geq \frac{1}{2^{1-s}}.$$

Inserting this into (A.12) we finally obtain

$$(-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) \geq g(r(x_0, x)) \underbrace{\int_K \bar{C}_4 \frac{e^{(N-1)r(x_0,y)}}{2^{1-s}} \psi(y, t) d\mu_{\mathbb{H}^N}(y)}_{=\underline{C}(t) \in C_c^1(0,T)} \quad \text{for all } r(x_0, x) > 3\gamma.$$

Summarising, the proof of (A.10) follows by taking $\hat{R} := \max\{1 + \gamma, 3\gamma\}$.

Proof of (A.11). As above, let K be the compact support of ψ with respect to x and let $\gamma > 0$ be such that $r(x_0, y) \leq \gamma$ for all $y \in K$. Let $R > 0$, for all $r(x_0, x) \leq R$ we have $r(x, y) \leq r(x, x_0) + r(x_0, y) \leq R + \gamma$. Hence, by exploiting the estimate (A.8), we infer

$$\begin{aligned} (-\Delta_{\mathbb{H}^N})^{-s} \psi(x, t) &= \int_K \mathbb{G}_{\mathbb{H}^N}^s(x, y) \psi(y, t) d\mu_{\mathbb{H}^N}(y) \\ &\leq \|\psi(x, t)\|_{L^\infty(\mathbb{H}^N)} \int_{B_{R+\gamma}(x)} \frac{\bar{C}_5}{(r(x, y))^{N-2s}} d\mu_{\mathbb{H}^N}(y) \end{aligned}$$

and the proof follows by setting $D(t) := \|\psi(x, t)\|_{L^\infty(\mathbb{H}^N)} \omega_N \int_0^{R+\gamma} \frac{\overline{C}_5}{r^{N-2s}} (\sinh(r))^{N-1} dr$, where ω_N is the volume of the N dimensional unit sphere. \square

Acknowledgments. The second author is partially supported by Project MTM2017-85757-P (Spain), and by the E.U. H2020 MSCA programme, grant agreement 777822. The first and fourth author are partially supported by the PRIN project 201758MTR2 “Direct and inverse problems for partial differential equations: theoretical aspects and applications” (Italy) and they are members of the Gruppo Nazionale per l’Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM). The research of the third author is supported in part by an INSPIRE faculty fellowship (IFA17-MA98).

REFERENCES

- [1] L. Ambrosio, E. Bruè, D. Semola, “Notes on Gradient Flows”, Book in preparation.
- [2] L. Ambrosio, N. Gigli, G. Savare, “Gradient Flows In Metric Spaces and in the Space of Probability Measures”, Lectures in Mathematics. ETH Zürich, 2008, Birkhäuser Basel, ISBN 978-3-7643-8721-1.
- [3] I. Athanasopoulos, L. Caffarelli, *Continuity of the temperature in boundary heat control problems*, Adv. Math. **224** (2010), 293–315.
- [4] C. Bandle, M.d.M. González, M.A. Fontelos, N. Wolanski, *A nonlocal diffusion problem on manifolds*, Comm. Partial Differential Equations **43** (2018), 652–676.
- [5] V. Banica, M.d.M González, M. Sáez, *Some constructions for the fractional Laplacian on noncompact manifolds*, Rev. Mat. Iberoam. **31** (2015), 681-712.
- [6] P. Bénilan, M. G. Crandall, *Regularizing effects of homogeneous evolution equations*, Contributions to Analysis and Geometry (suppl. to Amer. Jour. Math.), Johns Hopkins Univ. Press, Baltimore, Md., 1981. Pp. 23-39.
- [7] P. Bénilan, M. G. Crandall, A. Pazy, “Nonlinear evolution equations in Banach spaces”, Unpublished Book.
- [8] M. Bonforte, A. Figalli, X. Ros-Oton, *Infinite speed of propagation and regularity of solutions to the fractional porous medium equation in general domains*, Comm. Pure Appl. Math. **70** (2017), 1472–1508.
- [9] M. Bonforte, A. Figalli, J. L. Vázquez, *Sharp global estimates for local and nonlocal porous medium-type equations in bounded domains*, Anal. PDE **11** (2018), 945–982.
- [10] M. Bonforte, F. Gazzola, G. Grillo, J. L. Vázquez, *Classification of radial solutions to the Emden-Fowler equation on the hyperbolic space*, Calc. Var. Partial Differential Equations **46** (2013), 375–401.
- [11] M. Bonforte, G. Grillo, J. L. Vázquez, *Fast diffusion flow on manifolds of nonpositive curvature*, J. Evol. Equ. **8** (2008), 99–128.
- [12] M. Bonforte, Y. Sire, J. L. Vázquez, *Existence, uniqueness and asymptotic behaviour for fractional porous medium equations on bounded domains*, Discrete Contin. Dyn. Syst. **35** (2015), 5725–5767.
- [13] M. Bonforte, J. L. Vázquez, *Quantitative local and global a priori estimates for fractional nonlinear diffusion equations*, Adv. Math. **250** (2014), 242–284.
- [14] M. Bonforte, J. L. Vázquez, *A priori estimates for fractional nonlinear degenerate diffusion equations on bounded domains*, Arch. Ration. Mech. Anal. **218** (2015), no. 1, 317–362.
- [15] M. Bonforte, J. L. Vázquez, *Fractional nonlinear degenerate diffusion equations on bounded domains part I. Existence, uniqueness and upper bounds*, Nonlinear Anal. **131** (2016), 363–398.
- [16] H. Brezis. *Monotonicity methods in Hilbert spaces and some applications to nonlinear partial differential equations*, Proc. Symp. Nonlinear Funct. Anal., Madison, Acad. Press (1971), 101–156.
- [17] H. Brezis, “Opérateurs maximaux monotones et semi-groupes de contractions dans les espaces de Hilbert”, North-Holland, 1973.
- [18] L. Caffarelli, L. Silvestre, *An extension problem related to the fractional Laplacian*, Comm. Partial Differential Equations **32** (2007), 1245–1260.
- [19] M. G. Crandall, T.M. Liggett, *Generation of semi-groups of nonlinear transformations on general Banach spaces*, Amer. J. Math. **93** (1971) 265–298.
- [20] M. G. Crandall, M. Pierre, *Regularizing Effectd for $u_t = A\phi(u)$ in L^1* , J. Funct. Anal. **45** (1982) 194–212.
- [21] E. B. Davies, *Heat kernels and spectral theory*, Cambridge Tracts in Mathematics, 92. Cambridge University Press, Cambridge, 1989. x+197 pp.
- [22] A. de Pablo, F. Quirós, A. Rodríguez, J.L. Vázquez, *A fractional porous medium equation*, Adv. Math. **226** (2011), 1378–1409.

- [23] A. de Pablo, F. Quirós, A. Rodríguez, J.L. Vázquez, *A general fractional porous medium equation*, Comm. Pure Appl. Math. **65** (2012), 1242–1284.
- [24] E. Giusti, “Direct methods in the calculus of variations”, World Scientific Publishing Co., Inc., River Edge, NJ, 2003. viii+403 pp. ISBN: 981-238-043-4.
- [25] M.d.M. González, M. Sáez, *Fractional Laplacians and extension problems: the higher rank case*, Trans. Amer. Math. Soc. **370** (2018), 8171–8213.
- [26] A. Grigor’yan, *Analytic and geometric background of recurrence and non-explosion of the Brownian motion on Riemannian manifolds*, Bull. Amer. Math. Soc. (N.S.) **36** (1999), 135–249.
- [27] A. Grigor’yan, *Heat kernels on weighted manifolds and applications*, Cont. Math. **398** (2006), 93–191.
- [28] A. Grigor’yan, “Heat Kernel and Analysis on Manifolds”, AMS/IP Studies in Advanced Mathematics, 47, 2013.
- [29] G. Grillo, K. Ishige, M. Muratori, *Nonlinear characterizations of stochastic completeness*, J. Math. Pures Appl. (9) **139** (2020), 63–82.
- [30] G. Grillo, M. Muratori, *Radial fast diffusion on the hyperbolic space*, Proc. London Math. Soc. **109** (2014), 283–317.
- [31] G. Grillo, M. Muratori, *Smoothing effects for the porous medium equation on Cartan-Hadamard manifolds*, Nonlin. Anal. **131** (2016), 346–362.
- [32] G. Grillo, M. Muratori, F. Punzo, *Weighted fractional porous media equations: existence and uniqueness of weak solutions with measure data*, Calc. Var. Partial Diff. Equ. **54** (2015), 3303–3335.
- [33] G. Grillo, M. Muratori, F. Punzo, *The porous medium equation with measure data on negatively curved Riemannian manifolds*, J. European Math. Soc. **20** (2018), 2769–2812.
- [34] G. Grillo, M. Muratori, F. Punzo, *The porous medium equation with large initial data on negatively curved Riemannian manifolds*, J. Math. Pures Appl. **113** (2018), 195–226.
- [35] G. Grillo, M. Muratori and J. L. Vázquez, *The porous medium equation on Riemannian manifolds with negative curvature: the superquadratic case*, Math. Ann. **373** (2019), 119–153.
- [36] G. Grillo, M. Muratori, J. L. Vázquez, *The porous medium equation on Riemannian manifolds with negative curvature. The large-time behaviour*, Adv. Math. **314** (2017), 328–377.
- [37] Y. Komura, *Nonlinear semi-groups in Hilbert space*, J. Math. Soc. Japan **19** (1967), 493–507.
- [38] N. Roidos, Y. Shao, *The fractional porous medium equation on manifolds with conical singularities*, preprint arXiv:1908.06915.
- [39] N. Roidos, E. Schrohe, *Existence and maximal L^p -regularity of solutions for the porous medium equation on manifolds with conical singularities*, Comm. Partial Differential Equations **41** (2016), 1441–1471.
- [40] N. Roidos, E. Schrohe, *Smoothness and long time existence for solutions of the porous medium equation on manifolds with conical singularities*, Comm. Partial Differential Equations **43** (2018), no. 10, 1456–1484.
- [41] J.L. Vázquez, “The Porous Medium Equation. Mathematical Theory”, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2007.
- [42] J.L. Vázquez, “Smoothing and decay estimates for nonlinear diffusion equations. Equations of porous medium type”, Oxford Lecture Series in Mathematics and its Applications, 33. Oxford University Press, Oxford, 2006.
- [43] J.L. Vázquez, *Barenblatt solutions and asymptotic behaviour for a nonlinear fractional heat equation of porous medium type*, J. Eur. Math. Soc. **16** (2014), 769–803.
- [44] J.L. Vázquez, *Fundamental solution and long time behaviour of the porous medium equation in hyperbolic space*, J. Math. Pures Appl. **104** (2015), 454–484.
- [45] J.L. Vázquez, A. de Pablo, F. Quirós, A. Rodríguez, *Classical solutions and higher regularity for nonlinear fractional diffusion equations*, J. Eur. Math. Soc. (JEMS) **19** (2017), 1949–1975.

DIPARTIMENTO DI SCIENZE MATEMATICHE,
POLITECNICO DI TORINO,
CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY.

E-mail address: `elvise.berchio@polito.it`

DEPARTAMENTO DE MATEMÁTICAS, UNIVERSIDAD AUTÓNOMA DE MADRID, AND
ICMAT - INSTITUTO DE CIENCIAS MATEMÁTICAS, CSIC-UAM-UC3M-UCM
CAMPUS DE CANTOBLANCO, 28049 MADRID, SPAIN

E-mail address: `matteo.bonforte@uam.es`

DEPARTMENT OF MATHEMATICS,
INDIAN INSTITUTE OF TECHNOLOGY DELHI,
IIT CAMPUS, HAUZ KHAS, NEW DELHI
DELHI 110016, INDIA.

E-mail address: `debdipmath@gmail.com`

DIPARTIMENTO DI MATEMATICA,
POLITECNICO DI MILANO,
PIAZZA LEONARDO DA VINCI 32, 20133 MILANO, ITALY.

E-mail address: `gabriele.grillo@polimi.it`