

Lowest order stabilization free Virtual Element Method for the Poisson equation

Original

Lowest order stabilization free Virtual Element Method for the Poisson equation / Berrone, Stefano; Borio, Andrea; Marcon, Francesca. - ELETTRONICO. - (2021).

Availability:

This version is available at: 11583/2881864 since: 2021-04-01T15:27:11Z

Publisher:

Published

DOI:

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Lowest order stabilization free Virtual Element Method for the Poisson equation

Stefano Berrone, Andrea Borio, Francesca Marcon *

Abstract

We introduce and analyse the first order Enlarged Enhancement Virtual Element Method (E²VEM) for the Poisson problem. The method has the interesting property of allowing the definition of bilinear forms that do not require a stabilization term. We provide a proof of well-posedness and optimal order a priori error estimates. Numerical tests on convex and non-convex polygonal meshes confirm the theoretical convergence rates.

1 Introduction

In recent years, the study of polygonal methods for solving partial differential equations has received a huge attention. The main reason for this great interest relies in the flexibility of polygonal meshes to discretize domains with high geometrical complexity. A large number of families of polygonal/polyhedral methods has been developed, among them we can list Discontinuous Galerkin Methods [26, 37, 33], Polygonal Finite Elements (PFEM) [41], Mimetic Finite Difference Methods (MFD) [8, 22, 42], Hybrid High Order Methods (HHO) [27, 28, 29], Gradient Discretisation Methods [31, 30], CutFEM [24], other methods that help in circumventing geometrical complexities are Extended FEMs (XFEM) [35], Generalised FEMs (GFEM) [38, 40, 39] as well as Fictitious Domain Methods [32, 5], Immersed Boundary Methods [36], PDE-constrained Optimization Methods [18, 17, 19] and many others. One of the most recent developments in this field is the family of the Virtual Element Methods (VEM). These methods were first introduced in primal conforming form in [6] and were later on applied to most of

*The authors wish to thank professor Paolo Tilli for his insightful comments. The three authors are members of the Gruppo Nazionale Calcolo Scientifico (GNCS) at Istituto Nazionale di Alta Matematica (INdAM). The authors kindly acknowledge partial financial support by INdAM-GNCS Projects 2020, by the Italian Ministry of Education, University and Research (MIUR) through the “Dipartimenti di Eccellenza” Programme (2018–2022) – Department of Mathematical Sciences “G. L. Lagrange”, Politecnico di Torino (CUP:E11G18000350001) and through the PRIN 2017 project (No. 201744KLJL.004).

the relevant problems of interest in applications, such as advection-diffusion-reaction equations [7, 13, 15], elastic and inelastic problems [9], plate bending problems [23], parabolic and hyperbolic problems [44, 43], simulations in fractured media [14, 12, 11].

Standard VEM discrete bilinear forms are the sum of a singular part maintaining consistency on polynomials and a stabilizing form enforcing coercivity. In the literature, the stabilization term has been extensively studied, for instance in [10], and remains a somehow arbitrarily chosen component of the method with several possible effects on the stability and conditioning of the method. Moreover, the stabilization term causes issues in many theoretical contexts. The first one that we mention is the derivation of a posteriori error estimates [25, 15], where the stabilization term is always at the right-hand side when bounding the error in terms of the error estimator, both from above and from below. Moreover, the isotropic nature of the stabilization term becomes an issue when devising SUPG stabilizations [13, 16], or in the derivation of anisotropic a posteriori error estimates [3]. Finally, other contexts in which the stabilization may induce problems are multigrid analysis [4] and complex non-linear problems [34].

In this work, we introduce a new family of VEM, that we call Enlarged Enhancement Virtual Element Methods (E^2 VEM), designed to avoid the need of the stabilization term. The method is based on the use of higher order polynomial projections in the discrete bilinear form with respect to the standard one [7] and on a modification of the VEM space to allow the computation of such projections. In particular, we extend the enhancement property that is used in the definition of the VEM space ([1], [7]). Indeed, the name of the method comes from this enlarged enhancement property. The degree of polynomial enrichment is chosen locally on each polygon, such that the discrete bilinear form is continuous and coercive, and depends on the number of vertices of the polygon. The resulting discrete functional space has the same set of degrees of freedom of the one defined in [7].

The proof of well-posedness is quite elaborate, thus in this paper we choose to deal only with the lowest order formulation and, for the sake of simplicity, we focus on the two dimensional Poisson's problem with homogeneous Dirichlet boundary conditions, the extension to general boundary conditions being analogous to what is done for classical VEM.

The outline of the paper is as follows. In section 2 we state our model problem. In section 3 we introduce the approximation functional spaces and projection operators and we state the discrete problem. Section 4 contains the discussion about the well-posedness of the discrete problem under suitable sufficient conditions on the local projections. In section 5 we prove optimal order H^1 and L^2 a priori error estimates. Section 6 contains some numerical results assessing the rates of convergence of the method.

Throughout the work, we denote by $(\cdot, \cdot)_\omega$ the standard L^2 scalar product defined on a generic $\omega \subset \mathbb{R}^2$, by $\gamma^{\partial\omega}$ the trace operator, that restricts on the

boundary $\partial\omega$ an element of a space defined over $\omega \subset \mathbb{R}^2$. Inside the proofs, we decide to use a single character C for constants, independent of the mesh size, that appear in the inequalities, which means that we suppose to take at each step the maximum of the constants involved.

2 Model Problem

Let $\Omega \subset \mathbb{R}^2$ be a bounded open set. We are interested in solving the following problem:

$$\begin{cases} -\Delta U = f & \text{in } \Omega, \\ U = 0 & \text{on } \partial\Omega. \end{cases} \quad (1)$$

Defining $a: H_0^1(\Omega) \times H_0^1(\Omega) \rightarrow \mathbb{R}$ such that,

$$a(U, W) := (\nabla U, \nabla W)_\Omega \quad \forall U, W \in H_0^1(\Omega), \quad (2)$$

then, given $f \in L^2(\Omega)$, the variational formulation of (1) is given by: find $U \in H_0^1(\Omega)$ such that,

$$a(U, W) = (f, W)_\Omega \quad \forall W \in H_0^1(\Omega). \quad (3)$$

3 Discrete formulation

In order to define the discrete form of (3), we denote by \mathcal{M}_h a conforming polygonal tessellation of Ω and by E a generic polygon of \mathcal{M}_h . We denote by $\#\mathcal{M}_h$ the number of polygons of \mathcal{M}_h and by h the maximum diameter of all the polygons in \mathcal{M}_h . Let $\{x_i\}_{i=1}^{N_E^V}$ be the N_E^V vertices of E , \mathcal{E}_E the set of its edges and $\mathbf{n}^e = (n_x^e, n_y^e)$ the outward-pointing unit normal vector to the edge e of E . We assume that \mathcal{M}_h satisfies the standard mesh assumptions for VEM (see for instance [10, 21]), i.e. $\exists \kappa > 0$ such that

1. for all $E \in \mathcal{M}_h$, E is star-shaped with respect to a ball of radius $\rho \geq \kappa h_E$, where h_E is the diameter of E ;
2. for all edges $e \subset \partial E$, $|e| \geq \kappa h_E$.

Notice that the above conditions imply that, denoting by N_E^V the number of vertices of E , it holds

$$\exists N_{\max}^V > 0: \forall E \in \mathcal{M}_h, N_E^V \leq N_{\max}^V. \quad (4)$$

For any given $E \in \mathcal{M}_h$, let $\mathbb{P}_k(E)$ be the space of polynomials of degree k defined on E . Let $\Pi_{1,E}^\nabla: H^1(E) \rightarrow \mathbb{P}_1(E)$ be the $H^1(E)$ -orthogonal operator, defined up to a constant by the orthogonality condition: $\forall u \in H^1(E)$,

$$(\nabla(\Pi_{1,E}^\nabla u - u), \nabla p)_E = 0 \quad \forall p \in \mathbb{P}_1(E). \quad (5)$$

In order to define $\Pi_{1,E}^\nabla$ uniquely, we choose any continuous and linear projection operator $P_0 : H^1(E) \rightarrow \mathbb{P}_0(E)$, whose continuity constant in H^1 -norm is independent of h_E , and we impose $\forall u \in H^1(E)$,

$$P_0(\Pi_{1,E}^\nabla u - u) = 0. \quad (6)$$

Remark 1. A suitable choice for P_0 is the integral mean on the boundary of E , i.e.

$$P_0(u) := \frac{1}{|\partial E|} \int_{\partial E} \gamma^{\partial E}(u) ds \quad \forall u \in H^1(E).$$

Notice that this is a common choice, see for instance [7].

For any given $E \in \mathcal{M}_h$, let $l \in \mathbb{N}$ be given, as detailed in the next section. Let $\mathcal{EN}_{1,l}^E$ be the set of functions $v \in H^1(E)$ satisfying

$$(v, p)_E = (\Pi_{1,E}^\nabla v, p)_E \quad \forall p \in \mathbb{P}_{l+1}(E). \quad (7)$$

We define the Enlarged Enhancement Virtual Space of order 1 as

$$\mathcal{V}_{1,l}^E := \{v \in \mathcal{EN}_{1,l}^E : \Delta v \in \mathbb{P}_{l+1}(E), \gamma^e(v) \in \mathbb{P}_1(e) \quad \forall e \in \mathcal{E}_E, v \in C^0(\partial E)\}.$$

We define as degrees of freedom of this space the values of functions at the vertices of E (see [6, 7]).

Moreover, let $\ell \in \mathbb{N}^{\#\mathcal{M}_h}$ be a vector and denote by $\ell(E)$ the element corresponding to the polygon E , we define the global discrete space as

$$\mathcal{V}_{1,\ell} := \{v \in H_0^1(\Omega) : v|_E \in \mathcal{V}_{1,l}^E, \text{ where } l = \ell(E)\}.$$

Note that $v \in \mathcal{V}_{1,\ell}$ is a continuous function that is a polynomial of degree 1 on each edge of the mesh.

To define our discrete bilinear form, let $\Pi_{l,E}^0 \nabla : \mathcal{V}_{1,l}^E \rightarrow [\mathbb{P}_l(E)]^2$ be the $L^2(E)$ -projection operator of the gradient of functions in the Enlarged Enhancement VEM Space, defined, $\forall u \in \mathcal{V}_{1,l}^E$, by the orthogonality condition

$$(\Pi_{l,E}^0 \nabla u, \mathbf{p})_E = (\nabla u, \mathbf{p})_E \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2. \quad (8)$$

Remark 2. The above projection is computable given the degrees of freedom of $u \in \mathcal{V}_{1,l}^E$, applying the Gauss-Green formula and exploiting (7).

Let $a_h^E : \mathcal{V}_{1,l}^E \times \mathcal{V}_{1,l}^E \rightarrow \mathbb{R}$ be defined as

$$a_h^E(u, v) := (\Pi_{l,E}^0 \nabla u, \Pi_{l,E}^0 \nabla v)_E \quad \forall u, v \in \mathcal{V}_{1,l}^E,$$

and $a_h : \mathcal{V}_{1,\ell} \times \mathcal{V}_{1,\ell} \rightarrow \mathbb{R}$ as

$$a_h(u, v) := \sum_{E \in \mathcal{M}_h} a_h^E(u, v) \quad \forall u, v \in \mathcal{V}_{1,\ell}. \quad (9)$$

We can state the discrete problem as: find $u \in \mathcal{V}_{1,\ell}$ such that

$$a_h(u, v) = \sum_{E \in \mathcal{M}_h} (f, \Pi_{0,E}^0 v)_E \quad \forall v \in \mathcal{V}_{1,\ell}, \quad (10)$$

where, $\forall E \in \mathcal{M}_h$, $\Pi_{0,E}^0: \mathcal{V}_{1,l}^E \rightarrow \mathbb{R}$ is the $L^2(E)$ -projection, defined by

$$\Pi_{0,E}^0 v := \frac{1}{|E|} (v, 1)_E \quad \forall v \in \mathcal{V}_{1,\ell}^E.$$

The above projection is computable exploiting (7).

4 Well-posedness

This section is devoted to prove the well-posedness of the discrete problem stated by (10), under suitable sufficient conditions on ℓ . The main result is given by Theorem 1, that induces the existence of an equivalent norm on $\mathcal{V}_{1,\ell}$, which implies the well-posedness of (10).

Theorem 1. *Let $E \in \mathcal{M}_h$, $u \in \mathcal{V}_{1,l}^E$ and $l \in \mathbb{N}$ such that*

$$(l+1)(l+2) \geq N_E^V - 1, \quad (11)$$

then

$$\Pi_{l,E}^0 \nabla u = 0 \implies \nabla u|_E = 0. \quad (12)$$

We omit in the following the proof of the case of triangles ($N_E^V = 3$ and $l = 0$), indeed this case can be led back to classical results. Then, for technical reasons, the proof of Theorem 1 in the case $N_E^V > 3$ is split into two results, described in Section 4.1 and in Section 4.2, respectively. The proof relies on an auxiliary inf-sup condition that is proved by constructing a suitable Fortin operator, whose existence is guaranteed under condition (11).

4.1 Auxiliary inf-sup condition

In this section, after some auxiliary results, we prove through Proposition 1 that (12) is satisfied if the auxiliary inf-sup condition (26) holds true.

Lemma 1. *Let $u \in \mathcal{V}_{1,l}^E$, with $l \geq 1$. Then*

$$\Pi_{l,E}^0 \nabla u = 0 \implies \Pi_{1,E}^\nabla u \in \mathbb{P}_0(E).$$

Proof. Applying (8), we have

$$\Pi_{l,E}^0 \nabla u = 0 \implies (\nabla u, \mathbf{p})_E = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2,$$

that implies

$$(\nabla u, \nabla p)_E = 0 \quad \forall p \in \mathbb{P}_1(E), \quad (13)$$

thanks of the relation $\nabla \mathbb{P}_1(E) \subseteq \nabla \mathbb{P}_{l+1}(E) \subseteq [\mathbb{P}_l(E)]^2$.

Given (13) and (5),

$$\begin{aligned} (\nabla \Pi_{1,E}^\nabla u, \nabla p)_E = 0 \quad \forall p \in \mathbb{P}_1(E) &\implies \nabla \Pi_{1,E}^\nabla u = 0 \\ &\implies \Pi_{1,E}^\nabla u \in \mathbb{P}_0(E). \end{aligned}$$

□

Lemma 2. *Let $u \in \mathcal{V}_{1,l}^E$. If $\Pi_{l,E}^0 \nabla u = 0$, then (7) can be rewritten as*

$$(u, p)_E = P_0(u) \cdot (1, p)_E \quad \forall p \in \mathbb{P}_{l+1}(E), \quad (14)$$

where P_0 is the projection operator chosen in Section 3.

Proof. Applying Lemma 1 and (6),

$$\Pi_{l,E}^0 \nabla u = 0 \implies \Pi_{1,E}^\nabla u = P_0(u).$$

Then, (7) provides (14). □

We now need to introduce some notations and definitions. First, we denote by \mathcal{T}_E the triangulation of E obtained linking each vertex of E to the centre of the ball with respect to which E is star-shaped, denoted by x_C . Let us define the set of internal edges of the triangulation \mathcal{T}_E as $\mathcal{I}_{\mathcal{E}_E}$. For any $i = 1, \dots, N_E^V$, let $\tau_i \in \mathcal{T}_E$ be the triangle whose vertices are x_i, x_{i+1} and x_C . We denote by e_i the edge $\overline{x_C x_i} \in \mathcal{I}_{\mathcal{E}_E}$ and by \mathbf{n}^{e_i} the outward-pointing unit normal vector to the edge e_i of τ_i .

Definition 1. *Let $\mathbf{H}_{\mathcal{T}}^1(E)$ be the broken Sobolev space*

$$\mathbf{H}_{\mathcal{T}}^1(E) := \bigcup_{\tau \in \mathcal{T}_E} \mathbf{H}^1(\tau).$$

Let $u \in \mathbf{H}_{\mathcal{T}}^1(E)$, we define $\forall e_i \in \mathcal{I}_{\mathcal{E}_E}$ the jump function $[[\cdot]]_{e_i} : \mathbf{H}_{\mathcal{T}}^1(E) \rightarrow \mathbf{L}^2(e_i)$ such that

$$[[u]]_{e_i} := \gamma^{e_i}(u|_{\tau_i}) - \gamma^{e_i}(u|_{\tau_{i-1}}).$$

Moreover, $[[u]]_{\mathcal{I}_{\mathcal{E}_E}}$ denotes the vector containing the jumps of u on each $e_i \in \mathcal{I}_{\mathcal{E}_E}$. We endow $\mathbf{H}_{\mathcal{T}}^1(E)$ with the following seminorm and norm : $\forall u \in \mathbf{H}_{\mathcal{T}}^1(E)$,

$$|u|_{\mathbf{H}_{\mathcal{T}}^1(E)}^2 := \sum_{\tau \in \mathcal{T}_E} \|\nabla u\|_{\mathbf{L}^2(\tau)}^2 + \sum_{i=1}^{N_E^V} \|[u]_{e_i}\|_{\mathbf{L}^2(e_i)}^2, \quad (15)$$

$$\|u\|_{\mathbf{H}_{\mathcal{T}}^1(E)}^2 := |u|_{\mathbf{H}_{\mathcal{T}}^1(E)}^2 + \sum_{\tau \in \mathcal{T}_E} \|u\|_{\mathbf{L}^2(\tau)}^2. \quad (16)$$

Definition 2. Let us define $V \subset H_{\mathcal{T}}^1(E)$ given by

$$V := \{v \in H_{\mathcal{T}}^1(E) : \forall e_i \in \mathcal{I}_{\mathcal{E}_E}, \llbracket v \rrbracket_{e_i} \in L^\infty(e_i)\}.$$

Then $\forall v \in V$, we define its seminorm and its norm:

$$\begin{aligned} |v|_V^2 &:= \sum_{\tau \in \mathcal{T}_E} \|\nabla v\|_{[L^2(\tau)]^2}^2 + \left\| \llbracket v \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})}^2, \\ \|v\|_V^2 &:= |v|_V^2 + \sum_{\tau \in \mathcal{T}_E} \|v\|_{L^2(\tau)}^2, \end{aligned}$$

where

$$\left\| \llbracket v \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} := \max_{i \in \{1, \dots, N_E^V\}} \left\| \llbracket v \rrbracket_{e_i} \right\|_{L^\infty(e_i)}.$$

Remark 3. Let us observe that

$$[\mathbb{P}_l(E)]^2 \subset \bigcup_{\tau \in \mathcal{T}_E} [\mathbb{P}_l(\tau)]^2 \subset [V]^2.$$

Hence, we can use $\|\cdot\|_{[V]^2}$ as a norm for $[\mathbb{P}_l(E)]^2$. Notice that, since $[\mathbb{P}_l(E)]^2 \subset [C^0(E)]^2$, $\left\| \llbracket \mathbf{p} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} = 0$, $\forall \mathbf{p} \in [\mathbb{P}_l(E)]^2$.

Definition 3. Let $\mathcal{Q}(\partial E)$ be the vector space

$$\mathcal{Q}(\partial E) := \{q : q \in \mathbb{P}_1(e) \ \forall e \in \mathcal{E}_E, q \in C^0(\partial E), P_0(q) = 0\}. \quad (17)$$

Let $\{\varphi_j\}_{j=1}^{N_E^V-1} \in \mathcal{Q}(\partial E)$ be the set of basis functions of $\mathcal{Q}(\partial E)$ defined such that

$$\varphi_j(x_i) := \begin{cases} 1 & \text{if } i = j \\ c_j : P_0(\varphi_j) = 0 & \text{if } i = j + 1, \quad \forall j = 1, \dots, N_E^V - 1. \\ 0 & \text{otherwise} \end{cases}$$

Definition 4. Let $\mathcal{R}_{\mathcal{Q}}(E)$ be the vector space, lifting of $\mathcal{Q}(\partial E)$ on E , given by:

$$\mathcal{R}_{\mathcal{Q}}(E) := \left\{ \bar{q} \in \mathbb{P}_1(\tau) \ \forall \tau \in \mathcal{T}_E, \gamma^{\partial E}(\bar{q}) \in \mathcal{Q}(\partial E), \bar{q}(x_C) = 0 \right\}. \quad (18)$$

We note that $\mathcal{R}_{\mathcal{Q}}(E) \subset H_{\mathcal{T}}^1(E) \cap C^0(E)$. Hence, we use the norm $\|\cdot\|_{H_{\mathcal{T}}^1(E)}$ defined in (16) as a norm for $\mathcal{R}_{\mathcal{Q}}(E)$. Notice that $\sum_{i=1}^{N_E^V} \left\| \llbracket \bar{q} \rrbracket_{e_i} \right\|_{L^2(e_i)} = 0$ and $\nabla \bar{q} \in [V]^2$, $\forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E)$. We can consider as a basis of $\mathcal{R}_{\mathcal{Q}}(E)$ the set of functions $\{r_j\}_{j=1}^{N_E^V-1} \in \mathcal{R}_{\mathcal{Q}}(E)$:

$$\gamma^{\partial E}(r_j) = \varphi_j, \quad \forall j = 1, \dots, N_E^V - 1. \quad (19)$$

Now, we can introduce the bilinear form b which is crucial for Proposition 1.

Definition 5. Let $b : \mathcal{R}_{\mathcal{Q}}(E) \times [V]^2 \rightarrow \mathbb{R}$, such that $\forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E), \forall \mathbf{v} \in [V]^2$

$$b(\bar{q}, \mathbf{v}) := \int_{\partial E} \bar{q} \mathbf{v} \cdot \mathbf{n}^{\partial E} dx. \quad (20)$$

Applying the divergence theorem, we can rewrite the form b :

$$b(\bar{q}, \mathbf{v}) = \sum_{\tau \in \mathcal{T}_E} \int_{\tau} [\nabla \bar{q} \mathbf{v} + \bar{q} \nabla \cdot \mathbf{v}] dA - \sum_{i=1}^{N_E^V} \int_{e_i} \gamma^{e_i}(\bar{q}) \llbracket \mathbf{v} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i} dx. \quad (21)$$

In order to prove the continuity of b , we have to present a preliminary result.

Lemma 3. Let $\bar{q} \in \mathcal{R}_{\mathcal{Q}}(E)$, we have that $\exists C > 0$, independent of h_E , such that

$$\sum_{i=1}^{N_E^V} \bar{q}(x_i) \leq C \sqrt{\sum_{\tau \in \mathcal{T}_E} \|\nabla \bar{q}\|_{L^2(\tau)}^2}. \quad (22)$$

Proof. By Hölder inequality, we have

$$\sum_{i=1}^{N_E^V} \bar{q}(x_i) \leq \sqrt{N_E^V} \sqrt{\sum_{i=1}^{N_E^V} \bar{q}^2(x_i)}.$$

Moreover, we can apply the property $\exists C > 0$ such that

$$\sqrt{\bar{q}^2(x_i) + \bar{q}^2(x_{i+1})} \leq C \|\nabla \bar{q}\|_{L^2(\tau_i)}, \quad (23)$$

which comes from the equivalence of norms on finite dimensional vector spaces. Finally, recalling the mesh assumption (4) we prove (22). Notice that the constant C of (23) does not depend on h_E by a standard scaling argument. \square

The following lemma proves the continuity of the bilinear form b .

Lemma 4. Let b be given by (20). Then b is a bilinear form and, for h_E sufficiently small,

$$\exists C > 0 : b(\bar{q}, \mathbf{v}) \leq C \|\bar{q}\|_{H^1_\tau(E)} \|\mathbf{v}\|_{[V]^2} \quad \forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E), \forall \mathbf{v} \in [V]^2 .$$

Proof. Let $\bar{q} \in \mathcal{R}_{\mathcal{Q}}(E)$ and $\mathbf{v} \in [V]^2$ be given. Starting from (21) and applying the triangular inequality, we have

$$|b(\bar{q}, \mathbf{v})| \leq \left| \sum_{\tau \in \mathcal{T}_E} \int_{\tau} [\nabla \bar{q} \mathbf{v} + \bar{q} \nabla \cdot \mathbf{v}] dA \right| + \left| \sum_{i=1}^{N_E^V} \int_{e_i} \gamma^{e_i}(\bar{q}) [\mathbf{v}]_{e_i} \cdot \mathbf{n}^{e_i} dx \right|. \quad (24)$$

Let us consider separately the two terms involved in the inequality.

The first part can be analysed applying the properties,

$$\begin{aligned} \forall \mathbf{v} \in [V]^2, \quad \|\nabla \cdot \mathbf{v}\|_{[L^2(\tau)]^2}^2 &\leq 2 \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4}^2 \\ \forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E), \quad \sum_{\tau \in \mathcal{T}_E} \|\bar{q}\|_{L^2(\tau)} + \|\nabla \bar{q}\|_{[L^2(\tau)]^2} &\leq \sqrt{2N_E^V} \|\bar{q}\|_{H_{\gamma}^1(E)} \end{aligned}$$

and the mesh assumption (4), as follows

$$\begin{aligned} \left| \sum_{\tau \in \mathcal{T}_E} \int_{\tau} [\nabla \bar{q} \mathbf{v} + \bar{q} \nabla \cdot \mathbf{v}] dA \right| &\leq \sum_{\tau \in \mathcal{T}_E} \|\nabla \bar{q}\|_{[L^2(\tau)]^2} \|\mathbf{v}\|_{[L^2(\tau)]^2} + \\ &\quad + \sum_{\tau \in \mathcal{T}_E} \|\bar{q}\|_{L^2(\tau)} \|\nabla \cdot \mathbf{v}\|_{L^2(\tau)} \\ &\leq \sum_{\tau \in \mathcal{T}_E} \|\nabla \bar{q}\|_{[L^2(\tau)]^2} \left(\|\mathbf{v}\|_{[L^2(\tau)]^2} + \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4} \right) + \\ &\quad + \sum_{\tau \in \mathcal{T}_E} \|\bar{q}\|_{L^2(\tau)} \left(\|\mathbf{v}\|_{[L^2(\tau)]^2} + \sqrt{2} \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4} \right) \\ &\leq C \sum_{\tau \in \mathcal{T}_E} \left(\|\mathbf{v}\|_{[L^2(\tau)]^2} + \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4} \right) \times \\ &\quad \times \left(\|\nabla \bar{q}\|_{L^2(\tau)} + \|\bar{q}\|_{L^2(\tau)} \right) \\ &\leq C \|\bar{q}\|_{H_{\gamma}^1(E)} \sum_{\tau \in \mathcal{T}_E} \left(\|\mathbf{v}\|_{[L^2(\tau)]^2} + \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4} \right). \end{aligned}$$

Moreover, let us consider the second term of (24), computing exactly the term $\|\gamma^{e_i}(\bar{q})\|_{L^2(e_i)}$ and applying the properties

$$\begin{aligned} \forall \mathbf{v} \in [V]^2, \quad \sum_{i=1}^{N_E^V} \|\llbracket \mathbf{v} \rrbracket_{e_i}\|_{L^2(e_i)} &\leq \sqrt{2N_E^V} \sqrt{\sum_{i=1}^{N_E^V} \|\llbracket \mathbf{v} \rrbracket_{e_i}\|_{L^2(e_i)}^2}, \\ \sum_{i=1}^{N_E^V} \|\llbracket \mathbf{v} \rrbracket_{e_i}\|_{L^2(e_i)}^2 &\leq Ch_E \|\llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}}\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})}^2, \end{aligned} \quad (25)$$

we have

$$\begin{aligned}
\left| \sum_{i=1}^{N_E^V} \int_{e_i} \gamma^{e_i}(\bar{q}) \llbracket \mathbf{v} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i} dx \right| &\leq \sum_{i=1}^{N_E^V} \|\gamma^{e_i}(\bar{q})\|_{L^2(e_i)} \|\llbracket \mathbf{v} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i}\|_{L^2(e_i)} \\
&\leq \sum_{i=1}^{N_E^V} \frac{\sqrt{h_{e_i}}}{\sqrt{3}} |\bar{q}(x_i)| \|\llbracket \mathbf{v} \rrbracket_{e_i}\|_{[L^2(e_i)]^2} \\
&\leq Ch_E \|\llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}}\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \sum_{i=1}^{N_E^V} |\bar{q}(x_i)| \\
&\leq Ch_E \|\llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}}\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \|\bar{q}\|_{H^1_\tau(E)},
\end{aligned}$$

where we apply Lemma 3 in the last step.

Finally, substituting into (24), we obtain

$$\begin{aligned}
|b(\bar{q}, \mathbf{v})| &\leq C \|\bar{q}\|_{H^1_\tau(E)} \left(\sum_{\tau \in \mathcal{T}_E} \left(\|\mathbf{v}\|_{[L^2(\tau)]^2} + \|\nabla \mathbf{v}\|_{[L^2(\tau)]^4} \right) + h_E \|\llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}}\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \right) \\
&\leq C \max(1, h_E) \|\bar{q}\|_{H^1_\tau(E)} \|\mathbf{v}\|_{[V]^2}.
\end{aligned}$$

□

The following proposition is the first step towards the proof of Theorem 1.

Proposition 1. *Let $u \in \mathcal{V}_{1,l}^E$ and $N_E^V > 3$, let b the continuous bilinear form defined by (20). If $\exists \beta > 0$, independent of h_E , such that*

$$\forall \bar{q} \in \mathcal{R}_Q(E), \quad \sup_{\mathbf{p} \in [\mathbb{P}_l(E)]^2} \frac{b(\bar{q}, \mathbf{p})}{\|\mathbf{p}\|_{[V]^2}} \geq \beta \|\bar{q}\|_{H^1_\tau(E)}, \quad (26)$$

then (12) holds true.

Proof. Given (8),

$$\Pi_{l,E}^0 \nabla u = 0 \implies (\nabla u, \mathbf{p})_E = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2.$$

Applying Gauss-Green formula, the previous relation becomes

$$(\nabla u, \mathbf{p})_E = \left(\gamma^{\partial E}(u), \mathbf{p} \cdot \mathbf{n}^{\partial E} \right)_{\partial E} - (u, \nabla \cdot \mathbf{p})_E = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2.$$

Since $\nabla \cdot \mathbf{p} \in \mathbb{P}_{l-1}(E)$ we apply (14) and we obtain

$$\left(\gamma^{\partial E}(u), \mathbf{p} \cdot \mathbf{n}^{\partial E} \right)_{\partial E} - P_0(u) \cdot (1, \nabla \cdot \mathbf{p})_E = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2.$$

Then we can apply the divergence theorem and find the relation

$$\left(\gamma^{\partial E}(u) - P_0(u), \mathbf{p} \cdot \mathbf{n}^{\partial E} \right)_{\partial E} = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2. \quad (27)$$

We have $q = \gamma^{\partial E}(u) - P_0(u) \in \mathcal{Q}(\partial E)$ ($\mathcal{Q}(\partial E)$ defined in (17)). Let $\bar{q} \in \mathcal{R}_{\mathcal{Q}}(E)$ be the lifting of q ($\mathcal{R}_{\mathcal{Q}}(E)$ defined in (18)), then the relation (27) is

$$b(\bar{q}, \mathbf{p}) = 0 \quad \forall \mathbf{p} \in [\mathbb{P}_l(E)]^2.$$

Then, since b is a continuous bilinear form, (26) implies $q \equiv 0$. Finally, since $u \in \mathcal{V}_{1,l}^E$, then $u = P_0(u)$. \square

4.2 Proof of the inf-sup condition

In this section we show that (26) holds with β independent of h_E . The proof relies on the technique known as Fortin trick [20], that consists in the following two classical results.

Proposition 2 ([20, Proposition 5.4.2]). *Assume that there exists an operator $\Pi_E : [V]^2 \rightarrow [\mathbb{P}_l(E)]^2$ that satisfies*

$$b(\bar{q}, \Pi_E \mathbf{v} - \mathbf{v}) = 0 \quad \forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E)$$

and assume that there exists a constant $C_{\Pi} > 0$, independent of h_E , such that

$$\|\Pi_E \mathbf{v}\|_{[V]^2} \leq C_{\Pi} \|\mathbf{v}\|_{[V]^2} \quad \forall \mathbf{v} \in [V]^2.$$

Assume moreover that $\exists \eta > 0$, independent of h_E such that

$$\inf_{q \in \mathcal{R}_{\mathcal{Q}}(E)} \sup_{\mathbf{v} \in [V]^2} \frac{b(q, \mathbf{v})}{\|q\|_{\mathbb{H}_T^1(E)} \|\mathbf{v}\|_{[V]^2}} \geq \eta. \quad (28)$$

Then the discrete inf-sup condition (26) is satisfied, with $\beta = \frac{\eta}{C_{\Pi}}$.

Remark 4. *The inf-sup constant β in (26) has to be independent of the mesh size in order to guarantee that the constant in (44), involved in the coercivity of the bilinear form of (10), is independent of the mesh size.*

Remark 5. *The operator Π_E defined in the following is such that the constant C_{Π} depends on N_{\max}^V and on the constant of continuity of P_0 , both are bounded independently of h_E by assumption.*

Proposition 3 ([20, Proposition 5.4.4]). *Let $\Pi_1, \Pi_2 \in \mathcal{L}([V]^2, [\mathbb{P}_l(E)]^2)$ be such that $\exists c_1, c_2 > 0$,*

$$\|\Pi_1 \mathbf{v}\|_{[V]^2} \leq c_1 \|\mathbf{v}\|_{[V]^2} \quad \forall \mathbf{v} \in [V]^2, \quad (29a)$$

$$b(\bar{q}, \Pi_2 \mathbf{v} - \mathbf{v}) = 0 \quad \forall \bar{q} \in \mathcal{R}_{\mathcal{Q}}(E), \forall \mathbf{v} \in [V]^2, \quad (29b)$$

$$\|\Pi_2 (\mathbf{I} - \Pi_1) \mathbf{v}\|_{[V]^2} \leq c_2 \|\mathbf{v}\|_{[V]^2} \quad \forall \mathbf{v} \in [V]^2. \quad (29c)$$

Then, the operator $\Pi_E := \Pi_2 (\mathbf{I} - \Pi_1) + \Pi_1$ satisfies the hypothesis of Proposition 2.

Following the above results, we have to prove (28) and to show the existence of two operators Π_1, Π_2 satisfying (29a), (29b) and (29c). In the following proposition we achieve the first task.

Proposition 4. *Let $b: \mathcal{R}_Q(E) \times [V]^2 \rightarrow \mathbb{R}$ be defined by (20). Then, for h_E sufficiently small, the inf-sup condition (28) holds true.*

Proof. Let $\bar{q} \in \mathcal{R}_Q(E)$ be given. Recall that $\nabla \bar{q} \in [V]^2$. Notice that, since $\nabla \bar{q} \in \bigcup_{\tau \in \mathcal{T}_E} \mathbb{P}_0(\tau)$,

$$\|\nabla \bar{q}\|_{[V]^2}^2 = \|\nabla \bar{q}\|_{[L^2(E)]^2}^2 + \left\| \llbracket \nabla \bar{q} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})}^2.$$

Since $\|\nabla \bar{q}\|_{[V]^2}^2 = 0 \iff \nabla \bar{q} = 0 \iff \bar{q} = 0$, we deduce that, $\forall \bar{q} \in \mathcal{R}_Q(E)$, $\|\nabla \bar{q}\|_{[V]^2}$ is a norm on $\mathcal{R}_Q(E)$. The same holds for $\|\nabla \bar{q}\|_{[L^2(E)]^2}$. Then, by equivalence of norms on a finite dimensional space and standard scaling arguments, we have

$$\exists C > 0: \|\nabla \bar{q}\|_{[L^2(E)]^2} \geq C \|\nabla \bar{q}\|_{[V]^2}. \quad (30)$$

Moreover, using (21), we get

$$b(\bar{q}, \nabla \bar{q}) = \|\nabla \bar{q}\|_{[L^2(E)]^2}^2 - \sum_{i=1}^{N_E^V} \int_{e_i} \gamma^{e_i}(\bar{q}) \llbracket \nabla \bar{q} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i}, \quad (31)$$

and, since $\llbracket \nabla \bar{q} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i} \in \mathbb{P}_0(e_i) \forall e_i \in \mathcal{I}_{\mathcal{E}_E}$ and $\bar{q}(x_C) = 0$, we get

$$\int_{e_i} \gamma^{e_i}(\bar{q}) \llbracket \nabla \bar{q} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i} = (\llbracket \nabla \bar{q} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i}) \int_{e_i} \gamma^{e_i}(\bar{q}) = (\llbracket \nabla \bar{q} \rrbracket_{e_i} \cdot \mathbf{n}^{e_i}) \frac{|e_i|}{2} \bar{q}(x_i).$$

Then, using Lemma 3 and $|\mathbf{n}^{e_i}| = 1 \forall i = 1, \dots, N_E^V$,

$$\begin{aligned} \sum_{e \in \mathcal{I}_{\mathcal{E}_E}} \int_e \gamma^e(\bar{q}) \llbracket \nabla \bar{q} \rrbracket_e \cdot \mathbf{n}^e &\leq \left\| \llbracket \nabla \bar{q} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \sum_{i=1}^{N_E^V} \frac{|e_i|}{2} \bar{q}(x_i) \\ &\leq \frac{h_E}{2} \left\| \llbracket \nabla \bar{q} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \sum_{i=1}^{N_E^V} \bar{q}(x_i) \\ &\leq Ch_E \left\| \llbracket \nabla \bar{q} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \|\nabla \bar{q}\|_{[L^2(E)]^2}. \end{aligned}$$

Then, from (31) we get,

$$b(\bar{q}, \nabla \bar{q}) \geq \|\nabla \bar{q}\|_{[L^2(E)]^2}^2 \left(\|\nabla \bar{q}\|_{[L^2(E)]^2} - Ch_E \left\| \llbracket \nabla \bar{q} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})} \right).$$

Finally, the term in the parentheses can be bounded from below exploiting (30), as follows:

$$\|\nabla\bar{q}\|_{[L^2(E)]^2} - Ch_E \left\| \llbracket \nabla\bar{q} \rrbracket_{\mathcal{I}_{\varepsilon_E}} \right\|_{L^\infty(\mathcal{I}_{\varepsilon_E})} \geq C_*(1 - h_E) \|\nabla\bar{q}\|_{[V]^2} .$$

Choosing $\mathbf{v}^\star = \nabla\bar{q}$, we obtain the thesis $\forall h_E \leq h_0$, for any $h_0 < 1$. \square

Now, let us focus on the operator Π_1 of Proposition 3. This is a best-approximation operator satisfying the Poincaré-type inequality (35). In order to prove it, let us consider the following lemma.

Lemma 5. *Let $P : \mathbf{H}_{\mathcal{T}}^1(E) \rightarrow \mathbb{P}_0(E) \subset \mathbf{H}_{\mathcal{T}}^1(E)$, such that $\forall v \in \mathbf{H}_{\mathcal{T}}^1(E)$,*

$$Pv := \frac{1}{|E|} \int_E v \, dA .$$

Then $\exists C > 0 : \forall v \in \mathbf{H}_{\mathcal{T}}^1(E)$,

$$\|v - Pv\|_{L^2(E)} \leq C |v|_{\mathbf{H}_{\mathcal{T}}^1(E)} , \quad (32)$$

where C depends on h_E .

Proof. By contradiction, suppose

$$\forall C > 0, \exists v \in \mathbf{H}_{\mathcal{T}}^1(E) : \|v - Pv\|_{L^2(E)} > C |v|_{\mathbf{H}_{\mathcal{T}}^1(E)} .$$

Then, it is possible to define a sequence $w_k \in \mathbf{H}_{\mathcal{T}}^1(E)$ such that, $\forall k \in \mathbb{N}$,

$$\|w_k - Pw_k\|_{L^2(E)} > k |w_k|_{\mathbf{H}_{\mathcal{T}}^1(E)} \quad \|w_k - Pw_k\|_{L^2(E)} = 1 ,$$

which means that

$$|w_k|_{\mathbf{H}_{\mathcal{T}}^1(E)} < \frac{1}{k} \Rightarrow |w_k|_{\mathbf{H}_{\mathcal{T}}^1(E)} \rightarrow 0 . \quad (33)$$

If we define $u_k = w_k - Pw_k$, we have, since Pw_k is constant,

$$|u_k|_{\mathbf{H}_{\mathcal{T}}^1(E)} = |w_k|_{\mathbf{H}_{\mathcal{T}}^1(E)} \rightarrow 0 . \quad (34)$$

The sequence u_k is bounded in $\mathbf{H}_{\mathcal{T}}^1(E)$ by (33), (34) and by the fact that $\|u_k\|_{L^2(E)} = 1$. Thus, by Banach-Alaoglu theorem, it converges weakly in $\mathbf{H}_{\mathcal{T}}^1(E)$ to a function u^\star up to sub-sequences, i.e.

$$u_{k_j} \xrightarrow{\mathbf{H}_{\mathcal{T}}^1(E)} u^\star .$$

Moreover, $\mathbf{H}_{\mathcal{T}}^1(E)$ is contained in the space of functions of bounded variations on E , thus it is compactly embedded in $L^2(E)$ (see [2, Corollary 3.49]).

Then, u_{k_j} converges to a function u^{**} strongly in $L^2(E)$, up to sub-sequences, and by uniqueness of the limit we have $u^{**} = u^*$. Let $u_{\tilde{k}} = u_{k_{j_l}}$ be the sub-sequence such that

$$u_{\tilde{k}} \xrightarrow{H^1_\gamma(E)} u^*, \quad u_{\tilde{k}} \xrightarrow{L^2(E)} u^*.$$

By (34), $|u^*|_{H^1_\gamma(E)} = 0$, thus u^* is constant on E . Since $\|u^*\|_{L^2(E)} = 1$ then $u^* = |E|^{-\frac{1}{2}}$. This is a contradiction because we have $P(u_{\tilde{k}}) = P(w_{\tilde{k}} - Pw_{\tilde{k}}) = 0 \forall \tilde{k}$, then $P(u^*)$ should be zero by continuity and linearity of P . \square

Proposition 5. Let $\Pi_1 : [V]^2 \rightarrow [\mathbb{P}_l(E)]^2$ be the operator defined $\forall \mathbf{v} \in [V]^2$ by

$$\Pi_1 \mathbf{v} := \begin{pmatrix} \frac{1}{|E|} \int_E v_1 dA \\ \frac{1}{|E|} \int_E v_2 dA \end{pmatrix}.$$

Π_1 satisfies the condition (29a) and the following inequality: $\exists C > 0$ such that $\forall \mathbf{v} \in [V]^2$

$$\|\mathbf{v} - \Pi_1 \mathbf{v}\|_{[L^2(E)]^2} \leq Ch_E |\mathbf{v}|_{[V]^2}. \quad (35)$$

Proof. Since $\Pi_1 \mathbf{v} \in [\mathbb{P}_0(E)]^2$, we have

$$\begin{aligned} \|\Pi_1 \mathbf{v}\|_{[V]^2}^2 &= \|\Pi_1 \mathbf{v}\|_{[L^2(E)]^2}^2 = (\Pi_1 \mathbf{v}, \Pi_1 \mathbf{v})_{[L^2(E)]^2} = (\Pi_1 \mathbf{v}, \mathbf{v})_{[L^2(E)]^2} \\ &\leq \|\Pi_1 \mathbf{v}\|_{[L^2(E)]^2} \|\mathbf{v}\|_{[L^2(E)]^2} \leq \|\Pi_1 \mathbf{v}\|_{[V]^2} \|\mathbf{v}\|_{[V]^2}. \end{aligned}$$

The condition (29a) is satisfied.

Since Π_1 satisfies the hypothesis of Lemma 5, we can apply (32) to each component of $\hat{\mathbf{v}} - \hat{\Pi}_1 \hat{\mathbf{v}}$ and, by standard scaling argument, we get

$$\|\mathbf{v} - \Pi_1 \mathbf{v}\|_{[L^2(E)]^2}^2 \leq Ch_E^2 \left(\|\nabla \mathbf{v}\|_{[L^2(E)]^4}^2 + h_E^{-1} \sum_{i=1}^{N_E^V} \|[\mathbf{v}]_{e_i}\|_{[L^2(e_i)]^2}^2 \right).$$

Finally, applying the property (25), inequality (35) is proved. \square

In the following, assuming (11), we prove the existence of an operator Π_2 satisfying (29b). First, we need some auxiliary results.

Definition 6. Let $\{r_i\}_{i=1}^{N_E^V-1}$ be the basis of $\mathcal{R}_Q(E)$, defined in (19). Let us define the set of linear operators $D_i : [V]^2 \rightarrow \mathbb{R}$ such that $\forall \mathbf{v} \in [V]^2$

$$D_i(\mathbf{v}) := \int_{\partial E \cap \text{supp}(r_i)} (\mathbf{v} \cdot \mathbf{n}^{\partial E}) r_i dx, \quad \forall i = 1, \dots, N_E^V - 1.$$

Lemma 6. *If $(l+1)(l+2) \geq N_E^V - 1$, there exists a set of linearly independent functions $\pi_j \in [\mathbb{P}_l(E)]^2$ defined by*

$$D_i(\pi_j) = \delta_{ij} \quad \forall i, j = 1, \dots, N_E^V - 1. \quad (36)$$

Proof. Let $\pi_j \in [\mathbb{P}_l(E)]^2$. The relations (36) are $N_E^V - 1$ independent conditions because the supports of two basis functions r_{i_1}, r_{i_2} never coincide $\forall i_1 \neq i_2$. Then, since $(l+1)(l+2)$ is the dimension of $[\mathbb{P}_l(E)]^2$, the assumption $(l+1)(l+2) \geq N_E^V - 1$ implies that π_j can satisfy all the conditions. \square

In the following proposition we provide a definition of Π_2 and prove an approximation result that is used in Proposition 7.

Proposition 6. *Under the hypothesis of Theorem 1, let us define $\Pi_2 : [V]^2 \rightarrow [\mathbb{P}_l(E)]^2$ such that $\forall \mathbf{v} \in [V]^2$*

$$\Pi_2 \mathbf{v} := \sum_{i=1}^{N_E^V - 1} D_i(\mathbf{v}) \pi_i,$$

where π_i satisfy (36).

Then Π_2 satisfies (29b) and the property $\exists C > 0 : \forall \mathbf{v} \in [V]^2$

$$\|\Pi_2 \mathbf{v}\|_{[V]^2} \leq C \left((1 + h_E^{-1}) \|\mathbf{v}\|_{[\mathbb{L}^2(E)]^2} + (h_E + 1) |\mathbf{v}|_{[V]^2} \right). \quad (37)$$

Proof. Since

$$\forall \mathbf{v} \in [V]^2, \quad D_i(\Pi_2 \mathbf{v}) = D_i(\mathbf{v}) \quad \forall i = 1, \dots, N_E^V - 1,$$

let us check that Π_2 satisfies (29b), indeed by construction $\forall r_i \in \mathcal{R}_{\mathcal{Q}}(E)$, $i = 1, \dots, N_E^V - 1$, $\forall \mathbf{v} \in [V]^2$:

$$b(r_i, \Pi_2 \mathbf{v} - \mathbf{v}) = \int_{\partial E} r_i (\Pi_2 \mathbf{v} - \mathbf{v}) \cdot n^{\partial E} dx = D_i(\Pi_2 \mathbf{v} - \mathbf{v}) = 0.$$

Furthermore, applying Lemma 4 on the reference polygon \hat{E} , we have

$$D_i(\widehat{\Pi_2 \mathbf{v}}) = \int_{\partial \hat{E} \cap \overline{\text{supp}(\hat{r}_i)}} (\hat{\mathbf{v}} \cdot n^{\partial \hat{E}}) \hat{r}_i d\hat{x} = b(\hat{r}_i, \hat{\mathbf{v}}) \leq C \|\hat{r}_i\|_{\mathbb{H}_7^1(\hat{E})} \|\hat{\mathbf{v}}\|_{[\hat{V}]^2}. \quad (38)$$

Then, we want to prove the continuity of $\widehat{\Pi_2 \mathbf{v}}$, i.e.

$$\begin{aligned} \left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\hat{V}]^2} &\leq \sum_{i=1}^{N_E^V - 1} \left| D_i(\widehat{\Pi_2 \mathbf{v}}) \right| \|\widehat{\pi}_i\|_{[\hat{V}]^2} \\ &\leq (N_E^V - 1) \max_i \left| D_i(\widehat{\Pi_2 \mathbf{v}}) \right| \max_i \|\widehat{\pi}_i\|_{[\hat{V}]^2}. \end{aligned}$$

Since $\widehat{\boldsymbol{\pi}}_i \in [\mathbb{P}_l(\widehat{E})]^2$, $\|\widehat{\boldsymbol{\pi}}_i\|_{[\widehat{V}]^2} \leq C \forall i = 1, \dots, N_E^V$ and applying the mesh assumption (4), we have

$$\left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\widehat{V}]^2} \leq CN_{\max}^V \max_i \left| D_i \left(\widehat{\Pi_2 \mathbf{v}} \right) \right|.$$

Applying (38) we obtain

$$\left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\widehat{V}]^2} \leq C \|\widehat{\mathbf{v}}\|_{[\widehat{V}]^2} \max_i \|\widehat{r}_i\|_{\mathbb{H}_T^1(\widehat{E})}.$$

Finally, since r_i is a piecewise linear polynomial on \widehat{E} we have that $\forall i \|\widehat{r}_i\|_{\mathbb{H}_T^1(\widehat{E})} \leq C$, where C depends on the continuity constant of P_0 , that is bounded independently of h_E by assumption. It results,

$$\left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\widehat{V}]^2} \leq C \|\widehat{\mathbf{v}}\|_{[\widehat{V}]^2}. \quad (39)$$

Then, since $\Pi_2 \mathbf{v} \in C^0(E)$, we have

$$\|\Pi_2 \mathbf{v}\|_{[V]^2}^2 = \|\Pi_2 \mathbf{v}\|_{[L^2(E)]^2}^2 + \sum_{\tau \in \mathcal{T}} \|\nabla \Pi_2 \mathbf{v}\|_{[L^2(\tau)]^2}^2. \quad (40)$$

Applying (39) and a standard scaling argument, we can analyse the second term as follows:

$$\begin{aligned} \sum_{\tau \in \mathcal{T}} \|\nabla \Pi_2 \mathbf{v}\|_{[L^2(\tau)]^2}^2 &= \sum_{\widehat{\tau} \in \widehat{\mathcal{T}}} \left\| \widehat{\nabla \Pi_2 \mathbf{v}} \right\|_{[L^2(\widehat{\tau})]^2}^2 \leq \left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\widehat{V}]^2}^2 \leq C \|\widehat{\mathbf{v}}\|_{[\widehat{V}]^2}^2 \\ &= C \left(h_E^{-2} \|\mathbf{v}\|_{[L^2(E)]^2}^2 + \|\nabla \mathbf{v}\|_{[L_T^2(E)]^4}^2 + \left\| \llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})}^2 \right). \end{aligned} \quad (41)$$

Moreover, applying similar arguments to the term $\|\Pi_2 \mathbf{v}\|_{[L^2(E)]^2}^2$, we have

$$\begin{aligned} \|\Pi_2 \mathbf{v}\|_{[L^2(E)]^2}^2 &= h_E^2 \left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[L^2(\widehat{E})]^2}^2 \leq h_E^2 \left\| \widehat{\Pi_2 \mathbf{v}} \right\|_{[\widehat{V}]^2}^2 \\ &\leq Ch_E^2 \left(h_E^{-2} \|\mathbf{v}\|_{[L^2(E)]^2}^2 + \|\nabla \mathbf{v}\|_{[L_T^2(E)]^4}^2 + \left\| \llbracket \mathbf{v} \rrbracket_{\mathcal{I}_{\mathcal{E}_E}} \right\|_{L^\infty(\mathcal{I}_{\mathcal{E}_E})}^2 \right). \end{aligned} \quad (42)$$

Applying (41) and (42) to (40), we prove (37). \square

Finally, we show that the operators Π_1 and Π_2 defined above satisfy (29c).

Proposition 7. *Let $\Pi_1, \Pi_2 \in \mathcal{L}([V]^2, [\mathbb{P}_l(E)]^2)$ be given according to Proposition 5 and 6 respectively, then (29c) is satisfied.*

Proof. Applying (37), we have

$$\|\Pi_2(I - \Pi_1)\mathbf{v}\|_{[V]^2} \leq C \left((1 + h_E^{-1}) \|(I - \Pi_1)\mathbf{v}\|_{[L^2(E)]^2} + (h_E + 1) |(I - \Pi_1)\mathbf{v}|_{[V]^2} \right).$$

Then, applying (35) to the first term and the property

$$\Pi_1\mathbf{v} \in [\mathbb{P}_0(E)]^2 \implies |(I - \Pi_1)\mathbf{v}|_{[V]^2} = |\mathbf{v}|_{[V]^2},$$

to the second one, we have, for h_E sufficiently small,

$$\|\Pi_2(I - \Pi_1)\mathbf{v}\|_{[V]^2} \leq C(1 + h_E) |\mathbf{v}|_{[V]^2} \leq C |\mathbf{v}|_{[V]^2} \leq C \|\mathbf{v}\|_{[V]^2}.$$

□

4.3 Coercivity of the discrete bilinear form

In this section we prove the coercivity of the discrete problem defined by (10) with respect to the standard $H_0^1(\Omega)$ norm, denoted by

$$\|V\|_{H_0^1(\Omega)} = \|\nabla V\|_{[L^2(\Omega)]^2} \quad \forall V \in H_0^1(\Omega).$$

Let

$$\|v\|_{\boldsymbol{\ell}} := \left(\sum_{E \in \mathcal{M}_h} \left\| \Pi_{\boldsymbol{\ell}(E), E}^0 \nabla v \right\|_{[L^2(E)]^2}^2 \right)^{\frac{1}{2}} \quad \forall v \in \mathcal{V}_{1, \boldsymbol{\ell}}.$$

We have the following result.

Proposition 8. *Suppose $\boldsymbol{\ell}$ satisfies (11) $\forall E \in \mathcal{M}_h$. Then, $\|\cdot\|_{\boldsymbol{\ell}}$ is a norm on $\mathcal{V}_{1, \boldsymbol{\ell}}$.*

Proof. Let $v \in \mathcal{V}_{1, \boldsymbol{\ell}}$ be given. It is clear from its definition that $\|v\|_{\boldsymbol{\ell}}$ is a semi-norm. Applying Theorem 1 and since $v \in H_0^1(\Omega)$, we have that

$$\|v\|_{\boldsymbol{\ell}} = 0 \implies \|v\|_{H_0^1(\Omega)} = 0 \implies v = 0.$$

□

Lemma 7. *We have that*

$$\|v\|_{\boldsymbol{\ell}} \leq \|v\|_{H_0^1(\Omega)} \quad \forall v \in \mathcal{V}_{1, \boldsymbol{\ell}}. \quad (43)$$

Moreover, if $\boldsymbol{\ell}(E)$ satisfies (11) $\forall E \in \mathcal{M}_h$, then

$$\exists c_* > 0: \|v\|_{\boldsymbol{\ell}} \geq c_* \|v\|_{H_0^1(\Omega)} \quad \forall v \in \mathcal{V}_{1, \boldsymbol{\ell}}, \quad (44)$$

where c_* does not depend on h .

Proof. The relation (43) follows by the definition of $\Pi_{l,E}^0$ and an application of the Cauchy-Schwarz inequality. Moreover, (44) follows from the equivalence of norms on finite dimensional spaces. By standard scaling arguments, we see that c_* is independent of h . \square

In the following theorem, we provide a proof of the continuity and the coercivity of the discrete bilinear form. The coercivity property follows from Lemma 7.

Theorem 2. *Let a_h be the bilinear form defined by (9). Then,*

$$a_h(w, v) \leq \|w\|_{\mathbb{H}_0^1(\Omega)} \|v\|_{\mathbb{H}_0^1(\Omega)} \quad \forall w, v \in \mathcal{V}_{1,\ell}. \quad (45)$$

Moreover, suppose $\ell(E)$ satisfies (11) $\forall E \in \mathcal{M}_h$. Then,

$$\exists C > 0, \text{ independent of } h: a_h(w, w) \geq C \|w\|_{\mathbb{H}_0^1(\Omega)}^2 \quad \forall w \in \mathcal{V}_{1,\ell}. \quad (46)$$

Proof. Let $w, v \in \mathcal{V}_{1,\ell}$ be given. Applying the Cauchy-Schwarz inequality and (43) we get

$$\begin{aligned} a_h(w, v) &= \sum_{E \in \mathcal{M}_h} \left(\Pi_{\ell(E),E}^0 \nabla w, \Pi_{\ell(E),E}^0 \nabla v \right)_E \\ &\leq \sum_{E \in \mathcal{M}_h} \left\| \Pi_{\ell(E),E}^0 \nabla w \right\|_{[L^2(E)]^2} \left\| \Pi_{\ell(E),E}^0 \nabla v \right\|_{[L^2(E)]^2} \\ &\leq \|w\|_{\ell} \|v\|_{\ell} \leq \|w\|_{\mathbb{H}_0^1(\Omega)} \|v\|_{\mathbb{H}_0^1(\Omega)}. \end{aligned}$$

Moreover, assuming that $\ell(E)$ satisfies (11) $\forall E \in \mathcal{M}_h$, we can apply the lower bound in (44) and get

$$a_h(w, w) = \|w\|_{\ell}^2 \geq (c_*)^2 \|w\|_{\mathbb{H}_0^1(\Omega)}^2.$$

\square

This theorem implies that the bilinear form a_h of the problem (10) satisfies the hypothesis of Lax-Milgram theorem, then the problem admits a unique solution.

5 A priori error estimates

In this section we derive error estimates for the proposed method, in \mathbb{H}_0^1 norm and in the standard L^2 norm. Then, we recall classical results for Virtual Element Methods concerning the interpolation error and the polynomial projection error (see [6, 7]). For each element $E \in \mathcal{M}_h$ we denote by dof_i the operator that maps each sufficiently smooth function U on the i -th local degree of freedom in $\mathcal{V}_{1,\ell}$. According to the definition of the degrees of

freedom, it is true that for every $U \in \mathbf{H}^2(E)$ there exists a unique $U_I \in \mathcal{V}_{1,\ell}$ such that

$$U_I = \sum_{i=1}^{\dim \mathcal{V}_{1,\ell}} \text{dof}_i(U) \xi_i, \quad (47)$$

where ξ_i is the related basis function, defined such that $\text{dof}_j(\xi_i) = \delta_{ij} \quad \forall j = 1, \dots, \dim \mathcal{V}_{1,\ell}$. The following results hold.

Lemma 8 ([6, Proposition 4.3]). *Let U be a smooth enough function, then there exists $C > 0$ such that $\forall h, \exists U_I \in \mathcal{V}_{1,\ell}$, defined as (47), the following relation holds:*

$$\|U - U_I\|_{\mathbf{L}^2(\Omega)} + h \|U - U_I\|_{\mathbf{H}_0^1(\Omega)} \leq Ch^2 |U|_2. \quad (48)$$

Lemma 9 ([7, Lemma 5.1]). *Let U be a smooth enough function, there exist $C_1, C_2 > 0$ such that*

$$\|\Pi_\ell^0 \nabla U - \nabla U\|_{\mathbf{L}^2(\Omega)} \leq C_1 h |U|_2, \quad (49)$$

and

$$\|\Pi_0^0 U - U\|_{\mathbf{L}^2(\Omega)} \leq C_2 h \|U\|_{\mathbf{H}_0^1(\Omega)}. \quad (50)$$

Theorem 3. *Let $U \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$ and $f \in \mathbf{L}^2(\Omega)$ be the solution and the right-hand side of (1), respectively. For h sufficiently small, $\exists C > 0$ such that the unique solution $u \in \mathcal{V}_{1,\ell}$ of problem (10) satisfies the following error estimate:*

$$\|U - u\|_{\mathbf{H}_0^1(\Omega)} \leq Ch \left(|U|_2 + \|f\|_{\mathbf{L}^2(\Omega)} \right). \quad (51)$$

Proof. Let U_I be given by (47). Applying the triangle inequality, we have

$$\|U - u\|_{\mathbf{H}_0^1(\Omega)} \leq \|U - U_I\|_{\mathbf{H}_0^1(\Omega)} + \|U_I - u\|_{\mathbf{H}_0^1(\Omega)}. \quad (52)$$

We deal with the two terms separately. The first one can be bounded applying (48), i.e.

$$\|U - U_I\|_{\mathbf{H}_0^1(\Omega)} \leq Ch |U|_2. \quad (53)$$

On the other hand, in order to deal with the second term of (52) let us denote by $\varepsilon = U_I - u$. First, applying the coercivity of the bilinear form a_h (46) and the discrete problem (10), we have that $\exists C > 0$:

$$\begin{aligned} C \|\varepsilon\|_{\mathbf{H}_0^1(\Omega)}^2 &\leq a_h(\varepsilon, \varepsilon) = a_h(U_I, \varepsilon) - a_h(u, \varepsilon) \\ &= a_h(U_I, \varepsilon) - \sum_{E \in \mathcal{M}_h} (f, \Pi_{0,E}^0 \varepsilon)_E. \end{aligned} \quad (54)$$

Applying the definition of the L^2 projectors and adding and subtracting terms $(\Pi_{l,E}^0 \nabla U, \nabla U)$, we have

$$\begin{aligned}
a_h(\varepsilon, \varepsilon) &= a_h(U_I - U, \varepsilon) + a_h(U, \varepsilon) - \sum_{E \in \mathcal{M}_h} (\Pi_{0,E}^0 f, \varepsilon)_E \\
&= a_h(U_I - U, \varepsilon) + \sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla U - \nabla U, \nabla \varepsilon)_E + (\nabla U, \nabla \varepsilon)_E - (\Pi_{0,E}^0 f, \varepsilon)_E \\
&= a_h(U_I - U, \varepsilon) + \sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla U - \nabla U, \nabla \varepsilon)_E + (f - \Pi_{0,E}^0 f, \varepsilon)_E.
\end{aligned}$$

Let us consider the last three terms separately. The first one can be bounded applying (45) and (48), i.e.

$$a_h(U_I - U, \varepsilon) \leq C \|U_I - U\|_{\mathbf{H}_0^1(\Omega)} \|\varepsilon\|_{\mathbf{H}_0^1(\Omega)} \leq Ch |U|_2 \|\varepsilon\|_{\mathbf{H}_0^1(\Omega)}. \quad (55)$$

Applying the Cauchy-Schwarz inequality and (49), the second term can be bounded as follows:

$$\begin{aligned}
\sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla U - \nabla U, \nabla \varepsilon)_E &\leq \sum_{E \in \mathcal{M}_h} \|\Pi_{l,E}^0 \nabla U - \nabla U\|_{L^2(E)} \|\varepsilon\|_{\mathbf{H}_0^1(E)} \\
&\leq Ch |U|_2 \|\varepsilon\|_{\mathbf{H}_0^1(\Omega)}.
\end{aligned} \quad (56)$$

The last term can be bounded applying the definition of $\Pi_{0,E}^0$, the Cauchy-Schwarz inequality and (50), i.e.

$$\begin{aligned}
\sum_{E \in \mathcal{M}_h} (f - \Pi_{0,E}^0 f, \varepsilon)_E &= \sum_{E \in \mathcal{M}_h} (f, \varepsilon - \Pi_{0,E}^0 \varepsilon)_E \\
&\leq \sum_{E \in \mathcal{M}_h} \|f\|_{L^2(E)} \|\varepsilon - \Pi_{0,E}^0 \varepsilon\|_{L^2(E)} \leq Ch \|f\|_{L^2(\Omega)} \|\varepsilon\|_{\mathbf{H}_0^1(\Omega)}.
\end{aligned} \quad (57)$$

Finally, applying together (55), (56) and (57) into (54) and simplifying, we have

$$\|\varepsilon\|_{\mathbf{H}_0^1(\Omega)} \leq Ch \left(|U|_2 + \|f\|_{L^2(\Omega)} \right). \quad (58)$$

Considering together (53) and (58) we prove (51). \square

Theorem 4. *Let $U \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$ and $f \in \mathbf{H}^1(\Omega)$ be the solution and the right-hand side of (1), respectively. For h sufficiently small, $\exists C > 0$ such that the unique solution $u \in \mathcal{V}_{1,\ell}$ of problem (10) satisfies the following error estimate:*

$$\|U - u\|_{L^2(\Omega)} \leq Ch^2 \left(|U|_2 + \|f\|_{\mathbf{H}_0^1(\Omega)} \right). \quad (59)$$

Proof. Let us define the auxiliary problem: let $\Psi \in \mathbf{H}^2(\Omega) \cap \mathbf{H}_0^1(\Omega)$ the solution of $a(V, \Psi) = (U - u, V)_\Omega \quad \forall V \in \mathbf{H}_0^1(\Omega)$. From the definition of Ψ , we get:

$$\exists C > 0 : \quad |\Psi|_2 \leq C \|U - u\|_{\mathbf{L}^2(\Omega)}, \quad (60)$$

and

$$\exists C > 0 : \quad \|\Psi\|_{\mathbf{H}_0^1(\Omega)} \leq C \|U - u\|_{\mathbf{L}^2(\Omega)}. \quad (61)$$

Let us denote by Ψ_I the interpolant of Ψ according to (47). Applying the auxiliary problem, the discrete problem (10) and the definition of the bilinear form a (2), we have

$$\begin{aligned} \|U - u\|_{\mathbf{L}^2(\Omega)}^2 &= (U - u, U - u)_\Omega = a(U - u, \Psi) \\ &= a(U, \Psi - \Psi_I) + a(U, \Psi_I) - a(u, \Psi) \\ &= a(U, \Psi - \Psi_I) + (f, \Psi_I)_\Omega - a(u, \Psi) \\ &= a(U, \Psi - \Psi_I) + (f, \Psi_I)_\Omega - \left(\sum_{E \in \mathcal{M}_h} (f, \Pi_{0,E}^0 \Psi_I)_E \right) + \\ &\quad + a_h(u, \Psi_I) - a(u, \Psi) \pm a(u, \Psi_I) \\ &= a(U - u, \Psi - \Psi_I) + \left(\sum_{E \in \mathcal{M}_h} (f, \Psi_I - \Pi_{0,E}^0 \Psi_I)_E \right) + \\ &\quad + a_h(u, \Psi_I) - a(u, \Psi_I). \end{aligned} \quad (62)$$

Let us consider the terms of the previous relation separately. First, applying the Cauchy-Schwarz inequality, (48), (50) and (60), we have, for the first term,

$$\begin{aligned} a(U - u, \Psi - \Psi_I) &\leq \|U - u\|_{\mathbf{H}_0^1(\Omega)} \|\Psi - \Psi_I\|_{\mathbf{H}_0^1(\Omega)} \\ &\leq Ch \|U - u\|_{\mathbf{H}_0^1(\Omega)} |\Psi|_2 \leq Ch \|U - u\|_{\mathbf{H}_0^1(\Omega)} \|U - u\|_{\mathbf{L}^2(\Omega)}, \end{aligned} \quad (63)$$

and, for the second one,

$$\begin{aligned} \sum_{E \in \mathcal{M}_h} (f, \Psi_I - \Pi_{0,E}^0 \Psi_I)_E &= \sum_{E \in \mathcal{M}_h} (f - \Pi_{0,E}^0 f, \Psi_I - \Pi_{0,E}^0 \Psi_I)_E \\ &\leq \sum_{E \in \mathcal{M}_h} \|f - \Pi_{0,E}^0 f\|_{\mathbf{L}^2(E)} \|\Psi_I - \Pi_{0,E}^0 \Psi_I\|_{\mathbf{L}^2(E)} \\ &\leq Ch |f|_{\mathbf{H}^1(\Omega)} \sum_{E \in \mathcal{M}_h} \|\Psi_I - \Pi_{0,E}^0 \Psi_I\|_{\mathbf{L}^2(E)}. \end{aligned} \quad (64)$$

Applying the property

$$\forall E \in \mathcal{M}_h, \quad \|\Psi_I - \Pi_{0,E}^0 \Psi_I\|_{\mathbf{L}^2(E)} \leq \|\Psi_I - \Pi_{0,E}^0 \Psi\|_{\mathbf{L}^2(E)},$$

(48) and (50) to (64), we obtain

$$\begin{aligned}
\sum_{E \in \mathcal{M}_h} (f, \Psi_I - \Pi_{0,E}^0 \Psi_I)_E &\leq Ch |f|_{\mathbf{H}^1(\Omega)} \sum_{E \in \mathcal{M}_h} \|\Psi_I - \Pi_{0,E}^0 \Psi\|_{L^2(E)} \\
&\leq Ch |f|_{\mathbf{H}^1(\Omega)} \sum_{E \in \mathcal{M}_h} \left(\|\Psi_I - \Psi\|_{L^2(E)} + \|\Psi - \Pi_{0,E}^0 \Psi\|_{L^2(E)} \right) \\
&\leq Ch |f|_{\mathbf{H}^1(\Omega)} \left(h |\psi|_2 + \|\Psi\|_{\mathbf{H}_0^1(\Omega)} \right). \tag{65}
\end{aligned}$$

We can omit higher order terms and apply (61), obtaining

$$\sum_{E \in \mathcal{M}_h} (f, \Psi_I - \Pi_{0,E}^0 \Psi_I)_E \leq Ch |f|_{\mathbf{H}^1(\Omega)} \|U - u\|_{L^2(\Omega)}. \tag{66}$$

Finally, we have to bound $a_h(u, \Psi_I) - a(u, \Psi_I)$. Then, applying the orthogonality property of $\Pi_{l,E}^0$, adding and subtracting terms, we have

$$\begin{aligned}
a_h(u, \Psi_I) - a(u, \Psi_I) &= \sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla u, \nabla \Psi_I)_E - (\nabla u, \nabla \Psi_I)_E \\
&= \sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla u - \nabla u, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E \\
&= \sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla u - \Pi_{l,E}^0 \nabla U, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E + \\
&\quad + (\Pi_{l,E}^0 \nabla U - \nabla U, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E + \\
&\quad + (\nabla U - \nabla u, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E. \tag{67}
\end{aligned}$$

Notice that, applying (48) and (49), we have the property $\forall E \in \mathcal{M}_h$:

$$\|\nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I\|_{L^2(E)} \leq \|\nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi\|_{L^2(E)} \leq Ch |\Psi|_{2,E}.$$

Therefore, applying the continuity of the projection operator and (60), the first and the last term of (67) can be bounded as

$$\begin{aligned}
\sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla u - \Pi_{l,E}^0 \nabla U, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E + (\nabla U - \nabla u, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E \\
\leq Ch \|U - u\|_{\mathbf{H}_0^1(\Omega)} \|U - u\|_{L^2(\Omega)}. \tag{68}
\end{aligned}$$

Similarly, the second term is bounded as

$$\sum_{E \in \mathcal{M}_h} (\Pi_{l,E}^0 \nabla U - \nabla U, \nabla \Psi_I - \Pi_{l,E}^0 \nabla \Psi_I)_E \leq Ch^2 |U|_2 \|U - u\|_{L^2(\Omega)}. \tag{69}$$

Finally, applying (63),(66),(68) and (69) to (62) and simplifying, we obtain

$$\|U - u\|_{L^2(\Omega)} \leq C \left(h \|U - u\|_{\mathbf{H}_0^1(\Omega)} + h^2 |f|_{\mathbf{H}^1(\Omega)} + h^2 |U|_2 \right).$$

Applying the \mathbf{H}^1 -estimate (Theorem 3) we obtain the relation (59). \square

6 Numerical Results

Let us consider problem (1) on the unit square with homogeneous Dirichlet boundary conditions and the right-hand side defined such that the exact solution is

$$U_{ex} = \sin(2\pi x) \sin(2\pi y).$$

In the following, we show, in log-log scale plots, the convergence curves of the L^2 and H^1 errors that we measure respectively as follows,

$$\begin{aligned} L^2 \text{ error} &= \sqrt{\sum_{E \in \mathcal{M}_h} \left\| \Pi_{1,E}^\nabla u - U_{ex} \right\|_{L^2(E)}^2}, \\ H^1 \text{ error} &= \sqrt{\sum_{E \in \mathcal{M}_h} \left\| \Pi_{l,E}^0 \nabla u - \nabla U_{ex} \right\|_{L^2(E)}^2}, \end{aligned}$$

where u is the discrete solution of (10). Then, for each polygon $E \in \mathcal{M}_h$ we choose l such that the sufficient condition (11) is satisfied (see Table 1 for some choices of l).

Table 1: Sufficient l for polygons that have up to 20 edges.

N_E^V	l
3	0
from 4 to 7	1
from 8 to 13	2
from 14 to 20	3

6.1 Meshes

We consider four sequences of meshes for the convergence test. The first sequence, labeled *Hexagonal*, is a tessellation made by hexagons and triangles, as it is shown in Figure 1a. The second sequence, shown in Figure 1b and labeled *Octagonal*, is made by octagons, squares and triangles. Then, the third sequence, labeled *Hexadecagonal*, is made by hexadecagons and concave pentagons, as it is shown in Figure 1c. Finally, the last sequence, labeled *Star Concave*, is a non-convex tessellation made by octagons and nonagons, as it is shown in Figure 1d.

In each case we start from a mesh of $\#\mathcal{M}_h$ polygons then we refine it, obtaining meshes made by $4\#\mathcal{M}_h$, $16\#\mathcal{M}_h$ and $64\#\mathcal{M}_h$ polygons. The first and the third sequence start with $\#\mathcal{M}_h$ equal to 320, the second and the fourth with $\#\mathcal{M}_h$ equal to 164 and 192 respectively.




Figure 1: Meshes

6.2 Convergence results

For the four mesh sequences, we report the trend of the L^2 and the H^1 errors in Figure 2 and in Figure 3, respectively, decreasing the maximum diameter of the polygons. In the legends, we report the computed convergence rates with respect to h , denoted by α . We see that we get the expected values for all the meshes, as obtained in (51) and (59).

References

- [1] B. Ahmad, A. Alsaedi, F. Brezzi, L. D. Marini, and A. Russo. Equivalent projectors for virtual element methods. *Computers & Mathematics with Applications*, 66:376–391, September 2013.




Figure 2: Logarithmic plot of the L^2 error.




Figure 3: Logarithmic plot of the H^1 error.

- [2] L. Ambrosio, N. Fusco, and D. Pallara. *Functions of bounded variation and free discontinuity problems*. Oxford mathematical monographs. Clarendon, Oxford, 2000.
- [3] P. F. Antonietti, S. Berrone, A. Borio, A. D'Auria, M. Verani, and S. Weisser. Anisotropic a posteriori error estimate for the virtual element method. *IMA Journal of Numerical Analysis*, 02 2021.
- [4] P. F. Antonietti, L. Mascotto, and M. Verani. A multigrid algorithm

for the p -version of the virtual element method. *ESAIM: Mathematical Modelling and Numerical Analysis*, 52:337–364, 03 2017.

- [5] I. Babuška, E. G. Podnos, and G. J. Rodin. New fictitious domain methods: formulation and analysis. *Mathematical Models and Methods in Applied Sciences*, 15(10):1575–1594, 2005.
- [6] L. Beirão da Veiga, F. Brezzi, A. Cangiani, G. Manzini, L. D. Marini, and A. Russo. Basic principles of virtual element methods. *Mathematical Models and Methods in Applied Sciences*, 23(01):199–214, 2013.
- [7] L. Beirão da Veiga, F. Brezzi, L. D. Marini, and A. Russo. Virtual element methods for general second order elliptic problems on polygonal meshes. *Mathematical Models and Methods in Applied Sciences*, 26(04):729–750, 2015.
- [8] L. Beirão da Veiga, K. Lipnikov, and G. Manzini. *The mimetic finite difference method for elliptic problems*. Ms&A Modeling, simulation and applications 11. Springer, Cham, 2014.
- [9] L. Beirão da Veiga, C. Lovadina, and D. Mora. A virtual element method for elastic and inelastic problems on polytope meshes. *Computer Methods in Applied Mechanics and Engineering*, 295:327–346, 2015.
- [10] L. Beirão da Veiga, C. Lovadina, and A. Russo. Stability analysis for the virtual element method. *Mathematical Models and Methods in Applied Sciences*, 27(13):2557–2594, 2017.
- [11] M. F. Benedetto, S. Berrone, and A. Borio. The Virtual Element Method for underground flow simulations in fractured media. In *Advances in Discretization Methods*, volume 12 of *SEMA SIMAI Springer Series*, pages 167–186. Springer International Publishing, Switzerland, 2016.
- [12] M. F. Benedetto, S. Berrone, A. Borio, S. Pieraccini, and S. Scialò. A hybrid mortar virtual element method for discrete fracture network simulations. *J. Comput. Phys.*, 306:148–166, 2016.
- [13] M. F. Benedetto, S. Berrone, A. Borio, S. Pieraccini, and S. Scialò. Order preserving SUPG stabilization for the virtual element formulation of advection-diffusion problems. *Comput. Methods Appl. Mech. Engrg.*, 311:18 – 40, 2016.
- [14] M. F. Benedetto, S. Berrone, and S. Scialò. A globally conforming method for solving flow in discrete fracture networks using the virtual element method. *Finite Elem. Anal. Des.*, 109:23–36, 2016.

- [15] S. Berrone and A. Borio. A residual a posteriori error estimate for the virtual element method. *Mathematical Models and Methods in Applied Sciences*, 27(08):1423–1458, 2017.
- [16] S. Berrone, A. Borio, and G. Manzini. SUPG stabilization for the nonconforming virtual element method for advection–diffusion–reaction equations. *Computer Methods in Applied Mechanics and Engineering*, 340:500 – 529, 2018.
- [17] S. Berrone, S. Pieraccini, and S. Scialò. On simulations of discrete fracture network flows with an optimization-based extended finite element method. *SIAM J. Sci. Comput.*, 35(2):A908–A935, 2013.
- [18] S. Berrone, S. Pieraccini, and S. Scialò. A PDE-constrained optimization formulation for discrete fracture network flows. *SIAM J. Sci. Comput.*, 35(2):B487–B510, 2013.
- [19] S. Berrone, S. Pieraccini, and S. Scialò. An optimization approach for large scale simulations of discrete fracture network flows. *J. Comput. Phys.*, 256:838–853, 2014.
- [20] D. Boffi, F. Brezzi, and M. Fortin. *Approximation of Saddle Point Problems*, chapter 5, pages 265–335. Springer Berlin Heidelberg, Berlin, Heidelberg, 2013.
- [21] Susanne C. Brenner and Li-Yeng Sung. Virtual element methods on meshes with small edges or faces. *Mathematical Models and Methods in Applied Sciences*, 28(07):1291–1336, 2018.
- [22] F. Brezzi, K. Lipnikov, and V. Simoncini. A family of mimetic finite difference methods on polygonal and polyhedral meshes. *Mathematical Models and Methods in Applied Sciences*, 15(10):1533–1551, 2005.
- [23] F. Brezzi and L. D. Marini. Virtual element methods for plate bending problems. *Computer Methods in Applied Mechanics and Engineering*, 253:455–462, 2013.
- [24] E. Burman, S. Claus, P. Hansbo, M. G. Larson, and A. Massing. Cut-FEM: Discretizing geometry and partial differential equations. *International Journal for Numerical Methods in Engineering*, 104(7):472–501, 2015.
- [25] A. Cangiani, E. H. Georgoulis, T. Pryer, and O. J. Sutton. A posteriori error estimates for the virtual element method. *Numerische Mathematik*, 137(4):857–893, Dec 2017.
- [26] D. A. Di Pietro and A. Ern. *Mathematical Aspects of Discontinuous Galerkin Methods*. Springer Berlin Heidelberg, Berlin, Heidelberg, 2012.

- [27] D. A. Di Pietro and A. Ern. A hybrid high-order locking-free method for linear elasticity on general meshes. *Computer Methods in Applied Mechanics and Engineering*, 283:1–21, 2015.
- [28] D. A. Di Pietro and A. Ern. Hybrid high-order methods for variable-diffusion problems on general meshes. *Comptes Rendus Mathématique*, 353(1):31–34, 2015.
- [29] D. A. Di Pietro, A. Ern, and S. Lemaire. An arbitrary-order and compact-stencil discretization of diffusion on general meshes based on local reconstruction operators. *Computational Methods in Applied Mathematics*, 14(4):461–472, 2014.
- [30] J. Droniou, R. Eymard, T. Gallou et, C. Guichard, and R. Herbin. *The Gradient Discretisation Method*. Mathématiques et Applications. Springer, Cham, 2018.
- [31] J. Droniou, R. Eymard, T. Gallou et, and R. Herbin. Gradient Schemes: a generic framework for the discretisation of linear, nonlinear and nonlocal elliptic and parabolic equations. *Mathematical Models and Methods in Applied Sciences*, 23(13):2395–2432, 2013.
- [32] R. Glowinski, T.-W. Pan, and J. Periaux. A fictitious domain method for Dirichlet problem and applications. *Computer Methods in Applied Mechanics and Engineering*, 111(3):283–303, 1994.
- [33] J. S Hesthaven and T. Warburton. *Nodal Discontinuous Galerkin methods: algorithms, analysis, and applications*. Texts in applied mathematics 54. Springer Science & Business Media, New York, 2008.
- [34] B. Hudobivnik, F. Aldakheel, and P. Wriggers. A low order 3D virtual element formulation for finite elasto–plastic deformations. *Computational Mechanics*, 63:253–269, 02 2019.
- [35] N. Moës, J. Dolbow, and T. Belytschko. A finite element method for crack growth without remeshing. *International Journal for Numerical Methods in Engineering*, 46(1):131–150, 1999.
- [36] C. S. Peskin. The immersed boundary method. *Acta Numerica*, 11:479–517, 2002.
- [37] B. Rivière. *Discontinuous Galerkin Methods for Solving Elliptic and Parabolic Equations*. Society for Industrial and Applied Mathematics, 2008.
- [38] T. Strouboulis, I. Babuška, and K. Copps. The design and analysis of the Generalized Finite Element Method. *Computer Methods in Applied Mechanics and Engineering*, 181(1):43–69, 2000.

- [39] T. Strouboulis, K. Copps, and I. Babuška. The generalized finite element method: an example of its implementation and illustration of its performance. *International Journal for Numerical Methods in Engineering*, 47(8):1401–1417, 2000.
- [40] T. Strouboulis, K. Copps, and I. Babuška. The generalized finite element method. *Computer Methods in Applied Mechanics and Engineering*, 190(32):4081–4193, 2001.
- [41] N. Sukumar and A. Tabarraei. Conforming polygonal finite elements. *International Journal for Numerical Methods in Engineering*, 61(12):2045–2066, 2004.
- [42] V. Tishkin, A. A. Samarskii, A. P. Favorskii, and M. Shashkov. Operational finite-difference schemes. *Differential Equations*, 17:854–862, 07 1981.
- [43] G. Vacca. Virtual element methods for hyperbolic problems on polygonal meshes. *Computers & Mathematics with Applications*, 74(5):882 – 898, 2017. SI: SDS2016 – Methods for PDEs.
- [44] G. Vacca and L. Beirão da Veiga. Virtual element methods for parabolic problems on polygonal meshes. *Numerical Methods for Partial Differential Equations*, 31(6):2110–2134, 2015.