



POLITECNICO DI TORINO
Repository ISTITUZIONALE

Additive bases and Niven numbers

Original

Additive bases and Niven numbers / Sanna, Carlo. - In: BULLETIN OF THE AUSTRALIAN MATHEMATICAL SOCIETY.
- ISSN 0004-9727. - STAMPA. - (2021), pp. 1-8. [10.1017/S0004972721000186]

Availability:

This version is available at: 11583/2883056 since: 2021-04-02T18:07:07Z

Publisher:

Cambridge University Press

Published

DOI:10.1017/S0004972721000186

Terms of use:

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

ADDITIVE BASES AND NIVEN NUMBERS

CARLO SANNA[†]

ABSTRACT. Let $g \geq 2$ be an integer. A natural number is said to be a *base- g Niven number* if it is divisible by the sum of its base- g digits. Assuming Hooley's Riemann Hypothesis, we prove that the set of base- g Niven numbers is an additive basis, that is, there exists a positive integer C_g such that every natural number is the sum of at most C_g base- g Niven numbers.

1. INTRODUCTION

One of the principal problems of additive number theory is to determine, given a set of natural numbers \mathcal{A} , if there exists a positive integer k such that every natural number (resp., every sufficiently large natural number) is the sum of at most k elements of \mathcal{A} . In such a case, \mathcal{A} is said to be an *additive basis* (resp., an *asymptotic additive basis*) of order k .

Probably the most famous result of additive number theory is Lagrange's theorem, proved by Lagrange in 1770, which says that the set of perfect squares is an additive basis of order 4. More generally, Waring's problem asks whether the set of perfect h -th powers is an additive basis, which was answered in the affirmative by Hilbert in 1909. Furthermore, in 1937, Vinogradov proved that every sufficiently large odd number is the sum of three prime numbers, which implies that the set of prime numbers is an asymptotic basis of order 4. For an introduction to these classic results see, e.g., Nathanson's book [17].

Let $g \geq 2$ be an integer. Recently, some authors considered additive basis of sets of natural numbers whose base- g representations are restricted in certain ways. For example, Cilleruelo, Luca, and Baxter [4], improving a result of Banks [1], proved that, for $g \geq 5$, the set of natural numbers whose base- g representations are palindrome is an additive basis of order 3. Rajasekaran, Shallit, and Smith [18] showed that the same is true for $g = 3, 4$ but not for $g = 2$; and they proved that the binary palindromes are an additive basis of order 4. Moreover, Madhusudan, Nowotka, Rajasekaran, and Shallit [15] proved that the set of natural numbers whose binary representations consist of two identical repeated blocks is an asymptotic basis of order 4, while Kane, Sanna, and Shallit [14] gave a generalization regarding k repeated blocks. For other results of this kind see also [2, 3].

A natural number is said to be a *base- g Niven number* if it is divisible by the sum of its base- g digits. De Koninck, Doyon, and Kátai [7], and (independently) Mauduit, Pomerance, and Sárközy [16], proved that the number of base- g Niven numbers not exceeding x is asymptotic to $c_g x / \log x$, as $x \rightarrow +\infty$, where $c_g > 0$ is an explicit constant (see [5] for a generalization). Also, De Koninck, Doyon, and Kátai [6, 8] studied gaps between Niven numbers, and runs of consecutive Niven numbers.

Our result is the following:

Theorem 1.1. *Let $g \geq 2$ be an integer. Assuming Hooley's Riemann Hypothesis, we have that the set of base- g Niven numbers is an additive basis.*

Hooley's Riemann Hypothesis for an integer a (HRH(a) for short) states that, for all square-free positive integers m , the Dedekind zeta function Z_K of the number field $K := \mathbb{Q}(\zeta_m, \sqrt[m]{a})$, where ζ_m is a primitive m -th root of unity, satisfies the Riemann hypothesis, that is, if

2010 *Mathematics Subject Classification.* Primary: 11B13, Secondary: 11A63.

Key words and phrases. additive basis; Niven number; sum of digits.

[†]C. Sanna is a member of GNSAGA of INdAM and of CrypTO, the group of Cryptography and Number Theory of Politecnico di Torino.

$Z_K(s) = 0$ for some $s \in \mathbb{C}$, with $0 < \operatorname{Re}(s) < 1$, then $\operatorname{Re}(s) = 1/2$. We assumed $\operatorname{HRH}(g_0)$, where g_0 is an integer depending on g , to use some deep results of Frei, Koymans, and Sofos [12] (Theorem 2.7 and Theorem 2.8 below) concerning sums of three prime numbers with prescribed primitive roots. Except for that, our proof of Theorem 1.1 employs only elementary methods.

Finding an unconditional proof of Theorem 1.1 and determining the order of the additive basis of the set of base- g Niven numbers are two natural problems. We checked that every natural number not exceeding 10^9 is the sum of at most two base-10 Niven numbers.

A related problem stems from considering the multiplicative analog of Niven numbers. A natural number is said to be a *base- g Zuckerman number* if it is divisible by the product of its base- g digits. De Koninck and Luca [9] (see also [10] for the correction of a numerical error in [9]), and Sanna [19] gave upper and lower bounds for the number of base- g Zuckerman numbers not exceeding x . In particular, there are at least $x^{0.122}$, and at most $x^{0.717}$, base-10 Zuckerman numbers not exceeding x , for every sufficiently large x . A question is whether the set of base- g Zuckerman numbers is an additive basis. We checked that every natural number $n \neq 106$ not exceeding 10^9 is the sum of at most four base-10 Zuckerman numbers.

2. PRELIMINARIES

Throughout this section, let $g \geq 2$ be a fixed integer. For every positive integer n , there are uniquely determined $d_1, \dots, d_\ell \in \{0, \dots, g-1\}$, with $d_\ell \neq 0$, such that $n = \sum_{i=1}^\ell d_i g^{i-1}$. We let $[n]_g := d_1, \dots, d_\ell$ (a string), $s_g(n) := \sum_{i=1}^\ell d_i$, and $\ell_g(n) := \ell$. Moreover, for two strings a and b , we write $a \leq b$ if a is a substring of b , and we let $a | b$ be the concatenation of a and b .

We begin with two simple lemmas.

Lemma 2.1. *Let n and s_1, \dots, s_v be positive integers such that $s_g(n) = s_1 + \dots + s_v$ and $s_v > (g-2)(v-1)$. Then there exist positive integers n_1, \dots, n_v such that $[n]_g = [n_1]_g | \dots | [n_v]_g$ and $|s_g(n_i) - s_i| \leq (g-2)(v-1)$ for $i = 1, \dots, v$.*

Proof. If $v = 1$ then the claim follows by picking $n_1 := n$. Hence, assume that $v \geq 2$. Let $n = \sum_{j=1}^\ell d_j g^{j-1}$, where $d_1, \dots, d_\ell \in \{0, \dots, g-1\}$, with $d_\ell \neq 0$. We construct n_1, \dots, n_v in the following way: $n_1 := \sum_{j=1}^{\ell_1} d_j g^{j-1}$, where ℓ_1 is the minimal integer in $[1, \ell]$ such that $\sum_{j=1}^{\ell_1} d_j \geq s_1$; then $n_2 := \sum_{j=\ell_1+1}^{\ell_2} d_j g^{j-\ell_1-1}$, where ℓ_2 is the minimal integer in $(\ell_1, \ell]$ such that $\sum_{j=\ell_1+1}^{\ell_2} d_j \geq s_2$; and so on, up to $n_{v-1} := \sum_{j=\ell_{v-2}+1}^{\ell_{v-1}} d_j g^{j-\ell_{v-2}-1}$, where ℓ_{v-1} is the minimal integer in $(\ell_{v-2}, \ell]$ such that $\sum_{j=\ell_{v-2}+1}^{\ell_{v-1}} d_j \geq s_{v-1}$; Finally, $n_v := \sum_{j=\ell_{v-1}+1}^\ell d_j g^{j-\ell_{v-1}-1}$. From this construction, it follows that $s_i \leq s_g(n_i) \leq s_i + g - 2$ and, by induction, that

$$(1) \quad \sum_{j=i+1}^v s_j - (g-2)i \leq \sum_{j=\ell_i+1}^\ell d_j \leq \sum_{j=i+1}^v s_j,$$

for $i = 1, \dots, v-1$. In fact, the first inequality in (1) and $s_v \geq (g-2)(v-2)$ ensure that each $\ell_1, \dots, \ell_{v-1}$ is well defined. Moreover, (1) with $i = v-1$ yields $s_v - (g-2)(v-1) \leq s_g(n_v) \leq s_v$. In particular, $n_v > 0$ since $s_v > (g-2)(v-1)$. Lastly, by the minimality of each ℓ_i , it follows that $d_{\ell_i} \neq 0$, so that $[n_i]_g = d_{\ell_{i-1}+1}, \dots, d_{\ell_i}$, for $i = 1, \dots, v$, where $\ell_0 := 0$ and $\ell_v := \ell$. Consequently, $[n]_g = [n_1]_g | \dots | [n_v]_g$ and the proof is complete. \square

Lemma 2.2. *Let n and n_1, \dots, n_v be positive integers such that $[n]_g = [n_1]_g | \dots | [n_v]_g$ and n_i is the sum of t_i base- g Niven numbers for $i = 1, \dots, v$. Then n is the sum of $t_1 + \dots + t_v$ base- g Niven numbers.*

Proof. The claim follows easily after noticing that if m is a base- g Niven number then $g^i m$ is a base- g Niven number for every integer $i \geq 0$. \square

The next theorem is a result of additive combinatorics due to Dias da Silva and Hamidoune [11]. For every integer $h \geq 1$ and every subset \mathcal{A} of an additive abelian group, let $h \wedge \mathcal{A}$ denote the set of the sums of h pairwise distinct elements of \mathcal{A} , that is, $h \wedge \mathcal{A} := \{\sum_{a \in \mathcal{A}'} a : \mathcal{A}' \subseteq \mathcal{A}, |\mathcal{A}'| = h\}$.

Theorem 2.3. *Let h be a positive integer, let p be a prime number, and let $\mathcal{A} \subseteq \mathbb{F}_p$. Then*

$$|h^\wedge \mathcal{A}| \geq \min\{p, h|\mathcal{A}| - h^2 + 1\}.$$

In particular, if $|\mathcal{A}| \geq \frac{1}{h}(p-1) + h$ then $h^\wedge \mathcal{A} = \mathbb{F}_p$.

The next lemma shows that every positive integer whose sum of digits satisfies certain properties can be written as the sum of a bounded number of Niven numbers. Hereafter, write $g = g_0^{2^u}$, where $g_0 \geq 2$ is a nonsquare integer and $u \geq 0$ is an integer.

Lemma 2.4. *If n is a positive integer such that:*

- (i) $s_g(n) = p + h$ for a prime number p and an integer $h \in [4g, 8g]$;
- (ii) g_0 is a primitive root modulo p ; and
- (iii) $s_g(n) > \max\{\frac{g-1}{3} \ell_g(n), 140g^3\}$;

then n is the sum of at most $8g + 1$ base- g Niven numbers.

Proof. Put $\ell := \ell_g(n)$, $s := s_g(n)$, and write $n = \sum_{i=0}^{\ell-1} d_i g^i$, where $d_0, \dots, d_{\ell-1} \in \{0, \dots, g-1\}$. Also, let $\mathcal{I} := \{i \in \{0, \dots, \ell-1\} : d_i \neq 0\}$ and $\mathcal{I}' := \{i \bmod t : i \in \mathcal{I}\}$, where t is the multiplicative order of g modulo p . By (ii) and recalling that $g = g_0^{2^u}$, we get that $t \geq (p-1)/2^u > (p-1)/g$. Hence,

$$|\mathcal{I}| = \sum_{i' \in \mathcal{I}'} |\{i \in \mathcal{I} : i \equiv i' \pmod{t}\}| < |\mathcal{I}'| \left(\frac{\ell}{t} + 1\right) < |\mathcal{I}'| \left(\frac{g\ell}{p-1} + 1\right).$$

Since g modulo p has order t , letting $\mathcal{A} := \{g^i \bmod p : i \in \mathcal{I}\} \subseteq \mathbb{F}_p$ we have

$$\begin{aligned} |\mathcal{A}| &= |\mathcal{I}'| > \frac{|\mathcal{I}|}{\frac{g\ell}{p-1} + 1} \geq \frac{\frac{s}{g-1}}{\frac{g\ell}{p-1} + 1} > \frac{\frac{s}{g-1}}{\frac{3gs}{(g-1)(p-1)} + 1} \\ &= \frac{(p-1)s}{3gs + (g-1)(p-1)} > \frac{p-1}{4g-1} > \frac{p-1}{4g} + 8g \geq \frac{p-1}{h} + h, \end{aligned}$$

where we used the inequalities $|\mathcal{I}| \geq \frac{s}{g-1}$, $\ell < \frac{3}{g-1}s$, $s > p-1 > 128g^3$; of which the last three are consequences of (i) and (iii). Hence, Theorem 2.3 yields that $h^\wedge \mathcal{A} = \mathbb{F}_p$. In particular, there exists $\mathcal{J} \subseteq \mathcal{I}$ such that $|\mathcal{J}| = h$ and $\sum_{i \in \mathcal{J}} g^i \equiv n \pmod{p}$. As a consequence, letting $m := n - \sum_{i \in \mathcal{J}} g^i$, it follows easily that $s_g(m) = s - h = p$ and $m \equiv 0 \pmod{p}$, so that m is a base- g Niven number. Thus $n = m + \sum_{i \in \mathcal{J}} g^i$ is the sum of $h + 1$ base- g Niven numbers and, recalling that $h \leq 8g$, the proof is complete. \square

For all integers $q > 0$ and r , let $\mathcal{S}_{q,r}$ be the set of of positive integers n such that:

- (S1) $s_g(n) \equiv r \pmod{q}$;
- (S2) for all positive integers m such that $[m]_g \leq [n]_g$ and $\ell_g(m) \geq 36 \log \ell_g(n)$, we have $s_g(m) > \frac{q-1}{3} \ell_g(m)$.

Recall that the *lower asymptotic density* of a set of positive integers \mathcal{A} is defined as the limit infimum of $|\mathcal{A} \cap [1, x]|/x$, as $x \rightarrow +\infty$.

Lemma 2.5. *Let $q > 0$ and r be integers. Then $\mathcal{S}_{q,r}$ has positive lower asymptotic density.*

Proof. Let $\ell > q$ be an integer and let n be a uniformly distributed random integer in $\{0, \dots, g^\ell - 1\}$. Then $n = \sum_{i=1}^{\ell} d_i g^{i-1}$, where d_1, \dots, d_ℓ are independent uniformly distributed

random variables in $\{0, \dots, g-1\}$. On the one hand, we have

$$\begin{aligned} P_1 &:= \Pr \left[\ell_g(n) = \ell \text{ and } s_g(n) \equiv r \pmod{q} \right] = \Pr \left[d_\ell \neq 0 \text{ and } \sum_{i=1}^{\ell} d_i \equiv r \pmod{q} \right] \\ &= \sum_{s=0}^{q-1} \Pr \left[d_\ell \neq 0 \text{ and } \sum_{i=\ell-q+1}^{\ell} d_i \equiv r-s \pmod{q} \right] \cdot \Pr \left[\sum_{i=1}^{\ell-q} d_i \equiv s \pmod{q} \right] \\ &\geq \frac{1}{g^q} \sum_{s=0}^{q-1} \Pr \left[\sum_{i=1}^{\ell-q} d_i \equiv s \pmod{q} \right] \geq \frac{1}{g^q}. \end{aligned}$$

On the other hand, by Hoeffding's inequality [13, Theorem 2], we have

$$\Pr \left[\sum_{j=1}^k d_{i+j} \leq \frac{g-1}{3}k \right] \leq e^{-k/18},$$

for all $k \in \{1, \dots, \ell\}$ and $i \in \{0, \dots, \ell-k\}$. Hence, letting $y := 36 \log \ell$, we get

$$\begin{aligned} P_2 &:= \Pr \left[\exists m \in \mathbb{N} \text{ s.t. } [m]_g \leq [n]_g, \ell_g(m) \geq y, s_g(m) \leq \frac{g-1}{3}\ell_g(m) \right] \\ &\leq \sum_{\substack{y \leq k \leq \ell \\ i \in \{0, \dots, \ell-k\}}} \Pr \left[\sum_{j=1}^k d_{i+j} \leq \frac{g-1}{3}k \right] \leq \ell \sum_{k \geq y} e^{-k/18} \leq \frac{\ell e^{-y/18}}{1 - e^{-1/18}} < \frac{20}{\ell} \rightarrow 0, \end{aligned}$$

as $\ell \rightarrow +\infty$. Therefore, for every sufficiently large $x > 0$, letting ℓ be the greatest integer such that $g^\ell \leq x$, we obtain

$$\frac{|\mathcal{S}_{q,r} \cap [1, x]|}{x} > \frac{|\mathcal{S}_{q,r} \cap [1, g^\ell]|}{g^{\ell+1}} \geq \frac{P_1 - P_2}{g} > \frac{1}{2g^{q+1}}.$$

Hence, $\mathcal{S}_{q,r}$ has positive lower asymptotic density. \square

The next result is an easy consequence of an important theorem of Schnirelmann.

Theorem 2.6. *Let \mathcal{A} be a set of positive integers such that $1 \in \mathcal{A}$. If \mathcal{A} has positive lower asymptotic density, then \mathcal{A} is an additive basis.*

Proof. Since $1 \in \mathcal{A}$ and $\liminf_{n \rightarrow +\infty} \frac{|\mathcal{A} \cap [1, n]|}{n} > 0$, it follows that $\inf_{n \geq 1} \frac{|\mathcal{A} \cap [1, n]|}{n} > 0$, that is, \mathcal{A} has positive *Schnirelmann density*. Consequently, by Schnirelmann's Theorem [17, Theorem 7.7], it follows that \mathcal{A} is an additive basis. (Note that in [17] they say that \mathcal{A} is a *basis of finite order* if there exists a positive integer k such that every natural number is the sum of *exactly* k elements of \mathcal{A} , and that [17, Theorem 7.7] has to be applied to $\mathcal{A} \cup \{0\}$.) \square

The following deep results of Frei, Koymans, and Sofos [12, Theorem 1.1 and Theorem 1.7] are crucial to the proof of Theorem 1.1.

Theorem 2.7. *Let $\mathbf{a} = (a_1, a_2, a_3) \in \mathbb{Z}^3$ such that no a_i is -1 or a square. Assuming $\text{HRH}(a_i)$ for $i = 1, 2, 3$, we have*

$$\sum_{\substack{n = p_1 + p_2 + p_3 \\ a_i \text{ primitive root mod } p_i, \\ \text{for } i = 1, 2, 3.}} \prod_{i=1}^3 \log p_i \sim \mathcal{A}_{\mathbf{a}}(n) n^2 \quad \text{as } n \rightarrow +\infty,$$

with an explicit factor $\mathcal{A}_{\mathbf{a}}(n) \geq 0$ that satisfies $\mathcal{A}_{\mathbf{a}}(n) \geq A_{\mathbf{a}}$, whenever $\mathcal{A}_{\mathbf{a}}(n) > 0$, where $A_{\mathbf{a}} > 0$ is a constant depending only on \mathbf{a} .

Theorem 2.8. *Let $a \neq -1$ be a nonsquare integer. Then $\mathcal{A}_{(a, a, a)}(n) > 0$ if and only if $(n \bmod 420) \in \mathcal{R}_a$, where \mathcal{R}_a is a nonempty set of residues modulo 420 depending only on a .*

As a consequence, we obtain the following:

Corollary 2.9. *Let $a \neq -1$ be a nonsquare integer and let $\delta \in (0, 1)$. Assuming $\text{HRH}(a)$, for every sufficiently large natural number n such that $(n \bmod 420) \in \mathcal{R}_a$ there exist prime numbers $p_1, p_2, p_3 > n^\delta$ such that $n = p_1 + p_2 + p_3$ and a is a primitive root modulo each of p_1, p_2, p_3 .*

Proof. The claim follows easily from Theorem 2.7 and Theorem 2.8 by noticing that

$$\sum_{\substack{n=p_1+p_2+p_3 \\ p_1 \leq n^\delta}} \prod_{i=1}^3 \log p_i \leq \sum_{\substack{p_1 \leq n^\delta \\ p_2 \leq n}} (\log n)^3 \leq n^{\delta+1} (\log n)^3 = o(n^2),$$

as $n \rightarrow +\infty$. □

3. PROOF OF THEOREM 1.1

Let $g \geq 2$ be an integer, write $g = g_0^{2^u}$, where $g_0 \geq 2$ is a nonsquare integer and $u \geq 0$ is an integer, and assume that $\text{HRH}(g_0)$ holds. Put $q := 420$ and $r := r' + 18g$, where r' is any fixed element of \mathcal{R}_{g_0} , and let $\mathcal{A} := \mathcal{S}_{q,r}$ so that, thanks to Lemma 2.5, \mathcal{A} has positive lower asymptotic density. By Theorem 2.6, we have that $\mathcal{A} \cup \{1\}$ is an additive basis.

Now, in order to prove Theorem 1.1, it suffices to show that every sufficiently large (depending only on g) element of \mathcal{A} is the sum of a bounded number (depending only on g) of base- g Niven numbers. Let $n \in \mathcal{A}$ be sufficiently large, and let $\ell := \ell_g(n)$, $s := s_g(n)$, and $s' := s - 18g$.

Clearly, $\ell \rightarrow +\infty$ as $n \rightarrow +\infty$. In particular, $\ell \geq 36 \log \ell$ for every sufficiently large n . Hence, from (S2) with $m = n$, we get that $s > \frac{g-1}{3}\ell$ and $s' > \frac{g-1}{3}\ell - 18g$. Consequently, in what follows, we can assume that ℓ, s, s' are sufficiently large.

By (S1), we have $s' \equiv s - 18g \equiv r - 18g \equiv r' \pmod{q}$, so that $(s' \bmod 420) \in \mathcal{R}_{g_0}$ and s' is sufficiently large. Hence, by Corollary 2.9, there exist prime numbers $p_1, p_2, p_3 > \sqrt{s'}$ such that $s' = p_1 + p_2 + p_3$ and g_0 is a primitive root modulo each of p_1, p_2, p_3 .

As a consequence, $s = s_1 + s_2 + s_3$ where $s_i := p_i + 6g$ for $i = 1, 2, 3$. Hence, by Lemma 2.1, there exist positive integers n_1, n_2, n_3 such that $[n]_g = [n_1]_g \mid [n_2]_g \mid [n_3]_g$ and $|s_g(n_i) - s_i| \leq 2(g-2)$ for $i = 1, 2, 3$. In particular, $s_g(n_i) = p_i + h_i$ for some integer $h_i \in [4g, 8g]$. Note that $[n_i]_g \leq [n]_g$ and

$$\ell_g(n_i) \geq \frac{s_g(n_i)}{g-1} > \frac{p_i}{g-1} > \frac{\sqrt{s'}}{g-1} \geq \frac{\sqrt{\frac{g-1}{3}\ell - 9g}}{g-1} > 36 \log \ell.$$

Therefore, from (S2) it follows that $s_g(n_i) > \frac{g-1}{3}\ell_g(n_i)$.

Thus we have proved that each n_i satisfies the hypotheses of Lemma 2.4, and consequently each n_i is the sum of at most $8g + 1$ base- g Niven numbers. Then, from Lemma 2.2, it follows that n is the sum of at most $24g + 3$ base- g Niven numbers.

The proof is complete.

Remark 3.1. An inspection of the proof of Theorem 1.1, in particular Lemma 2.4, shows that, actually, we proved a stronger result: Assuming $\text{HRH}(g_0)$, the union of $\{g^i : i = 0, 1, \dots\}$ and the set of base- g Niven numbers m such that $p = s_g(m)$ is a prime number and g_0 is a primitive root modulo p is an additive basis.

4. ACKNOWLEDGEMENTS

The computational resources were provided by `hpc@polito` (<http://www.hpc.polito.it>). The author thanks Daniele Mastrostefano (University of Warwick) for suggestions that improved the paper.

REFERENCES

- [1] W. D. Banks, *Every natural number is the sum of forty-nine palindromes*, *Integers* **16** (2016), Paper No. A3, 9.
- [2] J. Bell, K. Hare, and J. Shallit, *When is an automatic set an additive basis?*, *Proc. Amer. Math. Soc. Ser. B* **5** (2018), 50–63.
- [3] J. P. Bell, T. F. Lidbetter, and J. Shallit, *Additive number theory via approximation by regular languages*, *Developments in language theory, Lecture Notes in Comput. Sci.*, vol. 11088, Springer, Cham, 2018, pp. 121–132.
- [4] J. Cilleruelo, F. Luca, and L. Baxter, *Every positive integer is a sum of three palindromes*, *Math. Comp.* **87** (2018), no. 314, 3023–3055.
- [5] R. Daileda, J. Jou, R. Lemke-Oliver, E. Rossolimo, and E. Treviño, *On the counting function for the generalized Niven numbers*, *J. Théor. Nombres Bordeaux* **21** (2009), no. 3, 503–515.
- [6] J.-M. De Koninck and N. Doyon, *Large and small gaps between consecutive Niven numbers*, *J. Integer Seq.* **6** (2003), no. 2, Article 03.2.5, 8.
- [7] J.-M. De Koninck, N. Doyon, and I. Kátai, *On the counting function for the Niven numbers*, *Acta Arith.* **106** (2003), no. 3, 265–275.
- [8] J.-M. De Koninck, N. Doyon, and I. Kátai, *Counting the number of twin Niven numbers*, *Ramanujan J.* **17** (2008), no. 1, 89–105.
- [9] J.-M. De Koninck and F. Luca, *Positive integers divisible by the product of their nonzero digits*, *Port. Math. (N.S.)* **64** (2007), no. 1, 75–85.
- [10] J.-M. De Koninck and F. Luca, *Corrigendum to “Positive integers divisible by the product of their nonzero digits”, portugaliae math. 64 (2007), 1: 75–85 [MR2298113]*, *Port. Math.* **74** (2017), no. 2, 169–170.
- [11] J. A. Dias da Silva and Y. O. Hamidoune, *Cyclic spaces for Grassmann derivatives and additive theory*, *Bull. London Math. Soc.* **26** (1994), no. 2, 140–146.
- [12] C. Frei, P. Koymans, and E. Sofos, *Vinogradov’s three primes theorem with primes having given primitive roots*, *Math. Proc. Cambridge Philos. Soc.* **170** (2021), no. 1, 75–110.
- [13] W. Hoeffding, *Probability inequalities for sums of bounded random variables*, *J. Amer. Statist. Assoc.* **58** (1963), 13–30.
- [14] D. M. Kane, C. Sanna, and J. Shallit, *Waring’s theorem for binary powers*, *Combinatorica* **39** (2019), no. 6, 1335–1350.
- [15] P. Madhusudan, D. Nowotka, A. Rajasekaran, and J. Shallit, *Lagrange’s theorem for binary squares*, 43rd International Symposium on Mathematical Foundations of Computer Science, LIPIcs. Leibniz Int. Proc. Inform., vol. 117, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, 2018, pp. Art. No. 18, 14.
- [16] C. Mauduit, C. Pomerance, and A. Sárközy, *On the distribution in residue classes of integers with a fixed sum of digits*, *Ramanujan J.* **9** (2005), no. 1-2, 45–62.
- [17] M. B. Nathanson, *Additive Number Theory: The Classical Bases*, Graduate Texts in Mathematics, vol. 164, Springer-Verlag, New York, 1996.
- [18] A. Rajasekaran, J. Shallit, and T. Smith, *Additive number theory via automata theory*, *Theory Comput. Syst.* **64** (2020), no. 3, 542–567.
- [19] C. Sanna, *On numbers divisible by the product of their nonzero base b digits*, *Quaest. Math.* **43** (2020), no. 11, 1563–1571.

POLITECNICO DI TORINO, DEPARTMENT OF MATHEMATICAL SCIENCES
 CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY
 Email address: carlo.sanna.dev@gmail.com