

On games with coordinating and anti-coordinating agents

*Original*

On games with coordinating and anti-coordinating agents / Vanelli, Martina; Arditti, Laura; Como, Giacomo; Fagnani, Fabio. - ELETTRONICO. - 53:(2020), pp. 10975-10980. (Intervento presentato al convegno 21st IFAC World Congress tenutosi a Berlin (Germany)) [10.1016/j.ifacol.2020.12.2848].

*Availability:*

This version is available at: 11583/2897494 since: 2021-04-28T19:03:59Z

*Publisher:*

Elsevier

*Published*

DOI:10.1016/j.ifacol.2020.12.2848

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Elsevier postprint/Author's Accepted Manuscript

© 2020. This manuscript version is made available under the CC-BY-NC-ND 4.0 license  
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:  
<http://dx.doi.org/10.1016/j.ifacol.2020.12.2848>

(Article begins on next page)

# On games with coordinating and anti-coordinating agents

Martina Vanelli\* Laura Arditti\* Giacomo Como\*  
 Fabio Fagnani\*

\* *Department of Mathematical Sciences “G.L. Lagrange”, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy*  
 (e-mail:  
 {martina.vanelli,laura.arditti,giacomo.como,fabio.fagnani}@polito.it).

**Abstract:** This work studies Nash equilibria for games where a mixture of coordinating and anti-coordinating agents, with possibly heterogeneous thresholds, coexist and interact through an all-to-all network. Whilst games with only coordinating or only anti-coordinating agents are potential, also in the presence of heterogeneities, this does not hold when both type of agents are simultaneously present. This makes their analysis more difficult and existence of Nash equilibria not guaranteed. Our main result is a checkable condition on the threshold distributions that characterizes the existence of Nash equilibria in such mixed games. When this condition is satisfied an explicit algorithm allows to determine the complete set of such equilibria. Moreover, for the special case when only one type of agents is present (either coordinating or anti-coordinating), our results allow an explicit computation of the cardinality of Nash equilibria. Copyright © 2020 The Authors. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0>)

**Keywords:** Game theory, coordination games, anti-coordination games, multi-agent systems, modelling and decision making in complex systems.

## 1. INTRODUCTION

In games of strategic complements, the best response action of a player is increasing in the action of the other players. Examples of such games include the adoption of a new technology, beliefs or behavioral attitudes in social influence systems, or cooperative interchanges among economical actors. In games of strategic substitutes just the opposite happens, the best response action of a player is decreasing in the action of the other players. Applications of such models include, for example, local public good provision, information gathering, firms interacting in competitive markets.

While these two classes of games have received considerable attention in recent literature (Jackson and Zenou (2015), Bramoullé (2007)), relatively unexplored are mixed models where these two strategic interactions coexist. However, examples of social or economic model where such behaviors coexist are rather frequent, e.g., collecting actions, interacting firms with cooperative and competitive features.

In this paper, we focus on a particular instance of such heterogeneous models. We consider games where actions are all binary  $\{-1, +1\}$  and players are split into two classes: one of coordinating agents and one of anti-coordinating agents. That of coordination and anti-coordination games are two basic examples of games with respectively strate-

gic complements and strategic substitutes. Coordinating agents have an incentive for action 1 when such action is played by a fraction of the remaining players above a certain threshold while anti-coordinating agents have an incentive for action 1 when such action is played by a fraction of the remaining players below a certain threshold. We will refer to such models as mixed coordination/anti-coordination (CAC) games.

Our aim is to study Nash equilibria in mixed CAC games. Pure coordination games and pure anti-coordination games always admit Nash equilibria as they are potential games (Monderer and Shapley (1996)) also when the thresholds are heterogeneous. Instead general mixed CAC games are no longer potential and the existence of Nash equilibria is not guaranteed. Furthermore, even when existence is guaranteed, the set of Nash equilibria is in general unknown. For the special case of coordinating agents with identical thresholds, a characterization of the Nash equilibria was proposed by Morris (2000).

There is a related literature where the best-response dynamics is analyzed, in particular its convergence to Nash equilibria. Ramazi et al. (2016) proved that the asynchronous best-response dynamics of pure coordination games and pure anti-coordination games will almost surely converge to Nash equilibria for every network topology and every set of thresholds. Granovetter (1978) studied the synchronous best-response dynamics of coordinating agents as a linear threshold model. Similar results have been generalized to configuration models by Rossi et al. (2017) and to games with a mixture of coordinating and anti-coordinating agents by Grabisch and Li (2019). Other results concern time of convergence of asynchronous best-

\* Giacomo Como is also with the Department of Automatic Control, Lund University, Sweden. This work was partially supported by MIUR grant Dipartimenti di Eccellenza 2018–2022 [CUP: E11G18000350001], the Swedish Research Council, and by the Compagnia di San Paolo.

response dynamics and its dependence on the network structure (Ellison (1993), Kandori et al. (1993), Montanari and Saberi (2010)).

Our contribution is a throughout analysis on the existence and the structure of pure strategy Nash equilibria for mixed coordination/anti-coordination (CAC) games with possibly heterogeneous activation thresholds. In particular we give the following contributions: (i) We establish a checkable necessary and sufficient condition involving the cumulative distribution functions of the thresholds for the existence of NE of a general mixed CAC game; and (ii) In the special case when only coordinating or anti-coordinating agents are present we classify all NE and determine their cardinality. While existence was well known in the literature, to the best of our knowledge such precise characterization was not yet presented.

The rest of the paper is organized as follows. Section 2 is devoted to the description of the model and Section 3 presents the main results. We conclude with Section 4 containing a summary and further work.

## 2. MODEL DESCRIPTION

In this section, we formally define the mixed CAC game, we explicit the best response function and we make some general observations on the existence of Nash equilibria. In order to give an intuition of the upcoming results, in the end we briefly study the case of an infinite population.

### 2.1 Definition of the game

Consider a finite set of agents  $\mathcal{V} = \{1, \dots, n\}$  partitioned into two classes:  $\mathcal{V}_c \cup \mathcal{V}_a = \mathcal{V}$ ,  $\mathcal{V}_c \cap \mathcal{V}_a = \emptyset$ . Each agent  $i \in \mathcal{V}$  is described by two elements: the type  $\delta_i$  describing the class it belongs to

$$\delta_i := \begin{cases} 1 & \text{if } i \in \mathcal{V}_c \\ -1 & \text{if } i \in \mathcal{V}_a \end{cases}$$

and its personal weight  $d_i \in \mathbb{R}$ .

The action set of all agents is the binary set  $\mathcal{A} := \{-1, +1\}$ . We denote with  $x_i \in \mathcal{A}$  the action chosen by agent  $i$  and with  $x \in \mathcal{X} := \mathcal{A}^{\mathcal{V}}$  the vector assembling the actions of all players (action profile). As usual in game theory,  $x_{-i} \in \mathcal{A}^{\mathcal{V} \setminus \{i\}}$  is the vector obtained from the action profile  $x \in \mathcal{X} := \mathcal{A}^{\mathcal{V}}$  by removing its  $i^{\text{th}}$  entry. The utility of agent  $i$ , when she plays action  $x_i$  and the remaining players  $x_{-i}$ , is given by

$$u_i(x_i, x_{-i}) := \delta_i \left( \sum_{j \neq i} x_i x_j - d_i x_i \right), \quad i \in \mathcal{V} \quad (1)$$

Agents in  $\mathcal{V}_c$  ( $\delta_i = 1$ ) are *coordinating agents* (their utility increases with the number of agents playing their same action), while agents in  $\mathcal{V}_a$  ( $\delta_i = -1$ ) are *anti-coordinating agents* (their utility decreases with the number of agents playing their same action).

The weight  $d_i$  represents the tendency of agent  $i$  in choosing an action over the other. Indeed, the sign of weight  $d_i$ , along with  $\delta_i$ , determines which is the risk dominant action, i.e., which is the best action for agent

$i$  in the absence of any external influences. If  $d_i = 0$  the two actions are called *risk-neutral*.

*Definition.* The *mixed coordination/anti-coordination* (CAC) game is a game with agent set  $\mathcal{V}$ , action configuration space  $\mathcal{X}$  and utilities  $\{u_i\}_{i \in \mathcal{V}} : \mathcal{X} \rightarrow \mathbb{R}$  in (1).

In the special cases when only one type of agents is present, namely  $\mathcal{V}_c = \mathcal{V}$  or  $\mathcal{V}_a = \mathcal{V}$ , we will simply call it, *coordination game* or, respectively, *anti-coordination game*.

### 2.2 Best response and Nash equilibria

In strategic games, agents are assumed to be rational, i.e., they choose their action with the aim of maximizing the utility. Given the actions of the others, the *best response* (BR) function returns the set of the best actions for agent  $i$ , that is, the actions that achieve the highest utility. Formally, the BR is given by

$$\mathcal{B}_i(x_{-i}) = \operatorname{argmax}_{x_i \in \mathcal{A}} u_i(x_i, x_{-i}).$$

For the purpose of our study, it is convenient to express the best response function in terms of the number of opponents playing action +1, that is,

$$n_i^+(x) = |\{j \in \mathcal{V} \setminus \{i\} \mid x_j = +1\}|.$$

From (1) and the fact that  $\sum_{j \neq i} x_j = 2n_i^+(x) - (n-1)$ , it follows that the BR of an agent  $i \in \mathcal{V}$  can be written in the following way:

$$\mathcal{B}_i(x_{-i}) = \begin{cases} \{+1\} & \text{if } \delta_i(n_i^+(x) - r_i(n-1)) > 0 \\ \{-1\} & \text{if } \delta_i(n_i^+(x) - r_i(n-1)) < 0 \\ \{\pm 1\} & \text{if } \delta_i(n_i^+(x) - r_i(n-1)) = 0 \end{cases} \quad (2)$$

where  $r_i := \frac{1}{2} + \frac{d_i}{2(n-1)}$ .

In words, when  $\delta_i = 1$ , the best action is +1 if more than  $r_i(n-i)$  opponents are playing +1, whereas, when  $\delta_i = -1$ , the best action is +1 if less than  $r_i(n-i)$  opponents are playing +1. Hence, the best response changes depending if the fraction of opponents playing action +1, that is,  $n_i^+(x)/(n-1)$ , is above or below the *threshold*  $r_i$ . Agents having  $r_i > 1$  or  $r_i < -1$  are called *stubborn agents*, as their best response function is constantly equal to either  $\{-1\}$  or  $\{+1\}$ , independently of the actions of others.

A *Nash equilibria* is a configuration  $x \in \mathcal{X}$  such that  $x_i \in \mathcal{B}_i(x_{-i})$  for any agent  $i \in \mathcal{V}$ . In the following we will denote by  $\mathcal{N}$  the set of Nash equilibria of the game.

Both the coordination game and the anti-coordination game, for any possible choice of the weights, are potential games (Monderer and Shapley (1996)). This fact is very well known in case of homogeneous thresholds, but actually holds in general. Indeed, if we define the two functions

$$\Phi_c(x) = \frac{1}{2} \sum_{\substack{i,j \in \mathcal{V} \\ i \neq j}} x_i x_j - \sum_{i \in \mathcal{V}} d_i x_i \quad (3)$$

$$\Phi_a(x) = -\Phi_c(x) \quad (4)$$

we can derive from (1) that, depending if the game is a coordination or an anti-coordination one, it respectively holds

$$\begin{aligned} u_i(y_i, x_{-i}) - u_i(x_i, x_{-i}) &= \Phi_c(y_i, x_{-i}) - \Phi_c(x_i, x_{-i}) \\ u_i(y_i, x_{-i}) - u_i(x_i, x_{-i}) &= \Phi_a(y_i, x_{-i}) - \Phi_a(x_i, x_{-i}) \end{aligned} \quad (5)$$

for all  $x_i, y_i \in \mathcal{A}$ ,  $x_{-i} \in \mathcal{A}^{\mathcal{V} \setminus \{i\}}$ .

The potential property guarantees the existence of at least one Nash equilibrium, which is a maximum point of the potential function. If there are no stubborn agents, the two consensus configurations, namely  $\mathbf{1}$  and  $-\mathbf{1}$ , are NE of the coordination game for any set of weights. This is a peculiar property of coordination games and, in general, an explicit characterization of all the NE is hard to find.

It is enough for a coordinating agent to interact with an anti-coordinating agent to lose the potential property.

*Proposition 1.* If  $\mathcal{V}_c \neq \emptyset$  and  $\mathcal{V}_a \neq \emptyset$  the mixed CAC game is *not* a potential game.

Hence, the study of Nash equilibria for the mixed CAC game is a challenging problem and not even the existence of such configuration is guaranteed, as the following simple example shows.

*Example 1.* Let us consider the very simple case of a two-player game with  $\mathcal{V}_c = \{1\}$  and  $\mathcal{V}_a = \{2\}$  where both agents have weights  $d_i = 0$  (Fig. 1). This is known as the *discoordination game* or *matching pennies* game, since it describes the situation where one agent aims at coordinating while the other wants to play the opposite action to that of her opponent. The discoordination game admits no Nash equilibria.




Fig. 1. The discoordination game.

### 2.3 The formalism of distribution functions

We now set some notation that will be used throughout the rest of the paper.

- $n_c = |\mathcal{V}_c|$  and  $n_a = |\mathcal{V}_a|$  are the number of coordinating and anti-coordinating agents respectively, while  $\alpha = n_c/n$  is the fraction of coordinating agents.
- Given a configuration  $x \in \mathcal{X}$ , we denote by  $z(x)$ ,  $z_c(x)$ ,  $z_a(x)$  the fraction of agents that in  $x$  are playing action 1 in, respectively, the total population, in the set of coordinating agents, and in the set of anti-coordinating agents:

$$\begin{aligned} z(x) &:= n^{-1} |\{i \in \mathcal{V} \mid x_i = +1\}|, \\ z_c(x) &:= n_c^{-1} |\{i \in \mathcal{V}_c \mid x_i = +1\}|, \\ z_a(x) &:= n_a^{-1} |\{i \in \mathcal{V}_a \mid x_i = +1\}| \end{aligned} \quad (6)$$

It holds

$$z(x) = \alpha z_c(x) + (1 - \alpha) z_a(x)$$

Furthermore, let

$$F(z) := \frac{1}{n} |\{i \in \mathcal{V} \mid r_i \leq z\}| \quad (7)$$

be the *threshold cumulative distribution function* (CDF) that returns the fraction of agents having threshold *less* than or *equal* to  $z$ .

In general, the threshold CDF is non-decreasing, piece-wise constant and continuous to the right with discontinuities occurring at points  $r_i$ ,  $i \in \mathcal{V}$ . We denote by

$$F^-(z) := \lim_{a \rightarrow z^-} F(a) = \frac{1}{n} |\{i \in \mathcal{V} \mid r_i < z\}|$$

the left limit of  $F$  in  $z$ .

Similarly, the *threshold complementary cumulative distribution function* (CCDF) is given by

$$G(z) := \frac{1}{n} |\{i \in \mathcal{V} \mid r_i > z\}| \quad (8)$$

and returns the fraction of agents having threshold *greater* than a given value. Note that  $G(z) = 1 - F(z)$ . Accordingly, the threshold CCDF is non-increasing, piece-wise constant and continuous to the right with discontinuities occurring at points  $r_i$ ,  $i \in \mathcal{V}$ . We denote by  $G^-(z) := \lim_{a \rightarrow z^-} G(a)$  the left limit of  $G$  in  $z$ .

Throughout the paper, we denote the threshold CDF of the coordinating agents by  $F_c$  and threshold CCDF of the anti-coordinating agents by  $G_a$ . Our goal is to characterize Nash equilibria in terms of the thresholds CDF and CCDF. Part of the difficulty is due to the fact that in our finite size setting, the fraction variables  $z$ ,  $z_c$ , and  $z_a$  are not continuous and variations restricted to multiples of  $1/n$ . To give an intuition of the results that will be presented in the next section, below we briefly analyze the case of an infinite population of agents where such technical issues disappear.

### 2.4 Infinite population

We assume an infinite continuous population of agents with a percentage  $\alpha \in [0, 1]$  of coordinating agents, while the remaining agents are anti-coordinating. Furthermore, we assume that  $F_c : \mathbb{R} \rightarrow [0, 1]$  and  $G_a : \mathbb{R} \rightarrow [0, 1]$  are *continuous* functions.

A configuration corresponding to a global fraction of 1 given by  $z^*$  is a Nash if and only if the players in  $\mathcal{V}_c$  playing action 1 are exactly those having threshold above  $z^*$  and all players in  $\mathcal{V}_a$  playing action 1 are exactly those having threshold below  $z^*$ . Indicating with  $z_c^*$ ,  $z_a^*$  the fraction of agents playing action 1 in respectively  $\mathcal{V}_c$  and  $\mathcal{V}_a$ , Nash equilibria can then be characterized by the equations

$$\begin{cases} z_c^* = F_c(z^*) \\ z_a^* = G_a(z^*) \\ z^* = \alpha z_c^* + (1 - \alpha) z_a^* \end{cases} \quad (9)$$

If we define

$$H_\alpha(z) := \alpha F_c(z) + (1 - \alpha) G_a(z), \quad (10)$$

we can see that all the solutions of (9) can be found by first solving

$$z^* = H_\alpha(z^*) \quad (11)$$

and then considering  $z_c^* = F_c(z^*)$  and  $z_a^* = G_a(z^*)$ . Notice that since  $H_\alpha : [0, 1] \rightarrow [0, 1]$  is continuous, solutions to (11) always exist. This shows that the mixed CAC game for an infinite continuous population and assuming that  $F_c$  and  $G_a$  are continuous, always admits Nash equilibria.

*Example 2.* Let the thresholds of the coordinating agents be *uniformly distributed* in  $[0, 1]$ , as well as the thresholds of the anti-coordinating agents, i.e.  $F_c(z) = z \mathbf{1}_{[0,1]}(z) + \mathbf{1}_{(1,\infty)}(z)$  and  $G_a(z) = 1 - F_c(z)$ . Then

$$H_\alpha(z) = (2\alpha - 1)z + (1 - \alpha)$$

and its unique fixed point is  $z^* = \frac{1}{2}$  (Fig. 2). The NE is given by  $z_c^* = F_c(\frac{1}{2}) = \frac{1}{2}$  and  $z_a^* = G_a(\frac{1}{2}) = \frac{1}{2}$ .




Fig. 2. We study the fixed point equation (11) for Ex. 2. On the left we consider  $\alpha < \frac{1}{2}$ , on the right  $\alpha > \frac{1}{2}$ .

### 3. NASH EQUILIBRIA OF MIXED CAC GAMES

In this section, we investigate the existence, the uniqueness and the characterization of NE of mixed CAC games.

Our main result is to provide a *necessary* and *sufficient* condition for an action configuration to be a Nash equilibrium of the mixed CAC game. The condition, which is a slight modification of (9), only involves the fractions of agents playing action +1 in the whole population and in the subsets  $\mathcal{V}_c$  and  $\mathcal{V}_a$ .

As we shall see, the condition is checkable and it reduces the complexity of the search of all NE to analyze  $n + 1$  triples of candidate fractions. In the special case where only coordinating or anti-coordinating agents are present, we will be able to give more explicit results.

#### 3.1 Formal statement of the main result

In this section, we formally state the main result. Given a Nash equilibrium  $x^* \in \mathcal{X}$ , we shortly denote  $z^* := z(x^*)$ ,  $z_c^* := z_c(x^*)$  and  $z_a^* := z_a(x^*)$ .

**Theorem 1.** Consider the mixed CAC game.

- (i) A Nash equilibrium  $x^* \in \mathcal{X}$  satisfies

$$\begin{cases} F_c^- \left( \frac{n}{n-1} z^* \right) = z_c^* = F_c \left( \frac{n}{n-1} (z^* - \frac{1}{n}) \right) \\ G_a^- \left( \frac{n}{n-1} (z^* - \frac{1}{n}) \right) \geq z_a^* \geq G_a \left( \frac{n}{n-1} z^* \right) \\ z^* = \alpha z_c^* + (1 - \alpha) z_a^* \end{cases} \quad (12)$$

- (ii) Given  $z^* \in \{0, \frac{1}{n}, \dots, 1\}$ ,  $z_c^* \in \{0, \frac{1}{n_c}, \dots, 1\}$ ,  $z_a^* \in \{0, \frac{1}{n_a}, \dots, 1\}$  satisfying (12), there exists at least one Nash equilibrium  $x^* \in \mathcal{X}$  such that  $z(x^*) = z^*$ ,  $z_c(x^*) = z_c^*$  and  $z_a(x^*) = z_a^*$ .

**Remark 1.** The Nash equilibrium whose existence is guaranteed by (ii) can be built in the following way. First, we set the actions of the coordinating agents  $i \in \mathcal{V}_c$  accordingly to the following formula

$$x_i^* = \begin{cases} +1 & \text{if } r_i \leq \frac{n}{n-1} \left( z^* - \frac{1}{n} \right) \\ -1 & \text{if } r_i > \frac{n}{n-1} z^* \end{cases} \quad (13)$$

Then, we force the actions of the anti-coordinating agents  $i \in \mathcal{V}_a$  to satisfy

$$x_i^* = \begin{cases} -1 & \text{if } r_i < \frac{n}{n-1} \left( z^* - \frac{1}{n} \right) \\ +1 & \text{if } r_i > \frac{n}{n-1} z^* \end{cases} \quad (14)$$

Finally, we set the actions of the anti-coordinating agents  $i \in \mathcal{V}_a$  having thresholds  $r_i \in [\frac{n}{n-1} (z^* - \frac{1}{n}), \frac{n}{n-1} z^*]$  in such a way that the condition  $z^*(x^*) = z^*$  is met. In general, the associated Nash equilibrium is not unique.

Theorem 1 gives a tool to investigate the existence of Nash equilibria of the mixed CAC game and it permits to characterize all the NE of the game. Indeed, once we find all triples of fractions  $(z^*, z_c^*, z_a^*)$  satisfying (12), then we can trivially find all the Nash equilibria of the game.

**Remark 2.** Consider a triple  $(z^*, z_c^*, z_a^*)$  satisfying (12). The values of  $z_c^*$  and  $z_a^*$  are uniquely determined from  $z^*$  through the 1<sup>st</sup> and 3<sup>rd</sup> equations. Hence, there are at most  $n + 1$  triples satisfying the conditions in (12).

#### 3.2 Nash equilibria of the coordination game

In this section, we focus on the coordination game, i.e.,  $\mathcal{V}_c = \mathcal{V}$ . A straightforward substitution of  $\alpha = 1$  in (12) leads to the following characterization of NE for coordination games.

**Corollary 2.** Consider the coordination game.

- (i) A Nash equilibrium  $x^* \in \mathcal{X}$  satisfies

$$F_c^- \left( \frac{n}{n-1} z^* \right) = z^* = F_c \left( \frac{n}{n-1} (z^* - \frac{1}{n}) \right) \quad (15)$$

- (ii) Given  $z^* \in \{0, \frac{1}{n}, \dots, 1\}$  satisfying (15), there exists a Nash equilibrium  $x^* \in \mathcal{X}$  such that  $z(x^*) = z^*$ .

In this case, we can strengthen the analysis as reported in the result below.

**Proposition 2.** The coordination game always admits Nash equilibria. Moreover, the number of NE coincide with the number of solutions  $z^*$  of the equations (15) and this number never exceeds  $n + 1$ .

We comment on this result. Existence follows from the fact that the game is in this case potential, but can also directly derived from condition (15). Notice that, if there are no stubborn agents, that is,  $F(0) = 0$  and  $F(1) = 1$ , then  $z^0 = 0$  and  $z^1 = 1$  are solutions of (15). They correspond to the two *consensus configurations*. On the other hand, the fact that a NE  $x^*$  is completely determined by the corresponding value of  $z^*$  is an immediate consequence of the characterization expressed in formula (13).

The following example shows a case where equations (15) has exactly  $n + 1$  distinct solutions.

**Example 3.** Consider the thresholds  $r = [0, \frac{1}{n-1}, \dots, 1]$ . The condition in (15) admits the  $n + 1$  solutions  $0, \frac{1}{n}, \dots, 1$  as shown in Fig. 3.

Finally, notice that a fixed point  $z^* = F_c^- (\frac{n}{n-1} z^*)$  is a solution of (15) only when there is a flat interval preceding the fixed point large at least  $\frac{1}{n}$ . In Fig. 3, we provide an example where a fixed point does not satisfy (15).

#### 3.3 Nash equilibria of the anti-coordination game

In this section, we focus on the anti-coordination game, i.e.,  $\mathcal{V}_a = \mathcal{V}$ . A straightforward substitution of  $\alpha = 0$  in (12) leads to the following characterization of NE for anti-coordination games.






Fig. 3. On the left, for thresholds in Ex. 3, (15) admits  $n + 1$  solutions. On the right, we consider an example where  $z^0 = 0$  and  $z^1 = 1$  are the unique solutions of (15). Even if  $z^* = \frac{n-1}{2n}$  is a fixed point, the CDF is not flat in the interval  $[z^* - \frac{1}{n}, z^*]$ .

**Corollary 3.** Consider the anti-coordination game.

- (i) A Nash equilibrium  $x^* \in \mathcal{X}$  satisfies

$$G_a^-\left(\frac{n}{n-1}(z^* - \frac{1}{n})\right) \geq z^* \geq G_a\left(\frac{n}{n-1}z^*\right) \quad (16)$$

- (ii) Given  $z^* \in \{0, \frac{1}{n}, \dots, 1\}$  satisfying (16), there exists a Nash equilibrium  $x^* \in \mathcal{X}$  such that  $z(x^*) = z^*$ .

A deeper insight is possible also for this case, though it has peculiar differences with respect to the pure coordination case. We first note that a *fixed point*  $z^* \in \{0, \frac{1}{n}, \dots, 1\}$  of the threshold CCDF, namely  $z^* = G_a(\frac{n}{n-1}z^*)$ , satisfies (16). Such fixed points however not necessarily exist and, in any case, other solutions to (16) may show up. In the following result, we propose a sufficient existence condition for NE that encompasses the fixed point condition above.

**Proposition 3.** Consider the inequalities

$$\begin{cases} z^* - \frac{1}{n} < G_a\left(\frac{n}{n-1}(z^* - \frac{1}{n})\right) \\ z^* \geq G_a\left(\frac{n}{n-1}z^*\right) \end{cases} \quad (17)$$

Then, any  $z^* \in \{0, \frac{1}{n}, \dots, 1\}$  satisfying (17) is a solution of (16).

Since  $G_a$  is non-increasing with codomain  $[0, 1]$ , solutions to (17) always exist. An example is shown in Fig. 4. This argument in particular proves the existence of NE for anti-coordination game, fact that could also be derived from the fact that such games are potential. Notice that, concretely, starting from  $z^*$  a corresponding NE  $x^*$  can be constructed setting the actions of part of the agents as in (14) and the remaining actions in such a way that the condition  $z^*(x^*) = z^*$  is met. The associated Nash equilibrium is, in general, *not* unique.

The next result further refines our analysis of the NE of the anti-coordination game and includes an analytical computation of its cardinality.

**Proposition 4.** For the anti-coordination game, the following facts hold true.

- (1) The number of solutions of the conditions (16) is either 1 or 2.
- (2) Indicated with  $\mathcal{Z}$  the set of solutions of (16), the number of NE is given by

$$|\mathcal{N}| = \sum_{z \in \mathcal{Z}} \binom{n(\zeta_2 - \zeta_1)}{n(z - \zeta_1)} \quad (18)$$




Fig. 4. On the left, an example when inequality (16) admits the unique solution  $z^* = \frac{9}{18}$ . On the right, we consider  $n = 19$  and  $r_i = \frac{1}{2}$ , for all  $i \in \mathcal{V}$ . The inequality (16) admits the two solutions  $z^* = \frac{9}{19}$  and  $z^{**} = \frac{10}{19}$ .

where

$$\zeta_1 := G_a\left(\frac{n}{n-1}z\right), \zeta_2 := G_a^-\left(\frac{n}{n-1}(z - \frac{1}{n})\right).$$

We briefly comment on the statement in item 1.: the possibility of a second solution in (16) come from the fact that if  $z^*$  is a solution and there is a sufficiently big jump between  $G_a(z^*)$  and  $G_a^-(z^*)$ , then  $z^{**} = z^* + \frac{1}{n}$  is another solution. No other choice is possible as  $G_a$  is by definition non-increasing. This phenomenon is shown in Fig. 4.

### 3.4 Nash Equilibria of the mixed CAC game

In this section, we make some final considerations on the general mixed case. The difficulty in analyzing the existence condition (12) is due to the simultaneous presence of the two threshold functions that give rise to two coupled conditions. In the infinite continuous population case this was essentially overcome by considering the function  $H_\alpha$  defined in (10) and reducing the problem to a fixed point investigation. Similar considerations can also be done in the finite size case and, even if they do not lead to a complete characterization, they do shed some light on the problem. To this aim, consider any  $(z^*, z_c^*, z_a^*)$  satisfying (12) and makes the extra assumption that

$$\begin{aligned} F_c^-\left(\frac{n}{n-1}z^*\right) &= F_c\left(\frac{n}{n-1}z^*\right), \\ G_a^-\left(\frac{n}{n-1}(z^* - \frac{1}{n})\right) &= G_a\left(\frac{n}{n-1}(z^* - \frac{1}{n})\right), \end{aligned}$$

Then,  $z^*$  also satisfies the pair of inequalities

$$H_\alpha\left(\frac{n}{n-1}(z^* - \frac{1}{n})\right) \geq z^* \geq H_\alpha\left(\frac{n}{n-1}z^*\right). \quad (19)$$

This can be seen as an approximation of the fixed point relation  $z^* = H_\alpha(z^*)$  we had in the infinite case. This argument says that, in determining the solutions of (12), all the possible candidates for the fraction  $z^*$  are:

- (i) the solutions of (19) that are continuous points of both  $F_c(\frac{n}{n-1}*)$  and  $G_a(\frac{n}{n-1}(* - \frac{1}{n}))$ .
- (ii) The discontinuous points of either  $F_c(\frac{n}{n-1}*)$  and  $G_a(\frac{n}{n-1}(* - \frac{1}{n}))$ .

An example that shows how such argument can be concretely used is presented below.

**Example 4 (Mixed majority/minority game).** Consider  $r_i = \frac{1}{2}$  for all agents, and  $\alpha \in (0, 1)$ . Then, we have




Fig. 5. We consider Ex. 4 for  $\alpha \geq \frac{1}{2} + \frac{1}{n}$ . The pair of inequalities in (19) admits the two solutions  $z^* = \alpha$  and  $z^{**} = 1 - \alpha$ .

$$H_\alpha \left( \frac{n}{n-1} z \right) = (1 - \alpha) \mathbb{1}_{[0, \frac{n-1}{2n}]}(z) + \alpha \mathbb{1}_{[\frac{n-1}{2n}, 1]}(z).$$

Notice that  $F_c(\frac{n}{n-1} *)$  is discontinuous in  $\frac{n-1}{2n}$ , whereas  $G_\alpha(\frac{n}{n-1}(* - \frac{1}{n}))$  is discontinuous in  $\frac{n+1}{2n}$ .

CASE 1: If  $\alpha \geq \frac{1}{2} + \frac{1}{n}$ , the condition in (19) admits the two solutions  $z^* = \alpha$  and  $z^{**} = 1 - \alpha$  (Fig. 5). Hence, we have 4 candidates:  $z^*$ ,  $z^{**}$ , and the 2 discontinuous points.

If we substitute the first candidate  $z^* = \alpha$  in the 1<sup>st</sup> and 3<sup>rd</sup> equations of (12), we get  $z_c^* = 1$  and  $z_a^* = 0$ . Note that the triple  $(\alpha, 1, 0)$  does satisfy (12). The triple associated to  $z^{**}$  is  $(1 - \alpha, 0, 1)$  and it satisfies (12) as well. The two solutions correspond to the NE where all coordinating agents play action 1, or action  $-1$ , respectively, while anti-coordinating agents play the opposite one.

Notice that there are no other NE as the triples associated to  $\frac{n-1}{2n}$  and  $\frac{n+1}{2n}$  do not satisfy (12).

CASE 2: If  $\alpha < \frac{1}{2} + \frac{1}{n}$  and  $n$  is odd, (19) admits the two solutions  $z^* = \frac{n-1}{2n}$  and  $z^{**} = \frac{n+1}{2n}$  (Fig. 6), which are also discontinuous points.

The fractions associated to  $z^*$  are  $z_c^* = F_c^-(\frac{n}{n-1} \frac{n-1}{2n}) = F_c^-(\frac{1}{2}) = 0$  and  $z_a^* = z^*/(1 - \alpha)$ . Also in this case, the triple  $(z^*, 0, z^*/(1 - \alpha))$  satisfies (12), and the same holds for the triple  $(z^{**}, 1, (z^{**} - \alpha)/(1 - \alpha))$ .

CASE 3: If  $\alpha < \frac{1}{2} + \frac{1}{n}$  and  $n$  is even, the unique solution of (19) is  $z^* = \frac{1}{2}$  (Fig. 6). Notice that

$$F_c^-\left(\frac{n}{n-1} \frac{1}{2}\right) = 1 \neq F_c\left(\frac{n}{n-1} \left(\frac{1}{2} - \frac{1}{n}\right)\right) = 0.$$

Hence,  $z^*$  does not induce a triple that solves (12). By checking the discontinuous points, one can prove that, in this case, the mixed CAC game admits no NE.

#### 4. CONCLUDING REMARKS

We considered mixed ensembles of agents playing coordinating and anti-coordinating agents with possibly different activation thresholds. We found a necessary and sufficient condition for the existence of Nash equilibria in terms of the distribution of the thresholds in the ensemble and an explicit characterization of Nash equilibria when do exist. When only coordinating or anti-coordinating agents are present, we were capable to determine NE's cardinality.

Current work includes the extension of this analysis to the case when agents are confined to interact through




Fig. 6. We consider Ex. 4 for  $\alpha < \frac{1}{2} + \frac{1}{n}$ . If  $n$  is odd (left), the solutions of (19) are  $z^* = \frac{n-1}{2n}$  and  $z^{**} = \frac{n+1}{2n}$ . If  $n$  is even (right), the unique solution is  $z^* = \frac{1}{2}$ .

a network. Preliminary results in this direction can be found in Vanelli (2019) and Arditti et al. (2020), where we determined graph-theoretical sufficient conditions for the existence of a pure strategy Nash equilibria for mixed network CAC games. Also, we are interested in the behavior of evolutionary dynamics associated to these games, such as the the best response dynamics.

#### REFERENCES

- Arditti, L., Como, G., Fagnani, F., and Vanelli, M. (2020). Robustness of Nash equilibria in network games. *Preprint*. <https://arxiv.org/abs/2004.12869>.
- Bramoullé, Y. (2007). Anti-coordination and social interactions. *Games and Economic Behavior*, 58, 30–49.
- Ellison, G. (1993). Learning, local interaction, and coordination. *Econometrica*, 61(5), 1047–1071.
- Grabisch, M. and Li, F. (2019). Anti-conformism in the threshold model of collective behavior. *Dynamic Games and Applications*, 9, 1–34.
- Granovetter, M. (1978). Threshold models of collective behavior. *American Journal of Sociology*, 83(6), 1420–1443.
- Jackson, M. and Zenou, Y. (2015). Games on networks. *Handbook of Game Theory with Economic Applications*, 4.
- Kandori, M., Mailath, G.J., and Rob, R. (1993). Learning, mutation, and long run equilibria in games. *Econometrica*, 61(1), 29.
- Monderer, D. and Shapley, L.S. (1996). Potential games. *Games and Economic Behavior*, 14(1), 124–143.
- Montanari, A. and Saberi, A. (2010). The spread of innovations in social networks. *Proceedings of the National Academy of Sciences*, 107(47), 20196–20201.
- Morris, S. (2000). Contagion. *The Review of Economic Studies*, 67(1), 57–78.
- Ramazi, P., Riehl, J., and Cao, M. (2016). Networks of conforming and nonconforming individuals tend to reach satisfactory decisions. *Proceedings of the National Academy of Sciences*, 113(46), 12985–12990.
- Rossi, W.S., Como, G., and Fagnani, F. (2017). Threshold models of cascades in large-scale networks. *IEEE Transactions on Network Science and Engineering*, 6(2), 158 – 172.
- Vanelli, M. (2019). *On network games with coordinating and anti-coordinating agents*. Master's thesis, Politecnico di Torino. <https://webthesis.biblio.polito.it/11996>.