

Managing floodplains using naturebased solutions to support multiple ecosystem functions and services

Original

Managing floodplains using naturebased solutions to support multiple ecosystem functions and services / Jakubínský, Jií; Prokopová, Marcela; Raška, Pavel; Salvati, Luca; Bezak, Nejc; Cudlín, Ondej; Cudlín, Pavel; Purkyt, Jan; Vezza, Paolo; Camporeale, Carlo; Dank, Jan; Pástor, Michal; Lepeška, Tomáš. - In: WIRES. WATER. - ISSN 2049-1948. - ELETTRONICO. - e1545:(2021). [10.1002/wat2.1545]

Availability:

This version is available at: 11583/2911314 since: 2021-07-06T17:58:45Z

Publisher:

John Wiley & Sons

Published

DOI:10.1002/wat2.1545



Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Managing floodplains using nature-based solutions to support multiple ecosystem functions and services

Jiří Jakubínský¹  | Marcela Prokopová¹ | Pavel Raška² | Luca Salvati³ | Nejc Bezak⁴ | Ondřej Cudlín¹ | Pavel Cudlín¹ | Jan Purkyt¹ | Paolo Vezza⁵ | Carlo Camporeale⁵ | Jan Daněk^{1,6} | Michal Pástor⁷ | Tomáš Lepeška⁸ 

¹Global Change Research Institute CAS, Brno, Czech Republic

²Jan Evangelista Purkyně University in Ústí nad Labem, Ústí nad Labem, Czech Republic

³Department of Economics and Law, University of Macerata, Marche, Italy

⁴University of Ljubljana, Faculty of Civil and Geodetic Engineering, Ljubljana, Slovenia

⁵Department of Environment, Land and Infrastructure Engineering (DIATI), Politecnico di Torino, Torino, Italy

⁶Faculty of Humanities, Charles University, Praha, Czech Republic

⁷National Forest Center, Banská Bystrica, Slovakia

⁸Faculty of Ecology and Environmental Sciences, Technical University in Zvolen, Zvolen, Slovakia

Correspondence

Jiří Jakubínský, Global Change Research Institute CAS, Bělidla 986/4a, 603 00 Brno, Czech Republic
Email: jakubinsky.j@czechglobe.cz

Funding information

European Cooperation in Science and Technology, Grant/Award Number: CA16209; Javna Agencija za Raziskovalno Dejavnost RS, Grant/Award Number: P2-0180; Ministerstvo Školství, Mládeže a Tělovýchovy, Grant/Award Number: LTC18025

Edited by: Stuart N. Lane, Editor-in-Chief

Abstract

Floodplains include unique environments shaped over a long time horizon along rivers and smaller streams and formed by alluvial sediments. As floodplains are flat, often with highly fertile and well-accessible land, they have become the intrinsic focus of human society—while providing a variety of goods and ecosystem services. Intensive land use of floodplains is degrading their natural values and significantly reducing their ecosystem functions and services. A significant part of these key services is related with the ability of floodplains to retain water and nutrients, which can be understood as a flood control and a water-retention function. Although these ecosystems serve a number of other basic functions, the importance of floodplains as a place for water retention during extreme discharges caused by intense rainfall or snowmelt and the supply of water in times of drought are essential under conditions of global change. In order to increase the ability of floodplains to perform these functions, it is increasingly required to preserve the connectivity of rivers with surrounding floodplains and adapt human activities to maintain and restore river ecosystems. This article reviews the recent understanding of floodplain delimitation, the most common causes of disturbance, the ecosystem functions being performed, discussing in turn the measures being considered to mitigate the frequency and magnitude of hydrologic extremes resulting from ongoing environmental changes.

This article is categorized under:

Water and Life > Conservation, Management, and Awareness
Engineering Water > Planning Water

This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

© 2021 The Authors. *WIREs Water* published by Wiley Periodicals LLC.

1 | INTRODUCTION

Floodplains are specific parts of the natural landscape, whose formation and existence are due to their association with a watercourse, allowing the exchange of flow, sediment, nutrients, and organisms (Amoros and Bornette, 2002; Benjankar, Egger, Jorde, Goodwin, & Glenn, 2011). Mostly defined as an area along the watercourse, floodplains are usually formed by alluvial sediments deposited during floods of varying magnitude and associated geomorphological processes (e.g., Lewin, 1978; Nardi, Vivoni, & Grimaldi, 2006). Flooding, with a wide variety of discharge magnitudes and events ranging from extreme low flow events to infrequent high flow events (Poff et al., 1997; Whiting, 2002), underpin floodplain ecosystem dynamics and influences a variety of biophysical and territorial features (e.g., fluvial landforms, soil hydrology, or vegetation patterns), being a crucial process for ecosystem functioning. The timing, duration, and magnitude of floods influence the structure of riparian communities (Auble & Scott, 1998; Scott, Shafroth, Auble, & Eggleston, 1997), as well as ecosystem functions and services. At the same time, flooding can be a dangerous phenomenon when floodplains are intensively developed (de Martino, de Paola, Fontana, Marini, & Ranucci, 2012). The strength of connectivity between a river and the surrounding terrestrial environment (i.e., the floodplain) varies depending on the hydrogeomorphic control of the downstream flux of water and materials—both dissolved and particulate matter (Stanford, Lorang, & Hauer, 2005). A river should retain a flow regime with sufficient variability to encompass the flow levels and events that support important floodplain processes (Opperman et al., 2010). Lateral exchange between river channels and their floodplains, known as hydrologic connectivity, has been identified as a key variable in biodiversity and composition of aquatic (e.g., Desjonquères, Rybak, Castella, Llusia, & Sueur, 2018; Leigh & Sheldon, 2009; Paillex, Dolédec, Castella, Mérigoux, & Aldridge, 2013) and terrestrial (e.g., Casco, Neiff, & de Neiff, 2010; Souter, Wallace, Walter, & Watts, 2014) biological communities. Hydrological connectivity also represents a key feature supporting ecosystem processes such as nutrient turnover and geomorphic change (Hein, Baranyi, Reckendorfer, & Schiemer, 2004; Schönbrunner, Preiner, & Hein, 2012; Welti, Bondar-Kunze, Tritthart, Pinay, & Hein, 2012 or Park, 2020).

Floodplain ecosystems are unique in terms of their constantly recurring hydrological dynamics (Funk et al., 2019; Schindler et al., 2014; 2016), which result from the interaction of geomorphic, hydrological, and biological processes (Tomscha et al., 2017). Together with the outstanding ability of floodplains to retain water (Getirana et al., 2017), such intrinsic dynamics modify morphology and water conditions and ensure the high diversity of natural conditions, as well as their temporal variability (Tockner et al., 2000), maintaining a highly diversified mosaic of habitats, from open soils to deciduous forests (Fischer et al., 2019), with a marked variability of aquatic, semi-aquatic and terrestrial habitats (Hughes et al., 2005). Worldwide, with some of the most distinctive examples found in Europe, floodplains are threatened by the loss of floodplain and riparian habitats, as well as by pollution and alteration of hydromorphological conditions (Funk et al., 2020; Habersack et al., 2016; Vörösmarty et al., 2010). The most obvious effect of human activities is urbanization, which increases the proportion of impermeable surfaces in floodplains, degrades landform diversity, and affect sediment balance by altering runoff regimes (Chin, 2006; Booth and Bledsoe, 2009; Raška et al., 2019). Additionally, levee construction along rivers often exacerbates the downcutting of riparian forests or plant communities and increases in bank height, reflecting the urgent need for channel stabilization measures following changes in flow or sediment regimes (Zachary et al., 2003). Higginson et al. (2020) considers water resource development to be one of the main causes of floodplain degradation, which has led to a decline in floodplain ecological condition. The embankment and isolation of rivers from their floodplains, which allows their intensive use for agriculture, settlements or traffic routes, are among the most common interventions in Central and Southern European floodplains (Hein, Schwarz, et al., 2016). The floodplain that remains active are altered due to these changes and habitat regeneration is hindered (Díaz-Redondo et al., 2017). Where floodplain forests remain, they are mostly converted from naturally regenerating stands to stands managed for forestry. The extensive hardwood forests of Central European floodplains are of particular economic importance (Klimo and Hager, 2008). Wetlands are being replaced by plant communities adapted to the water regime of reservoir shorelines (Keddy, 2010), or even with forestry systems using non-native tree species (Hughes et al., 2012). Recreation in floodplains is increasing in many parts of Europe, being a threat to conservation goals as well as a chance for a better public appreciation of the value of floodplains and rivers (Hughes et al., 2012). Channel stabilization and peak flow reduction in turn, disconnect a river channel from a floodplain, reducing both the channel migration rates and the channel avulsion (rapid channel shift during floods; Shields Jr et al., 2000, Zachary et al., 2003). Approximately 70–90% of Europe's current floodplain area is estimated to be ecologically degraded due to human activities over the centuries, especially since the early 1950s (EEA, 2018), thus we can summarize that Europe is the continent most affected by disconnection of floodplains from rivers (Nilsson et al. 2005; Schindler et al. 2016). Tockner and

Stanford (2002) suggested that approximately 46% of floodplains in North America (excluding northern Canada and Alaska) were intensively cultivated, and 11% of floodplains across Africa were farmed at the beginning of the 21st century. According to Erwin (2009), 90% of floodplains in North America are described as “cultivated” and non-functional. Furthermore, climate projections in many parts of Europe, as well as other regions of the world, indicate an increasing occurrence of intense rainfall and prolonged droughts, which would affect the condition of floodplains (e.g. Moradkhani et al., 2010; Schneider et al., 2011; Politti et al., 2014 or O’Brian, 2019). This article brings an overview of ecosystem functions and services provided by floodplains and focuses on (a) operational frameworks for defining floodplains, (b) major causes of their disturbance, and (c) more general approaches for protecting and restoring these ecosystems. The issue of floodplain protection is outlined in this study with examples of the current situation in several Central and Southern European countries.

2 | FLOODPLAIN DEFINITION

2.1 | Floodplain as a soil phenomenon

Different approaches have been adopted over the time to distinguish the floodplains from other landscape types. Derived from the definition of a floodplain as an area along a watercourse formed by alluvial sediments deposited during floods, gathering information on the soil properties or spatial extent of the inundation area is the most relevant approach. A necessary prerequisite for this approach is data available at a relatively detailed scale (spatial resolution) and comparable across larger territorial units or countries (Jakubínský et al., 2020). Using soil data, a floodplain is usually defined based on the spatial distribution of hydromorphic soils, characterized by the temporary or permanent wetting of soil pores. Fluvisol is the most widespread floodplain soil type, formed by erosion of sediments in the upland zone and deposited in lowland sites with flat valley bottoms in the transfer (piedmont) zone (WRB, 2015). Another soil type that commonly found within a floodplain is Gleysol (WRB, 2015), the formation of which is influenced by periodic recurring or persistent excess moisture in the near-surface soil layers. In addition, a much less widespread soil type within a floodplain is Phaeozems (Fluvisol Phaeozems according to WRB [2015]), in the form of deep semi-hydromorphic soils. The spatial extent of certain soil types represents a stable component of the landscape. In fact, even after a possible loss of floodplain connectivity with the watercourse due to anthropogenic interventions, the soil types remain for a long period of time, although the floodplain itself loses its natural functions.

2.2 | Hydrologically conceived floodplain

Hydrological and hydraulic data, mostly the results of modeling based on digital elevation models (DEMs), are often used to define floodplains, in particular using the extent of inundated areas of 100-year flood (e.g., Omer et al., 2003 or EEA, 2018). However, other values have been frequently used. For instance, Witner (1966) proposed an area of alluvial soils corresponding to 50-year floods. Within this hydrological approach, the identification of floodplains relied upon the creation of flood hazard maps, produced through detailed hydraulic modeling techniques (Grimaldi et al., 2013; Noman et al., 2001). Existing methods for delineating inundated areas using hydraulic simulations were reviewed by Noman et al. (2001) and Horritt and Bates (2002). Although these techniques and models can result in highly accurate delineation of a floodplain, they can be computationally expensive and time-consuming to run, requiring the calibration of a large number of variables and model parameters (e.g., Horritt and Bates, 2002; Liu and Gupta, 2007). Additionally, adequate input data (e.g., river cross-section or floodplain LiDAR data) are needed to obtain reliable results. The significant refinement and facilitation of modeling is currently related to the availability of very accurate elevation data provided by remote sensing technologies such as LiDAR (Light Detection and Ranging) methods in a number of affluent countries (e.g. Bezak et al., 2018; Rak et al., 2018; Ureta et al., 2020). In contrast to soil data, operational definitions by flooded area are highly dependent on valley floor terrain characteristics and any anthropogenic intervention can directly affect the extent of inundated areas and, thus, floodplains defined in this manner.

Since hydrologically and hydraulically defined floodplains depend on terrain characteristics, this delineation approach is also understood as a “hydrogeomorphic method” (Nardi et al., 2006). This method uses GIS technologies and hydrologic modeling techniques (e.g., HEC-RAS software; Ackerman et al., 2009; Patel et al., 2016) to delineate floodplains as buffers at a specified distance from the watercourse (Entwistle et al., 2019), depending on the elevation of

the terrain above the river water level. Floodplain delineation approaches (based on hydrologic–hydraulic and soils data) often achieve very similar results in terms of the spatial extent of the defined floodplains in non-urban areas, as evidenced in Figure 1. In urban areas, on the other hand, both approaches can be very different in terms of the location of the borderline between the floodplain and the surrounding landscape, due to anthropogenic influences (e.g., the presence of levees, flood protection walls and road or railway embankments).

In the Figure 2a, a broad active floodplain, which is limited by the slope edge adjacent to the valley floor, is depicted. In contrast, Figure 2b shows the influence of anthropogenic landforms (presence of levees) affecting the extent of the active floodplain, which is inundated during regular flood events. Behind the levee, the pedologically defined floodplain (also referred to as the “geologic floodplain” according to Fuller [2018]), i.e., alluvial soil types formed in the past, actually shows intermittent connectivity with the riverbed in the presence.

2.3 | Floodplain according to specific vegetation

As floodplains represent a specific environment formed by a long-term connectivity with a watercourse, in the case of less-significant anthropogenic interventions and a minor modification of the natural environment, they can also be defined on the basis of specific vegetation cover. These are almost always communities of azonal vegetation that do not




FIGURE 1 Comparison of floodplain areas defined on the basis of hydrological and pedological data along the Dřevnice River in the Czech Republic, in urban (a) and forest-agricultural (b) landscape.

Source: authors, based on data provided by T. G. Masaryk Water Research Institute and Research Institute for Soil and Water Conservation




FIGURE 2 (a) Floodplain area covering the entire valley floor in the case of a near-natural landscape (the Berounka River floodplain, Czech Republic); (b) urban floodplain limited by the presence of levees (the Vltava River floodplain in České Budějovice, Czech Republic).

Source: authors

occur in any other landscape type than in floodplains, depending on the specific location relative to the channel axis (Tockner and Stanford, 2002). Spatiotemporal variability in surface and groundwater hydrology, microclimate, geomorphology, and soils, combined with inter- and intraspecific competition, results in distinctive floodplain biodiversity (Hughes et al., 2005; Meli et al., 2014; Robinson et al., 2002; Salo et al., 1986) that is most evident along aquatic–terrestrial boundaries (Bunn and Arthington, 2002). Dynamic fluvial processes cause habitat rejuvenation and succession (Hohensinner and Drescher, 2008), resulting in habitat heterogeneity (Opperman et al., 2010), which supports a floodplain with a shifting mosaic of habitat patches in terms of species, age classes, and physical structure (Ward et al., 2002). Many authors (e.g., Evette et al., 2014; Sponseller et al., 2013; Corenblit et al., 2014 or McShane et al., 2015) identify hydro-climatic constraints such as climate, soil moisture availability and fluvial disturbance as a major factor influencing vegetation near watercourses. Gurnell et al. (2016) propose the conceptual model of vegetation—hydrogeomorphology interactions using so-called dynamic zones within river corridors where different hydrogeomorphological processes dominate so that plants and hydrogeomorphological processes interact in different ways.

3 | THE ECOSYSTEM FUNCTIONS AND SERVICES OF A FLOODPLAIN

Floodplains represent one of the most productive ecosystems on Earth (Opperman et al., , 2010b). The high degree of biodiversity and level of primary productivity of floodplains exceed the production of either purely terrestrial or aquatic ecosystems (Tockner and Stanford, 2002). The dynamics and naturally high biodiversity of floodplains are responsible for their high multifunctionality (Meli et al., 2014; Funk et al., 2020). Focusing on the value of global ecosystem services, Costanza et al. (1997) found that floodplains are the second best ranked ecosystem type, behind estuaries, in terms of their per hectare value to society. Despite representing <2% of the Earth's terrestrial land surface area, floodplains provide approximately 25% of all “terrestrial” (i.e., non-marine) ecosystem service benefits, with the regulation of disturbance (i.e., attenuation of flood flows) providing the greatest value (Akanbi et al., 1999). Floodplains contribute to a wide range of ecosystem functions by controlling the regional hydrologic cycle and the retention and transformation of nutrients in river systems (Sanon et al., 2012; Schindler et al., 2014; Weigelhofer and Hein 2015). Their connectivity patterns are also crucial for the provisioning of ecosystem services, including floodwater retention (e.g., Clilverd et al., 2016; Habersack et al., 2015; Schober et al., 2015), nutrient retention (e.g., Hein et al., 2004; Natho et al., 2013; Newcomer Johnson et al., 2016), or greenhouse gas emission retention (e.g., Audet et al., 2013; Funk et al., 2020). Other ecosystem services provided by floodplains include surface water filtration (Mitsch et al., 2001; Noe and Hupp, 2005), groundwater recharge (Hein et al., 2004; Jolly, 1996), water purification (Hein, van Koppen, et al., 2016), and provision of food and fiber (e.g., fish, timber, and other plant resources; Welcomme, 1979). Fisheries supported by floodplain productivity provide one of the most tangible examples of an economically and socially valuable ecosystem service (Opperman et al., 2010). Furthermore, recent efforts to quantify the cultural ecosystem services provided by floodplains and river ecosystems point to a range of non-material benefits (Funk et al., 2020; Hale et al., 2019). Traditionally, floodplains provide various water-related recreational opportunities, including swimming, boating, angling, and ice skating (Funk et al., 2020). River landscapes may also be valued for their aesthetic quality and cultural or heritage significance (Ghermandi et al., 2020; Thiele et al., 2020; Tieskens et al., 2018). A list of the key ecosystem functions and services provided by floodplains can be found in Table 1.

Floodplain ecosystem services are inextricably linked to hydrology (Morris et al., 2009); the hydrologic regime of a floodplain determines what will grow there and how it can be used (Posthumus et al., 2010). Forest ecosystems in particular depend on natural hydrological and biological diversity (Turner et al., 2016), which are very important for the delivery of regulation, provisioning, and cultural services (Mamat et al., 2018; Xu et al., 2017). Our understanding of ecosystem services can benefit greatly from drawing on classic river–floodplain principles that recognize both longitudinal and lateral connectivity (Tomscha et al., 2017).

Floodplains host a unique suite of habitats, and their composition and condition can serve as indicators of the ecosystem services they are able to provide (Burkhard et al., 2012; Podschun et al., 2018). Riparian systems (vegetation in close proximity to the watercourse) are those of a highest importance; due to their spatial location and connectivity with stream channels, they are inundated periodically and play an important role in water infiltration and aquifer recharge (Gonzalez del Tanago et al., 2011), as well as flood attenuation and hydrological risk reduction (Horn and Richards, 2006). Anderson et al. (2006) reported that these are especially smaller floods (with an average recurrence interval of 2 or 5 years) that are more sensitive to the riparian vegetation conditions, and in these cases riparian

TABLE 1 The frequently cited ecosystem functions and services of floodplains and their measured parameters/indicators (a nonexhaustive list)

Category of service (CICES section)	Ecosystem function	Class of ecosystem service (CICES v5)	Parameter/indicator
Provisioning	Production (biomass)	Fibers and other materials from cultivated (and wild) plants, fungi, algae and bacteria for direct use or processing (excluding genetic materials); Animals reared for nutritional purposes	Gross output ¹ ; Annual biomass increase ² ; Nutritive productivity ²
	Water-retention and evapotranspiration	Surface water used as a material (non-drinking purposes); Ground and subsurface water for drinking	Difference between water rainfall and evapotranspiration ^{3,4,5} ; Potential for water provision ⁶
Regulation and Maintenance	Water-retention	Hydrological cycle and water flow regulation (Including flood control, and coastal protection)	Time to fill water capacity; Curve number (CN) ⁷ ; Quality of land cover, slope, soil permeability and flow length ⁸ ; Floodplain water storage volume ^{9,10} ; Effective retention volume ^{11,12} ; Net supply of water remaining after evapotranspiration losses ⁴ ; Water holding capacity ¹³ ; Retaining coefficients for forest management ⁵
	Self-cleaning processes of water	Mediation by other chemical or physical means (e.g. via Filtration, sequestration, storage or accumulation)	Nutrient leaching ¹ ; Nitrogen leaching from floodplain area ⁴ ; Total nitrogen and total phosphorus removed from water ³ ; Phosphorus load in the river modeled by InVEST ¹⁵ ; Water quality index ⁶ ; Diversity of the instream macroinvertebrates ¹⁶
	Evapotranspiration and Condensation	Regulation of chemical composition of atmosphere and ocean (e.g., greenhouse gases concentration, isotopic variance in atmospheric moisture); Carbon sequestration by terrestrial ecosystems	Inverse values of daily temperatures range for land use types ² ; Evapotranspiration ^{17,18,19} ; Albedo of land use types ¹⁷ ; Extent of vegetation cover ¹⁴ ; Evapotranspiration and heat exchange based on functional plant traits ²⁰ ; Land surface temperature ²¹
	Carbon capture	Carbon sequestration by terrestrial ecosystems	Carbon sequestration by plants ² ; Above ground carbon storage ²² ; Global warming potential ¹ ; Carbon stock ⁶ ; Fluxes of greenhouse gases for land use types ^{10,4}
	Mineralization and accumulation of organic matter, storage and recycling of nutrients	Decomposition and fixing processes and their effect on soil quality	Soil carbon stock ¹ ; Organic matter layer and total nitrogen in top soil ² ; Floodplain connectivity ¹⁴
	Functions of species composition and diversity	Maintaining nursery populations and habitats (Including gene pool protection)	A species value indicator ¹ ; Habitat-conservation value ¹ ; Riparian quality index ^{2,23} ; Habitat provision index ²⁴ ; Fish capacity index ²² ; Diversity of the instream macroinvertebrates ¹⁶ ; Proportion of natural land cover weighted by a condition index ⁴ ; Presence of threatened species ¹⁴ ; Habitat value according to Habitat Valuation Method ²⁵

TABLE 1 (Continued)

Category of service (CICES section)	Ecosystem function	Class of ecosystem service (CICES v5)	Parameter/indicator
Cultural	Recreation	Characteristics of living systems that enable activities promoting health and wellbeing, recuperation or enjoyment through active or immersive interactions; Characteristics of living systems that enable activities promoting health, recuperation or enjoyment through passive or observational interactions	Possibility to experience the terrain, Presence of protected areas, Water surface area, Presence of sandbanks and meanders, Visibility depth, Minimum width for (non-) motorized boating ²⁶ ; Frequency of tourists per year ²⁷ ; Content of geotagged photographs uploaded to social media sites ^{28,31,32} ; Diversity of potential for nature experiences ²⁹ ; Recreation potential, Recreation opportunity spectrum ³⁰
	Heritage	Characteristics of living systems that are resonant in terms of culture or heritage	Density of monuments and cultural-historical facilities, Density of archeological and natural monuments ²⁶ ; Content of geotagged photographs uploaded to social media sites ^{28,32}
	Aesthetic values	Characteristics of living systems that enable aesthetic experiences	Landscape diversity, naturalness and uniqueness ^{26,33} ; Level of aesthetic value ²⁷ ; Content of geotagged photographs uploaded to social media sites ^{28,32}

Note: Source: authors. ¹Posthumus et al. (2010); ²Felipe-Lucia et al. (2014); ³Boithias et al. (2016); ⁴Ausseil et al. (2013); ⁵Morri et al. (2014) ⁶Larsen et al. (2012); ⁷Fu et al. (2013); ⁸Nin et al. (2016); ⁹Grygoruk et al. 2013; ¹⁰Peh et al. (2014); ¹¹Pithart et al. (2010); ¹²Karpack et al. (2020); ¹³Ghaley et al. (2014); ¹⁴Peters (2016); ¹⁵Johnson et al. (2012); ¹⁶Ncube et al. (2018); ¹⁷West et al. (2011), ¹⁸Smith et al. (2013); ¹⁹Serna-Chavez et al. (2017); ²⁰de Bello et al. (2010); ²¹Alkama and Cescatti (2016); ²²Tomscha et al. 2017; ²³Gonzalez del Tanago et al. 2011; ²⁴Fischer et al. 2019; ²⁵Seják et al. (2010); ²⁶Thiele et al. 2020; ²⁷Ajwang' Ondiek et al. 2016; ²⁸Ghermandi et al. 2020; ²⁹Funk et al. (2020); ³⁰Grizzetti et al. (2019); ³¹Tieskens et al. (2018); ³²Hale et al. (2019); ³³Thiele et al. (2020); ³³Thiele et al. (2019). CICES stands for the Common International Classification of Ecosystem Services developed by the European Environment Agency (EEA).

vegetation affects the magnitude of a flood event and the resulting damage. As an important landform agent and flow resistance factor, riparian vegetation is responsible for the majority of energy losses in fluvial systems by controlling sediment erosion, transport, and deposition in both the channel and floodplain (Corenblit et al., 2008; Gonzalez del Tanago et al., 2011). The most important ecological functions of the riparian zone include providing a habitat and refuge for aquatic and terrestrial species, facilitating biological connections across the landscape, maintaining plant diversity, providing organic material to aquatic food chains, and controlling stream water temperature (Forman, 1999; Gonzalez del Tanago and Garcia de Jalon, 2011). These functions are all related to the dimensions, longitudinal continuity and vegetation structure of riparian corridors (Gurnell et al., 2016), becoming particularly important in regions with a Mediterranean-climate (Stella et al., 2013). In addition, riparian vegetation provides many other aesthetic and economic benefits, including food resources (Pusey and Arthington, 2003).

4 | DISTURBANCE, PROTECTION, AND RESTORATION OF FLOODPLAINS IN CENTRAL AND SOUTHERN EUROPE

Almost all large rivers in Central and Southern Europe are affected by dikes and other flood-protection measures, in major catchments, such as the Rhine, Elbe, Danube and Oder, only 10–20% of the former floodplains are left as inundation areas (Brunotte et al., 2009). Only 10% of the original extent of European floodplain forests has been preserved, most of which are located in Eastern Europe (Hughes et al., 2012). Areas with the best-preserved floodplain forests remained in Croatia along the Danube and Sava Rivers (Anić, 2008). Many floodplain areas were drained in the past in

order to intensify agricultural use, for example in Slovakia (Holubová et al., 2003), in the Czech Republic (Brázdil et al., 2011) or in Hungary in large parts of the Tisza lowland basin (Szmańda et al., 2008). Recently, urbanization and the development of new transport routes along river valleys with associated channelization works are another threat to floodplains, particularly in some Mediterranean countries and in the more recent accession countries of Central Europe (Hughes et al., 2012). The construction of dams also has a significant impact on floodplains, not only affecting the area inundated by the construction of a waterworks, but the sediment starvation downstream of dams has perhaps the greatest potential to impact on floodplain development. According to Marren et al. (2014), we can identify several ways in which floodplains could potentially be affected by dams, with varying degrees of confidence, including a distinction between passive impacts (floodplain disconnection) and active impacts (changes in geomorphological processes and functions). According to EEA (2018), 15% of the European population lives in floodplains, rising to 25% in Austria, Slovakia and Slovenia. In Southern Europe, problems with water supply to agriculture have led to an intensive river regulation strategy, including the construction of reservoirs in upland valleys (Hughes et al., 2012). Many lowland rivers have been realigned to maximize agricultural production. These rivers are among the most regulated in the world (Magdaleno and Fernández, 2011), which is reflected in the poor ecological condition of their floodplains. The conclusion of the study by Kuiper et al. (2014), based on a meta-analysis of the scientific literature, was that altering of a natural flow regime reduces mean species abundance (MSA) of floodplains by more than 50% on average, and species richness by more than 25%. The effects on species richness and abundance tend to be related to the degree of hydrologic alteration. A list of the most common causes of floodplain ecosystem degradation and corresponding mitigation measures, usually taken in Central and Southern Europe can be found in Table 2.

As floodplains are valuable ecosystems that provide a range of ecosystem functions and services, it is necessary to address their protection and implement selected measures to ensure that their environmental values are maintained or restored (Hughes et al., 2008). According to EEA (2018), the protection and restoration of European floodplains is promoted within environmental policy, but only indirectly required—i.e., by the Water Framework Directive—WFD (2000/60/EC), the Floods Directive (2007/60/EC), the Habitat (1992/43/EEC) and Birds—HBD (1979/409/EEC) Directives, the EU 2020 Biodiversity Strategy, the Green Infrastructure initiative, the EU Climate Change Adaptation Strategy, and the Ramsar Convention. As the protection of such human altered floodplains along large European rivers is one of the objectives of the WFD and HBD (Funk et al., 2019), achieving these objectives require detailed planning of various compromise solutions that are ecologically, commercially, and socially acceptable (Rouquette et al., 2011). In addition to European legislation, the floodplains or at least some of the ecosystems found in Europe (e.g., floodplain forests or wetlands) are subject to legal protection at the level of individual countries. To illustrate the different approaches to floodplain management, here we outline the current situation in several Central and Southern European countries. In the Czech Republic floodplains are protected by law; according to Act No. 114/1992 Coll., a floodplain area is considered a “significant landscape element.” From a practical point of view, this form of floodplain protection is not completely effective, as it is not clear how to approach the protection of already partially degraded floodplains. The

TABLE 2 The most common causes of floodplain degradation in central and southern Europe and appropriate measures to mitigate the effects of degradation (Source: authors)

Cause of floodplain degradation	Mitigation measure(s)
Drainage and riverbed reinforcement due to agricultural activities	River restoration (reestablish morphological river type and lateral connectivity), set initial measures for type-specific self-development of the river/the floodplain system, detention ponds construction, adopt ad hoc crop rotations and agricultural practices (tillage systems, soil cover management, etc.), check and rebuild old drainage systems.
Urbanization and transport infrastructure development	Develop urban green infrastructure and stormwater drainage management.
Technical/structural flood protection measures limiting lateral floodplain connectivity (embankment)	Creation of room for river (polders and areas suitable for periodic flooding) to alleviate floods where possible.
Dam construction and construction of torrent controls	Mitigate hydropeaking, implement fish bypasses and modify migration barriers in order to reestablish longitudinal continuum; remove or reconstruct torrent controls in headwater areas.
Land-use/land cover changes	Floodplain habitat restoration, modify land-use (afforestation of unused land a pastures).

same law provides for better protection of floodplains only if they are located in core zones of large-scale protected areas or in small-scale protected areas. In addition, floodplain protection is hampered by the absence of datasets that capture the precise spatial extent of floodplain ecosystems. Although floodplains in the Czech Republic represent a significant part of the landscape, covering approximately 10% of the total area of the country (Štěrba, 2008), ineffective legal protection has led to extensive degradation in the past and it is currently difficult to implement large-scale restorations that would improve the ecological condition of floodplains. Instead of complex restoration of floodplain ecosystems, these areas are more often subjected to construction of various flood protection measures, aimed at protecting residential and industrial infrastructure located directly in flooded areas (Loučková, 2014). Whether it is a structural measure located in the urban area of a floodplain (e.g., artificial levee or solid concrete wall) or outside urban areas (e.g., polder), all these measures affect biodiversity and the quality of ecosystem functions and services provided. While flood control measures implemented outside urban areas usually have a positive effect on several different ecosystem functions, on the contrary, measures implemented in built-up areas usually degrade many ecosystem functions. The poor condition of riparian habitats in the Czech Republic is also related to the practical maintenance of the River Basin Management Authorities based on the “Water Act” (No. 254/2001 Coll.), which aims to ensure sufficient space for the water flowing in the riverbed by cutting down forests on the riverbanks.

More often than entire floodplain areas, their individual parts (specific ecosystems or habitats) are protected by law; for example, in Slovakia, where wetlands, bogs or peat bogs, wet meadows, natural flowing waters and natural standing (lentic) waters are protected under Act No. 543/2002. There are no specific maps of floodplain areas in Slovakia; however, background materials are available that can be considered as proxy data of floodplain distribution—e.g., maps of Quaternary deposits (Maglay et al., 2009), soil maps (Hraško et al., 1993), or potential primary vegetation (Michalko et al., 1986). One of the most important localities where the protection of natural values of the floodplain is addressed in Slovakia is the great floodplain of the Danube River. An example of suitable measures to increase the environmental values of the floodplains is the LIFE project “Conservation and Management of Danube Floodplain Forests”, which focuses on the conservation of the last remaining natural floodplain forests in the Slovak part of the Danube floodplain and the introduction of sustainable forest management in this area (BROZ, 2003). Restoration measures such as reconnection of meanders with the river system, increase of flow dynamics, excavation of sediment deposits from the meanders, and a special mowing scheme have been proposed and partially implemented to conserve the natural values and the derived human benefits (Holubová et al., 2003). Restoration of the Danube floodplain in Slovakia is perceived as an important factor in improving environmental values, as it is a highly anthropogenically modified area. Major interventions in this area include, for example, the construction of the Gabčíkovo waterworks, leading to a slow degradation of rare and endangered habitats of softwood floodplain forests in the Danube inland delta—see Figure 3 (Petrášová-Šibíková et al., 2017).

In Slovenia, conditions and limitations related to construction and activities in floodplain areas are defined by the “Decree on conditions and limitations for constructions and activities on flood risk areas” (PISRS, 2020). This regulation mentions the flooding and erosion processes of surface water and sea. The methodology that should be used to define endangered floodplain is determined by the “Rules on methodology to define flood risk areas and erosion areas connected to floods and classification of plots into risk classes.” These rules use the concept of 10-year, 100-year, and 500-year return periods. Moreover, in relation to flood risk, the “Water Act” should also be mentioned. As a result of this legislation, flood hazard maps for various parts of the country have been prepared. However, since the adoption of this law about 10 years ago, a lot of construction has already taken place in floodplains, especially near the larger cities such as Ljubljana and Celje (e.g., Glavan et al., 2020). The floodplain area, defined based on the extent of flooding during the 100-year return period, where much of the urban development of the city of Ljubljana in Slovenia is located, is shown in Figure 4. The strong dependence of the floodplain extent on the terrain characteristics is a major drawback of this hydrological (hydraulic) approach, since in the case of urban areas or localities with traffic embankments, the course of the floodplain borderlines is intensively modified.

River lateral connectivity with floodplains is essential to create and maintain habitats for animals and plants, ensure ecosystem services and integrity, enhance carbon sequestration and storage (Wohl et al., 2017). A case study of the Orco River (North-Western Italy) is used to demonstrate how maintaining an adequate width of the river corridor and sustaining lateral river migration can be used as an effective solution to (a) mitigate flood risk, (b) minimize damage to transport infrastructure, (c) support the objectives of the EU Water Framework, Floods and Habitat Directives. The Orco river basin has a total area of about 910 km², of which 78% is located in the Alpine mountain range and 22% in the Po Valley plain. The Orco River flows on the southern slope of the Gran Paradiso massif, where an area of 11 km² is currently occupied by glaciers. In the Po Valley plain (between the municipality of Cuornè and the confluence with



FIGURE 3 Danube floodplain forest near the Gabčíkovo waterworks, Slovakia
(Source: Jaroslav Jankovič)




FIGURE 4 Significantly urbanized floodplain area (defined as the extent of flooding of the 100-year return period) of Ljubljanica River and its tributaries in the eastern part of Ljubljana, Slovenia
(Source: Geoportal ARSO, 2020)

the Po River near Chivasso municipality), the river is characterized by a wandering morphology with a significant lateral migration of bends, due to the intense sediment transport during the autumn and spring floods. Thanks to moderate anthropogenic pressures, the Orco River corridor still preserves a riparian forest in the floodplain, located between the river banks and the terraces. In late October 2019, a significant meander chute-cut occurred near San Benigno Canavese village (Figure 5), with a consecutive evolution of the chute channel in 2020. The chute channel has been formed in the floodplain peninsula enclosed by the meander loop, resulting in gradual closure of the abandoned channel. The chute channel is currently continuing to incise and widen into the floodplain. This process naturally preempted and avoided the realization of a river engineering measure envisioned by the authorities to reduce lateral bank migration towards the highway. The formation of this new channel did indeed reduce flood risk across the right river bank, minimizing possible damages to the highway and to the San Benigno Canavese village (Figure 5). The river bank protection by riprap at the outer bend of the meander is also currently not necessary because, after 16 months, the chute channel is acting as the main channel, conveying the discharge during the majority of time. The shortening of the river



FIGURE 5 A meander chute cut-off on the Orco River near San Benigno Canavese village, Italy (Source: Paolo Maschio, Politecnico di Torino)

centerline has also resulted in increase in water gradient, with the consequent generation of erosion waves slowly migrating upstream, which will likely affect the hydraulic geometry upstream of the chute.

Water regulation in Italy provides for a very extensive stratification of planning competence, involving the national level, the regional administrative level and intermediary authorities whose sphere of influence extends beyond traditional administrative boundaries, such as the basin authorities (Salvati et al., 2012). This complex management framework often collides with the urgency of controlling, managing and regulating water flows in a very complex socio-economic context, such as in Italy and, more generally, Mediterranean Europe (e.g. Chelleri et al., 2015). Informal settlements spreading in the floodplain areas without considering sufficient buffer zones have sometimes led to severe conditions not only for ecosystems, but also for human health and life (Chelli et al., 2016; Ciommi et al., 2017; Gigliarano and Chelli, 2016). These situations are exacerbated by erosive processes due to land use changes, fires, landslides and abandonment of marginal lands (Salvati and Zitti, 2009; Salvati et al., 2011). Regional planning has mainly acted through integrated tools, which involved regional landscape plans, provincial coordination plans, river basin plans and planning at a more detailed scale of intervention, allowing both water regulation in contexts of particularly intense meteoric inflows, and emergency water management under drought conditions to be organized fairly effectively (e.g. Bajocco et al., 2012). Environmental policies at national and regional level have privileged the protection of relict floodplains in northern Italy, especially in flat areas, allowing the creation of habitats with high biodiversity, representing the natural extension of rivers with an alpine water regime (Smiraglia et al., 2016).

5 | DISCUSSION AND CONCLUSIONS

This article provided comparative insight into different approaches to floodplain delineation and outlined a conceptual nexus between floodplains and ecosystem functions, both illustrated with case studies. It is worth reiterating that direct human intervention (e.g., Entwistle et al., 2019; Lewin, 2013; Westra and de Wulf, 2006) can be seen as a major cause of floodplain ecosystem degradation, most evident in the urban landscape. There, the floodplain area is built-up and natural ecosystem functions are reduced due to loss of connectivity with the watercourse caused by river bank fortification and construction of levees along the river (e.g. Hein, van Koppen, et al., 2016 or Amoateng et al., 2018). In the agricultural landscape, the quality of ecosystem functions performed by the floodplain area is negatively affected by significant human-induced channel incision and narrowing, resulting in a lowering of the water table. As cropland is the most prevalent land-use category in floodplains in the Central and Southern European countries (cropland occupies between 40% and 60% of floodplain area according to EEA, 2018), the above anthropogenic interventions are the most

common cause of reduced ability of floodplains to perform ecosystem functions. Drainage and replacement of floodplain forests by agriculture, primarily fields and meadows, is most common (Klimo et al. 2008), resulting in loss of inundation areas and increasing sediment and nutrient delivery to the river (van Andel and Aronson 2012).

Based on the above examples, it can be concluded that floodplain protection is at a different stage of development in each country and has different preferences. Since all floodplains in Central and Southern Europe are very intensively anthropogenically used and no change can be expected, multifunctional floodplain management can be seen as a solution to ensure the sustainable use of these areas. Multifunctional floodplain management can be defined as a management approach that aims at a balanced provision of multiple ecosystem services that serve the needs of local residents. Existing trade-offs imply the need of provisioning services reduction to decrease their dominance (Schindler et al., 2014). Landscapes can be enhanced by adding (or maintaining) semi-natural landscape features designed to provide multiple ecosystem services (Lovell and Johnston, 2009). The importance of investing in natural ecosystems, in particular urban green spaces, floodplains and areas for recreation, is recognized as a source of economic development in EU regional and cohesion policies (COM, 2011). In Germany, for example multifunctionality is to some extent included in legal regulations – the Federal Water Resources Act requires water managers to preserve, protect and even enhance natural habitats in order to manage water resources sustainably (Schindler et al., 2016). The solution to these ecologically unsatisfactory conditions, coupled with increased flood risk, is possible through the restoration of a watercourse or an entire floodplain (Keesstra et al., 2018). In order to find the optimal combination of spatially distinct large-scale and small-scale measures to increase habitat availability for all relevant species, detailed spatial planning is an important component of floodplain restoration (Remm et al., 2019). Weigelhofer et al. (2020) consider a combination of multiple-species (aiming at restoring natural hydrological dynamics) and single-species approaches (focusing on the conservation status of individual species) as a sound basis for decision-making processes in floodplain restoration in accordance with the EU Water Framework Directive and the Birds and Habitat Directives, as well as local legislation.

To ensure both ecologically and socially viable restoration efforts, future research should explore the following uncertainties and trade-offs. First, a proper delineation of current floodplains should take into account the legacy of Late Holocene climatic oscillations that influence the magnitude of sediment fluxes in floodplains (Stacke et al., 2014), as well as the legacy of past human activities (Swinnen et al., 2020, 2020). Decoupling the cause–effect feedback between these processes is difficult (Hoffman and Rohde, 2011), but crucial for establishing historical baselines of floodplain restoration to improve ecosystem functions. Second, given the uncertainties associated with ongoing climate change and its spatio-temporally varying impacts, floodplain management must consider the different scales at which socio-ecological systems are transformed (Liu et al., 2007), including (a) variations in floodplain adjustment processes (Chin, 2006), (b) the existing mismatches between the scales of ecohydrological processes and those of planning and policy interventions (Raška et al., 2019), and (c) complicated and dynamic property rights and tenure systems (Hartmann, 2009) and the social importance of floodplains (Richards et al., 2017). These issues point out the necessity to balance ecosystem functions with the livelihood benefits of direct floodplain use (Juarez Lucas and Kibler, 2016). Awareness of the social significance of floodplain ecosystems is likely to be a key element in improving the overall ecological conditions of floodplains and ensuring their sustainability, as only an awareness of this fact can lead to restoration efforts being supported by floodplain landowners.

ACKNOWLEDGMENTS

The authors would like to thank Paolo Maschio and Jaroslav Jankovič for providing a photo of the Orco River meander in Italy and the Danube floodplain forest in Slovakia. We are also grateful to the anonymous reviewers for their valuable comments and feedback, which helped to improve the quality of the article. This article contributes to the EU COST Action network “LAND4FLOOD: Natural Flood Retention on Private Land” (CA16209, <http://www.land4flood.eu>) and the complementary national Inter-COST Project (MŠMT LTC18025). N. Bezak work was partially supported by the Slovene Research Agency (ARRS) through grant P2-0180.

CONFLICT OF INTEREST

The authors declare no conflicts of interest.

AUTHOR CONTRIBUTIONS

Jiří Jakubínský: Conceptualization; funding acquisition; methodology; supervision; visualization; writing-original draft; writing-review & editing. **Marcela Prokopová:** Data curation; resources; writing-original draft; writing-review & editing. **Pavel Raska:** Conceptualization; methodology; writing-original draft; writing-review & editing. **Luca Salvati:**

Methodology; resources; supervision; writing-original draft; writing-review & editing. **Nejc Bezak**: Data curation; methodology; visualization; writing-original draft; writing-review & editing. **Ondřej Cudlín**: Data curation; methodology; resources; writing-original draft; writing-review & editing. **Pavel Cudlín**: Conceptualization; methodology; resources; supervision; writing-original draft; writing-review & editing. **Jan Purkyt**: Data curation; resources; visualization. **Paolo Vezza**: Data curation; writing-original draft. **Carlo Camporeale**: Data curation; writing-original draft. **Jan Daněk**: Data curation; writing-original draft; writing-review & editing. **Michal Pástor**: Data curation; resources; writing-original draft; writing-review & editing. **Tomáš Lepeška**: Data curation; resources; writing-original draft; writing-review & editing.

DATA AVAILABILITY STATEMENT

The data that support the findings of this study are available from the corresponding author upon reasonable request.

ORCID

Jiří Jakubínský  <https://orcid.org/0000-0002-7461-2611>

Tomáš Lepeška  <https://orcid.org/0000-0003-0385-3482>

RELATED WIREs ARTICLES

[Developing and using geomorphic condition assessments for river rehabilitation planning, implementation and monitoring](#)

[Natural flood management](#)

[Monitoring the effectiveness of floodplain habitat restoration: A review of methods and recommendations for future monitoring](#)

FURTHER READING

- FAO (1988). *Soil map of the world. Revised legend*. World Soil Resources Report, No. 60. Rome, FAO-UNESCO-ISRIC.
- Fausch, K. D., Torgersen, C. E., Baxter, C. V., & Li, H. W. (2002). Landscapes to riverscapes: Bridging the gap between research and conservation of stream fishes: A continuous view of the river is needed to understand how processes interacting among scales set the context for stream fishes and their habitat. *Bioscience*, 52(6), 483–498.
- Gonzalez del Tanago, M., & Garcia de Jalon, D. (2011). Riparian Quality Index (RQI): A methodology for characterising and assessing the environmental conditions of riparian zones. *Limnetica*, 30(2), 235–254.
- Hoffmann, T., Erkens, G., Gerlach, R., Klostermann, J., & Lang, A. (2009). Trends and controls of Holocene floodplain sedimentation in the Rhine catchment. *Catena*, 77, 96–106.
- Kuiper, J. J., Janse, J. H., Teurlincx, S., Verhoeven, J. T., & Alkemade, R. (2014). The impact of river regulation on the biodiversity intactness of floodplain wetlands. *Wetlands Ecology and Management*, 22(6), 647–658.
- Leopold, L. B., & O'Brien Marchand, M. (1968). On the quantitative inventory of the riverscape. *Water Resources Research*, 4(4), 709–717.
- Patel, C. G., & Gundaliya, P. J. (2016). Floodplain delineation using HEC-RAS model—A case study of Surat City. *Open Journal of Modern Hydrology*, 6(1), 34–42. <http://dx.doi.org/10.4236/ojmh.2016.61004>.
- Pithart, D., Žaloudík, J., Dostál, T., Valentová, J., Valenta, P., Weyskrabová, J., & Dušek, J. (2010). Ecosystem services of natural floodplain segment-Lužnice River, Czech Republic. *WIT Transactions on Ecology and the Environment*, 133, 129–139.
- Richards, K., & Hughes, F. (2008). Floodplains in Europe: The case for restoration. In T. Moss & J. Monstadt (Eds.), *Restoring floodplains in Europe. Policy contexts and project experiences* (pp. 16–46). IWA.
- Wiens, J. A. (2002). Riverine landscapes: Taking landscape ecology into the water. *Freshwater Biology*, 47(4), 501–515.

REFERENCES

- Ackerman, C. T. (2009). *HEC-GeoRAS GIS tools for support of HEC-RAS using ArcGIS user's manual version 4.2*. US Army Corps of Engineers Institute for Water Resources Hydrologic Engineering Center (HEC).
- Ajwang' Ondiek, R., Kitaka, N., & Omondi Odour, S. (2016). Assessment of provisioning and cultural ecosystem services in natural wetlands and rice fields in Kano floodplain, Kenya. *Ecosystem Services*, 21, 166–173.
- Akanbi, A. A., Lian, Y. Q., Soong, T. W. (1999). *An analysis on managed flood storage options for selected levees along the lower Illinois river for enhancing flood protection*. Report No 4: Flood Storage Reservoirs and Flooding on the Lower Illinois River. Illinois State Water Survey. Contract Report 645.
- Alkama, R., & Cescatti, A. (2016). Biophysical climate impacts of recent changes in global forest cover. *Science*, 351(6273), 600–604.
- Amoateng, P., Finlayson, C. M., Howard, J., & Wilson, B. (2018). Dwindling rivers and floodplains in Kumasi, Ghana: A socio-spatial analysis of the extent and trend. *Applied Geography*, 90, 82–95.
- Amoros, C., & Bornette, G. (2002). Connectivity and biocomplexity in waterbodies of riverine floodplains. *Freshwater Biology*, 47(4), 761–776.

- Anderson, B. G., Rutherford, I. D., & Western, A. W. (2006). An analysis of the influence of riparian vegetation on the propagation of flood waves. *Environmental Modelling & Software*, 21(9), 1290–1296.
- Anić, I. (2008). Floodplain forests of Croatia. In E. Klimo & H. Hager (Eds.), *The floodplain forests in Europe: Current situation and perspectives*, European Forest Institute research report no. 10. Koninklijke Brill NV.
- Auble, G. T., & Scott, M. L. (1998). Fluvial disturbance patches and cottonwood recruitment along the upper Missouri River, Montana. *Wetlands*, 18(4), 546–556.
- Audet, J., Elsgaard, L., Kjaergaard, C., Larsen, S. E., & Hoffmann, C. C. (2013). Greenhouse gas emissions from a Danish riparian wetland before and after restoration. *Ecological Engineering*, 57, 170–182.
- Ausseil, A. G., Dymond, J. R., Kirschbaum, M. U. F., Andrew, R. M., & Parfitt, R. L. (2013). Assessment of multiple ecosystem services in New Zealand at the catchment scale. *Environmental Modelling & Software*, 43, 37–48.
- Bajocco, S., de Angelis, A., & Salvati, L. (2012). A satellite-based green index as a proxy for vegetation cover quality in a Mediterranean region. *Ecological Indicators*, 23, 578–587.
- Benjankar, R., Egger, G., Jorde, K., Goodwin, P., & Glenn, N. F. (2011). Dynamic floodplain vegetation model development for the Kootenai River, USA. *Journal of Environmental Management*, 92(12), 3058–3070.
- Bezák, N., Šraj, M., Rusjan, S., & Mikoš, M. (2018). Impact of the rainfall duration and temporal rainfall distribution defined using the huff curves on the hydraulic flood modelling results. *Geosciences*, 8, 69.
- Boithias, L., Terrado, M., Corominas, L., Ziv, G., Kumar, V., Marqués, M., Schuhmacher, M., & Acuña, V. (2016). Analysis of the uncertainty in the monetary valuation of ecosystem services: A case study at the river basin scale. *Science of the Total Environment*, 543, 683–690.
- Booth, D. B., & Bledsoe, B. P. (2009). Streams and urbanization. In L. Baker (Ed.), *The water environment of cities*. Springer.
- Brázdil, R., Máčka, Z., Řezníčková, L., Soukalová, E., Dobrovolný, P., & Grygar, M. (2011). Floods and floodplain changes of the river Morava, the Strážnické Pomoraví region (Czech Republic) over the past 130 years. *Hydrological Sciences Journal*, 56(7), 1166–1185.
- BROZ. (2003). *Conservation and management of Danube floodplain forests* (p. 16). BROZ: Regional Association for Nature Conservation and Sustainable Development.
- Brunotte, E., Dister, E., Guenther-Diringer, D., Koenzen, U., & Mehl, D. (2009). Flussauen in Deutschland: Erfassung und Bewertung des Auenzustandes (Riparian floodplains in Germany: survey and evaluation of floodplain conditions). In *Naturschutz und Biologische Vielfalt* (p. 87). BfN.
- Bunn, S. E., & Arthington, A. H. (2002). Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity. *Environmental Management*, 30(4), 492–507.
- Burkhard, B., Kroll, F., Nedkov, S., & Müller, F. (2012). Mapping ecosystem service supply, demand and budgets. *Ecological Indicators*, 21, 17–29.
- Casco, S. L., Neiff, J. J., & de Neiff, A. P. (2010). Ecological responses of two pioneer species to a hydrological connectivity gradient in riparian forests of the lower Paraná River. *Plant Ecology*, 209(1), 167–177.
- Chelleri, L., Schuetze, T., & Salvati, L. (2015). Integrating resilience with urban sustainability in neglected neighborhoods: Challenges and opportunities of transitioning to decentralized water management in Mexico City. *Habitat International*, 48, 122–130.
- Chelli, F. M., Ciommi, M., Emili, A., Gigliarano, C., & Taralli, S. (2016). Assessing the equitable and sustainable well-being of the Italian provinces. *International Journal of Uncertainty, Fuzziness and Knowledge-Based Systems*, 24, 39–62.
- Chin, A. (2006). Urban transformation of river landscapes in a global context. *Geomorphology*, 79, 460–487.
- Ciommi, M., Gentili, A., Ermini, B., Chelli, F. M., & Gallegati, M. (2017). Have your cake and eat it too: The well-being of the Italians (1861–2011). *Social Indicators Research*, 134(2), 473–509.
- Ciliverd, H. M., Thompson, J. R., Heppell, C. M., Sayer, C. D., & Axmacher, J. C. (2016). Coupled hydrological/hydraulic modelling of river restoration impacts and floodplain hydrodynamics. *River Research and Applications*, 32(9), 1927–1948.
- COM (2011). Regional policy contributing to sustainable growth in Europe 2020. European Commission, Brussels. COM 17, 2011. http://ec.europa.eu/regional_policy/sources/docoffic/official/communic/sustainable/comm2011_17_en.pdf
- Corenblit, D., Gurnell, A. M., Steiger, J., & Tabacchi, E. (2008). Reciprocal adjustments between landforms and living organisms: Extended geomorphic evolutionary insights. *Catena*, 73(3), 261–273.
- Corenblit, D., Steiger, J., Tabacchi, E., González, E., & Planty-Tabacchi, A. M. (2014). Ecosystem engineers modulate exotic invasions in riparian plant communities by modifying hydrogeomorphic connectivity. *River Research and Applications*, 30(1), 45–59.
- Costanza, R., d'Arge, R., de Groot, R., Farber, S., Grasso, M., Hannon, B., Limburg, K., Naeem, S., O'Neill, R. V., Paruelo, J., Raskin, R. G., Sutton, P., & van den Belt, M. (1997). The value of the world's ecosystem services and natural capital. *Nature*, 387(387), 253–260.
- de Bello, F., Lavorel, S., Díaz, S., Harrington, R., Cornelissen, J. H. C., Bardgett, R. D., Berg, M. P., Cipriotti, P., Feld, C. K., Hering, D., da Silva, P. M., Potts, S. G., Sandin, L., Sousa, J. P., Storkey, J., Wardle, D. A., & Harrison, P. A. (2010). Towards an assessment of multiple ecosystem processes and services via functional traits. *Biodiversity and Conservation*, 19, 2873–2893.
- de Martino, G., de Paola, F., Fontana, N., Marini, G., & Ranucci, A. (2012). Experimental assessment of level pool routing in preliminary design of floodplain storage. *Science of the Total Environment*, 416, 142–147.
- Desjonquères, C., Rybak, F., Castella, E., Llusia, D., & Sœur, J. (2018). Acoustic communities reflects lateral hydrological connectivity in riverine floodplain similarly to macroinvertebrate communities. *Scientific Reports*, 8(1), 1–11.
- Diaz-Redondo, M., Egger, G., Marchamalo, M., Hohensinner, S., & Dister, E. (2017). Benchmarking fluvial dynamics for process-based river restoration: The upper Rhine River (1816–2014). *River Research and Applications*, 33(3), 403–414.

- EEA (2018) European freshwater: Why should we care about floodplains? Briefing No. 14/2018. doi:<https://doi.org/10.2800/548993>
- Entwistle, N. S., Heritage, G. L., Schofield, L. A., & Williamson, R. J. (2019). Recent changes to floodplain character and functionality in England. *Catena*, *174*, 490–498.
- Erwin, K. L. (2009). Wetlands and global climate change: The role of wetland restoration in a changing world. *Wetlands Ecology and Management*, *17*(1), 71–84.
- Evette, A., Zanetti, C., Cavaillé, P., Dommanget, F., Mériaux, P., & Vennetier, M. (2014). The paradox when managing the riparian zones of rivers with engineered embankments in The French prealps. *Revue de géographie alpine*, 102–4, <http://dx.doi.org/10.4000/rga.2373>.
- Felipe-Lucia, M. R., Comín, F. A., & Bennett, E. M. (2014). Interactions among ecosystem services across land uses in a floodplain agroecosystem. *Ecology and Society*, *19*(1). <http://dx.doi.org/10.5751/es-06249-190120>.
- Fischer, C., Damm, C., Foeckler, F., Gelhaus, M., Gerstner, L., Harris, R., Hoffmann, T. G., Iwanowski, J., Kasperidus, H., Mehl, D., Podschum, S. A., Rumm, A., Stammel, B., & Scholz, M. (2019). The “habitat provision” index for assessing floodplain biodiversity and restoration potentials as an ecosystem service: Method and application. *Frontiers in Ecology and Evolution*, *7*, 483.
- Forman, R. T. T. (1999). *Land mosaics. The ecology of landscapes and regions* (p. 632). Cambridge University Press.
- Fu, B., Wang, Y. K., Xu, P., & Yan, K. (2013). Mapping the flood mitigation services of ecosystems: A case study in the Upper Yangtze River basin. *Ecological Engineering*, *52*, 238–246.
- Fuller, J.E. (2018). *Defining ordinary and natural conditions for state navigability determinations*. Arizona Geological Survey Contributed Report CR-18-B, 135p.
- Funk, A., Martínez-López, J., Borgwardt, F., Trauner, D., Bagstad, K. J., Balbi, S., & Hein, T. (2019). Identification of conservation and restoration priority areas in the Danube River based on the multi-functionality of river-floodplain systems. *Science of the Total Environment*, *654*, 763–777.
- Funk, A., Tschikof, M., Grüner, B., Böck, K., Hein, T., & Bondar-Kunze, E. (2020). Analysing the potential to restore the multi-functionality of floodplain systems by considering ecosystem service quality, quantity and trade-offs. *River Research and Applications*, *37*, 221–232.
- Getirana, A., Kumar, S., Giroto, M., & Rodell, M. (2017). Rivers and floodplains as key components of global terrestrial water storage variability. *Geophysical Research Letters*, *44*, 10359–10368.
- Ghaley, B. B., Vesterdal, L., & Porter, J. R. (2014). Quantification and valuation of ecosystem services in diverse production systems for informed decision-making. *Environmental Science & Policy*, *39*, 139–149.
- Ghermandi, A., Camacho-Valdez, V., & Trejo-Espinosa, H. (2020). Social media-based analysis of cultural ecosystem services and heritage tourism in a coastal region of Mexico. *Tourism Management*, *77*, 104002.
- Gigliarano, C., & Chelli, F. M. (2016). Measuring inter-temporal intragenerational mobility: An application to the Italian labour market. *Quality and Quantity*, *50*(1), 89–102.
- Glavan, M., Cvejić, R., Zupanc, V., Knapič, M., & Pintar, M. (2020). Agricultural production and flood control dry detention reservoirs: Example from lower Savinja Valley, Slovenia. *Environmental Science and Policy*, *114*, 394–402.
- Gonzalez del Tanago, M., & Garcia de Jalon, D. (2011). Riparian quality index (RQI): A methodology for characterising and assessing the environmental conditions of riparian zones. *Limnetica*, *30*(2), 235–254.
- Grimaldi, S., Petroselli, A., Arcangeletti, E., & Nardi, F. (2013). Flood mapping in ungauged basins using fully continuous hydrologic-hydraulic modeling. *Journal of Hydrology*, *487*, 39–47.
- Grizzetti, B., Liqueste, C., Pistocchi, A., Vigiak, O., Zuiliana, G., Bouraoui, F., de Roo, A., & Cardoso, A. C. (2019). Relationship between ecological condition and ecosystem services in European rivers lakes and coastal waters. *Science of the Total Environment*, *671*, 452–465.
- Grygoruk, M., Mirosław-Świątek, D., Chrzanowska, W., & Ignar, S. (2013). How much for water? Economic assessment and mapping of floodplain water storage as a catchment-scale ecosystem service of wetlands. *Water*, *5*(4), 1760–1779.
- Gurnell, A. M., Corenblit, D., García de Jalón, D., González del Tánago, M., Grabowski, R. C., O'hare, M. T., & Szewczyk, M. (2016). A conceptual model of vegetation–hydrogeomorphology interactions within river corridors. *River Research and Applications*, *32*(2), 142–163.
- Habersack, H., Schober, B., & Hauer, C. (2015). Floodplain evaluation matrix (FEM): An interdisciplinary method for evaluating river floodplains in the context of integrated flood risk management. *Natural Hazards*, *75*(1), 5–32.
- Habersack, H., Hein, T., Stanica, A., Liska, I., Mair, R., Jäger, E., & Bradley, C. (2016). Challenges of river basin management: Current status of, and prospects for, the river Danube from a river engineering perspective. *Science of the Total Environment*, *543*, 828–845.
- Hale, R. L., Cook, E. M., & Beltrán, B. J. (2019). Cultural ecosystem services provided by rivers across diverse social-ecological landscapes: A social media analysis. *Ecological Indicators*, *107*, 105580.
- Hartmann, T. (2009). Clumsy floodplains and the law: Towards a responsive land policy for extreme floods. *Built Environment*, *35*(4), 531–544.
- Hein, L., van Koppen, C. K., van Ierland, E. C., & Leidekker, J. (2016). Temporal scales, ecosystem dynamics, stakeholders and the valuation of ecosystems services. *Ecosystem Services*, *21*, 109–119.
- Hein, T., Baranyi, C., Reckendorfer, W., & Schiemer, F. (2004). The impact of surface water exchange on the nutrient and particle dynamics in side-arms along the River Danube, Austria. *Science of the Total Environment*, *328*, 207–218.
- Hein, T., Schwarz, U., Habersack, H., Nichersu, I., Preiner, S., Willby, N., & Weigelhofer, G. (2016). Current status and restoration options for floodplains along the Danube River. *Science of the Total Environment*, *543*, 778–790.
- Higginson, W., Higginson, B., Powell, M., Driver, P., & Dyer, F. (2020). Impacts of water resource development on hydrological connectivity of different floodplain habitats in a highly variable system. *River Research and Applications*, *36*(4), 542–552.

- Hoffman, M. T., & Rohde, R. F. (2011). Rivers through time: Historical changes in the riparian vegetation of the semi-arid, winter rainfall region of South Africa in response to climate and land use. *Journal of the History of Biology*, 44(1), 59–80.
- Hohensinner, S., & Drescher, A. (2008). *Historical change of European floodplains: The Danube River in Austria. The floodplain forests of temperate zone of Europe*. Lesnická práce.
- Holubová, K., Hey, R. D., & Lisicky, M. J. (2003). Middle Danube tributaries: Constraints and opportunities in lowland river restoration. *Large Rivers*, 15(1–4), 507–519.
- Horn, R., & Richards, J. S. (2006). Flow-vegetation interactions in restored floodplain environments. In P. J. Wood, D. M. Hannah & J. P. Sadler (Eds.), *Hydroecology and Ecohydrology: Past, Present and Future* (pp. 269–294). Chichester, England: John Wiley & Sons, Ltd.
- Horritt, M. S., & Bates, P. D. (2002). Evaluation of 1D and 2D numerical models for predicting river flood inundation. *Journal of Hydrology*, 268(1–4), 87–99.
- Hraško, J., Linkeš, V., Šály, R., & Šurina, B. (1993). *Pôdna mapa Slovenska. 1: 400 000*. VÚPÚ/Slov.
- Hughes, F. M., Moss, T., & Richards, K. S. (2008). Uncertainty in riparian and floodplain restoration. In S. Darby & D. Sear (Eds.), *River restoration: managing the uncertainty in restoring physical habitat*. (Vol. 79, pp. 94–119). Chichester, England: John Wiley & Sons, Ltd.
- Hughes, F. M., del Tánago, M. G., & Mountford, J. O. (2012). Restoring floodplain forests in Europe. In *A goal-oriented approach to forest landscape restoration* (pp. 393–422). Springer.
- Hughes, F. M. R., Colston, A., & Mountford, J. O. (2005). Restoring riparian ecosystems: The challenge of accommodating variability and designing restoration trajectories. *Ecology and Society*, 10(1). <http://www.ecologyandsociety.org/vol10/iss1/art12/>.
- Jakubínský, J., Herber, V., & Cudlín, P. (2020). A comparison of four approaches to river landscape delineation: The case of small water-courses in The Czech Republic. *Moravian Geographical Reports*, 27(4), 229–240.
- Johnson, K. A., Polasky, S., Nelson, E., & Pennington, D. (2012). Uncertainty in ecosystem services valuation and implications for assessing land use tradeoffs: An agricultural case study in the Minnesota River basin. *Ecological Economics*, 79, 71–79.
- Jolly, I. D. (1996). The effects of river management on the hydrology and Hydroecology of arid and semi-arid floodplains. In M. G. Anderson, D. E. Walling, & P. D. Bates (Eds.), *Floodplain processes* (pp. 577–609). John Wiley & Sons Ltd.
- Juarez Lucas, A. M., & Kibler, K. M. (2016). Integrated flood management in developing countries: Balancing flood risk, sustainable livelihoods, and ecosystem services. *International Journal of River Basin Management*, 14(1), 19–31.
- Karpack, M. N., Morrison, R. R., & McManamay, R. A. (2020). Quantitative assessment of floodplain functionality using an index of integrity. *Ecological Indicators*, 111, 106051.
- Keddy, P. A. (2010). *Wetland ecology: Principles and conservation*. Cambridge University Press.
- Keesstra, S., Nunes, J., Novara, A., Finger, D., Avelar, D., Kalantari, Z., & Cerdà, A. (2018). The superior effect of nature based solutions in land management for enhancing ecosystem services. *Science of the Total Environment*, 610, 997–1009.
- Klimo, E., Hager, H., Machar, I., Buček, A., & Schmalfus, R. (2008). Revitalization and protection of floodplain forests. Floodplain forests of the temperate zone of Europe. *Lesnická práce*, 87, 301–323.
- Larsen, F. W., Turner, W. R., & Brooks, T. M. (2012). Conserving critical sites for biodiversity provides disproportionate benefits to people. *PLoS One*, 7(5), e36971.
- Leigh, C., & Sheldon, F. (2009). Hydrological connectivity drives patterns of macroinvertebrate biodiversity in floodplain rivers of the Australian wet/dry tropics. *Freshwater Biology*, 54(3), 549–571.
- Lewin, J. (1978). Floodplain geomorphology. *Progress in Physical Geography*, 2(3), 408–437.
- Lewin, J. (2013). Enlightenment and the GM floodplain. *Earth Surface Processes and Landforms*, 38(1), 17–29.
- Liu, J., Dietz, T., Carpenter, S. R., Alberti, M., Folke, C., Moran, E., & Taylor, W. W. (2007). Complexity of coupled human and natural systems. *Science*, 317, 1513–1516.
- Liu, Y., & Gupta, H. V. (2007). Uncertainty in hydrologic modeling: Toward an integrated data assimilation framework. *Water Resources Research*, 43, W07401. <https://doi.org/10.1029/2006WR005756>
- Loučková, B. (2014). Eastern European perspective on the environmental aspects in current flood risk management: The example of The Czech Republic. In A. Bhadurri, J. Bogardi, J. Leentvaar, & S. Marx (Eds.), *The global water system in the Anthropocene. Challenges for science and governance* (pp. 183–196). Springer.
- Lovell, S. T., & Johnston, D. M. (2009). Creating multifunctional landscapes: How can the field of ecology inform the design of the landscape? *Frontiers in Ecology and the Environment*, 7(4), 212–220.
- Magdaleno, F., & Fernández, J. A. (2011). Hydromorphological alteration of a large Mediterranean river: Relative role of high and low flows on the evolution of riparian forests and channel morphology. *River Research and Applications*, 27(3), 374–387.
- Maglay, J., Pristaš, J., Kučera, M., & Ábelová, M. (2009). *Geologická mapa kvartéru Slovenska, Genetické typy kvartérnych uloženin 1:500 000*. MŽP SR a ŠGÚDŠ.
- Mamat, Z., Halik, Ü., Keyimu, M., Keram, A., & Nurmamat, K. (2018). Variation of the floodplain forest ecosystem service value in the lower reaches of Tarim River, China. *Land Degradation & Development*, 29(1), 47–57.
- Marren, P. M., Grove, J. R., Webb, J. A., & Stewardson, M. J. (2014). The potential for dams to impact lowland meandering river floodplain geomorphology. *The Scientific World Journal*, 2014, 1–24. <http://dx.doi.org/10.1155/2014/309673>.
- McShane, R. R., Auerbach, D. A., Friedman, J. M., Auble, G. T., Shaforth, P. B., Merigliano, M. F., & Poff, N. L. (2015). Distribution of invasive and native riparian woody plants across the western USA in relation to climate, river flow, floodplain geometry and patterns of introduction. *Ecography*, 38(12), 1254–1265.

- Meli, P., Benayas, J. M. R., Balvanera, P., & Ramos, M. M. (2014). Restoration enhances wetland biodiversity and ecosystem service supply, but results are context-dependent: A meta-analysis. *PLoS One*, 9(4), e93507.
- Michalko, J., Berta, J., & Magic, D. (1986). *Geobotanická mapa ČSSR (SSR): textová a mapová část. 1. vyd* (p. 162). Veda.
- Mitsch, W. J., Day, J. W., Gilliam, J. W., Groffman, P. M., Hey, D. L., Randall, G. W., & Wang, N. (2001). Reducing nitrogen loading to the Gulf of Mexico from the Mississippi River basin: Strategies to counter a persistent ecological problem: Ecotechnology—The use of natural ecosystems to solve environmental problems—Should be a part of efforts to shrink the zone of hypoxia in the Gulf of Mexico. *Bioscience*, 51(5), 373–388.
- Moradkhani, H., Baird, R. G., & Wherry, S. A. (2010). Assessment of climate change impact on floodplain and hydrologic ecotones. *Journal of Hydrology*, 395(3–4), 264–278.
- Morri, E., Pruscini, F., Scolozzi, R., & Santolini, R. (2014). A forest ecosystem services evaluation at the river basin scale: Supply and demand between coastal areas and upstream lands (Italy). *Ecological Indicators*, 37, 210–219.
- Morris, J., Posthumus, H., Hess, T., Gowing, D., & Rouquette, J. (2009). Watery land: The management of lowland floodplains in England. In M. Winter & M. Loble (Eds.), *What is land for* (pp. 135–166). New York: Earthscan.
- Nardi, F., Vivoni, E. R., & Grimaldi, S. (2006). Investigating a floodplain scaling relation using a hydrogeomorphic delineation method. *Water Resources Research*, 42(9), 1–15.
- Natho, S., Venohr, M., Henle, K., & Schulz-Zunkel, C. (2013). Modelling nitrogen retention in floodplains with different degrees of degradation for three large rivers in Germany. *Journal of Environmental Management*, 122, 47–55.
- Ncube, S., Visser, A., & Beavers, L. (2018). A framework for assessing Instream supporting ecosystem services based on Hydroecological Modelling. *Water*, 10(9), 1247.
- Newcomer Johnson, T. A., Kaushal, S. S., Mayer, P. M., Smith, R. M., & Svirich, G. M. (2016). Nutrient retention in restored streams and rivers: A global review and synthesis. *Water*, 8(4), 116.
- Nilsson, C., Reidy, C. A., Dynesius, M., & Revenga, C. (2005). Fragmentation and flow regulation of the world's large river systems. *Science*, 308, 405–408.
- Nin, M., Soutullo, A., Rodríguez-Gallego, L., & di Minin, E. (2016). Ecosystem services-based land planning for environmental impact avoidance. *Ecosystem Services*, 17, 172–184.
- Noe, G. B., & Hupp, C. R. (2005). Carbon, nitrogen and phosphorus accumulation in floodplains of Atlantic coastal plain rivers, USA. *Ecological Applications*, 15, 1178–1190.
- Noman, N. S., Nelson, E. J., & Zundel, A. K. (2001). Review of automated floodplain delineation from digital terrain models. *Journal of Water Resources Planning and Management*, 127(6), 394–402.
- O'Briain, R. (2019). Climate change and European rivers: An eco-hydromorphological perspective. *Ecology*, 12(5), e2099.
- Omer, C. R., Nelson, E. J., & Zundel, A. K. (2003). Impact of varied data resolution on hydraulic modeling and floodplain delineation. *Journal of the American Water Resources Association (JAWRA)*, 39(2), 467–475.
- Opperman, J. J., Luster, R., McKenney, B. A., Roberts, M., & Meadows, A. W. (2010b). Ecologically functional floodplains: Connectivity, flow regime, and scale 1. *JAWRA Journal of the American Water Resources Association*, 46(2), 211–226.
- Paillex, A., Dolédec, S., Castella, E., Mérigoux, S., & Aldridge, D. C. (2013). Functional diversity in a large river floodplain: Anticipating the response of native and alien macroinvertebrates to the restoration of hydrological connectivity. *Journal of Applied Ecology*, 50(1), 97–106.
- Park, E. (2020). Characterizing channel-floodplain connectivity using satellite altimetry: Mechanism, hydrogeomorphic control, and sediment budget. *Remote Sensing of Environment*, 243, 111783.
- Peh, K. S. H., Balmford, A., Field, R. H., Lamb, A., Birch, J. C., Bradbury, R. B., Brown, C., Butchart, S. H. M., Lester, M., Morrison, R., Sedgwick, I., Soans, C., Stattersfield, A. J., Stroh, P. A., Swetnam, R. D., Thomas, D. H. L., Walpole, M., Warrington, S., & Hughes, F. M. R. (2014). Benefits and costs of ecological restoration: Rapid assessment of changing ecosystem service values at a U.K. wetland. *Ecology and Evolution*, 4, 3875–3886.
- Peters, G. (2016). Identifying and valuing the functions of floodplains. Paper presented at The 2016 Floodplain Management Association National Conference (pp. 1–16).
- Petrášová-Šibíková, M., Matečný, I., Uherčíková, E., Pišút, P., Kubalová, S., Valachovič, M., & Medvecká, J. (2017). Effect of the Gabčíkovo waterworks (Slovakia) on riparian floodplain forest ecosystems in the Danube inland delta: Vegetation dynamics and trends. *Biologia*, 72(7), 722–734.
- PISRS. (2020). *Legal information system*. Legislation Office of the Government of the Republic of Slovenia. <http://www.pisrs.si/Pis.web/>
- Podschun, S. A., Thiele, J., Dehnhardt, A., Mehl, D., Hoffmann, T. G., Albert, C., & Costea, G. (2018). Das Konzept der Ökosystemleistungen-eine Chance für integratives Gewässermanagement. *Hydrologie und Wasserbewirtschaftung*, 62(6), 453–468.
- Poff, N. L., Allan, J. D., Bain, M. B., Karr, J. R., Prestegard, K. L., Richter, B. D., & Stromberg, J. C. (1997). The natural flow regime. *Bioscience*, 47(11), 769–784.
- Politti, E., Egger, G., Angermann, K., Rivaes, R., Blamauer, B., Klösch, M., Tritthart, M., & Habersack, H. (2014). Evaluating climate change impacts on alpine floodplain vegetation. *Hydrobiologia*, 737(1), 225–243.
- Posthumus, H., Rouquette, J. R., Morris, J., Gowing, D. J. G., & Hess, T. M. (2010). A framework for the assessment of ecosystem goods and services; a case study on lowland floodplains in England. *Ecological Economics*, 69(7), 1510–1523.
- Pusey, B. J., & Arthington, A. H. (2003). Importance of the riparian zone to the conservation and management of freshwater fish: A review. *Marine and Freshwater Research*, 54(1), 1–16.
- Rak, G., Grobljar, S., & Steinman, F. (2018). Flood modelling in urban areas. *Acta hydrotechnica*, 31(54), 21–33.

- Raška, P., Slavíková, L., & Sheehan, J. (2019). Scale in nature-based solutions for flood risk management. In T. Hartmann, L. Slavíková, & S. McCarthy (Eds.), *Nature-based flood risk management on private land* (pp. 9–20). Springer.
- Raška, P., Stehliková, M., Rybová, K., & Aubrechtová, T. (2019). Managing flood risk in shrinking cities: Dilemmas for urban development from the central European perspective. *Water International*, *44*(5), 520–538.
- Remm, L., Lohmus, A., Leibak, E., Kohv, M., Salm, J. O., Lohmus, P., Rosenvald, R., Runnel, K., Vellak, K., & Rannap, R. (2019). Restoration dilemmas between future ecosystem and current species values: The concept and a practical approach in Estonian mires. *Journal of Environmental Management*, *250*, 109439.
- Richards, D. R., Warren, P. H., Maltby, L., & Moggridge, H. L. (2017). Awareness of greater numbers of ecosystem services affects preferences for floodplain management. *Ecosystem Services*, *24*, 138–146.
- Robinson, C. T., Tockner, K., & Ward, J. V. (2002). The fauna of dynamic riverine landscapes. *Freshwater Biology*, *47*, 661–677.
- Rouquette, J. R., Posthumus, H., Morris, J., Hess, T. M., Dawson, Q. L., & Gowing, D. J. G. (2011). Synergies and trade-offs in the management of lowland rural floodplains: An ecosystem services approach. *Hydrological Sciences Journal*, *56*(8), 1566–1581.
- Salo, J., Kalliola, R., Hakkinen, L., Makinen, Y., Niemela, P., Puhakka, M., & Coley, P. D. (1986). River dynamics and the diversity of Amazon lowland forests. *Nature*, *322*, 254–258.
- Salvati, L., & Zitti, M. (2009). The environmental “risky” region: Identifying land degradation processes through integration of socio-economic and ecological indicators in a multivariate regionalization model. *Environmental Management*, *44*(5), 888.
- Salvati, L., Bajocco, S., Ceccarelli, T., Zitti, M., & Perini, L. (2011). Towards a process-based evaluation of land vulnerability to soil degradation in Italy. *Ecological Indicators*, *11*(5), 1216–1227.
- Salvati, L., Perini, L., Sabbi, A., & Bajocco, S. (2012). Climate aridity and land use changes: A regional-scale analysis. *Geographical Research*, *50*(2), 193–203.
- Sanon, S., Hein, T., Douven, W., & Winkler, P. (2012). Quantifying ecosystem service trade-offs: The case of an urban floodplain in Vienna, Austria. *Journal of Environmental Management*, *111*, 159–172.
- Schindler, S., Sebesvari, Z., Damm, C., Euller, K., Mauerhofer, V., Schneidergruber, A., Biró, M., Essl, F., Kanka, R., Lauwaars, S. G., Schulz-Zunkel, C., Sluis, T., Kropik, M., Gasso, V., Krug, A., Pusch, M. T., Zulka, K. P., Lazowski, W., Hainz-Renetzed, C., ... Wrbka, T. (2014). Multifunctionality of floodplain landscapes: Relating management options to ecosystem services. *Landscape Ecology*, *29*, 229–244.
- Schindler, S., O'Neill, F. H., Biró, M., Damm, C., Gasso, V., Kanka, R., Sluis, T., Krug, A., Lauwaars, S. G., Sebesvari, Z., Pusch, M., Baranovsky, B., Ehlert, T., Neukirchen, B., Martin, J. R., Euller, K., Mauerhofer, V., & Wrbka, T. (2016). Multifunctional floodplain management and biodiversity effects: A knowledge synthesis for six European countries. *Biodiversity and Conservation*, *25*(7), 1349–1382.
- Schneider, C., Flörke, M., Geerling, G., Duell, H., Grygoruk, M., & Okruszko, T. (2011). The future of European floodplain wetlands under a changing climate. *Journal of Water and Climate Change*, *2*(2–3), 106–122.
- Schober, B., Hauer, C., & Habersack, H. (2015). A novel assessment of the role of Danube floodplains in flood hazard reduction (FEM method). *Natural Hazards*, *75*(1), 33–50.
- Schönbrunner, I. M., Preiner, S., & Hein, T. (2012). Impact of drying and re-flooding of sediment on phosphorus dynamics of river-floodplain systems. *Science of the Total Environment*, *432*, 329–337.
- Scott, M. L., Shafroth, P. B., Auble, G. T., & Eggleston, E. D. (1997). Flood dependency of cottonwood establishment along the Missouri River, Montana USA. *Ecological Applications*, *7*, 677–690.
- Seják, J., Cudlín, P., Pokorný, J., Zapletal, M., Petříček, V., Guth, J., & Vyskot, I. (2010). *Hodnocení funkcí a služeb ekosystémů České republiky* (198 p). Ústí nad Labem: Fakulta životního prostředí, Univerzita J. E. Purkyně.
- Serna-Chavez, H. M., Swenson, N. G., Weiser, M. D., van Loon, E. E., Bouten, W., Davidson, M. D., & Van Bodegom, P. M. (2017). Strong biotic influences on regional patterns of climate regulation services. *Global Biogeochemical Cycles*, *31*(5), 787–803.
- Shields, F. D., Jr., Simon, A., & Steffen, L. J. (2000). Reservoir effects on downstream river channel migration. *Environmental Conservation*, *27*(1), 54–66.
- Smiraglia, D., Ceccarelli, T., Bajocco, S., Salvati, L., & Perini, L. (2016). Linking trajectories of land change, land degradation processes and ecosystem services. *Environmental Research*, *147*, 590–600.
- Smith, P., Ashmore, M. R., Black, H. I., Burgess, P. J., Evans, C. D., Quine, T. A., & Orr, H. G. (2013). The role of ecosystems and their management in regulating climate, and soil, water and air quality. *Journal of Applied Ecology*, *50*(4), 812–829.
- Souter, N. J., Wallace, T., Walter, M., & Watts, R. (2014). Raising river level to improve the condition of a semi-arid floodplain forest. *Ecohydrology*, *7*(2), 334–344.
- Sponseller, R. A., Heffernan, J. B., & Fisher, S. G. (2013). On the multiple ecological roles of water in river networks. *Ecosphere*, *4*(2), 1–14.
- Stacke, V., Pánek, T., & Sedláček, J. (2014). Late Holocene evolution of the Bečva River floodplain (Outer Western Carpathians, Czech Republic). *Geomorphology*, *206*, 440–451.
- Stanford, J. A., Lorang, M. S., & Hauer, F. R. (2005). The shifting habitat mosaic of river ecosystems. *Internationale Vereinigung für theoretische und angewandte Limnologie: Verhandlungen*, *29*(1), 123–136.
- Stella, J. C., Rodríguez-González, P. M., Dufour, S., & Bendix, J. (2013). Riparian vegetation research in Mediterranean-climate regions: Common patterns, ecological processes, and considerations for management. *Hydrobiologia*, *719*(1), 291–315.
- Štěrba, O. (Ed.). (2008). *Říční krajina a její ekosystémy* (p. 391). Univerzita Palackého v Olomouci (in Czech).
- Swinnen, W., Broothaerts, N., Hoevers, R., & Verstraeten, G. (2020). Anthropogenic legacy effects control sediment and organic carbon storage in temperate river floodplains. *Catena*, *195*, 104897.

- Swinnen, W., Daniëls, T., Maurer, E., Broothaerts, N., & Verstraeten, G. (2020). Geomorphic controls on floodplain sediment and soil organic carbon storage in a Scottish mountain river. *Earth Surface Processes and Landforms*, 45(1), 207–223.
- Szmańda, J. B., Lehotský, M., & Novotný, J. (2008). Sedimental record of flood events from years 2002 and 2007 in the Danube river overbank deposits in Bratislava. *Moravian Geographical Reports*, 16, 2–8.
- Thiele, J., von Haaren, C., & Albert, C. (2019). Are river landscapes outstanding in providing cultural ecosystem services? An indicator-based exploration in Germany. *Ecological Indicators*, 101, 31–40.
- Thiele, J., Albert, C., Hermes, J., & von Haaren, C. (2020). Assessing and quantifying offered cultural ecosystem services of German river landscapes. *Ecosystem Services*, 42, 101080.
- Tieskens, K. F., van Zanten, B. T., Schulp, C. J., & Verburg, P. H. (2018). Aesthetic appreciation of the cultural landscape through social media: An analysis of revealed preference in the Dutch river landscape. *Landscape and Urban Planning*, 177, 128–137.
- Tockner, K., & Stanford, J. A. (2002). Riverine flood plains: Present state and future trends. *Environmental Conservation*, 29(3), 308–330.
- Tockner, K., Bunn, S. E., Gordon, C., Naiman, R. J., Quinn, G. P., & Stanford, J. A. (2000). Flood Plains: Critically threatened ecosystems. In C. Gordon, T. Tockner, S. E. Bunn, G. P. Quinn, & J. A. Stanford (Eds.), *Aquatic ecosystems - trends and global prospects* (pp. 45–61). Cambridge University Press.
- Tomscha, S. A., Gergel, S. E., & Tomlinson, M. J. (2017). The spatial organization of ecosystem services in river-floodplains. *Ecosphere*, 8(3), 1–18.
- Turner, K. G., Anderson, S., Gonzales, C. M., Costanza, R., Courville, S., Dalgaard, T., & Ratna, N. (2016). A review of methods, data, and models to assess changes in the value of ecosystem services from land degradation and restoration. *Ecological Modelling*, 319, 190–207. e01728.
- Ureta, J. C., Zurqani, H. A., Post, C. J., Ureta, J., & Motallebi, M. (2020). Application of nonhydraulic delineation method of flood hazard areas using LiDAR-based data. *Geosciences*, 10, 338.
- van Andel, J., & Aronson, J. (Eds.). (2012). *Restoration ecology: The new frontier*. John Wiley & Sons.
- Vörösmarty, C. J., McIntyre, P. B., Gessner, M. O., Dudgeon, D., Prusevich, A., Green, P., & Davies, P. M. (2010). Global threats to human water security and river biodiversity. *Nature*, 467(7315), 555.
- Ward, J. V., Tockner, K., Arscott, D. B., & Claret, C. (2002). Riverine landscape diversity. *Freshwater Biology*, 47, 517–539.
- Weigelhofer, G., & Hein, T. (2015). Efficiency and detrimental side effects of denitrifying bioreactors for nitrate reduction in drainage water. *Environmental Science and Pollution Research*, 22(17), 13534–13545.
- Weigelhofer, G., Feldbacher, E., Trauner, D., Pözl, E., Hein, T., & Funk, A. (2020). Integrating conflicting goals of the EC water framework directive and the EC habitats directives into floodplain restoration schemes. *Frontiers in Environmental Science*, 8, 225.
- Welcomme, R. L. (1979). *Fisheries ecology of floodplain Rivers*. Longman.
- Welti, N., Bondar-Kunze, E., Tritthart, M., Pinay, G., & Hein, T. (2012). Nitrogen dynamics in complex Danube River floodplain systems: Effects of restoration. *River Systems*, 20(1–2), 71–85.
- West, P. C., Narisma, G. T., Barford, C. C., Kucharik, C. J., & Foley, J. A. (2011). An alternative approach for quantifying climate regulation by ecosystems. *Frontiers in Ecology and the Environment*, 9(2), 126–133.
- Westra, T., & De Wulf, R. (2006). Monitoring floodplain dynamics in the Sahel region to detect land degradation processes. Paper presented at the Proceedings of the 1st international conference on remote sensing and geoinformation processing in the Assessment and monitoring of land degradation and desertification. University of Trier, Trier.
- Whiting, P. J. (2002). Streamflow necessary for environmental maintenance. *Annual Review of Earth and Planetary Sciences*, 30(1), 181–206.
- Witner, D. B. (1966). Soils and their role in planning a suburban community. In L. J. Bartelli, A. A. Klingebiel, J. V. Baird, & M. R. Heddleson (Eds.), *Soil surveys and land use planning*. Soil Science Society of America and American Society of Agronomy (p. 15–30). Soil Science Soc. of Amer. and Amer. Soc. of Agronomy.
- Wohl, E., Hall, R. O., Jr., Lininger, K. B., Sutfin, N. A., & Walters, D. M. (2017). Carbon dynamics of river corridors and the effects of human alterations. *Ecological Monographs*, 87, 379–409.
- WRB (2015). World Reference Base for soil resources 2014, update 2015. International soil classification system for naming soils and creating legends for soil maps. World soil resources reports no. 106. FAO, Rome.
- Xu, M., Dong, X., Yang, X., Wang, R., Zhang, K., Zhao, Y., & Jeppesen, E. (2017). Using palaeolimnological data and historical records to assess long-term dynamics of ecosystem services in typical Yangtze shallow lakes (China). *Science of the Total Environment*, 584, 791–802.
- Zachary, H. B., Ken, D. B., & Terry, J. W. (2003). Effects of flow regulation on shallow-water habitat dynamics and floodplain connectivity. *Transactions of the American Fisheries Society*, 132, 809–823.

How to cite this article: Jakubínský, J., Prokopová, M., Raška, P., Salvati, L., Bezak, N., Cudlín, O., Cudlín, P., Purkyt, J., Vezza, P., Camporeale, C., Daněk, J., Pástor, M., & Lepeška, T. (2021). Managing floodplains using nature-based solutions to support multiple ecosystem functions and services. *Wiley Interdisciplinary Reviews: Water*, e1545. <https://doi.org/10.1002/wat2.1545>