POLITECNICO DI TORINO Repository ISTITUZIONALE

Recent results for the one-proton transfer reaction in the 18O+48Ti collision at 275 MeV

Original

Recent results for the one-proton transfer reaction in the 18O+48Ti collision at 275 MeV / Sgouros, O.; Cavallaro, M.; Cappuzzello, F.; Carbone, D.; Agodi, C.; Altana, C.; Brischetto, G. A.; Burrello, S.; Calabrese, S.; Calvo, D.; Capirossi, V.; Chávez Lomelí, E. R.; Ciraldo, I.; Cutuli, M.; De Gregorio, G.; Delaunay, F.; Djapo, H.; Eke, C.; Finocchiaro, P.; Fisichella, M.; Foti, A.; Gargano, A.; Hacisalihoglu, A.; Iazzi, F.; La Fauci, L.; Linares, R.; Lubian, J.; Medina, N. H.; Moralles, M.; Oliveira, J. R. B.; Pakou, A.; Pandola, L.; Pinna, F.; Russo, G.; Guazzelli, M. A.; Soukeras, V.; Souliotis, G.; Spatafora, A.; Torresi, D.; Yildirim, A.; Zagatto, V. A. B.. - In: EPJ WEB OF CONFERENCES. - ISSN 2100-014X. -EVENTRONICO. - 252 (2021), p. 04002 [10.1051/epiconf/202125204002] This version is available at: 11583/2931628 since: 2021-10-15110:24:112

Publisher: EDP Sciences

Published DOI:10.1051/epjconf/202125204002

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Recent results for the one-proton transfer reaction in the $^{18}\text{O+}^{48}\text{Ti}$ collision at 275 MeV

O. Sgouros^{1,*}, *M*. Cavallaro¹, *F*. Cappuzzello^{1,2}, *D*. Carbone¹, *C*. Agodi¹, *C*. Altana¹, *G*. *A*. Brischetto^{1,2}, *S*. Burrello^{3,4}, *S*. Calabrese¹, *D*. Calvo⁵, *V*. Capirossi^{5,6}, *E*. *R*. Chávez Lomelí⁷, *I*. Ciraldo^{1,2}, *M*. Cutuli^{1,2}, *G*. De Gregorio^{8,9}, *F*. Delaunay^{1,2,10}, *H*. Djapo¹¹, *C*. Eke¹², *P*. Finocchiaro¹, *M*. Fisichella¹, *A*. Foti¹³, *A*. Gargano⁸, *A*. Hacisalihoglu¹⁴, *F*. Iazzi^{5,6}, *L*. La Fauci^{1,2}, *R*. Linares¹⁵, *J*. Lubian¹⁵, *N*. *H*. Medina¹⁶, *M*. Moralles¹⁷, *J*. *R*. *B*. Oliveira¹⁵, *A*. Pakou¹⁸, *L*. Pandola¹, *F*. Pinna^{5,6}, *G*. Russo^{2,13}, *M*. *A*. Guazzelli¹⁹, *V*. Soukeras¹, *G*. Souliotis²⁰, *A*. Spatafora^{1,2}, *D*. Torresi¹, *A*. Yildirim²¹, and *V*. *A*. *B*. Zagatto¹⁵ for the NUMEN collaboration

¹INFN - Laboratori Nazionali del Sud, Catania, Italy

- ²Dipartimento di Fisica e Astronomia "Ettore Majorana", Università di Catania, Catania, Italy
- ³Université Paris-Saclay, CNRS/IN2P3, IJCLab, Orsay, France
- ⁴Technische Universität Darmstadt, Institut für Kernphysik, Darmstadt, Germany
- ⁵INFN Sezione di Torino, Torino, Italy
- ⁶DISAT Politecnico di Torino, Torino, Italy
- ⁷Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico
- ⁸INFN Sezione di Napoli, Napoli, Italy
- ⁹Dipartimento di Matematica e Fisica, Università della Campania "Luigi Vanvitelli", Caserta, Italy

¹⁰LPC Caen, Normandie Université, ENSICAEN, UNICAEN, CNRS/IN2P3, Caen, France

¹¹Ankara University, Institute of Accelerator Technologies, Turkey

- ¹²Department of Mathematics and Science Education, Faculty of Education, Akdeniz University, Antalya, Turkey
- ¹³INFN Sezione di Catania, Catania, Italy
- ¹⁴Institute of Natural Sciences, Karadeniz Teknik Universitesi, Trabzon, Turkey
- ¹⁵Instituto de Física, Universidade Federal Fluminense, Niterói, Brazil
- ¹⁶Instituto de Física, Universidade de São Paulo, São Paulo, Brazil
- ¹⁷Instituto de Pesquisas Energeticas e Nucleares IPEN/CNEN, São Paulo, Brazil

¹⁸Department of Physics, University of Ioannina and Hellenic Institute of Nuclear Physics, Ioannina, Greece

- ¹⁹Centro Universitario FEI, São Bernardo do Campo, Brazil
- ²⁰Department of Chemistry, University of Athens and Hellenic Institute of Nuclear Physics, Athens, Greece

²¹Department of Physics, Akdeniz Universitesi, Antalya, Turkey

Abstract. The ¹⁸O+⁴⁸Ti reaction was studied at the energy of 275 MeV for the first time under the NUMEN and NURE experimental campaigns with the aim to investigate the complete net of reaction channels potentially involved in the ⁴⁸Ca \rightarrow ⁴⁸Ti double charge exchange transition. Such a transition is of great interest because of its relevance to the extraction of ⁴⁸Ca \rightarrow ⁴⁸Ti double beta decay nuclear matrix element. The relevant experiment was carried out at the MAGNEX facility of INFN-LNS in Catania. Angular distribution measurements for the various reaction products were performed by using the MAGNEX large acceptance magnetic spectrometer. The present contribution is focused on the analysis of the one-proton transfer channel with emphasis on the particle identification technique and the estimation of background contaminations.

1 Introduction

The interest of the Nuclear Physics community for the Double Charge Exchange (DCE) reactions is still vivid due to their possible relation with the double beta ($\beta\beta$) decay [1]. Into this context, a novel idea has been recently conceived at Istituto Nazionale di Fisica Nucleare – Laboratori Nazionali del Sud (INFN-LNS), under the NUMEN (NUclear Matrix Elements for Neutrinoless double beta decay) project [2]. This is to use for the first time DCE reactions induced by heavy-ion beams as a mean for the

determination of the $\beta\beta$ decay Nuclear Matrix Elements (NMEs). This unique opportunity is based on the fact that the two processes present some interesting similarities. Among them, both processes probe the same initial and final nuclear wave functions and the operators connecting them have a similar spin – isospin mathematical structure [2–4]. Thus, even if the two processes are mediated by different interactions, the involved NMEs could be connected and the determination of the DCE reaction cross-sections may provide an important piece of information on the $\beta\beta$ matrix elements. In the present study, which is a part of the

^{*}e-mail: onoufrios.sgouros@lns.infn.it

NURE project [5], the ¹⁸O+⁴⁸Ti reaction was investigated by measuring not only the DCE channel, but also Single Charge Exchange (SCE) and multi-step transfer reactions which may lead to the same final states as the DCE one. It is very important to quantify possible contributions from other reaction channels to the DCE one for a precise determination of the absolute DCE cross sections, which may be the key for accessing the information of the NMEs of the $\beta\beta$ decay. In the present contribution, the data analysis of the one-proton transfer reaction ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc is presented. The experimental details are presented in Section 2, while the data reduction procedure is reported in Section 3. Our results are summarized in Section 4.

2 Experimental Setup

The experiment was performed at the MAGNEX facility [6] of INFN-LNS, where an ${}^{18}O^{8+}$ beam accelerated by the K800 Superconducting Cyclotron at the energy of 275 MeV impinged onto a TiO₂ target 510 μ g/cm² thick. An aluminum foil with a thickness of 216 μ g/cm² was used as a backing for the target material. In order to estimate the background originating from the different target components, two additional runs, one with a pure ²⁷Al target 226 μ g/cm² thick and a WO₃ one (284 μ g/cm²) evaporated on a thin aluminium foil, were also performed. The various reaction products were momentum analyzed by the MAG-NEX large acceptance magnetic spectrometer whose optical axis was set at $\theta_{opt} = 9^\circ$ with respect to the beam direction, spanning an angular range between 4° and 15° in the laboratory reference frame. The vertical slits of the MAG-NEX spectrometer, located 260 mm downstream the target position were set such as to cover a vertical angular range of $\pm 2^{\circ}$. The various reaction products emerging from the beam-target interaction were detected by the MAGNEX Focal Plane Detector (FPD) [7, 8].

The MAGNEX FPD is consisted of a gas section followed by a wall of 60 single silicon detectors. The use of the gas detector is twofold. It serves as a proportional drift chamber providing the energy loss signal (ΔE) of the ions inside the gas, but also as a mean to map the ions track. The tracker is divided in six sections each one having at the top a proportional wire (DC) in which the (ΔE) signal is measured. The sum of the 6 signals, ΔE_{tot} , is used in the particle identification process. Above the DC wires, a set of 6 segmented anode strips is located where each strip is composed of 224 induction pads allowing the measurement of the horizontal position (X_{foc}) and thus, the determination of the horizontal angle (θ_{foc}). Moreover, the electron drift time measurements inside the gas allow the determination of the vertical position (Y_{foc}) and angle (ϕ_{foc}). With the present gas tracker, the maximum rate that the detector can sustain is limited to a few kHz. However, this will be insufficient for the future NUMEN experiments at the MAGNEX facility with high intensity beams. To this extent, the development of a new configuration [9, 10] for the MAGNEX FPD is in progress based on the THick GEM (THGEM) technology for the new tracker.



Figure 1. Typical particle identification spectra for the ${}^{48}\text{Ti}({}^{18}\text{O}, {}^{19}\text{F}){}^{47}\text{Sc}$ reaction at the energy of 275 MeV. a) ΔE_{tot} as a function the of residual energy, E_{resid} , measured by one silicon detector of the FPD. A graphical selection of the fluorine ion events is illustrated by the solid black line. b) The horizontal position at the MAGNEX focal plane, X_{foc} , as a function of the residual energy gated by the selected fluorine ions of panel (a). The different loci correspond to ions with different \sqrt{m}/q . A graphical selection on the ${}^{19}\text{F}^{9+}$ events is shown by the solid black line.

3 Data Reduction

The data analysis relies on the accurate identification of the reaction channel of our interest. The particle identification (PID) is the first step of the data analysis which was performed following the prescription of Ref. [11]. The various ion species were well-discriminated using the conventional ΔE -E technique. A typical ΔE_{tot} - E_{resid} spectrum is presented in the top panel of Figure 1 for a single silicon detector. It is evident that the fluorine isotopes, highlighted with the black contour, are well separated from any other ion family. However, we are interested in identifying the one-proton pick up reaction products i.e. $^{19}F^{9+}$ so, the Z separation is not enough. To this direction, after the identification of the fluorine ions, the mass separation is feasible via the X_{foc} - E_{resid} (horizontal position versus



Figure 2. a) Correlation plot between the horizontal angle, θ_{foc} , and horizontal position, X_{foc} , measured by the focal plane detector of the MAGNEX spectrometer. b) A comparison between the experimental data of panel (a) with the simulated ones for three different reaction channels. The red, cyan and green points correspond to the simulated events originating from the one-proton transfer reaction on ²⁷Al, ⁴⁸Ti and ¹⁶O target, respectively. For reasons of clarity, the simulations were performed only for a few selective transitions for the one-proton transfer reaction on each target.

residual energy measured for each silicon detector) spectra. In a spectrometer, the position along the dispersive axis is analogous to the momentum of the ion. So, the position and the energy are related according to the following expression:

$$X_{foc} \propto \frac{\sqrt{m}}{q} \sqrt{E}_{resid},$$
 (1)

where m, q and E_{resid} are the ions' mass, charge state and residual energy, respectively. Ions possessing a different mass over charge state ratio lie on a different region at the X_{foc} - E_{resid} representation as it is illustrated in the lower panel of Figure 1. Looking at Figure 1 it is obvious that the mass resolution of MAGNEX is excellent. Thus, the ΔE -E technique in conjunction with mass separation method presented above, provide the root for an accurate identification of the one proton pick-up reaction channel.

Once the ¹⁹F⁹⁺ ions were identified, we have proceeded with the analysis of the the final phase space parameters. A typical θ_{foc} - X_{foc} spectrum obtained with the TiO₂+²⁷Al target is presented in Figure 2, where various loci are wellpronounced in the spectrum. Since a compound target was used in the current measurement, background events due to the reaction of the ¹⁸O beam with the oxygen and aluminium components of the target were also present in our spectra. In order to identify the fingerprint of each reaction, dedicated Monte Carlo simulations by taking into account the reaction kinematics but also the complete geometry of the spectrometer and the spatial distribution of the dipole and quadrupole fields were performed. The results of the simulations are compared to the experimental data in the lower panel of Figure 2. It is seen that the simulations describe in an excellent way the experimental data giving further support to the validity of the dipole and quadrupole fields which were adopted in the simulations. The different slope met both in experimental and simulated events is attributed partially to the different kinematics for each reaction and partially to the chromatic aberrations which are present in large-acceptance optical devices like MAGNEX. However, the latter are effectively compensated when a high-order software trajectory reconstruction [12] is applied to the data.



Figure 3. Reconstructed excitation energy spectrum for the ${}^{27}\text{Al}({}^{18}\text{O}, {}^{19}\text{F}){}^{26}\text{Mg}$ reaction at 275 MeV.

Having identified the background events, the analysis of the data obtained with the pure aluminum target and the WO₃+²⁷Al one was performed in order to estimate and subtract the contaminant events in the data set obtained with the TiO₂+²⁷Al target. A software ray reconstruction was applied to the data and the initial phase space parameters (e.g. θ_{lab} , kinetic energy) were reconstructed from the measured ones (e.g. θ_{foc} , ϕ_{foc} , X_{foc}). The excitation energy E_x was determined from the missing mass method [6] as:

$$E_x = Q_0 - Q, \tag{2}$$

where Q_0 is the ground state (g.s.) to g.s. transition. The reconstructed excitation energy spectrum for the data obtained with aluminum target is presented in Figure 3.



Figure 4. Bi-dimensional plot for the reconstructed scattering angle, θ_{lab} , as a function of the excitation energy, E_x , for the data obtained with the TiO₂+²⁷Al target. The inclined loci, highlighted with the dashed-green and long-dashed red curves, correspond to the one-proton transfer reaction of ¹⁸O with the oxygen and aluminum components of the target, respectively. The vertical loci correspond to the reconstructed events from the reaction on the ⁴⁸Ti target.

Various peaks associated to transition to the states of ¹⁹F and ²⁶Mg nuclei are well-formed in the spectrum up to the excitation energy of 10 MeV, where a rather continuum shape is observed due to the decay of ²⁶Mg nucleus ($^{26}Mg \rightarrow ^{22}Ne + \alpha$). A similar procedure was also followed for the data obtained with the $WO_3 + {}^{27}Al$ target and the excitation energy spectrum corresponding to the ¹⁶O(¹⁸O, ¹⁹F)¹⁵N reaction was deduced. After completing the analysis of the background runs, the same trajectory reconstruction technique was applied to the data obtained with the TiO₂+²⁷Al target. The reconstructed θ_{lab} - E_x plot is shown in Figure 4. Like in case of Figure 2, we can immediately identify in the spectrum the presence of the reaction contaminants. For reasons of clarity, we have pointed-out a few selected transitions. The inclined locus, highlighted with the red long-dashed curve, correspond to the g.s. of the ²⁷Al(¹⁸O,¹⁹F)²⁶Mg reaction, while the ones illustrated with the green dashed curves correspond to the g.s. and the 6.324 MeV $(3/2^{-})$ state of ¹⁵N from the ¹⁶O(¹⁸O,¹⁹F)¹⁵N reaction. The vertical oriented loci correspond to the ¹⁹F and ⁴⁷Sc states populated in the 48 Ti(18 O, 19 F) 47 Sc reaction. In order to isolate the energy spectrum for the one-proton transfer reaction on ⁴⁸Ti, after projecting the data of Figure 4 onto the E_x axis, we have superimposed to the mixed spectrum the previously analyzed data with the aluminum and oxygen targets, appropriately normalized. Subsequently, the background spectra were subtracted from the mixed one and the energy spectrum corresponding to the ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc reaction was deduced. The background subtraction procedure is presented in Figure 5. As it can be seen, the energy spectrum corresponding to the ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc reaction is rather structure-less as a result of the high density of states

of the ⁴⁷Sc nucleus. The data analysis for the reaction under study was recently completed and the results have been submitted for publication.



Figure 5. Decomposition of the excitation energy spectrum obtained with the $TiO_2+^{27}Al$ target. The total spectrum is presented with the solid black line. The blue-hatched area corresponds to the normalized background originating from the ²⁷Al backing material, while the brown solid line corresponds to the background arising from oxygen. The green solid line is the obtained excitation energy spectrum for the ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc reaction, after subtracting from the total spectrum the background contributions.

4 Summary

The study of the ¹⁸O+⁴⁸Ti reaction at 275 MeV was performed, for the first time, at INFN-LNS under the NU-MEN and NURE experimental campaigns. Angular distribution measurements for the ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc reaction were performed by using the MAGNEX large acceptance magnetic spectrometer. Complementary measurements with aluminum and $WO_3 + {}^{27}Al$ targets of the appropriate thickness were performed in order to estimate the background arising from the use of a TiO₂ target evaporated onto a thin aluminum foil. After subtracting the background contamination, the excitation energy spectrum corresponding to the ⁴⁸Ti(¹⁸O,¹⁹F)⁴⁷Sc reaction was deduced. Given the high density of states of the populated ⁴⁷Sc nucleus and the finite experimental energy resolution (about 500 keV FWHM) the measured excitation energy spectrum was rather structure-less. The results of this analysis will be the subject of a forthcoming publication.

Acknowledgments

The research leading to these results was partially funded by the European Research Council (ERC) under the European Union's Horizon 2020 Research and Innovation Programme (Grant Agreement No 714625).

References

- H. Ejiri, J. Suhonen and K. Zuber, Phys. Rep. 797, 1 (2019)
- [2] F. Cappuzzello, C. Agodi, M. Cavallaro et al., Eur. Phys. J. A 54, 72 (2018)
- [3] E. Santopinto, H. García-Tecocoatzi, R. I. Magaña Vsevolodovna et al., Phys. Rev. C 98, 061601(R) (2018)
- [4] H. Lenske, F. Cappuzzello, M. Cavallaro et al., Prog. Part. Nucl. Phys. 109, 103716 (2019)
- [5] M. Cavallaro, E. Aciksoz, L. Acosta et al., POS 302, (BORMIO2017) 015 (2017)
- [6] F. Cappuzzello, C. Agodi, D. Carbone et al., Eur. Phys. J. A 52, 167 (2016)

- [7] M. Cavallaro, F. Cappuzzello, D. Carbone et al., Eur. Phys. J. A 48, 59 (2012)
- [8] D. Torresi, O. Sgouros, V. Soukeras et al., Nucl. Instrum. Methods Phys. Res. A 989, 164918 (2021)
- [9] C. Agodi, A. D. Russo, L. Calabretta et al., Universe 7, 72 (2021)
- [10] F. Cappuzzello, L. Acosta, C. Agodi et al., Front. Astron. Space. Sci. 8, 668587 (2021)
- [11] F. Cappuzzello, M. Cavallaro, A. Cunsolo et al., Nucl. Instrum. Meth. A 621, 419 (2010)
- [12] F. Cappuzzello, D. Carbone, M. Cavallaro, Nucl. Instrum. Methods Phys. Res. A 638, 74 (2011)
- [13] M. Cavallaro, F.Cappuzzello, D.Carbone et al., Nucl. Instrum. Meth. A 637, 77 (2011)