

Higher order distance-like functions and Sobolev spaces

Original

Higher order distance-like functions and Sobolev spaces / Impera, Debora; Rimoldi, Michele; Veronelli, Giona. - In: ADVANCES IN MATHEMATICS. - ISSN 0001-8708. - 396:(2022), p. 108166. [10.1016/j.aim.2021.108166]

Availability:

This version is available at: 11583/2943972 since: 2021-12-09T23:29:25Z

Publisher:

Elsevier

Published

DOI:10.1016/j.aim.2021.108166

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.aim.2021.108166>

(Article begins on next page)

HIGHER ORDER DISTANCE-LIKE FUNCTIONS AND SOBOLEV SPACES

DEBORA IMPERA, MICHELE RIMOLDI, AND GIONA VERONELLI

ABSTRACT. On non-compact Riemannian manifolds, we construct distance-like functions with derivatives controlled up to some order k assuming bounds on the growth of the derivatives of the curvature up to order $k - 2$ and on the decay of the injectivity radius. This construction extends previously known results in various directions, permitting to obtain consequences which are (in a sense) sharp. As a first main application, we give refined conditions guaranteeing the density of compactly supported smooth functions in the Sobolev space $W^{k,p}$ on the manifold. Contrary to all previously known results this can be obtained also on manifolds with possibly unbounded geometry. In the particular case $p = 2$, making use of the Weitzenböck formula for a Lichnerowicz Laplacian acting on k -covariant totally symmetric tensor fields, we can weaken the assumptions needed to obtain the density property, avoiding any condition on the highest order derivatives of the curvature. Distance-like functions are also used to obtain new disturbed Sobolev inequalities, disturbed L^p -Calderón-Zygmund inequalities and the full Omori-Yau maximum principle for the Hessian under weak assumptions.

CONTENTS

1. Introduction and main results	2
1.1. Higher order distance-like functions	2
1.2. The density property for Sobolev spaces	4
1.3. Sampson-Lichnerowicz Laplacian and the special case $W^{k,2}(M)$	6
1.4. Other applications	7
1.5. Structure of the paper	7
2. Some notational conventions	7
3. Higher order distance-like functions	8
3.1. On the assumptions	8
3.2. A generalization of a result by Bianchi-Setti	10
3.3. Proof of Theorem 1.2: case (a)	13
3.4. Proof of Theorem 1.2: case (b)	21
4. k -th order cut-offs and application to the density problem	22
4.1. Proof of the density result	24
5. Case $p=2$	24
5.1. k -th order (rough) Laplacian cut-offs	24
5.2. Weitzenböck formulas	28
5.3. A Lichnerowicz Laplacian on symmetric $(0, k)$ -tensors	29
5.4. Proof of Theorem 1.7	29
6. Some sharp applications	34
6.1. Disturbed Sobolev inequalities	34
6.2. Calderón-Zygmund inequalities	35
6.3. Omori-Yau maximum principle	38
Appendix A. Some commutation formulas	39

Date: December 9, 2021.

2010 Mathematics Subject Classification. 46E35, 53C21.

Key words and phrases. Sobolev spaces on manifolds, Distance-like functions, Cut-off functions, Sampson-Lichnerowicz Laplacian, Calderón-Zygmund inequalities.

Appendix B. Weitzenböck formula for Δ_{Sym}	40
References	41

1. INTRODUCTION AND MAIN RESULTS

1.1. Higher order distance-like functions. Let (M, g) be a complete non-compact Riemannian manifold and let $r(x)$ be the distance from $x \in M$ to a fixed reference point $o \in M$. The behaviour of the function $r(x)$ detects how much the manifold differs from the Euclidean space. It turns out that in general $r(x)$ is 1-Lipschitz on M , but only a.e. differentiable. This happens as soon as the point o has a cut-locus, and in particular whenever M has non-trivial topology. This lack of regularity of r represents an obstacle to perform a certain kind of smooth analysis. In order to overcome this problem, it is often enough to use, in place of r , a suitable smooth approximation of the distance function. A first evidence in this direction is given by a well-known result by M. P. Gaffney [9] (often attributed in the literature to R. E. Greene and H. Wu, [11]), who showed the existence of a function $H(x)$ on M which is smooth, distance-like (i.e. $C^{-1}r(x) < H(x) < Cr(x)$ outside a compact set) and whose gradient has norm uniformly bounded by 1. This can be obtained by smoothing r in local coordinate charts by mollification.

In order to adapt to Riemannian manifolds certain techniques of C^k (Euclidean) analysis, one would like to have at disposal on M a smooth distance-like functions of which one controls not only the gradient, but also higher order (covariant) derivatives up to the order k . It is thus a natural question to what extent one can generalize Gaffney's result, providing sufficiently general geometric assumptions which guarantee the existence of a distance-like function with controlled higher order derivatives.

A first achievement in this direction is due to J. Cheeger and M. Gromov, [7] (see also [21] for a detailed proof), who remarked that a uniform double-sided bound on the sectional curvatures of M permits to get a uniform bound also on the norm of the second order derivatives of Gaffney's first-order distance-like function through the local smoothing process, thanks to the Hessian comparison. More recently, L. F. Tam provided in [28] a new completely different proof of Cheeger-Gromov's result: start with the first-order distance-like function by Gaffney-Greene-Wu and let it evolve by the heat flow of M ; the regularizing property of the heat flow, together with the sectional curvature bound, permits to control also the second order derivatives along the evolution, so that the evolution at a fixed time (say $t = 1$) gives the aimed second-order distance-like function. Contrary to the proof by Cheeger-Gromov, Tam's approach is more flexible, in the sense that it can be adapted to other sets of assumptions. This was first observed by the second and third authors in [25], where the existence of a second order-distance like function was obtained when the underlying manifold has (double-sided) bounded Ricci curvature and positive injectivity radius. For a further recent result, see also [19], where the assumption of positive injectivity radius is replaced by the request that the volumes of unitary balls are almost Euclidean. This seems a pretty strong assumption, which however a priori could not imply $\text{inj}_g(M) > 0$ in general.

As we will see later, for most applications the uniform bound on the second-order derivatives is a property stronger than what is really needed, in that we can actually allow derivatives to explode at infinity, provided that their growth is appropriately controlled. Starting from this observation, in the previous paper [20] we found conditions which ensure the existence of a distance-like function with uniformly bounded gradient and sub-linear growth of the second-order derivatives. Namely, this holds true provided that: a) the norm of the Riemann tensor grows sub-quadratically or b) the norm of the Ricci tensor grows sub-quadratically and the injectivity radius can possibly vanish at infinity, but no faster than $1/r(x)$. In order to obtain this improvement, we introduced a new strategy: the distance-like function is no more given by a solution of the linear homogeneous parabolic heat flow, but instead by the solution of a non-linear non-homogeneous elliptic equation.

Concerning the existence of higher order distance-like functions (i.e. with controlled derivatives up to the order $k > 2$), up to our knowledge, the only known achievement is due to L. F. Tam (see [8, Remark 26.50]) who proved the following

Proposition 1.1. *Let $k \geq 2$. Let (M^m, g) be a complete Riemannian manifold such that, for some $D > 0$, we have $|\nabla^j \text{Riem}| \leq D^2$ for $j = 0, \dots, k-2$. Then there exists a distance-like function $H \in C^\infty(M)$ such that $|\nabla^j H| \leq C$, for $j = 1, \dots, k$, with C a positive constant depending only on m, k, D .*

Let $\lambda : [0, +\infty) \rightarrow [0, +\infty)$ be a C^∞ function. Before stating the first main result of this paper, let us introduce the following assumptions:

(A1) There exists a constant $R_1 > 0$ such that λ is strictly positive and non-decreasing on $[R_1, +\infty)$.

(A2) For every $\delta > 0$, there exist constants $R_2 = R_2(\delta) > 0$ and $M_2 = M_2(\delta) > 1$ such that

$$\forall t > R_2, \quad M_2(\delta)^{-1} \leq \frac{\lambda(\delta t)}{\lambda(t)} \leq M_2(\delta), \quad \text{and} \quad M_2(\delta)^{-1} \leq \frac{\lambda'(\delta t)}{\lambda'(t)} \leq M_2(\delta).$$

(A3) There exist constants $R_3 > 0$ and $M_3 > 1$ such that

$$\forall t > R_3, \quad M_3^{-1} \leq \frac{t\lambda'(t)}{\lambda(t)} \leq M_3.$$

Moreover, for every integer $j \geq 1$ we introduce the assumptions

(A4(j)) There exist constants $R_{4(j)} > 0$ and $M_{4(j)} > 1$ such that

$$\forall t > R_{4(j)}, \quad \frac{t\lambda^{(j)}(t)}{\lambda^j(t)} \leq M_{4(j)}.$$

The first main theorem of this paper is the following.

Theorem 1.2. *Let (M, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}(x, o)$. Let $k \geq 2$ be an integer. Let λ satisfy the assumptions (A1), (A2), (A3) and (A4(j)), $j = 1, \dots, k$. Suppose that one of the following curvature assumptions holds*

(a) *for some $i_0 > 0$,*

$$|\nabla^j \text{Ric}|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{inj}_g(x) \geq \frac{i_0}{\lambda(r(x))} > 0 \quad \text{on } M.$$

(b)

$$|\nabla^j \text{Ric}|(x) \leq \lambda(r(x))^{2+j}, \quad 1 \leq j \leq k-2, \quad |\text{Sect}|(x) \leq D^2 \lambda(r(x))^2.$$

Then there exists an exhaustion function $H \in C^\infty(M)$ such that for some positive constant $C > 1$ independent of x and o , we have on M that

(i) $C^{-2}r(x) \leq H(x) \leq \max\{r(x), 1\}$;

(ii) $|\nabla H|(x) \leq 1$;

(iii) for $2 \leq j \leq k$, $|\nabla^j H|(x) \leq C \max\{\lambda(r(x))^{j-1}, 1\}$.

More comments about assumptions (A1) to (A4) will be given in Subsection 3.1. For the moment let us only point out that admissible choices for λ which are relevant in the applications are, for instance, constant multiples of

$$\lambda(r(x)) = r(x)^\eta, \quad \eta \geq 0, \quad \text{or} \quad \lambda(r(x)) = r(x) \prod_{i=1}^{\bar{i}} \ln^{[i]}(r(x))$$

where $\ln^{[i]}$ stands for the i -th iterated logarithm (e.g. $\ln^{[2]}(t) = \ln \ln t$, etc.) and \bar{i} is some positive integer. See Lemma 3.4 for more details.

Theorem 1.2 extends the previously known results in several directions. Namely note that:

- The Proposition 1.1 due to Tam is still true if we assume only $|\nabla^j \text{Ric}| \leq D^2$ and either a uniform bound on all the sectional curvatures or the positivity of the injectivity radius;
- as in [20], we can assume a controlled growth of the curvatures and/or a controlled decay of the injectivity radius and get an estimate for the growth of the derivatives, which is enough for some important applications, as for instance for proving the density of compactly supported functions in Sobolev spaces;
- we can admit more general (non-necessarily sub-polynomial) growth functions. This generalizes also the second order result obtained in [20], and permits to get consequences which are (in a sense) sharp. See for instance Remark 1.6 and Subsection 6.3.

We present here the strategy of the proof of Theorem 1.2 in the special easy case where $\lambda(r(x)) = r(x)$. Reasoning by induction, suppose that we have a distance-like function H_{k-1} which satisfies $|\nabla^j H_{k-1}| \leq r(x)^{j-1}$ outside a compact set for $j = 1, \dots, k-1$. According to (a generalization of) a theorem due to D. Bianchi and A. G. Setti, [6], we can prove the existence of a smooth function h such that

- (i) $C_h^{-1} r^2(x) \leq h(x) \leq C_h r^2(x)$;
- (ii) $|\nabla h| \leq C_h r(x)$;
- (iii) $\Delta h = |\nabla h|^2 - C_\theta H_{k-1}^2(x)$,

outside a compact set; see Theorem 3.5. Suppose that we already have obtained an estimate for $|\nabla^j h|$ for $j = 2, \dots, k-1$ (formally, this is done via a second induction process). Derivating the equation at (iii), we get

$$(1) \quad \nabla^{k-2} \Delta h = \nabla^{k-2} |\nabla h|^2 - C_\theta \nabla^{k-2} H_{k-1}^2(x).$$

The curvature and injectivity radius assumptions, together with an appropriate local rescaling of the metric, guarantee the existence of a harmonic atlas with respect to which the metric is uniformly controlled in $C^{k-1, \alpha}$. Accordingly, in any local harmonic coordinate chart Ω , equation (1) writes $\Delta_0 \nabla^{k-2} h = F$ on Ω , where Δ_0 is the Euclidean Laplacian and F a certain (vector-valued) function which depends on $\|g\|_{C^{k-1, \alpha}(\Omega)}$, $\|h\|_{C^{k-1}(\Omega)}$ and $\|H_{k-1}\|_{C^{k-1}(\Omega)}$. Since all these three norms are controlled, we are in the position to implement standard Euclidean elliptic theory and deduce a $C^{2, \alpha}$ control on $\nabla^{k-2} h$, that is, a $C^{k, \alpha}$ control on $\nabla^k h$. The desired function H is finally obtained letting $H = h^2$.

1.2. The density property for Sobolev spaces. As alluded to above, distance-like functions guarantee that the underlying manifold M is not too different from the Euclidean space (in a suitable sense). Namely, when M supports a distance-like function with controlled derivatives, certain properties and tools from classic analysis on \mathbb{R}^n can be proved to have a Riemannian counterpart on M . In this sense, one of the main applications is to density results of smooth compactly supported functions in Sobolev spaces, as probably first observed in [13], [16].

Given (M^m, g) a smooth, complete, possibly non-compact Riemannian manifold without boundary, k an integer, and $p \geq 1$, one can define the Sobolev space $W^{k, p}(M)$ as the space of functions on M all of whose (weak) derivatives of order 0 to k have finite L^p norm. By a generalised Meyers-Serrin-type theorem (see e.g. [12]) this coincides with the completion of the space

$$C_k^p(M) \doteq \left\{ u \in C^\infty(M) : \int_M |\nabla^j u|^p d\text{vol} < +\infty, \forall j = 0, \dots, k \right\}$$

with respect to the norm

$$\|u\|_{k, p} = \sum_{j=0}^k \left(\int_M |\nabla^j u|^p d\text{vol} \right)^{\frac{1}{p}}.$$

Moreover, we can define $W_0^{k, p}(M) \subseteq W^{k, p}(M)$ as the closure of the space of smooth compactly supported functions $C_c^\infty(M)$ with respect to the same norm.

In case $M = \mathbb{R}^m$, it is well known that $W_0^{k,p}(\mathbb{R}^m) = W^{k,p}(\mathbb{R}^m)$ for all k integer and $p \in [1, \infty)$, so that one could expect the same equivalence to hold true also on complete Riemannian manifold M . However, quite surprisingly, the problem seems to remain open in general so far; see for instance [18, p. 49].

Problem 1.3. *Given an arbitrary complete Riemannian manifold (M, g) , is it true that*

$$(2) \quad W_0^{k,p}(M) = W^{k,p}(M)$$

for all integer $k \geq 0$ and $p \in [1, \infty)$?

Of course, the answer is positive under certain additional assumptions. First, (2) is satisfied for all k and p when M is compact; see for instance [4, Theorem 2.9]. Concerning complete non-compact manifolds, it is a standard fact that $W_0^{0,p}(M) = W^{0,p}(M) = L^p(M)$, and with some effort one can also prove that $W_0^{1,p}(M) = W^{1,p}(M)$ for all k and $p \in [1, \infty)$; [3].

Regarding the first non-trivial order, i.e. $k = 2$, in our previous work [20] we showed that $W_0^{2,p}(M) = W^{2,p}(M)$, $p \in [1, +\infty)$, for complete manifolds with either a sub-quadratic growth of the norm of the Riemann curvature, or a subquadratic growth of both the norm of the Ricci curvature and the squared inverse of the injectivity radius. Previously known results, obtained in [18], [14], were assuming uniform constant bounds on either the Ricci tensor and the injectivity radius or the Riemann tensor. As far as concerns higher order Sobolev spaces the most up to date result is the following proposition due to E. Hebey, [18].

Proposition 1.4 (Proposition 3.2 in [18]). *Let (M, g) be a smooth, complete Riemannian manifold with positive injectivity radius, and let $k \geq 3$ be an integer. We assume that for $j = 0, \dots, k - 2$, $|\nabla^j \text{Ric}|$ is bounded. Then for any $p \in [1, \infty)$, $W_0^{k,p}(M) = W^{k,p}(M)$.*

As a first main application of Theorem 1.2, we will prove the density property on complete Riemannian manifolds with a suitable growth condition on the derivatives of the Ricci tensor and either the Riemannian curvature tensor or the inverse of the injectivity radius. The admissible growth rate depends explicitly on the order of the Sobolev space we are dealing with. Note that, contrary to Proposition 1.4, the following Theorem 1.5 concerns also manifolds with possibly unbounded geometry.

Theorem 1.5. *Let $k \geq 2$. In the assumptions of Theorem 1.2, if we further assume that $\lambda^{1-k} \notin L^1([1, +\infty))$, then we have that $W^{k,p}(M) = W_0^{k,p}(M)$ for all $p \in [1, +\infty)$.*

Remark 1.6. Choosing the function λ in Theorem 1.5 in essentially the best admissible way, we get that $W^{k,p}(M) = W_0^{k,p}(M)$ for all $p \in [1, +\infty)$ for instance if

$$|\nabla^j \text{Ric}|(x) \leq \left(r(x) \prod_{i=1}^{\bar{i}} \ln^{[i]}(r(x)) \right)^{(2+j)/(k-1)}, \quad 0 \leq j \leq k - 2,$$

and either

- (a) for some $i_0 > 0$, $\text{inj}(x) \geq i_0 \left(r(x) \prod_{i=1}^{\bar{i}} \ln^{[i]}(r(x)) \right)^{-1/(k-1)} > 0$, or
- (b) for some $D > 0$, $|\text{Riem}|(x) \leq D^2 \left(r(x) \prod_{i=1}^{\bar{i}} \ln^{[i]}(r(x)) \right)^{2/(k-1)}$.

Accordingly, this theorem improves also our previous [20, Theorem 1.4] when $k = 2$, permitting to achieve the density result $W_0^{2,p}(M) = W^{2,p}(M)$ on manifolds whose curvatures can grow more than quadratically.

To get a flavour of the proof of Theorem 1.5, consider here the easy case $k = 3$ and $\lambda(r) = r^{1/2}$ (so that $\lambda^{1-k} \notin L^1([1, +\infty))$). Following a standard construction, let $\phi_R \in C_c^\infty(\mathbb{R}_+)$ be a family

of cut-off functions such that $\phi_R \equiv 1$ on $[0, R]$ and $|\phi_R^{(j)}| \leq CR^{-j}\mathbb{1}_{[R, 2R]}$. Let H be the 3rd-order distance-like function given by Theorem 1.2 and define $\chi_R \doteq \phi_R \circ H$. Then $\{\chi_R\}_{R>1}$ is a family of 3rd-order cut-off functions, namely $|\nabla^j \chi_R|$ are uniformly bounded as $R \rightarrow \infty$ for $j = 1, 2, 3$. Once one has 3rd-order cut-off functions, it is a standard fact that $f \in W^{3,p}(M)$ can be approached by $\chi_R f$ in $W^{3,p}(M)$ -norm. In this process, the main term to control is of the form $|\phi_R'(H(x))\nabla^3 H(x)| \lesssim R^{-1}\lambda^2 = R^{-1}R = 1$. Here, this latter control was deduced from uniform estimates which we have separately on $\phi_R'(H(x))$ and on $\nabla^3 H(x)$ in $B_{2R}(o) \setminus B_R(o)$. One can look instead for a more careful point-wise control. Namely one imposes that $|\phi_R'(H(x))|$ is smaller and smaller as $H(x) \rightarrow \infty$ in order to compensate the growth of $\nabla^3 H(x)$. This leads us to a more careful construction of the real cut-offs ϕ_R , in such a way that $|\phi_R'(H(x))| \sim \frac{1}{\lambda^2(H(x))} \sim \frac{1}{|\nabla^3 H(x)|}$. This can be done as soon as $\lambda^{1-k} \notin L^1((1, +\infty))$, and permits us to get the results under the sharper assumptions which we described in Remark 1.6.

1.3. Sampson-Lichnerowicz Laplacian and the special case $W^{k,2}(M)$. In the particular case $p = 2$, the assumptions of Theorem 1.5 can be weakened. For $k = 2$, L. Bandara [5] proved that $W^{2,2}(M) = W_0^{2,2}(M)$ under the sole lower bound $\text{Ric}(x) \geq -C$ for some constant $C > 0$. In [20], we pointed out that in fact $\text{Ric}(x) \geq -Cr^2(x)$ suffices. The point is that in these weaker assumptions one can still prove the existence of a distance-like function H with $|\Delta H| \leq Cr^2$, and hence of Laplacian cut-offs with $|\Delta \chi_n|$ uniformly bounded, [6]. Given $f \in W^{2,2}(M)$, the Weitzenböck formula applied to the Hodge Laplacian acting on the (skew-symmetric) one form $d(\chi_n f)$ permit to control $\|f \nabla^2 \chi_n\|_{L^2}$ in terms of $\|f \Delta \chi_n\|_{L^2}$, and thus to prove that $\chi_n f \rightarrow f$ in $W^{2,2}(M)$. Generalizing this approach to higher order Sobolev spaces is non-trivial. Indeed, one is led to apply Bochner techniques to the $(0, k-1)$ -tensor $\nabla^{k-1}(\chi_n f)$, which is very far from being skew-symmetric when $k > 2$. Accordingly, the Bochner formula for the Hodge Laplacian acting on k -forms can not be exploited. However, it turns out that $\nabla^2(\chi_n f) = \text{Hess}(\chi_n f)$ is symmetric. This is no more true for $k > 3$, but $\nabla^{k-1}(\chi_n f)$ remains almost symmetric, in the sense that it has a symmetric principal term, plus lower order terms which can be controlled. Thus, one can apply an analogous technique to J. H. Sampson's Laplacian Δ_{Sym} , [26]. This latter is a Laplace operator acting on the space of symmetric $(0, k)$ -tensors which

- (a) is a Lichnerowicz Laplacian, i.e. it satisfies a Weitzenböck formula

$$\Delta_L = \Delta_B + c\mathfrak{Ric},$$

where c is a suitable (in this case negative) constant, Δ_B is the Bochner Laplacian and \mathfrak{Ric} is the Weitzenböck curvature operator (see Subsection 5.2 for more details).

- (b) has a Hodge-type decomposition $\Delta_{\text{Sym}} = D_S^* D_S - D_S D_S^*$, where D_S is the symmetrized covariant derivative, and D_S^* its formal adjoint.

Exploiting this, one can get a control on the L^2 -norm $\|f \nabla^k \chi_n\|_{L^2}$ in terms of $\|f \Delta \nabla^{k-2} \chi_n\|_{L^2}$. Choosing the cut-offs χ_n to be what we called k -th order (rough) Laplacian cut-offs, this latter term can be estimated under assumptions which are weaker than the assumptions of Theorem 1.5 (in particular we do not need to control the highest order derivatives of the curvature). More precisely, in Section 5 we will prove the following.

Theorem 1.7. *Let (M, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point $r(x) \doteq \text{dist}_g(x, o)$. Let $k \geq 3$ be an integer. Let λ satisfy assumptions (A1), (A2), (A3), and (A4(j)) for $j = 1, \dots, k-1$ above, and suppose that $\lambda^{1-k} \notin L^1([1, +\infty))$. Suppose that*

$$|\nabla^j \text{Riem}| \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-3.$$

Then we have that $W_0^{k,2}(M) = W^{k,2}(M)$.

1.4. Other applications. Beyond the density problem for Sobolev spaces, other results can be obtained by means of Theorem 1.2, notably when $k = 2$, i.e. for distance-like functions with controlled gradient and Hessian. Some of these applications were already observed in [20], including disturbed Sobolev inequalities, L^2 -Calderón-Zygmund inequalities and the full Omori-Yau maximum principle for the Hessian. Since Theorem 1.2 improves [20, Theorem 1.5] also when $k = 2$, by allowing for a wider class of admissible growth functions λ , the range of application of the results alluded to above is enlarged in this paper. This will be made explicit respectively in Theorem 6.1, Theorem 6.4 and Subsection 6.3. On the way, we will take also the occasion to point out a couple of new applications of the 2nd-order distance-like functions which were not contained in [20]. The first one is a higher order version of the disturbed Sobolev inequality, i.e.

$$\|\varphi\|_{L^{pm/(m-kp)}(M)} \leq A \sum_{r=-1}^{k-1} \left\| (\max\{1; r(x)\})^{\frac{\eta}{mp}(r+1)(2m-rp)} |\nabla^{k-r-1}\varphi| \right\|_{L^p(M)},$$

for all $\varphi \in C_c^\infty(M)$. Quite surprisingly, compared to the case $k = 1$, this does not require any additional assumption; see Proposition 6.3. The second new application is the following disturbed global L^p Calderón-Zygmund inequality for $p \neq 2$.

Theorem 1.8. *Let (M^m, g) be a smooth, complete non-compact Riemannian manifold without boundary. Let $o \in M$, $r(x) \doteq \text{dist}_g(x, o)$ and suppose that for some $\eta > 0$, some $D > 0$ and some $i_0 > 0$,*

$$|\text{Sect}|(x) \leq D^2(1 + r(x)^2)^\eta, \quad \text{inj}_g(x) \geq \frac{i_0}{D(1 + r(x))^\eta} > 0 \quad \text{on } M.$$

Then there exist constants $A > 0$ depending on m, η, D, i_0 and the constant C from Theorem 1.2, such that for all $\varphi \in C_c^\infty(M)$ it holds

$$\|\text{Hess } \varphi|_g\|_{L^p}^p \leq A \left[\|H^{2\eta}\varphi\|_{L^p}^p + \|\Delta\varphi\|_{L^p}^p \right],$$

where $H \in C^\infty(M)$ is the distance-like function given by Theorem 1.2, which satisfies in particular $H(x) \leq \max\{1; r(x)\}$.

Compared to the L^2 -case alluded to above, the main difficulty here is to control the injectivity radius of the manifold at hand under a conformal deformation of the metric.

1.5. Structure of the paper. The rest of the paper is structured as follows. In Section 2, for the reader's convenience, we list some notations which will be used throughout the paper. Section 3 is devoted to the construction of higher order distance-like functions described in Theorem 1.2. These will be then used in Section 4 to construct the cut-off functions needed for the proof of the density property for Sobolev spaces stated in Theorem 1.5 which will be proved in the final part of the section. In Section 5 we focus on the special case $p = 2$. We construct the k -th order rough Laplacian cut-off functions, introduce Sampson's Lichnerowicz Laplacian acting on the space of smooth sections of symmetric k -covariant tensors and its Weitzenböck formula, and give a proof of the density property stated in Theorem 1.7. The other applications of the higher order distance-like functions we mentioned above will be presented in Section 6, which contains in particular a proof of Theorem 1.8. For the sake of completeness, we end the paper with two appendices. In Appendix A, we give a proof of some commutation formulas which we use in Section 5, while in Appendix B we explicitly prove the Weitzenböck formula for Sampson's Laplacian.

2. SOME NOTATIONAL CONVENTIONS

The following are some notations and conventions which will be used throughout the paper.

For any $\beta > 0$, the Euclidean ball of radius β centered at the origin will be denoted by \mathbb{B}_β . Given a smooth enough function $\lambda : \mathbb{R} \rightarrow \mathbb{R}$, its j -th derivative will be denoted by $\lambda^{(j)}$. Moreover, using standard notation, given a smooth enough function $F : \mathbb{R}^m \rightarrow \mathbb{R}$ we will denote by DF the

Euclidean gradient of F and, given a multi-index $\gamma = (\gamma_1, \dots, \gamma_q) \in \{1, \dots, m\}^q$, we will denote $|\gamma| = q$ and $\partial_\gamma F = \partial_{\gamma_1 \dots \gamma_q} F \doteq \partial_{\gamma_1} \dots \partial_{\gamma_q} F$.

Given (M^m, g) an m -dimensional complete Riemannian manifold, we will denote by $\nabla_g, \Delta_g \doteq \operatorname{div}_g \nabla_g$ and ∇_g^j , respectively, the corresponding Riemannian gradient, the (negative definite) Laplace-Beltrami operator and the natural j -th covariant derivative, thus specifying in the subscript the metric we are considering. Given a smooth enough $f : M \rightarrow \mathbb{R}$, the Hessian of f , i.e. $\nabla_g^2 f$, will be also denoted by $\operatorname{Hess}_g f$. An analogous convention will be used for various concepts associated to the metric such as the geodesic distance $d_g(\cdot, \cdot)$, open geodesic balls $B_r^g(x)$ of radius r centered at a point $x \in M$, the injectivity radius $\operatorname{inj}_g(x)$ at a point $x \in M$, the global injectivity radius $\operatorname{inj}_g(M)$, and the volume measure $d\operatorname{vol}_g$. Likewise, we will denote the Riemann curvature tensor of (M, g) by Riem_g , the Ricci tensor by Ric_g and the Sectional curvature of a two dimensional subspace $\pi \subset T_x M$ by $\operatorname{Sect}_g(\pi)$. However, we will omit the subscript metric when the meaning is clear, as for instance for the fiber norms induced by g (as in the expression $|\nabla_g f|$), as well as in all the Section 5 and in the Appendix A and B, where we are considering a unique Riemannian metric g on M .

Note also that when writing an inequality of the form $|\operatorname{Sect}_g|(x) \leq a(x)$ on M we will mean that all the sectional curvatures at the point x are dominated, in absolute value, by $a(x)$ for every $x \in M$.

In some parts of the paper, especially in Section 5 and in the appendices, some of the formulae which we will need can be significantly simplified using the following "*" notation, used e.g. in [29]. We denote by $A * B$ any tensor field which is a real linear combination of tensor fields, each formed by starting with the tensor field $A \otimes B$, using the metric to switch the type of any number of T^*M components to TM components, or vice versa, taking any number of contractions, and switching any number of components in the product.

Finally, all over the paper, C will denote real constants, whose explicit value can possibly change from line to line. When it will seem appropriate we will specify some dependences of these constants at the subscript.

3. HIGHER ORDER DISTANCE-LIKE FUNCTIONS

3.1. On the assumptions. Let $\lambda : [0, +\infty) \rightarrow [0, +\infty)$ be a C^∞ function. In Section 1 we introduced the following assumptions:

(A1) There exists a constant $R_1 > 0$ such that λ is strictly positive and non-decreasing on $[R_1, +\infty)$.

(A2) For every $\delta > 0$, there exist constants $R_2 = R_2(\delta) > 0$ and $M_2 = M_2(\delta) > 1$ such that

$$\forall t > R_2, \quad M_2(\delta)^{-1} \leq \frac{\lambda(\delta t)}{\lambda(t)} \leq M_2(\delta), \quad \text{and} \quad M_2(\delta)^{-1} \leq \frac{\lambda'(\delta t)}{\lambda'(t)} \leq M_2(\delta).$$

(A3) There exist constants $R_3 > 0$ and $M_3 > 1$ such that

$$\forall t > R_3, \quad M_3^{-1} \leq \frac{t\lambda'(t)}{\lambda(t)} \leq M_3.$$

Moreover, for every integer $j \geq 1$ we have defined the property

(A4(j)) There exist constants $R_{4(j)} > 0$ and $M_{4(j)} > 1$ such that

$$\forall t > R_{4(j)}, \quad \frac{t\lambda^{(j)}(t)}{\lambda^j(t)} \leq M_{4(j)}.$$

Let $\theta(t) \doteq t\lambda(t)$. We have the validity of the following

Lemma 3.1. *Assumption (A3) implies that for all $t > R_3$,*

$$(1 + M_3^{-1})\lambda(t) \leq \theta'(t) \leq (1 + M_3)\lambda(t).$$

Let $j \geq 1$ be an integer. Since $\theta^{(j)} = t\lambda^{(j)}(t) + j\lambda^{(j-1)}(t)$, we have

Lemma 3.2. *Assumptions $(A_4(j-1))$ and $(A_4(j))$ imply that there exist constants $R_{5(j)} > 0$ and $M_{5(j)} > 1$ such that*

$$\forall t > R_{5(j)}, \quad \theta^{(j)}(t) \leq M_{5(j)} \lambda^j(t).$$

Now, let $j \geq 0$. Define $\Theta(t) \doteq (\theta'(t))^2 = (t\lambda'(t) + \lambda(t))^2$. Note that $\Theta^{(j)}$ is a linear combination of terms of the form $\theta^{(j_1)}\theta^{(j_2)}$, with j_1 and j_2 positive integers satisfying $j_1 + j_2 = j + 2$. Then we have also the following

Lemma 3.3. *Suppose that assumptions $(A_4(s))$, $s = 1, \dots, j+1$ are satisfied. Then there exist constants $R_{6(j)} > 0$ and $M_{6(j)} > 1$ such that*

$$\forall t > R_{6(j)}, \quad \Theta^{(j)}(t) \leq M_{6(j)} \lambda^{j+2}(t).$$

Lemma 3.4. (i) *The function $\lambda(t) \doteq t^\eta$ with $\eta > 0$ satisfies assumptions (A1), (A2), (A3) and $(A_4(j))$ for $j \geq 2$.*
(ii) *Suppose that*

$$\lambda(t) = \alpha t \prod_{j=1}^{\bar{j}} \ln^{[j]}(t)$$

for t large enough, where $\alpha > 0$ is a constant, \bar{j} a positive integer and we recall that $\ln^{[j]}$ stands for the j -th iterated logarithm. Then λ satisfies assumptions (A1), (A2), (A3) and $(A_4(j))$ for $j \geq 2$.

Proof. (i) (A1), (A2), (A3) are trivially satisfied. Concerning $(A_4(j))$ for $j \geq 2$, we have

$$\frac{t\lambda^{(j)}(t)}{\lambda^j(t)} = C \frac{t^{\eta-j+1}}{t^{j\eta}}$$

If $\eta \leq 1$, then $t^{\eta-j+1} \leq 1$ when $t > 1$, whence the thesis. For $\eta > 1$, we have $t^{\eta+1} \leq t^{\eta j}$ when $t > 1$, which permits to conclude.

(ii) (A1) is trivial. Concerning (A2) note that

$$\frac{\lambda(\delta t)}{\lambda(t)} = \delta \prod_{j=1}^{\bar{j}} \frac{\ln^{[j]}(\delta t)}{\ln^{[j]}(t)},$$

and $\frac{\ln^{[j]}(\delta t)}{\ln^{[j]}(t)} \rightarrow 1$ as $t \rightarrow \infty$ for every j . Similarly, $\lambda'(t) = \alpha + \alpha \sum_{q=1}^{\bar{j}} \prod_{j=q}^{\bar{j}} \ln^{[j]}(t)$, so that $\lambda'(t) \sim_{t \rightarrow \infty} \alpha \prod_{j=1}^{\bar{j}} \ln^{[j]}(t)$, and

$$\lim_{t \rightarrow \infty} \frac{\lambda'(\delta t)}{\lambda'(t)} = \lim_{t \rightarrow \infty} \frac{\prod_{j=1}^{\bar{j}} \ln^{[j]}(\delta t)}{\prod_{j=1}^{\bar{j}} \ln^{[j]}(t)} = 1.$$

Concerning (A3), we have that

$$\frac{t\lambda'(t)}{\lambda(t)} = \frac{\alpha t \left(1 + \sum_{q=1}^{\bar{j}} \prod_{j=q}^{\bar{j}} \ln^{[j]}(t)\right)}{\lambda(t)} \sim_{t \rightarrow \infty} \frac{\alpha t \prod_{j=1}^{\bar{j}} \ln^{[j]}(t)}{\lambda(t)} = 1$$

Concerning $(A_4(j))$ it is enough to remark that $\lim_{t \rightarrow \infty} \lambda^{(j)}(t) = 0$ and that $\lambda^j(t) \geq \alpha^j t^j \geq t$ when t is large enough. \square

3.2. A generalization of a result by Bianchi-Setti. Schoen and Yau proved in [27] the existence of families of Laplacian cut-off functions on manifolds with lower bounded Ricci curvature. In [6, Theorem 2.1], Bianchi and Setti generalized this result in two directions. On the one hand they got a better control on the Laplacian of the cut-offs when the negative part of the Ricci curvature vanishes at infinity. On the other hand they showed that one can get Laplacian cut-offs even when the negative part of the Ricci curvature explodes at infinity, provided that its growth is suitably controlled, i.e. $\text{Ric}_g \geq -Cr(x)^2$. In the following theorem, we extend this latter result to the assumption $\text{Ric}_g \geq -\lambda^2(r(x))$ for a larger class of growth functions λ . This in particular will permit to sharpen applications to the density problem for Sobolev spaces, see Subsection 4.1.

Theorem 3.5. *Let $\lambda : [0, +\infty) \rightarrow [0, +\infty)$ be a C^∞ function satisfying assumptions (A1), (A2), (A3). Let (M^m, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}_g(x, o)$. Suppose that for some $R_0 > 0$,*

$$\text{Ric}_g(x) \geq -\lambda^2(r(x)), \quad \text{on } M \setminus B_{R_0}(o).$$

Let $\tilde{r} \in C^\infty(M \setminus o)$ be an exhaustion function on M such that for some $C_{\tilde{r}} > 1$

$$\begin{cases} C_{\tilde{r}}^{-1}r(x) \leq \tilde{r}(x) \leq C_{\tilde{r}}r(x) \\ \tilde{r}(x) = r(x), \quad \text{on } B_2(o) \setminus B_{1/2}(o) \\ |\nabla_g \tilde{r}| \leq 1. \end{cases}$$

Then there exists $h \in C^\infty(M)$ such that

- (i) $C_h^{-1}r(x)\lambda(r(x)) \leq h(x) \leq C_h \max\{1; r(x)\lambda(r(x))\}$;
- (ii) $|\nabla_g h| \leq C_h \lambda(r(x))$ on M ;
- (iii) $\Delta_g h = |\nabla_g h|^2 - C_\theta(\theta'(\tilde{r}(x)))^2$ on $M \setminus B_{R_\theta}(o)$

for some constants $C_\theta > 0$, $R_\theta > 0$ and $C_h > 1$. Here θ is the function defined on $[0, +\infty)$ by $\theta(t) \doteq t\lambda(t)$.

Beyond the more restrictive choice for the growth function λ , in the original version of Theorem 3.5 given in [6, Theorem 2.1] the first-order distance-like function \tilde{r} was required to satisfy $\|\tilde{r} - r\|_{L^\infty(M)} < \epsilon$ for an ϵ arbitrarily small. This is the case for instance if \tilde{r} is a distance-like function provided by Gaffney's result [9]. Instead, our Theorem 3.5 works for general first-order distance-like functions; see Proposition 3.9 below. This represents a minor technical improvement, which will reveal however essential in proving Theorem 1.2.

Indeed, in order to prove the existence of a distance-like function H_k with a control on the derivatives up to order $k > 2$ we will first construct a distance-like function H_{k-1} which is controlled up to order $k - 1$ and then apply Theorem 3.5 with $\tilde{r} = H_{k-1}$ and set $h = \theta(H_k)$. The control on the k -th order derivatives is obtained by differentiating $k - 2$ times the equation appearing in (iii), and then using elliptic estimates in suitable harmonic coordinates; see Lemma 3.11. However, note that in order to get a second-order distance-like function (i.e. for $k = 2$) the exact form of equation (iii) is not needed since it suffices to have an estimate of the form $|\Delta_g h| \leq C\lambda^2(r(x))$.

Proof (of Theorem 3.5). The proof mimics the proof of Theorem 2.1 in [6]. The main difference is to modify the equation of which $\log h$ will be a solution, according to our more general set of assumptions. We will sketch the relevant changes. Also, if λ is bounded then the result is contained in [6, Theorem 2.1]. We can thus assume from now on that λ is unbounded and $\lambda(t) \rightarrow \infty$ as $t \rightarrow \infty$.

We start by pointing out the following lemma.

Lemma 3.6. *Let $\lambda : [0, +\infty) \rightarrow [0, +\infty)$ be a C^∞ function satisfying assumptions (A1) and (A2). Let (M^m, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq$*

$\text{dist}_g(x, o)$. Suppose that for some $R_0 > 0$,

$$\text{Ric}_g(x) \geq -\lambda^2(r(x)), \quad \text{on } M \setminus B_{R_0}(o).$$

Then there exists $\alpha > 1$, $C_{SY} > 0$ and $R_\alpha > 0$ such that for every $x \in M \setminus B_{R_\alpha}(o)$ it holds

$$\text{vol}_g(B_{1/4}(x)) \geq C_{SY} e^{-\alpha r(x)\lambda(r(x))}.$$

The original idea is due to Schoen and Yau, [27], while in [6, Proposition 2.11] the authors consider bounds given by (possibly negative) powers of $r(x)$.

Proof (of Lemma 3.6). Set $G_x \doteq \lambda^2(2r(x) + 1)$. By continuity, since $\lambda(t) \rightarrow \infty$ as $t \rightarrow \infty$, if $r(x)$ is large enough, then $\text{Ric}_g \geq -G_x$ on $B_{2r(x)+1}(o)$. By the Bishop-Gromov comparison,

$$\begin{aligned} \text{vol}_g(B_{1/4}(x)) &\geq \text{vol}_g(B_{r(x)+1}(x)) \frac{V_{G_x}(1/4)}{V_{G_x}(r(x)+1)} \geq \text{vol}_g(B_1(x)) \frac{V_{G_x}(1/4)}{V_{G_x}(r(x)+1)} \\ &\geq \text{vol}_g(B_1(x)) \frac{V_0(1/4)}{V_{G_x}(r(x)+1)}, \end{aligned}$$

where $V_{G_x}(t)$ represents the volume of a ball of radius t in the m -dimensional simply connected space-form of constant negative curvature $-G_x$, and $V_0(1/4)$ is the the volume of a ball of radius $1/4$ in \mathbb{R}^m . A standard computation using assumption (A2) shows that there exist constants $C_1 > 0$, $R_\alpha > 0$ and $\alpha > 1$ independent of x such that

$$V_{G_x}(r(x)+1) \leq C_1 e^{\alpha r(x)\lambda(r(x))},$$

as soon as $r(x) > R_\alpha$. Setting $C_{SY} = C_1^{-1} \text{vol}_g(B_1(x)) V_0(1/4)$ concludes the proof of the lemma. \square

Recall that $\theta(t) = t\lambda(t)$. Proceeding as in [6, Theorem 2.1] we can produce a function $\omega \in C^\infty(M \setminus B_1(o))$ solution of

$$(3) \quad \begin{cases} \Delta_g \omega = C_e^2 (\theta'(\tilde{r}(x)))^2 \omega, \\ \omega|_{\partial B_1} = 1, \\ 0 < \omega|_{M \setminus \bar{B}_1(o)} < 1, \end{cases}$$

with C_e a large enough constant to be fixed later. Computing as in [6, page 6] we get that for every $0 < \gamma < \sqrt{C_e^2}/2$ and every $x \in M \setminus B_2(o)$,

$$(4) \quad \int_{B_{1/4}(x)} (\theta'(\tilde{r}(y)))^2 e^{\gamma \theta(\tilde{r}(y))} \omega^2 d\text{vol}_g(y) \leq \int_{M \setminus B_1(o)} (\theta'(\tilde{r}(y)))^2 e^{\gamma \theta(\tilde{r}(y))} \omega^2 d\text{vol}_g(y) \leq \frac{A}{C_e^2 - 4\gamma^2},$$

where $A = e^{\lambda(1)} \int_{\partial B_1(o)} |\nabla_g \omega| d\text{vol}_{m-1}$. We need the following Li-Yau gradient estimate.

Lemma 3.7. *There exists a constant $C_{LY} > 0$ and $R_{LY} > 0$ such that for every $R > R_{LY}$ and for every $x \in B_{R+1}(o) \setminus B_{R_{LY}}(o)$,*

$$\frac{|\nabla_g \omega|^2}{\omega^2} \leq C_{LY}^2 C_e^2 \lambda^2(R).$$

Proof. A special case of Theorem 2.8 in [6] ensures the existence of a constant $C'_{LY} > 0$ such that

$$\frac{|\nabla_g \omega|^2}{\omega^2} \leq C'_{LY} \max\{\lambda^2(R); C_e^2 (\theta'(R))^2\}.$$

By Lemma 3.1, $\theta'(R) \leq (1 + M_3)\lambda(R)$ for R large enough, which proves the lemma. \square

Reasoning as in [6, page 6], thanks to Lemma 3.7, for every $x \in M \setminus B_3(o)$ and every $y \in B_{1/4}(x)$, we have

$$(5) \quad \omega(y) \geq \omega(x) e^{-\frac{1}{4} C_{LY} C_e \lambda(r(x))}.$$

Note also that there exist $\delta > 0$, $\kappa > 0$ and $R_\delta > 0$ such that for every $x \in M \setminus B_{R_\delta}(o)$ and every $y \in B_{1/4}(x)$,

$$(6) \quad (\theta'(\tilde{r}(y)))^2 e^{\gamma\theta(\tilde{r}(y))} \geq \delta (\theta'(\tilde{r}(x)))^2 e^{\kappa\gamma\theta(\tilde{r}(x))}.$$

This follows from assumption (A2) and the fact that $|\tilde{r}(x) - \tilde{r}(y)| < 1/4$ (because $|\nabla_g \tilde{r}| \leq 1$). Inserting (5) and (6) in (4) gives

$$\omega^2(x) \delta (\theta'(\tilde{r}(x)))^2 e^{\kappa\gamma\theta(\tilde{r}(x))} e^{-\frac{1}{2}C_{LY}C_e\lambda(r(x))} \text{vol}_g(B_{1/4}(x)) \leq \frac{A}{C_e^2 - 4\gamma^2}.$$

From Lemma 3.6, we deduce that

$$\omega^2(x) \leq C_{SY}^{-1} \delta^{-1} \frac{A}{C_e^2 - 4\gamma^2} (\theta'(\tilde{r}(x)))^{-2} e^{-\kappa\gamma\theta(\tilde{r}(x)) + \frac{1}{2}C_{LY}C_e\lambda(r(x)) + \alpha r(x)\lambda(r(x))}$$

whenever $r(x) > R_1 \doteq \max\{R_\alpha; R_\delta; R_{LY}; 3\}$. Recalling assumption (A1) and (A2), we get

$$\theta(\tilde{r}(x)) \geq \theta(C_{\tilde{r}}^{-1}r(x)) \geq \zeta\theta(r(x)),$$

with $\zeta = C_{\tilde{r}}^{-1}M_2(C_{\tilde{r}}^{-1})^{-1} > 0$ independent of x . Hence

$$-\kappa\gamma\theta(\tilde{r}(x)) + \frac{1}{2}C_{LY}C_e\lambda(r(x)) + \alpha r(x)\lambda(r(x)) \leq -[\kappa\gamma\zeta - \alpha]r(x)\lambda(r(x)) + \frac{1}{2}C_{LY}C_e\lambda(r(x)).$$

At this point we can make a choice of $\gamma > \frac{3\alpha}{\kappa\zeta}$, so that $\kappa\gamma\zeta - \alpha > 2\alpha$, and a choice of $C_e > 2\gamma$. There exists a radius $R_2 > 0$ such that

$$-[\kappa\gamma\zeta - \alpha]r(x)\lambda(r(x)) + \frac{1}{2}C_{LY}C_e\lambda(r(x)) \leq \alpha r(x)\lambda(r(x))$$

for all $x \in M \setminus B_{R_2}(o)$. Because of assumption (A1) and Lemma 3.1, $\theta'(t)$ is lower bounded. Hence we have obtained that

$$\omega^2(x) \leq C_{SY}^{-1} \delta^{-1} \frac{A}{C_e^2 - 4\gamma^2} e^{-\alpha r(x)\lambda(r(x))}$$

on $M \setminus B_{R_3}(o)$ for some $R_3 \geq R_2$. Define

$$h(x) \doteq (\phi(x) - 1) \ln(\omega(x)) + \phi(x),$$

where $\phi \in C^\infty(M)$ is such that $\phi(x) \equiv 1$ in $B_2(o)$ and $\phi(x) \equiv 0$ in $M \setminus B_3(o)$. We have that on $M \setminus B_{R_3}(o)$

$$h(x) = -\ln(\omega(x)) \geq \ln\left(C_{SY}\delta\frac{C_e^2 - 4\gamma^2}{A}\right) + \alpha r(x)\lambda(r(x)),$$

from which we deduce the existence of a radius $R_4 > R_3$ such that

$$(7) \quad h(x) \geq \frac{\alpha}{2}r(x)\lambda(r(x))$$

on $M \setminus B_{R_4}(o)$. On the other hand, since $h(o) = 1$, thanks to Lemma 3.7 we can compute

$$(8) \quad h(x) = h(o) + \int_0^{r(x)} |\nabla_g h(\sigma(s))| ds \leq 1 + R_{LY} \max_{B_{R_{LY}}} |\nabla_g h| + \int_{R_{LY}}^{r(x)} C_{LY}C_e\lambda(r(x)) ds \\ \leq \alpha' r(x)\lambda(r(x)),$$

for some constant α' independent of x . By definition of the smooth function h and up to choose $C_h > \max\{\alpha', 2/\alpha, C_{LY}C_e\}$ large enough, as a consequence of (7), (8) and Lemma 3.7, we have thus obtained the validity of the points (i) and (ii) in the statement of Theorem 3.5. The point (iii) follows from (3) and a direct computation. \square

3.3. Proof of Theorem 1.2: case (a). We are going to prove Theorem 1.2 in the set of assumptions (a), i.e.

Theorem 3.8. *Let (M^m, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}_g(x, o)$. Let $k \in \mathbb{N}^+$. If $k \geq 2$, suppose in addition that for some $i_0 > 0$,*

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{inj}_g(x) \geq \frac{i_0}{\lambda(r(x))} > 0 \quad \text{on } M,$$

where the function λ satisfies assumptions (A1), (A2), (A3), (A4(j)) for $j = 1, \dots, k$. Then there exists an exhaustion function $H = H_k \in C^\infty(M)$ such that for some positive constant $C_k > 1$ independent of x , we have on M that

- (i) $C_k^{-2} r(x) \leq H(x) \leq \max\{r(x), 1\}$;
- (ii) for $1 \leq j \leq k$, $|\nabla_g^j H|(x) \leq C_k^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$.

The proof is done by induction on k . For $k = 1$, the function λ is not involved in the statement, and this is the content of the well-known theorem by Gaffney, [9, 11]. Assume now that for some $k \geq 2$ and some $i_0 > 0$

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{inj}_g(x) \geq \frac{i_0}{\lambda(r(x))} > 0 \quad \text{on } M.$$

As induction hypothesis we suppose that the result holds true for $k-1$, i.e. there exists a distance-like function $H_{k-1} \in C^\infty(M)$ such that for some positive constant $C_{k-1} > 1$ independent of x , we have on M that

- (i) $C_{k-1}^{-2} r(x) \leq H_{k-1}(x) \leq \max\{r(x), 1\}$;
- (ii) for $1 \leq j \leq k-1$, $|\nabla_g^j H_{k-1}|(x) \leq C_{k-1}^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$.

By Theorem 3.5 we have the following

Proposition 3.9. *In the assumptions of Theorem 3.8, there exists a function $h = h_k \in C^\infty(M)$ and a constant $C_h > 1$ such that*

- (i) $C_h^{-1} r(x) \lambda(r(x)) \leq h(x) \leq C_h \max\{1; r(x) \lambda(r(x))\}$;
- (ii) $|\nabla_g h|_g \leq C_h \lambda(r(x))$ on M ;
- (iii) $\Delta_g h = |\nabla_g h|_g^2 - C_\theta (\theta'(H_{k-1}(x)))^2$ on $M \setminus B_{R_\theta}(o)$ for some constants $C_\theta > 0$ and $R_\theta > 0$.

Recall that a local coordinate system $\{x^i\}_{i=1}^m$ is said to be harmonic if for any i , $\Delta_g x^i = 0$. As a consequence of [2] we have the validity of the following

Proposition 3.10. *Let $\alpha \in (0, 1)$, $Q > 1$, $\delta > 0$. Let (M^m, g) be a smooth Riemannian manifold, and Ω an open subset of M . Set*

$$\Omega(\delta) = \{x \in M \quad \text{s.t.} \quad d_g(x, \Omega) < \delta\}.$$

Suppose that

$$|\nabla_g^j \text{Ric}_g|(x) \leq 1, \quad 0 \leq j \leq k-2 \quad \text{and} \quad \text{inj}_g(x) \geq i \quad \text{for all} \quad x \in \Omega(\delta),$$

then there exists a positive constant $C_{HR} = C_{HR}(m, Q, k, \alpha, \delta, i)$, such that on the geodesic ball $B_{C_{HR}}(x)$ of center x and radius C_{HR} , there is a centered harmonic coordinate chart such that the metric tensor is $C^{k-1, \alpha}$ controlled in these coordinates. In particular, if g_{ij} , $i, j = 1, \dots, m$, are the components of g in these coordinates, then

- (1) $Q^{-1} \delta_{ij} \leq g_{ij} \leq Q \delta_{ij}$ as bilinear forms;
- (2) $\sum_{1 \leq |\gamma| \leq k-1} r_H^{|\gamma|} \sup_y |\partial_\gamma g_{ij}(y)| \leq Q - 1$

By assumption, we have that

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{inj}_g(x) \geq \frac{i_0}{\lambda(r(x))} > 0 \quad \text{on } M.$$

Fix $x \in M \setminus B_2^g(o)$ and set

$$\lambda_1 = \lambda_1(x) \doteq \lambda(r(x) + 1).$$

Since λ is increasing, for all $y \in B_1^g(x)$ we have

$$|\nabla_g^j \text{Ric}_g|(y) \leq \lambda_1^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{and} \quad \text{inj}_g(y) \geq \frac{i_0}{\lambda_1}.$$

We define a new rescaled metric $g_\lambda \doteq \lambda_1^2 g$ on M , so that for all $y \in B_{\lambda_1}^{g_\lambda}(x)$ we have

$$|\nabla_{g_\lambda}^j \text{Ric}_{g_\lambda}|(y) \leq 1, \quad 0 \leq j \leq k-2, \quad \text{and} \quad \text{inj}_{g_\lambda}(y) \geq \lambda_1 \text{inj}_g(y) \geq i_0.$$

According to Proposition 3.10 (applied with $\delta = 1/2$) there exists a constant $C_{HR} = C_{HR}(m, Q, i_0, k)$ such that on $B_{C_{HR}}^{g_\lambda}(x)$ there exists a centered harmonic chart $\varphi_H = (y^1, \dots, y^m) : B_{C_{HR}}^{g_\lambda}(x) \rightarrow U \subset \mathbb{R}^m$ such that $\varphi_H(x) = 0 \in \mathbb{R}^m$ and, setting $\hat{g}_\lambda = g_\lambda \circ \varphi_H^{-1}$, it holds

- (1_{HC}) $Q^{-1} \delta_{ij} \leq (\hat{g}_\lambda)_{ij} \leq Q \delta_{ij}$ as bilinear forms;
- (2_{HC}) $\sum_{1 \leq |\gamma| \leq k-1} C_{HR}^{|\gamma|} \sup_y \left| \partial_\gamma (\hat{g}_\lambda)_{ij}(y) \right| \leq Q - 1,$

Since

$$\partial_q \hat{g}_\lambda^{ij} = -\partial_q (\hat{g}_\lambda)_{lk} \hat{g}_\lambda^{il} \hat{g}_\lambda^{kj},$$

we have also that

- (1'_{HC}) $Q^{-1} \delta^{ij} \leq \hat{g}_\lambda^{ij} \leq Q \delta^{ij};$
- (2'_{HC}) $\sum_{1 \leq |\gamma| \leq k-1} \sup_y \left| \partial_\gamma \hat{g}_\lambda^{ij}(y) \right| \leq C(Q),$

for some constant $C(Q)$, depending only on Q .

For $\beta \doteq C_{HR}/Q$, on \mathbb{B}_β we set $\hat{g} = g \circ \varphi_H^{-1}$, $\hat{h} = h \circ \varphi_H^{-1}$ and $\hat{H} = H_{k-1} \circ \varphi_H^{-1}$.

We need the following fundamental lemma.

Lemma 3.11. *Let i and j be integers in $\{1, \dots, m\}$. Let $0 \leq q \leq k-1$ and let $\gamma = (\gamma_1, \dots, \gamma_{q-1}) \in \{1, \dots, m\}^{q-1}$ be a multi-index (possibly empty for $q = 0, 1$). There exists a positive constant $C > 0$ such that*

$$(9) \quad \forall v \in \mathbb{B}_\beta, \quad |\partial_i \hat{h}|(v) \leq C,$$

and, if $q \geq 1$,

$$(10) \quad \forall v \in \mathbb{B}_{4^{-q}\beta}, \quad \left| \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{q-1}}^{q-1} \hat{h} \right|(v) \leq C.$$

Before proving Lemma 3.11, let us point out the following computational lemma which will be used repeatedly.

Lemma 3.12. *Let $F \in C^\infty(M)$. Set $\hat{F} \doteq F \circ \varphi_H^{-1}$ and let $v \in \mathbb{B}_\beta$. Let $d \leq k$. Then there exists a real positive constant C_1 independent of x such that*

- (1) *If $\left| \partial_{e_1 \dots e_n}^n \hat{F} \right|(v) \leq C_0$ for some constant $C_0 > 0$ (possibly depending on x), for all $n \in \{1, \dots, d\}$ and for all multi-index $(e_1, \dots, e_n) \in \{1, \dots, m\}^n$, then $\left| \nabla_{\hat{g}_\lambda}^n \hat{F} \right|_{\hat{g}_\lambda}(v) \leq C_1 C_0$ for all $n \in \{1, \dots, d\}$.*
- (2) *If $\left| \nabla_{\hat{g}_\lambda}^n \hat{F} \right|_{\hat{g}_\lambda}(v) \leq C_0$ for some constant $C_0 > 0$ (possibly depending on x) and for all $n \in \{1, \dots, d\}$, then $\left| \partial_{e_1 \dots e_n}^n \hat{F} \right|(v) \leq C_1 C_0$ for all $n \in \{1, \dots, d\}$ and for all multi-index $(e_1, \dots, e_n) \in \{1, \dots, m\}^n$.*

$$(3) \quad \lambda_1^d \left| \nabla_{\hat{g}_\lambda}^d \hat{F} \right|_{\hat{g}_\lambda} = \left| \nabla_{\hat{g}}^d \hat{F} \right|_{\hat{g}}.$$

$$(4) \quad \left| \nabla_{\hat{g}}^d \hat{F} \right|_{\hat{g}}(v) = \left| \nabla_g^d F \right|_g(\varphi_H^{-1}(v)).$$

Proof. As for (1) and (2) we proceed by induction on d . For $d = 0$, the result is trivial with $C_1 = 1$. Suppose that we have proved the assertions for some $d - 1$ and let us prove it for d . Clearly the only nontrivial case is $n = d$.

(1). By definition

$$\left| \nabla_{\hat{g}_\lambda}^d \hat{F} \right|_{\hat{g}_\lambda}^2 = \left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{e_1 \dots e_d} \left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{f_1 \dots f_d} \hat{g}_\lambda^{e_1 f_1} \dots \hat{g}_\lambda^{e_d f_d}$$

The expression in coordinates of $\left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{e_1 \dots e_d}$ can be written as a linear combination of terms, each of which is given by the product of factors of the form $\partial_\alpha \hat{F}$ with factors of the form $\partial_\beta (\hat{g}_\lambda \Gamma_{ij}^k)$, where the multi-indexes α and β have orders, respectively, $|\alpha| \leq d$ and $|\beta| \leq d - 2$. Note that $\partial^\beta (\hat{g}_\lambda \Gamma_{ij}^k)$ contains partial derivatives of \hat{g}_λ and \hat{g}_λ^{-1} of order at most $d - 1$, which are uniformly bounded as soon as $d \leq k$ by the properties of the harmonic chart φ_H .

(2). Reasoning as for the previous implication, we get

$$\begin{aligned} \left| \partial_{e_1 \dots e_n}^d \hat{F} \right| (v) &= \left| \left(D^d \hat{F} \right)_{e_1 \dots e_d} \right| (v) \\ &\leq \left| \left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{e_1 \dots e_d} \right| (v) + C \left(\left| \nabla_{\hat{g}_\lambda}^{d-1} \hat{F} \right| + \dots + \left| \nabla_{\hat{g}_\lambda}^1 \hat{F} \right| \right) \\ &\leq \left| \nabla_{\hat{g}_\lambda}^d \hat{F} \right|_{\hat{g}_\lambda} |\partial_{e_1}|_{\hat{g}_\lambda} \dots |\partial_{e_d}|_{\hat{g}_\lambda} + C \\ &\leq C. \end{aligned}$$

(3). Note that

$$\left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{e_1 \dots e_d} = \left(\nabla_{\hat{g}}^d \hat{F} \right)_{e_1 \dots e_d}.$$

Accordingly,

$$\begin{aligned} \left| \nabla_{\hat{g}_\lambda}^d \hat{F} \right|_{\hat{g}_\lambda}^2 &= \left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{e_1 \dots e_d} \left(\nabla_{\hat{g}_\lambda}^d \hat{F} \right)_{f_1 \dots f_d} \hat{g}_\lambda^{e_1 f_1} \dots \hat{g}_\lambda^{e_d f_d} \\ &= \left(\nabla_{\hat{g}}^d \hat{F} \right)_{e_1 \dots e_d} \left(\nabla_{\hat{g}}^d \hat{F} \right)_{f_1 \dots f_d} \lambda_1^{-2} \hat{g}^{e_1 f_1} \dots \lambda_1^{-2} \hat{g}^{e_d f_d} \\ &= \lambda_1^{-2d} \left| \nabla_{\hat{g}}^d \hat{F} \right|_{\hat{g}}^2. \end{aligned}$$

(4). By definition of \hat{g} . □

Proof (of Lemma 3.11). The property (9) is a direct consequence of (ii) in Proposition 3.9 and of Lemma 3.12. In particular, Lemma 3.11 is verified when $q = 0$. It remains to prove (10) for $q \geq 1$. Let us proceed by induction on q . Suppose that the lemma is proved for all the integers q between 0 and \bar{q} , with $0 \leq \bar{q} \leq k - 2$. Let us prove it for $\bar{q} + 1$.

Define $\Theta(t) \doteq (\theta'(t))^2 = (t\lambda'(t) + \lambda(t))^2$. By Proposition 3.9 (iii), h satisfies

$$\Delta_g h(x) = |\nabla_g h|^2(x) - C_\theta \Theta(H_{k-1}(x))$$

on $M \setminus B_{R_\theta}(o)$, which reads in harmonic coordinates as

$$\lambda_1^2 \hat{g}_\lambda^{ij} (v) \partial_i^2 \hat{h}(v) = \lambda_1^2 \hat{g}_\lambda^{ij} (v) \partial_i \hat{h}(v) \partial_j \hat{h}(v) - C_\theta \Theta(\hat{H}(v)), \quad \forall v \in \mathbb{B}_\beta.$$

Applying the differential operator $\partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}}$ to both sides, and multiplying by λ_1^{-2} , we get on \mathbb{B}_β

$$(11) \quad \hat{g}_\lambda^{ij} \partial_{ij}^2 \left(\partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \hat{h} \right) = \sum \left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) - C_\theta \lambda_1^{-2} \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\Theta(\hat{H})) \\ - \sum \left(\partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1 \dots b_{\bar{q}-n}}^{\bar{q}-n} \hat{g}_\lambda^{ij} \right) \\ \doteq \hat{f}.$$

Here the first summation is taken over all the (possibly empty) subsets $\{a_s\}_{s=1}^n \subset \{\gamma_s\}_{s=1}^{\bar{q}}$ and $\{b_s\}_{s=1}^l \subset \{\gamma_s\}_{s=1}^{\bar{q}} \setminus \{a_s\}_{s=1}^n$, and we have set $\{c_s\}_{s=1}^{\bar{q}-n-l} \doteq \{\gamma_s\}_{s=1}^{\bar{q}} \setminus (\{a_s\}_{s=1}^n \cup \{b_s\}_{s=1}^l)$. Similarly the second summation is taken over all the non-empty subsets $\{b_s\}_{s=1}^{\bar{q}-n} \subset \{\gamma_s\}_{s=1}^{\bar{q}}$ and we have set $\{a_s\}_{s=1}^n \doteq \{\gamma_s\}_{s=1}^{\bar{q}} \setminus \{b_s\}_{s=1}^{\bar{q}-n}$. Fix $w \in \mathbb{B}_{4-\bar{q}\beta}$. We need to prove that $\left| \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{\bar{q}-1}}^{\bar{q}-1} \hat{h} \right| (w) \leq C$ for some constant C independent of x and w . Define $\tilde{h} \in C^\infty(\mathbb{B}_\beta)$ by

$$\tilde{h}(v) \doteq \hat{h}(v) - \sum_{j=0}^{\bar{q}} \frac{D^j \hat{h}(w)(v, \dots, v)}{j!}.$$

Note that $\tilde{h} - \hat{h}$ is a polynomial of degree \bar{q} , so that

$$\hat{g}_\lambda^{ij} \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \tilde{h} = \hat{g}_\lambda^{ij} \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \hat{h} = \hat{f}.$$

Rescaling to $\mathbb{B}_{4-\bar{q}\beta}(w)$ the elliptic Schauder estimates given for instance in [17, Theorem 5.21], we know that there exists $C > 0$, depending on \bar{q} (hence on k), but not on w , such that

$$(12) \quad \left| \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \hat{h}(w) \right| = \left| \partial_{ij}^2 \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \tilde{h}(w) \right| \\ \leq C \left\{ \left\| \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \tilde{h} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))} + \left\| \hat{f} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))} + [\hat{f}]_{C^{0,\alpha}(\mathbb{B}_{4-\bar{q}\beta}(w))} \right\}.$$

Note that $\mathbb{B}_{4-\bar{q}\beta}(w) \subset \mathbb{B}_{4-\bar{q}\beta}(0)$. We will estimate the three terms at RHS of (12) separately.

1st term. *Estimating* $\left\| \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \tilde{h} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))}$.

Let $v \in \mathbb{B}_{4-\bar{q}\beta}(w)$. Note that

$$\partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \sum_{j=0}^{\bar{q}} \frac{D^j \hat{h}(w)(v, \dots, v)}{j!} = \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \frac{D^{\bar{q}} \hat{h}(w)(v, \dots, v)}{\bar{q}!} = D^{\bar{q}} \hat{h}(w)(\partial_{\gamma_1}, \dots, \partial_{\gamma_{\bar{q}}}).$$

Hence

$$\left| \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} \tilde{h} \right| (v) = \left| D^{\bar{q}} \tilde{h}(v)(\partial_{\gamma_1}, \dots, \partial_{\gamma_{\bar{q}}}) \right| \\ = \left| D^{\bar{q}} \hat{h}(v)(\partial_{\gamma_1}, \dots, \partial_{\gamma_{\bar{q}}}) - D^{\bar{q}} \hat{h}(w)(\partial_{\gamma_1}, \dots, \partial_{\gamma_{\bar{q}}}) \right| \\ \leq 2\beta \left\| D^{\bar{q}+1} \hat{h} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))} \\ \leq C$$

by the inductive hypothesis of the lemma.

2nd term. *Estimating* $\left\| \hat{f} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))}$.

Concerning the first addend in \hat{f} , since $n \leq k - 1$, by the property $(2'_{HC})$ of the harmonic chart we can compute

$$(13) \quad \begin{aligned} & \left\| \left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \\ & \leq C \left\| \partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \left\| \partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \\ & \leq C, \end{aligned}$$

where we have used the inductive hypothesis and the fact that $l + 1 \leq \bar{q} + 1$ and $\bar{q} - n - l + 1 \leq \bar{q} + 1$.

We now consider the second addend of \hat{f} and estimate $\left\| C_\theta \lambda_1^{-2} \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\Theta(\hat{H})) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))}$. If $\bar{q} = 0$, then

$$\left\| C_\theta \lambda_1^{-2} \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\Theta(\hat{H})) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} = \left\| C_\theta \lambda_1^{-2} \Theta(\hat{H}) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \leq C$$

as a direct consequence of Lemma 3.3. If $\bar{q} \geq 1$, then $\partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\Theta(\hat{H}))$ can be developed as a linear combinations of terms of the form

$$\Theta^{(j)}(\hat{H}) \prod_{s=1}^j \partial_{\gamma_{\sigma(e_{s-1}+1)} \dots \gamma_{\sigma(e_s)}}^{e_s - e_{s-1}} \hat{H},$$

where $e_0 = 0$, $j \in \{1, \dots, \bar{q}\}$, $\{e_s\}_{s=1}^j$ is an increasing subset of $\{1, \dots, \bar{q}\}$ with $e_j = \bar{q}$, and $\sigma \in \Pi_{\bar{q}}$ is a permutation of $\{1, \dots, \bar{q}\}$. By the inductive hypothesis of Theorem 3.8,

$$\left| \nabla_g^{e_s - e_{s-1}} H_{k-1} \Big|_g(v) \leq C \lambda(r(x))^{e_s - e_{s-1} - 1}.$$

Lemma 3.12 and assumption (A2) then imply that

$$\left| \partial_{\gamma_{\sigma(e_{s-1}+1)} \dots \gamma_{\sigma(e_s)}}^{e_s - e_{s-1}} \hat{H} \right| (v) \leq C \lambda_1^{-1},$$

and, using also Lemma 3.3,

$$\begin{aligned} \left\| \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\hat{H}^{2\eta}) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} & \leq C \max_{j, \{e_s\}_{s=1}^j, \sigma} \left\| \Theta^{(j)}(\hat{H}) \prod_{s=1}^j \partial_{\gamma_{\sigma(e_{s-1}+1)} \dots \gamma_{\sigma(e_s)}}^{e_s - e_{s-1}} \hat{H} \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \\ & \leq C \max_j \lambda(\hat{H})^{2+j} \prod_{s=1}^j \lambda_1^{-1} \\ & \leq C \max_j \lambda_1^{2+j-j} \\ & \leq C \lambda_1^2. \end{aligned}$$

Accordingly

$$(14) \quad \left\| C_\theta \lambda_1^{-2} \partial_{\gamma_1 \dots \gamma_{\bar{q}}}^{\bar{q}} (\Theta(\hat{H})) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \leq C$$

Finally, concerning the third addend in \hat{f} , since $n < \bar{q}$ and using again harmonic radius estimates we get

$$(15) \quad \begin{aligned} & \left\| \left(\partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1 \dots b_{\bar{q}-n}}^{\bar{q}-n} (\hat{g}_\lambda)_{ij} \right) \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \leq C \left\| \partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right\|_{L^\infty(\mathbb{B}_{4^{1-\bar{q}}\beta}(w))} \\ & \leq C. \end{aligned}$$

Combining (13), (14) and (15) we thus get

$$(16) \quad \left\| \hat{f} \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq C.$$

3rd term. *Estimating* $\left[\hat{f} \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))}$.

First, note that for any function $F \in C^1(\mathbb{B}_{4^{-\bar{q}}\beta}(w))$ it holds clearly that

$$(17) \quad [F]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq (2^{1-2\bar{q}}\beta)^{1-\alpha} \|F\|_{C^1(\mathbb{B}_{4^{-\bar{q}}\beta}(w))}.$$

Reasoning as for (14), we get that, for any $s \in \{1, \dots, m\}$,

$$\left\| C_\theta \lambda_1^{-2} \partial_s \partial_{\gamma_1 \dots \gamma_{\bar{q}}} (\Theta(\hat{H})) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq C,$$

from which, together with (17),

$$(18) \quad \left[C_\theta \lambda_1^{-2} \partial_{\gamma_1 \dots \gamma_{\bar{q}}} ((\Theta(\hat{H}))) \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq C.$$

Moreover, as long as either $n \in \{1, \dots, \bar{q}\}$ or $n = 0$ and $l \in \{1, \dots, \bar{q} - 1\}$ we have that

$$\begin{aligned} & \left\| \partial_s \left[\left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) \right] \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \leq \left\| \left(\partial_{sa_1 \dots a_n}^{n+1} \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \quad + \left\| \left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{sb_1 \dots b_l}^{l+1} \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \quad + \left\| \left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{sc_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l+1} \partial_j \hat{h} \right) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \leq C \end{aligned}$$

for any $s \in \{1, \dots, m\}$, where each addend has been estimated reasoning as for (13). Note that we used here the condition $n + 1 \leq \bar{q} + 1 \leq k - 1$ which permits to have the needed control in the harmonic chart. Combining with (17), we get

$$(19) \quad \left[\left(\partial_{a_1 \dots a_n}^n \hat{g}_\lambda^{ij} \right) \left(\partial_{b_1 \dots b_l}^l \partial_i \hat{h} \right) \left(\partial_{c_1 \dots c_{\bar{q}-n-l}}^{\bar{q}-n-l} \partial_j \hat{h} \right) \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq C,$$

Similarly, for any $s \in \{1, \dots, m\}$ and any $n \in \{1, \dots, \bar{q} - 2\}$, it holds

$$\begin{aligned} & \left\| \partial_s \left[\left(\partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1 \dots b_{\bar{q}-n}}^{\bar{q}-n} \hat{g}_\lambda^{ij} \right) \right] \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \leq \left\| \left(\partial_{sa_1 \dots a_n}^{n+1} \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1 \dots b_{\bar{q}-n}}^{\bar{q}-n} \hat{g}_\lambda^{ij} \right) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} + \left\| \left(\partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right) \left(\partial_{sb_1 \dots b_{\bar{q}-n}}^{\bar{q}-n+1} \hat{g}_\lambda^{ij} \right) \right\|_{L^\infty(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \\ & \leq C, \end{aligned}$$

from which

$$(20) \quad \left[\left(\partial_{a_1 \dots a_n}^n \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1 \dots b_{\bar{q}-n}}^{\bar{q}-n} \hat{g}_\lambda^{ij} \right) \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} \leq C.$$

Hence, in order to control $\left[\hat{f} \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))}$, it remains to estimate the LHS of (19) for $l = \bar{q}$ and $n = 0$, and the LHS of (20) for $n = \bar{q} - 1$. Namely, we have to bound

$$\left[\hat{g}_\lambda^{ij} \left(\partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \partial_i \hat{h} \right) \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))} + \left[\left(\partial_{a_1 \dots a_{\bar{q}-1}}^{\bar{q}-1} \partial_{ij}^2 \hat{h} \right) \left(\partial_{b_1}^1 \hat{g}_\lambda^{ij} \right) \right]_{C^{0,\alpha}(\mathbb{B}_{4^{-\bar{q}}\beta}(w))}.$$

By the properties of the harmonic chart, it is enough to obtain an upper bound for terms of the form

$$\left[\partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right]_{C^{0,\alpha}(\mathbb{B}_{4-\bar{q}\beta}(w))},$$

for some indexes $b_1, \dots, b_{\bar{q}+1} \in \{1, \dots, m\}$. Letting $p = \frac{m}{1-\alpha}$, by the Euclidean Sobolev embeddings (see e.g [1, p. 109]), we have

$$(21) \quad \left[\partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right]_{C^{0,\alpha}(\mathbb{B}_{4-\bar{q}\beta}(w))} \leq K_3 \left\{ \left\| \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right\|_{L^p(\mathbb{B}_{4-\bar{q}\beta}(w))}^p + \sum_{s=1}^m \left\| \partial_s \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right\|_{L^p(\mathbb{B}_{4-\bar{q}\beta}(w))}^p \right\}^{1/p},$$

with K_3 a positive constant depending only on m, α, β and k (via \bar{q}). Concerning the first term at the RHS of (21), by the inductive hypothesis we have

$$(22) \quad \left\| \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right\|_{L^p(\mathbb{B}_{4-\bar{q}\beta}(w))}^p \leq \left\| \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right\|_{L^\infty(\mathbb{B}_{4-\bar{q}\beta}(w))}^p \beta^m \kappa_m \leq C,$$

where κ_m is the volume of the m -dimensional unit ball $\mathbb{B}_1 \subset \mathbb{R}^m$. Concerning the second term at the RHS of (21), we note that $|\partial_s \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h}| \leq |D^2 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h}|$, where, given a C^2 function F on \mathbb{B}_β , $D^2 F$ denotes the Euclidean Hessian matrix of F and $|D^2 F|^2 \doteq \sum_{i,j=1}^m (\partial_i \partial_j F)^2$. According to the Euclidean Calderón-Zygmund inequality, [10, Theorem 9.11], we thus have

$$(23) \quad \left\| \partial_s \partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right\|_{L^p(\mathbb{B}_{4-\bar{q}\beta}(w))} \leq \left\| D^2 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right\|_{L^p(\mathbb{B}_{4-\bar{q}\beta}(w))} \\ \leq C_1 \left(\left\| \Delta_0 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right\|_{L^p(\mathbb{B}_{4^1-\bar{q}\beta}(w))} + \left\| \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right\|_{L^p(\mathbb{B}_{4^1-\bar{q}\beta}(w))} \right),$$

where $\Delta_0 = \sum_i \partial_i \partial_i$ is the Euclidean Laplacian and the positive constant C_1 depends on m, α (via p), β and k (via \bar{q}). We have that

$$(24) \quad \left\| \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right\|_{L^p(\mathbb{B}_{4^1-\bar{q}\beta}(w))} \leq C$$

by the inductive hypothesis. Concerning the other addend in (23), using the properties of the harmonic chart, we get

$$|\Delta_0 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h}| = \left| \sum_{s=1}^m \partial_{ss}^2 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right| \leq C |\hat{g}_\lambda^{ij} \partial_{ij}^2 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h}| = C |\hat{f}_b|,$$

where \hat{f}_b is defined as \hat{f} in (11), replacing the indices $\gamma_1, \dots, \gamma_{\bar{q}}$ in the definition of \hat{f} with $b_1, \dots, b_{\bar{q}}$. Accordingly, computing as for (16), we get

$$\left\| \Delta_0 \partial_{b_1 \dots b_{\bar{q}}}^{\bar{q}} \hat{h} \right\|_{L^p(\mathbb{B}_{4^1-\bar{q}\beta}(w))} \leq C \left\| \hat{f}_b \right\|_{L^p(\mathbb{B}_{4^1-\bar{q}\beta}(w))} \\ \leq C \left\| \hat{f}_b \right\|_{L^\infty(\mathbb{B}_{4^1-\bar{q}\beta}(w))} \\ \leq C.$$

Inserting this latter and (24) in (23), and combining with (21) and (22), we obtain

$$\left[\partial_{b_1 \dots b_{\bar{q}+1}}^{\bar{q}+1} \hat{h} \right]_{C^{0,\alpha}(\mathbb{B}_{4-\bar{q}\beta}(w))} \leq C.$$

Together with (18), (19) and (20), this gives

$$\left[\hat{f} \right]_{C^{0,\alpha}(\mathbb{B}_{4-\bar{q}\beta}(w))} \leq C.$$

Hence, all the three addends in the RHS of (12) are upper bounded by a positive constant C , so that we have shown the validity of (10) with $q = \bar{q} + 1$. This concludes the proof of Lemma 3.11. \square

We can now come back to the proof of Theorem 3.8. Note that by assumption (A1) $\theta(t) \doteq t\lambda(t)$ is smooth, increasing on $[R_1, +\infty)$ and $\lim_{t \rightarrow +\infty} \theta(t) = +\infty$, hence $\theta^{-1} : [\theta(R_1), +\infty) \rightarrow [R_1, +\infty)$ is well-defined, smooth and increasing. Define $H = H_k \in C^\infty(M \setminus B_{R_H}(o))$ by $H = \theta^{-1} \circ h$, where R_H is chosen large enough so that $h \geq \theta(R_1)$ on $M \setminus B_{R_H}(o)$. First, since θ^{-1} and λ are increasing, from $C_h^{-1}\theta(r) \leq h \leq C_h\theta(r)$ we deduce that

$$H(x) \geq \theta^{-1}(C_h^{-1}\theta(r(x))) \geq \theta^{-1}(C_h^{-1}r(x)\lambda(C_h^{-1}r(x))) = C_h^{-1}r(x),$$

and similarly $H(x) \leq C_h r(x)$. To finish the proof of Theorem 1.2 we are going to prove by induction on j that, for $1 \leq j \leq k$, we have that

$$|\nabla_g^j H|(x) \leq C^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}.$$

We first consider the case $j = 1$. Then $|\nabla_g h|(x) = \theta'(H(x))|\nabla_g H|(x)$, so that, using also Lemma 3.1,

$$\begin{aligned} |\nabla_g H|(x) &= \frac{|\nabla_g h|(x)}{\theta'(H(x))} \\ &\leq \frac{C_h \lambda(r(x))}{\theta'(H(x))} \\ &\leq C. \end{aligned}$$

Similarly, suppose that for some $1 \leq \bar{j} \leq k$ and all $1 \leq j \leq \bar{j}-1$, $|\nabla_g^j H|(x) \leq C^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$.

We have that $\nabla_g^{\bar{j}} h(x) - \nabla_g^{\bar{j}} H(x)\theta'(H)$ can be written as a linear combination of terms of the form $\theta^{(s)}(H) \otimes_{i=1}^s \nabla_g^{e_i} H$, with $2 \leq s \leq \bar{j}$ and $\sum_{i=1}^s e_i = \bar{j}$. By inductive hypothesis, using also Lemma 3.2, we have that

$$\begin{aligned} |\theta^{(s)}(H) \otimes_{i=1}^s \nabla_g^{e_i} H| &\leq |\theta^{(s)}(H)| \prod_{i=1}^s |\nabla_g^{e_i} H| \\ &\leq |\theta^{(s)}(H)| \prod_{i=1}^s \lambda(r(x))^{e_i-1} \\ &\leq C \lambda(H(x))^s \lambda(r(x))^{\bar{j}-s} \\ &\leq C \lambda(r(x))^{\bar{j}}. \end{aligned}$$

In particular, by Lemmas 3.11 and 3.12,

$$\begin{aligned} |\nabla_g^{\bar{j}} H(x)| &\leq C \frac{|\nabla_g^{\bar{j}} h(x)| + \sum_{s=2}^{\bar{j}} |\theta^{(s)}(H) \otimes_{i=1}^s \nabla_g^{e_i} H|}{\theta'(H(x))} \\ &\leq C \frac{\lambda_1^{\bar{j}}}{\lambda(r(x))} \\ &\leq C \lambda(r(x))^{\bar{j}-1}. \end{aligned}$$

We have thus proved that there exists a function $H \in C^\infty(M \setminus B_{R_H}(o))$ such that for any x outside a compact set K of M

- (i) $C_h^{-1}r(x) \leq H(x) \leq C_h r(x)$;
- (ii) for $1 \leq j \leq k$, $|\nabla_g^j H|(x) \leq C \lambda(r(x))^{j-1}$.

Replacing H with H/C_h , and choosing a suitable smooth continuation of $H|_{M \setminus K}$ inside the compact K , we get that, up to possibly increase the value of C , it holds

- (i) $C^{-2}r(x) \leq H(x) \leq \max\{r(x), 1\}$;
- (ii) for $1 \leq j \leq k$, $|\nabla_g^j H|(x) \leq C^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$;

as desired.

3.4. Proof of Theorem 1.2: case (b). In this section, we are going to prove Theorem 1.2 in the set of assumptions (b), i.e.

Theorem 3.13. *Let (M^m, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}_g(x, o)$. Let $k \in \mathbb{N}^+$. If $k \geq 2$, suppose in addition that for some $D > 0$,*

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad |\text{Sect}_g|(x) \leq D\lambda(r(x))^2 \quad \text{on } M,$$

where the function λ satisfies assumptions (A1), (A2), (A3), (A4(j)) for $j = 1, \dots, k$. Then there exists an exhaustion function $H = H_k \in C^\infty(M)$ such that for some positive constant $C_H > 1$ independent of x , we have on M that

- (i) $C_H^{-2}r(x) \leq H(x) \leq \max\{r(x), 1\}$;
- (ii) for $1 \leq j \leq k$, $|\nabla_g^j H|(x) \leq C_H^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$.

We proceed as in [20, Subsection 4.2]. Note that the first part of the proof of Theorem 3.8 does not require the control on the injectivity radius. In particular, we can suppose by induction that there exists a distance-like function $H_{k-1} \in C^\infty(M)$ such that for some positive constant $C_{k-1} > 1$ independent of x and o , we have on M that

- (i) $C_{k-1}^{-2}r(x) \leq H_{k-1}(x) \leq \max\{r(x), 1\}$;
- (ii) for $1 \leq j \leq k-1$, $|\nabla_g^j H_{k-1}|_g(x) \leq C_{k-1}^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$.

By Theorem 3.5 we get also in the present assumptions the validity of Proposition 3.9. In particular there exists $h = h_k \in C^\infty(M)$ such that

- (i) $C_h^{-1}r(x)\lambda(r(x)) \leq h(x) \leq C_h \max\{1; r(x)\lambda(r(x))\}$;
- (ii) $|\nabla_g h|_g \leq C_h \lambda(r(x))$ on M ;
- (iii) $\Delta_g h = |\nabla_g h|_g^2 - C_\theta(\theta'(H_{k-1}(x)))^2$ on $M \setminus B_{R_\theta}(o)$ for some constants $C_\theta > 0$ and $R_\theta > 0$.

Fix $R_0 \in \mathbb{R}^+$ such that $\lambda(2R_0 + 1) > \frac{\pi}{2R_0}$. This is always possible since λ is strictly positive and non-decreasing. If $x \in M$ satisfies $r(x) > 1 + R_0$, then on $B_{R_0}(x)$

$$|\text{Sect}_g| \leq D^2 \lambda^2(R_0 + r(x)) \doteq D^2 K_x^2.$$

By a localized version of the Cartan-Hadamard theorem (see e.g. [15, Lemma 2.7]) we have that, for every $0 < R < \min\left\{\frac{\pi}{DK_x}, R_0\right\}$, there exists a smooth complete Riemannian manifold (\bar{M}, \bar{g}) , $\bar{x} \in \bar{M}$ and a smooth surjective local isometry

$$F \doteq F_{g,x,R} : B_{\bar{R}}^{\bar{g}}(\bar{x}) \rightarrow B_R^g(x)$$

such that

- $F(\bar{x}) = x$;
- $\text{inj}_{\bar{g}}(\bar{x}) \geq R$;
- $|\text{Sect}_{\bar{g}}| \leq D^2 K_x^2$ on $B_{\bar{R}}^{\bar{g}}(\bar{x})$;
- $F(B_r^{\bar{g}}(\bar{x})) = B_r^g(x)$, for all $0 < r < R$.

In particular, for every $\bar{y} \in B_{\bar{R}/2}^{\bar{g}}(\bar{x})$, we have that

$$(25) \quad |\text{Sect}_{\bar{g}}|(\bar{y}) \leq D^2 K_x^2, \quad |\nabla_{\bar{g}}^j \text{Ric}_{\bar{g}}|(\bar{y}) \leq K_x^{2+j}, \quad \text{inj}_{\bar{g}}(\bar{y}) \geq d_{\bar{g}}(\bar{y}, \partial B_{\bar{R}}^{\bar{g}}(\bar{x})) \geq \frac{R}{2}.$$

Letting $\lambda_{R_0}^2 \doteq (m-1)K_x^2$, we set

$$\bar{g}_\lambda = \lambda_{R_0}^2 \bar{g}.$$

Then, by (25),

$$|\text{Ric}_{\bar{g}_\lambda}|(\bar{y}) \leq D^2, \quad |\nabla_{\bar{g}_\lambda}^j \text{Ric}_{\bar{g}_\lambda}|(\bar{y}) \leq 1, \quad \text{inj}_{\bar{g}_\lambda}(\bar{y}) \geq \lambda_{R_0} \frac{R}{2}.$$

We can choose $R = \frac{\pi}{2DK_x} < R_0$, obtaining that

$$\text{inj}_{\bar{g}_\lambda}(\bar{y}) \geq \frac{\sqrt{m-1}\pi}{4D} \doteq i_0.$$

Proposition 3.10 hence yields that there exists a constant $C_{HR} = C_{HR}(m, Q, i_0, k)$ independent of x such that on $B_{C_{HR}}^{\bar{g}_\lambda}(x) \subset \bar{M}$ there exists a centered harmonic chart $\varphi_H = (y^1, \dots, y^m) : B_{C_{HR}}^{\bar{g}_\lambda}(x) \rightarrow U \subset \mathbb{R}^m$ such that $\varphi_H(x) = 0$ and, setting $\hat{g}_\lambda = \bar{g}_\lambda \circ \varphi_H^{-1}$, it holds

$$\begin{aligned} (1_{HC}) \quad & Q^{-1}\delta_{ij} \leq (\hat{g}_\lambda)_{ij} \leq Q\delta_{ij} \text{ as bilinear forms;} \\ (2_{HC}) \quad & \sum_{1 \leq |\gamma| \leq k-1} C_{HR}^{|\gamma|} \sup_y \left| \partial_\gamma (\hat{g}_\lambda)_{ij}(y) \right| \leq Q - 1, \\ (1'_{HC}) \quad & Q^{-1}\delta^{ij} \leq \hat{g}_\lambda^{ij} \leq Q\delta^{ij}; \\ (2'_{HC}) \quad & \sum_{1 \leq |\gamma| \leq k-1} \sup_y \left| \partial_\gamma \hat{g}_\lambda^{ij}(y) \right| \leq C(Q), \end{aligned}$$

for some constant $C(Q)$, depending only on Q .

For a fixed x , we can define $\bar{h} : B_R^{\bar{g}}(\bar{x}) \rightarrow \mathbb{R}$ by $\bar{h} = h \circ F$ and $\hat{h} : \mathbb{B}_\beta \rightarrow \mathbb{R}$ by $\hat{h} = \bar{h} \circ \varphi_H^{-1}$.

Since F is a local isometry, at this stage, we can estimate the covariant derivatives of \hat{h} exactly as we did in the proof of Theorem 3.8. In order to get our conclusion, we deduce from these estimates a control on the covariant derivatives of \bar{h} , hence of h , and finally of $H = \theta^{-1} \circ h$ outside a compact set of M .

4. k -TH ORDER CUT-OFFS AND APPLICATION TO THE DENSITY PROBLEM

In the following corollary we notice that higher order exhaustion functions, as the ones which we obtained in the proof of Theorem 1.2, permit to construct k -th order cut-off functions. These, in turn, will allow us to conclude the proof of the density result Theorem 1.5.

Corollary 4.1. *Let (M, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}_g(x, o)$. Let $k \geq 2$ be an integer. Let λ satisfy assumption (A1), (A2), (A3) and (A4(j)), $j = 1, \dots, k$, and suppose that $\lambda^{1-k} \notin L^1([1, +\infty))$. Suppose that one of the following curvature assumptions holds*

(a) *for some $i_0 > 0$,*

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-2, \quad \text{inj}_g(x) \geq \frac{i_0}{\lambda(r(x))} > 0 \quad \text{on } M.$$

(b) *for some $D > 0$,*

$$|\nabla_g^j \text{Ric}_g|(x) \leq \lambda(r(x))^{2+j}, \quad 1 \leq j \leq k-2, \quad |\text{Sect}_g|(x) \leq D^2 \lambda(r(x))^2 \quad \text{on } M.$$

Then there exist a family of cut-off functions $\{\chi_R\}_{R>3} \subset C_c^\infty(M)$ and a constant $C_\chi > 0$ independent of R such that

- (1) $\chi_R = 1$ on $B_{C_H^{-1}(R-2)}(o)$, with C_H the constant appearing in Theorem 1.2;
- (2) $|\nabla_g^j \chi_R| \leq C_\chi$ for $j = 1, \dots, k$.

Proof. Let α_R be a positive constant and define the family $\{\psi_R\}_{R>3}$ of functions on \mathbb{R} by

$$\psi_R(t) \doteq \begin{cases} \max\{1 - \int_R^t \alpha_R \lambda^{1-k}(s) ds; 0\}, & \text{if } t > R \\ 1, & \text{if } t \leq R. \end{cases}$$

We note that $\psi_R \in C^0(\mathbb{R})$ and, by the assumptions on λ , for every R there exists a constant $T(R) \in \mathbb{R}$ depending on R such that $\psi_R(t) \equiv 0$ if and only if $t \in [T(R), \infty)$. We can choose the positive constant α_R small enough so that $T(R) - R > 4$. Note also that $\alpha_R \doteq \alpha$ can be chosen independent of R since λ is increasing. Clearly the ψ_R are not C^2 , so we want to regularize it in a neighborhood of R and $T(R)$, keeping the derivatives controlled.

Let $\zeta : \mathbb{R} \rightarrow \mathbb{R}$ be a C^∞ increasing function such that $\zeta \equiv 0$ in $(-\infty, -1]$, $\zeta \equiv 1$ in $[1, +\infty)$. There exists $q_R \in [-1, 1]$ such that

$$(26) \quad \int_0^{R+2} \frac{\zeta(t+R-q_R)}{\lambda^{k-1}(t)} dt = \int_0^{R+2} \frac{\mathbb{1}_{[R,+\infty)}(t)}{\lambda^{k-1}(t)} dt = \int_R^{R+2} \frac{1}{\lambda^{k-1}(t)} dt.$$

In fact, we have that

$$\int_0^{R+2} \frac{\zeta(t+R-1)}{\lambda^{k-1}(t)} dt \leq \int_0^{R+2} \frac{\mathbb{1}_{[R,+\infty)}(t)}{\lambda^{k-1}(t)} dt \leq \int_0^{R+2} \frac{\zeta(t+R+1)}{\lambda^{k-1}(t)} dt,$$

so that (26) follows by continuity. Similarly, we have also that there exists $Q_R \in [-1, 1]$ such that

$$(27) \quad \int_{R+2}^{T(R)+2} \frac{\zeta(-t+T(R)+Q_R)}{\lambda^{k-1}(t)} dt = \int_{R+2}^{T(R)+2} \frac{\mathbb{1}_{[0,T(R)]}(t)}{\lambda^{k-1}(t)} dt = \int_{R+2}^{T(R)} \frac{1}{\lambda^{k-1}(t)} dt.$$

Now for every $R > 1$, define the real smooth function

$$\mu_R(t) \doteq \frac{\zeta(-t+T(R)+Q_R)\zeta(t-R+q_R)}{\lambda^{k-1}(t)}$$

and

$$\phi_R(t) \doteq 1 - \alpha \int_{R-2}^t \mu_R(s) ds.$$

Then $\phi_R \in C^\infty(\mathbb{R})$, ϕ_R is decreasing, $\phi_R \equiv 1$ if $t < R-2$ and $\phi_R = \psi_R \equiv 0$ if $t > T(R)+2$ because of (26) and (27). Moreover

$$|\phi'_R(t)| = \alpha |\mu_R(t)| \leq \frac{\alpha}{\lambda^{k-1}(t)}.$$

Similarly, let $Z_R(t) \doteq \zeta(-t+T(R)+Q_R)\zeta(t-R+q_R)$ and $\Lambda(t) \doteq \lambda^{1-k}(t)$. Then, for $j = 2, \dots, k$

$$|\phi_R^{(j)}(t)| = \alpha |\mu_R^{(j-1)}(t)| \leq \alpha \sum_{s=1}^{j-1} \binom{j-1}{s} |Z_R^{(s)}(t)| |\Lambda^{(j-1-s)}(t)|.$$

We have that

$$|Z_R^{(s)}(t)| \leq 2^s \max_{0 \leq l \leq k} \|\zeta^{(l)}\|_\infty^2 < C$$

independently of R . Moreover $\Lambda^{(j-1-s)}(t)$ can be written as a linear combination of terms of the form $\lambda^{1-k-u}(t) \prod_{i=1}^u \lambda^{(e_i)}(t)$, with $1 \leq u \leq j-1-s$ and $\sum_{i=1}^u e_i = j-1-s$. Since, by assumption (A4(j)), $\lambda^{(e_i)}(t) \leq M_{4(e_i)} \lambda^{e_i}(t)$, we get

$$\left| \lambda^{1-k-u}(t) \prod_{i=1}^u \lambda^{(e_i)}(t) \right| \leq C \lambda^{1-k-u}(t) \lambda^{j-1-s}(t)$$

and

$$|\Lambda^{(j-1-s)}(t)| \leq C \lambda^{1-k}(t)$$

for every $s = 1, \dots, j-1$, so that

$$(28) \quad |\phi_R^{(j)}(t)| \leq C \lambda^{1-k}(t)$$

for every $j = 2, \dots, k$.

Now, define the family of cut-off functions $\{\chi_R\}_{R>3} \subset C_c^\infty(M)$ by $\chi_R \doteq \phi_R \circ H$, where H is the k -th order distance-like function given by Theorem 1.2. Note that if $r(x) \leq C_H^{-1}(R-2)$, then

$\chi_R(x) = 1$. Moreover, we have that for every $j = 1, \dots, k$, $\nabla_g^j \chi_R(x)$ can be written as a linear combination of terms of the form $\phi_R^{(s)}(H) \otimes_{i=1}^s \nabla_g^{e_i} H$, with $1 \leq s \leq j$ and $\sum_{i=1}^s e_i = j$. We have

$$\begin{aligned} |\phi_R^{(s)}(H) \otimes_{i=1}^s \nabla_g^{e_i} H| &\leq |\phi_R^{(s)}(H)| \prod_{i=1}^s |\nabla_g^{e_i} H| \\ &\leq |\phi_R^{(s)}(H)| \prod_{i=1}^s \lambda(r(x))^{e_i-1} \\ &\leq C \lambda(H(x))^{1-k} \lambda(r(x))^{j-s} \\ &\leq C, \end{aligned}$$

and thus

$$|\nabla_g^j \chi_R| \leq C$$

for every $j = 1, \dots, k$. □

4.1. Proof of the density result. We can now give the proof of the following result which we stated in Section 1 as Theorem 1.5.

Theorem 4.2. *In the assumptions of Corollary 4.1, we have that $W^{k,p}(M) = W_0^{k,p}(M)$ for all $p \in [1, +\infty)$.*

Proof (of Theorem 1.5). We can apply Corollary 4.1 and get the existence of a sequence of cut-off functions $\{\chi_n\}$ with uniformly bounded covariant derivatives up to order k . This suffices to get the desired density result. Indeed, first recall that $C^\infty(M) \cap W^{k,p}(M)$ is dense in $W^{k,p}(M)$ (see for instance [12]). Then, given a smooth function $f \in C^\infty(M) \cap W^{k,p}(M)$, define $f_n \doteq \chi_n f$. One obtains that

$$(29) \quad \|f_n - f\|_{L^p} = \|(1 - \chi_n)f\|_{L^p},$$

$$(30) \quad \|\nabla_g(f_n - f)\|_{L^p} \leq \|f \nabla_g \chi_n\|_{L^p} + \|(1 - \chi_n) \nabla_g f\|_{L^p},$$

$$(31) \quad \|\nabla_g^j(f_n - f)\|_{L^p} \leq \sum_{l=0}^{j-1} \left\| C_l |\nabla_g^{j-l} \chi_n| |\nabla_g^l f| \right\|_{L^p} + \|(1 - \chi_n) \nabla_g^j f\|_{L^p}, \quad j = 2, \dots, k,$$

where C_l are integer constants depending on j and l . Note that each of $(1 - \chi_n)$, $\{\nabla^j \chi_n\}_{j=1, \dots, k}$ is uniformly bounded and supported in $\text{supp}(1 - \chi_n)$. Moreover, given any compact set $K \subset M$, we have that $\text{supp}(1 - \chi_n) \subset M \setminus K$ for $n \gg 1$. since $f \in W^{k,p}(M)$ this permits to conclude that all the terms at the RHS of (29), (30), (31) tend to 0 as $n \rightarrow \infty$. More precisely we have that both the terms of the form

$$\left\| |\nabla_g^{j-l} \chi_n| |\nabla_g^l f| \right\|_{L^p} \leq C_\chi \left\| \nabla_g^l f \right\|_{L^p(\text{supp}(1 - \chi_n))}$$

and the terms of the form

$$\|(1 - \chi_n) \nabla_g^j f\|_{L^p} \leq \left\| \nabla_g^j f \right\|_{L^p(\text{supp}(1 - \chi_n))}$$

go to 0 as $n \rightarrow \infty$ since $f \in W^{k,p}(M)$. □

5. CASE P=2

5.1. k -th order (rough) Laplacian cut-offs. In this subsection we prove versions of Theorem 3.13 and Corollary 4.1 under weaker assumptions. Namely, we assume a control on the derivatives of the Ricci curvature up to a smaller order. As a price to pay, we do not get a control on the whole k -th order covariant derivative $\nabla^k H$ of the distance-like function H , but only on its trace, i.e. the rough Laplacian of the $(k - 2)$ -th covariant derivative of H . This result will be used in Corollary 5.2 to construct a family of k -th order (rough) Laplacian cut-offs. In the rest of this section

we will combine these (rough) Laplacian cut-offs and the Weitzenböck formula for the Sampson-Lichnerowicz Laplacian to get the density of smooth compactly supported function in $W^{k,2}(M)$.

Theorem 5.1. *Let (M, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point, $r(x) \doteq \text{dist}_g(x, o)$. Let $k \geq 3$ be an integer. Let λ satisfy assumption (A1), (A2), (A3) and (A4(j)), $j = 1, \dots, k-1$. Suppose that*

$$(32) \quad |\text{Sect}|(x) \leq \lambda(r(x))^2, \quad \text{and} \quad |\nabla^j \text{Ric}|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-3.$$

Then there exists an exhaustion function $H \in C^\infty(M)$ such that for some positive constant $C_\Delta > 1$ independent of x , we have on M that

- (i) $C_\Delta^{-2}r(x) \leq H(x) \leq \max\{r(x), 1\}$;
- (ii) $|\nabla^j H| \leq C_\Delta^{j-1} \max\{\lambda(r(x))^{j-1}, 1\}$, $1 \leq j \leq k-1$;
- (iii) $|\Delta \nabla^{k-2} H| \leq C_\Delta \max\{\lambda(r(x))^{k-1}, 1\}$.

When acting on sections of a tensor bundle, Δ denotes the rough Laplacian, i.e. $\Delta T = -\nabla^* \nabla T = \text{tr}_{12} \nabla^2 T$, for any tensor field T . Note that Δ is equal to minus the Bochner Laplacian Δ_B we will use in Subsection 5.2.

Proof. Note that in the proof of Theorem 3.13, the assumption $|\nabla^{k-2} \text{Ric}| \leq \lambda(r(x))^k$ was used only to control $|\nabla^k h|$. Accordingly, by the proof of Theorem 3.13 we already know that there exists a smooth exhaustion function $h \in C^\infty(M)$ such that

$$(33) \quad \begin{aligned} C^{-1}r(x)\lambda(r(x)) &\leq h(x) \leq C \max\{1; r(x)\lambda(r(x))\} \quad \forall x \in M \\ |\nabla^j h|(x) &\leq C\lambda^j(r(x)) \quad \forall x \in M \setminus \bar{B}_{\bar{\rho}}(o), \quad j = 1, \dots, k-1. \end{aligned}$$

Moreover, by construction

$$(34) \quad \Delta h = |\nabla h|^2 - C_\theta (\theta'(H_{k-2}(x)))^2 = |\nabla h|^2 - C_\theta \Theta(H_{k-2}(x)) \quad \text{on } M \setminus \bar{B}_{R_\theta}(o),$$

for some constants $C_\theta > 0$ and $R_\theta > 0$. Here H_{k-2} is such that for some positive constant $C > 1$ (independent of x and o) we have on M that

$$\begin{aligned} Cr(x) &\leq H_{k-2}(x) \leq \max\{r(x), 1\} \\ |\nabla H_{k-2}|(x) &\leq 1 \\ |\nabla^j H_{k-2}|(x) &\leq C \max\{\lambda^{j-1}(x), 1\}, \quad j = 2, \dots, k-2. \end{aligned}$$

Taking ∇^{k-2} of (34) we obtain that

$$\nabla^{k-2} \Delta h = 2\nabla^{k-3} (\nabla^2 h(\nabla h, \cdot)) - C_\theta \nabla^{k-2} (\Theta(H_{k-2})).$$

Hence

$$|\nabla^{k-2} \Delta h| \leq C \left[|\nabla^{k-1} h| |\nabla h| + |\nabla^{k-2} h| |\nabla^2 h| + \dots + |\nabla^2 h| |\nabla^{k-2} h| \right] + C_\theta |\mathcal{L}|.$$

Here \mathcal{L} is a linear combination of terms of the form

$$\Theta^{(j)}(H_{k-2}) \prod_{s=1}^j \nabla^{e_s - e_{s-1}} H_{k-2},$$

where $e_0 = 0$, $j \in \{1, \dots, k-2\}$ and $\{e_s\}_{s=1}^j$ is an increasing subset of $\{1, \dots, k-1\}$ with $e_j = k-2$. Using the above properties of h and H_{k-2} and Lemma 3.3 we hence get that

$$|\nabla^{k-2} \Delta h| \leq C\lambda^k(r(x)).$$

By Lemma A.1 in Appendix A, we thus obtain that

$$\begin{aligned}
(35) \quad |\Delta \nabla^{k-2} h| &= |\nabla^{k-2} \Delta h| + |\text{Riem} * \nabla^{k-2} h + \nabla \text{Ric} * \nabla^{k-3} h + \dots + \nabla^{k-3} \text{Ric} * \nabla h| \\
&\leq |\nabla^{k-2} \Delta h| + C \left(|\text{Riem}| |\nabla^{k-2} h| + |\nabla \text{Ric}| |\nabla^{k-3} h| + \dots + |\nabla^{k-3} \text{Ric}| |\nabla h| \right) \\
&\leq C \left(\lambda(r(x))^k + \sum_{j=0}^{k-3} \lambda(r(x))^{2+j} \lambda(r(x))^{k-2-j} \right) \\
&\leq C \lambda(r(x))^k.
\end{aligned}$$

For the definition of the notation $*$ appearing in the latter formula see Section 2.

As in the proof of Theorem 1.2 define $H \doteq H_{k-1} \in C^\infty(M \setminus B_{R_H}(o))$ by $H = \theta^{-1} \circ h$. Then H is distance-like and

$$|\nabla^j H| \leq C^{j-1} \max \{ \lambda(r(x))^{j-1}, 1 \} \quad 1 \leq j \leq k-1.$$

Moreover we have that

$$\Delta \nabla^{k-2} h = \Delta \left(\theta'(H) \nabla^{k-2} H \right) + \mathcal{L}_1,$$

where \mathcal{L}_1 is a linear combination of terms of the form $\Delta \left(\theta^{(s)}(H) \otimes \nabla^{e_i} H \right)$, with $2 \leq s \leq k-2$ and $\sum_{i=1}^s e_i = k-2$. Letting $\{E_i\}$ being a local orthonormal frame on M , note that

$$\begin{aligned}
\Delta \left(\theta'(H) \nabla^{k-2} H \right) &= \sum_i \nabla_{E_i} \left(\theta''(H) \nabla_{E_i} H \nabla^{k-2} H + \theta'(H) \nabla_{E_i} \nabla^{k-2} H \right) \\
&= \theta'''(H) |\nabla H|^2 \nabla^{k-2} H + \theta''(H) \Delta H \nabla^{k-2} H \\
&\quad + 2\theta''(H) \nabla_{\nabla H} \nabla^{k-2} H + \theta'(H) \Delta \nabla^{k-2} H,
\end{aligned}$$

and

$$\begin{aligned}
\Delta \left[\theta^{(s)}(H) \otimes_{i=1}^s \nabla^{e_i} H \right] &= \sum_i \nabla_{E_i} \left(\theta^{(s+1)}(H) \nabla_{E_i} H \otimes \bigotimes_{i=1}^s \nabla^{e_i} H + \theta^{(s)}(H) \nabla_{E_i} \left(\bigotimes_{i=1}^s \nabla^{e_i} H \right) \right) \\
&= \theta^{(s+2)}(H) |\nabla H|^2 \bigotimes_{i=1}^s \nabla^{e_i} H + \theta^{(s+1)}(H) \Delta H \bigotimes_{i=1}^s \nabla^{e_i} H \\
&\quad + 2\theta^{(s+1)}(H) \nabla_{\nabla H} \left(\bigotimes_{i=1}^s \nabla^{e_i} H \right) + \theta^{(s)}(H) \Delta \left(\bigotimes_{i=1}^s \nabla^{e_i} H \right).
\end{aligned}$$

Hence, using Lemma 3.1, Lemma 3.2, and proceeding as for the end of the proof of Theorem 3.8, we get that

$$\begin{aligned}
|\Delta \nabla^{k-2} H| &= \frac{C}{\theta'(H)} \left[|\Delta \nabla^{k-2} h| + |\theta'''(H)| |\nabla H|^2 |\nabla^{k-2} H| + |\theta''(H)| |\Delta H| |\nabla^{k-2} H| \right. \\
&\quad + |\theta''(H)| |\nabla_{\nabla H} \nabla^{k-2} H| + \max_{s=2, \dots, k-2} |\theta^{(s+2)}(H)| |\nabla H|^2 \left| \bigotimes_{i=1}^s \nabla^{e_i} H \right| \\
&\quad + \max_{s=2, \dots, k-2} |\theta^{(s+2)}| |\theta^{(s+1)}(H)| |\Delta H| \left| \bigotimes_{i=1}^s \nabla^{e_i} H \right| \\
&\quad \left. + \max_{s=2, \dots, k-2} \left(|\theta^{(s+1)}(H)| |\nabla_{\nabla H} \bigotimes_{i=1}^s \nabla^{e_i} H| + |\theta^{(s)}(H)| \left| \Delta \left(\bigotimes_{i=1}^s \nabla^{e_i} H \right) \right| \right) \right] \\
&\leq C \lambda(r)^{k-1}.
\end{aligned}$$

□

Using the function H coming from Theorem 5.1, we want now to produce a sequence of higher order (rough) Laplacian cut-off functions. This will be done in the following

Corollary 5.2. *Let (M, g) be a complete Riemannian manifold and $o \in M$ a fixed reference point $r(x) \doteq \text{dist}_g(x, o)$. Let $k \geq 3$ be an integer. Let λ satisfy assumptions (A1), (A2), (A3), and (A4(j)) for $j = 1, \dots, k-1$, and suppose that $\lambda^{1-k} \notin L^1([1, +\infty))$. Suppose that*

$$(36) \quad |\nabla^j \text{Riem}|(x) \leq \lambda(r(x))^{2+j}, \quad 0 \leq j \leq k-3.$$

Then there exists a family of cut-off functions $\{\chi_R\} \subset C_c^\infty(M)$, and a constant $C > 0$ independent of R such that,

- (1) $\chi_R = 1$ on $B_{C_H^{-1}(R-2)}(o)$;
- (2) $|\nabla^j \chi_R| \leq C \lambda^{-k+j}$, $j = 1, \dots, k-1$;
- (3) $|\Delta \nabla^{k-2} \chi_R| \leq C$,

Proof. As in Corollary 4.1, For each radius $R \gg 1$, define $\chi_R \doteq \phi_R \circ H$, with H the distance-like function given by Theorem 5.1 and ϕ_R defined as in the proof of Corollary 4.1. Properties (1) and (2) follow from the proof of Corollary 4.1. About (3), using Lemma A.1 we note that

$$(37) \quad |\Delta \nabla^{k-2} \chi_R| \leq |\nabla^{k-2} \Delta \chi_R| + |\text{Riem} * \nabla^{k-2} \chi_R + \nabla \text{Ric} * \nabla^{k-3} \chi_R + \dots \nabla^{k-3} \text{Ric} * \nabla \chi_R|.$$

About the first term on the RHS of (37), note that

$$\Delta(\phi_R \circ H) = \phi'_R \Delta H + \phi''_R |\nabla H|^2,$$

and hence

$$|\nabla^{k-2} \Delta \chi_R| = |\nabla^{k-2}(\phi'_R \Delta H)| + |\nabla^{k-2}(\phi''_R |\nabla H|^2)|.$$

We have that $\nabla^{k-2}(\phi'_R \Delta H)$ can be written as

$$(38) \quad \nabla^{k-2}(\phi'_R \Delta H) = \sum_{\mathbf{c}} C_{\mathbf{c}} \phi^{(c_1 + \dots + c_{k-3} + 1)} \left[\bigotimes_{t=1}^{k-3} (\nabla^t H)^{\otimes c_t} \right] \otimes \nabla^{c_{k-2}} \Delta H,$$

where \mathbf{c} varies among all $(k-2)$ -vectors of nonnegative integers such that $\sum_{t=1}^{k-3} t c_t + c_{k-2} = k-2$, and the $C_{\mathbf{c}}$ are positive integer constants.

With the same notations, $\nabla^{k-2}(\phi''_R |\nabla H|^2)$ can be written as

$$(39) \quad \begin{aligned} \nabla^{k-2}(\phi''_R |\nabla H|^2) &= \sum_{\mathbf{c}} C_{\mathbf{c}} \phi^{(c_1 + \dots + c_{k-3} + 2)} \left[\bigotimes_{t=1}^{k-3} (\nabla^t H)^{\otimes c_t} \right] \otimes \nabla^{c_{k-2}} |\nabla H|^2 \\ &= \sum_{\mathbf{c}} C_{\mathbf{c}} \phi^{(c_1 + \dots + c_{k-3} + 2)} \left[\bigotimes_{t=1}^{k-3} (\nabla^t H)^{\otimes c_t} \right] \otimes \nabla^{c_{k-2}-1} (\nabla^2 H(\nabla H, \cdot)). \end{aligned}$$

Inserting (28) in (38) and using Theorem 5.1 (also combined with Lemma A.1), we get

$$|\nabla^{k-2}(\phi'_R \Delta H)| \leq C \lambda^{1-k} \left[\lambda^{\sum_{t=1}^{k-3} (t c_t - 1) + c_{k-2} + 1} \right] \leq C \lambda^{1-k} \lambda^{k-2} = C \lambda^{-1} = C.$$

Analogously, recalling also that

$$|\nabla^{c_{k-2}-1} (\nabla^2 H(\nabla H, \cdot))| \leq C \left[|\nabla^{c_{k-2}+1} H| |\nabla H| + |\nabla^{c_{k-2}} H| |\nabla^2 H| + \dots + |\nabla^2 H| |\nabla^{c_{k-2}} H| \right],$$

by (39), we get that

$$|\nabla^{k-2}(\phi''_R |\nabla H|^2)| \leq C \lambda^{1-k} \left[\lambda^{\sum_{t=1}^{k-3} (t c_t - 1) + c_{k-2}} \right] \leq C \lambda^{1-k} \left[\lambda^{k-2+1} \right] \leq C.$$

About the second term on the RHS of (37), we have that by (36) and property (2) of the cut-off functions,

$$\begin{aligned} |\text{Riem}||\nabla^{k-2}\chi_R| &\leq C\lambda^2\lambda^{-k+k-2} = C, \\ |\nabla^l\text{Ric}||\nabla^{k-2-l}\chi_R| &\leq C\lambda^{2+l}\lambda^{-k+k-2-l} = C, \end{aligned}$$

for $l = 1, \dots, k-3$. This concludes the proof of property (3) of the cut-off functions and hence yields the validity of the Lemma. \square

5.2. Weitzenböck formulas. Fix a tensor bundle $E \rightarrow M$ with m -dimensional fibers, endowed with an inner product induced by the metric g and a compatible connection ∇ induced by the Levi-Civita connection on M . A Lichnerowicz Laplacian is a second order differential operator acting on the space of smooth sections $\Gamma(E)$ of the form

$$(40) \quad \Delta_L = \Delta_B + c\mathfrak{Ric},$$

for a suitable constant c . Here $\Delta_B = -\text{tr}_{12}(\nabla^2) = \nabla^*\nabla$ is the Bochner Laplacian (with ∇^* the formal L^2 -adjoint of ∇) and \mathfrak{Ric} is a smooth symmetric endomorphism of E known as Weitzenböck curvature operator. When T is a $(0, k)$ -tensor, the Weitzenböck curvature operator, takes the form

$$(41) \quad \mathfrak{Ric}(T)(X_1, \dots, X_k) = \sum_{i=1}^k \sum_j (R(E_j, X_i)T)(X_1, \dots, E_j, \dots, X_k),$$

where $\{E_i\}$ is a local orthonormal frame and

$$R(X, Y) \doteq \nabla_{X,Y}^2 - \nabla_{Y,X}^2 = \nabla_X\nabla_Y - \nabla_Y\nabla_X - \nabla_{[X,Y]},$$

which may be applied to any tensor field. Note that, by the classical Bochner-Weitzenböck formula, the Hodge Laplacian on exterior differential forms decomposes as in (40) with $c = 1$.

Since, obviously, $R(X, Y)$ vanishes on functions, for any $(0, k)$ -tensor T , we have that

$$\begin{aligned} \mathfrak{Ric}(T)(X_1, \dots, X_k) &= - \sum_{i=1}^k \left[\sum_j \sum_{p \neq i} T(X_1, \dots, R(E_j, X_i)X_p, \dots, E_j, \dots, X_k) \right. \\ &\quad \left. + \sum_j T(X_1, \dots, R(E_j, X_i)E_j, \dots, X_k) \right] \\ &= - \sum_{i=1}^k \left[\sum_j \sum_{p \neq i} T(X_1, \dots, \sum_l g(R(E_j, X_i)X_p, E_l)E_l, \dots, E_j, \dots, X_k) \right. \\ &\quad \left. - \sum_\nu T(X_1, \dots, \text{Ric}(E_\nu, X_i)E_\nu, \dots, X_k) \right] \\ &= - \sum_{i=1}^k \left[\sum_j \sum_{p \neq i} \sum_l T(X_1, \dots, E_l, \dots, E_j, \dots, X_k) \text{Riem}(E_j, X_i, X_p, E_l) \right. \\ &\quad \left. - \sum_\nu T(X_1, \dots, E_\nu, \dots, X_k) \text{Ric}(X_i, E_\nu) \right], \end{aligned}$$

where we are setting

$$\begin{aligned} \text{Riem}(X, Y, Z, W) &= g(\text{R}(X, Y)Z, W) \\ \text{Ric}(X, Y) &= \text{tr} \text{Riem}(X, \cdot, \cdot, Y). \end{aligned}$$

This curvature term has a quite complicated expression, but it can be estimated in terms of the curvature operator \mathcal{R} of M (the linear extension to $\Lambda^2 TM$ of the $(2, 2)$ -Riemann curvature tensor); see e.g. [23, Corollary 9.3.4].

Proposition 5.3. *Let T be a $(0, s)$ tensor. If the curvature operator \mathcal{R} satisfies $\mathcal{R} \geq \alpha$, for some constant $\alpha < 0$, then $\langle \mathfrak{Ric}(T), T \rangle \geq \alpha C |T|^2$, with C constant depending only on s .*

5.3. A Lichnerowicz Laplacian on symmetric $(0, k)$ -tensors. Another remarkable example of differential operator which can be rewritten as a Lichnerowicz Laplacian was introduced by J. H. Sampson in [26] on smooth sections of the bundle $S^{(0, k)}(M)$ of totally symmetric $(0, k)$ -tensors (see also [22]).

Namely, consider the symmetrization operator s_k , i.e. the projection of the full tensor bundle $T^{0, k}(M)$ on $S^{(0, k)}(M)$. Given h a $(0, k - 1)$ -tensor field, we can define the totally symmetric tensor

$$h^S(X_1, \dots, X_{k-1}) \doteq s_{k-1}(h)(X_1, \dots, X_{k-1}) = \frac{1}{(k-1)!} \sum_{\sigma \in \Pi_{k-1}} h(X_{\sigma(1)}, \dots, X_{\sigma(k-1)}).$$

Let us define the operator $D_S : \Gamma S^{(0, k-1)}(M) \rightarrow \Gamma S^{(0, k)}(M)$ by

$$(D_S h^S)(X_0, \dots, X_k) = k s_k(\nabla h^S)(X_0, \dots, X_k).$$

Its formal L^2 -adjoint $D_S^* : \Gamma S^{(0, k)}(M) \rightarrow \Gamma S^{(0, k-1)}(M)$ is then given by

$$(D_S^* h^S)(X_1, \dots, X_{k-2}) = - \sum_i (\nabla_{E_i} h^S)(E_i, X_1, \dots, X_{k-2}).$$

Letting $\Delta_{\text{Sym}} = D_S^* D_S - D_S D_S^*$, we have by [26] (see also Appendix B for a proof) that

$$\Delta_{\text{Sym}} h^S = \nabla^* \nabla h^S - \mathfrak{Ric}(h^S),$$

that is to say, Δ_{Sym} is of type (40) for $c = -1$.

In particular, one can compute that

$$\begin{aligned} (42) \quad \frac{1}{2} \Delta |h^S|^2 &= - \langle \nabla^* \nabla h^S, h^S \rangle + |\nabla h^S|^2 \\ &= - \langle \Delta_{\text{Sym}} h^S, h^S \rangle - \langle \mathfrak{Ric}(h^S), h^S \rangle + |\nabla h^S|^2 \end{aligned}$$

It is worth mentioning that for $(0, 2)$ -tensors, in [23, Chapter 9] it was introduced a different Lichnerowicz Laplacian acting on smooth sections of $S^{(0, 2)}(M)$, with particular focus on applications of the Bochner technique to this operator. However the operator Δ_{Sym} seems to conform better to our scope.

5.4. Proof of Theorem 1.7. In our assumptions, we know by Corollary 5.2 that there exists a sequence of cut-off functions $\{\chi_n\} \subset C_c^\infty(M)$, and a constant $C > 0$ independent of n such that,

- (1) $\chi_n = 1$ on $B_{C_H^{-1}(n-2)}(o)$;
- (2) $|\nabla^j \chi_n| \leq C \lambda^{-k+j}$, $j = 1, \dots, k - 1$;
- (3) $|\Delta \nabla^{k-2} \chi_n| \leq C$,

Since smooth functions are dense in $W^{k, 2}(M)$, to prove the density result it is sufficient to consider $f \in C^\infty(M) \cap W^{k, 2}(M)$; see for instance [12]. We want to prove that $\chi_n f$ converges to f in $W^{k, 2}(M)$. The lower order terms can be treated as in the proof of Theorem 4.2, by using the dominated convergence theorem and the properties of the cut-off functions. Here we prove that

$$\int_M |\nabla^k(\chi_n f) - \nabla^k f|^2 d\text{vol}_g \rightarrow 0,$$

as $n \rightarrow \infty$. Note that

$$\begin{aligned} \int_M |\nabla^k(\chi_n f) - \nabla^k f|^2 d\text{vol}_g &= \int_M \left| \left[\sum_{i=0}^k \binom{k}{i} \nabla^{k-i} \chi_n \otimes \nabla^i f \right] - \nabla^k f \right|^2 d\text{vol}_g \\ &\leq \int_M (1 - \chi_n)^2 |\nabla^k f|^2 + \sum_{i=0}^{k-1} \binom{k}{i} \int_M |\nabla^{k-i} \chi_n|^2 |\nabla^i f|^2 d\text{vol}_g \end{aligned}$$

Taking into account the properties of the cut-off functions, the only non-trivial term to study is

$$\int_M |f|^2 |\nabla^k \chi_n|^2$$

To prove that this goes to 0 as $n \rightarrow \infty$ we are going to use (42) with $h \doteq \nabla^{k-1} \chi_n$ and hence $h^S = s_{k-1}(\nabla^{k-1} \chi_n)$. First note that, in general,

$$\begin{aligned} \frac{1}{2} \text{div} \left(f^2 \nabla |h^S|^2 \right) &\leq f^2 \left[-\langle \Delta_{\text{Sym}} h^S, h^S \rangle - \langle \mathfrak{Ric}(h^S), h^S \rangle + |\nabla h^S|^2 \right] \\ &\quad + 2|f| |h^S| |\langle \nabla f, \nabla |h^S| \rangle|. \end{aligned}$$

Hence, by Young's and Kato's inequality, we get that, for any $0 < \eta < 1$,

$$\begin{aligned} \frac{1}{2} \text{div} \left(f^2 \nabla |h^S|^2 \right) + f^2 \langle \Delta_{\text{Sym}} h^S, h^S \rangle &\leq -f^2 \langle \mathfrak{Ric}(h^S), h^S \rangle + f^2 |\nabla h^S|^2 \\ &\quad + \eta f^2 |\nabla |h^S||^2 + \frac{1}{\eta} |\nabla f|^2 |h^S|^2 \\ &\leq -f^2 \langle \mathfrak{Ric}(h^S), h^S \rangle + (1 + \eta) f^2 |\nabla h^S|^2 + \frac{1}{\eta} |\nabla f|^2 |h^S|^2. \end{aligned}$$

Integrating, we get that

$$(43) \quad \begin{aligned} \int_M \langle \Delta_{\text{Sym}} h^S, f^2 h^S \rangle d\text{vol}_g &\leq - \int_M f^2 \langle \mathfrak{Ric}(h^S), h^S \rangle d\text{vol}_g \\ &\quad + (1 + \eta) \int_M f^2 |\nabla h^S|^2 d\text{vol}_g + \frac{1}{\eta} \int_M |\nabla f|^2 |h^S|^2 d\text{vol}_g, \end{aligned}$$

for any $\eta > 0$. Recall that $h^S = s_{k-1}(\nabla^{k-1} \chi_n)$. Note that, by Cauchy-Schwarz inequality,

$$(44) \quad |s_{k-1}(\nabla^{k-1} \chi_n)|^2 \leq |\nabla^{k-1} \chi_n|^2.$$

Hence, by the properties of the cut-off functions χ_n , the dominated convergence theorem, and the fact that $f \in W^{k,2}(M)$, we have that

$$\int_M |\nabla f|^2 |s_{k-1}(\nabla^{k-1} \chi_n)|^2 d\text{vol}_g \rightarrow 0$$

as $n \rightarrow \infty$. Furthermore, the curvature term in (43) can be controlled, under our assumptions, using Proposition 5.3. More precisely we have that

$$(45) \quad - \int_M f^2 \langle \mathfrak{Ric}(h^S), h^S \rangle d\text{vol}_g \leq (-\alpha) C \int_M f^2 |h^S|^2 d\text{vol}_g.$$

Let us now analyze the LHS of (43). We let

$$\begin{aligned} A &= \int_M \langle D_S^* D_S(s_{k-1}(\nabla^{k-1} \chi_n)), f^2 s_{k-1}(\nabla^{k-1} \chi_n) \rangle d\text{vol}_g \\ B &= \int_M \langle D_S D_S^*(s_{k-1}(\nabla^{k-1} \chi_n)), f^2 s_{k-1}(\nabla^{k-1} \chi_n) \rangle d\text{vol}_g, \end{aligned}$$

so that

$$(46) \quad \int_M \langle \Delta_{\text{Sym}} h^S, f^2 h^S \rangle d\text{vol}_g = A - B.$$

First, let us deal with the term B . Using (44) we get

$$(47) \quad \begin{aligned} B &= \int_M \langle D_S^*(s_{k-1}(\nabla^{k-1}\chi_n)), D_S^*(f^2 s_{k-1}(\nabla^{k-1}\chi_n)) \rangle d\text{vol}_g \\ &= \int_M \left[f^2 |D_S^*(s_{k-1}(\nabla^{k-1}\chi_n))|^2 - 2f \langle i_{\nabla f}(s_{k-1}(\nabla^{k-1}\chi_n)), D_S^*(s_{k-1}(\nabla^{k-1}\chi_n)) \rangle \right] d\text{vol}_g \\ &\leq 2 \int_M f^2 |D_S^*(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g + \int_M |\nabla f|^2 |s_{k-1}(\nabla^{k-1}\chi_n)|^2 d\text{vol}_g \\ &\leq 2 \int_M f^2 |D_S^*(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g + \int_M |\nabla f|^2 |\nabla^{k-1}\chi_n|^2 d\text{vol}_g. \end{aligned}$$

In the following we will use the "*" notation we defined in Section 2. Moreover, we will work in a normal orthonormal frame $\{E_i\}$ at $p \in M$, and in frame computations we will use the convention of lowering all indices, summing over repeated indices.

Note that

$$(48) \quad |D_S^*(s_{k-1}(\nabla^{k-1}\chi_n))|^2 = \sum_{i_2, \dots, i_{k-1}} \left[\frac{1}{(k-1)!} \sum_{i_1} \nabla_{i_1} \left(\sum_{\pi \in \Pi_{k-1}} \nabla_{i_{\pi(1)} \dots i_{\pi(k-1)}}^{k-1} \chi_n \right) \right]^2.$$

Recall that for any $(0, r)$ -tensor α , with $r \geq 1$, the standard commutation formula gives that

$$(49) \quad (\nabla_i \nabla_j - \nabla_j \nabla_i) \alpha_{i_1 \dots i_r} = \text{Riem} * \alpha$$

Hence for each of the terms in square parentheses on the RHS of (48) we can trace back to a rough Laplacian of a $(k-2)$ -th covariant derivative of χ_n . For instance, consider the term $\nabla_{i_1} \nabla_{i_2 \dots i_{k-1} i_1}^{k-1} \chi_n$. We can compute (at the point $p \in M$ about which we have selected the normal orthonormal frame):

$$\begin{aligned} \sum_{i_1} \nabla_{i_1} \nabla_{i_2 \dots i_{k-1} i_1}^{k-1} \chi_n &= \sum_{i_1} \nabla_{i_1} \left[\left(\nabla_{i_2 \dots i_{k-1} i_1}^{k-1} - \nabla_{i_2 \dots i_{k-2} i_1 i_{k-1}}^{k-1} \right) \chi_n \right. \\ &\quad \left. + \left(\nabla_{i_2 \dots i_{k-2} i_1 i_{k-1}}^{k-1} - \nabla_{i_2 \dots i_{k-3} i_1 i_{k-2} i_{k-1}}^{k-1} \right) \chi_n + \dots + \nabla_{i_1 i_2 \dots i_{k-1}}^{k-1} \chi_n \right] \\ &= \sum_{i_1} \nabla_{i_1} \left(\left(\nabla^{k-4} \text{Riem} * \nabla \chi_n + \dots + \text{Riem} * \nabla^{k-3} \chi_n \right)_{i_1 \dots i_{k-1}} + \nabla_{i_1 i_2 \dots i_{k-1}}^{k-1} \chi_n \right) \\ &= \left(\nabla^{k-3} \text{Riem} * \nabla \chi_n + \dots + \text{Riem} * \nabla^{k-2} \chi_n \right)_{i_2 \dots i_{k-1}} + \Delta \nabla_{i_2 \dots i_{k-1}}^{k-2} \chi_n. \end{aligned}$$

The other terms can be treated similarly. Hence, by (48) and Young's inequality, we get that

$$|D_S^*(s_{k-1}(\nabla^{k-1}\chi_n))|^2 \leq C(m, k) \left(|\Delta \nabla^{k-2} \chi_n|^2 + |\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right),$$

where (here and from now on) $C(m, k)$ is a constant, depending only on k and m , which can possibly change from line to line. Integrating, we get from (47) that

$$(50) \quad \begin{aligned} B &\leq \int_M 2C(m, k) f^2 \left(|\Delta \nabla^{k-2} \chi_n|^2 + |\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right) d\text{vol}_g \\ &\quad + \int_M |\nabla f|^2 |\nabla^{k-1} \chi_n|^2 d\text{vol}_g. \end{aligned}$$

On the other hand, by Young's inequality and reasoning as in (44), we have that

$$\begin{aligned}
(51) \quad A &= \int_M \langle D_S(s_{k-1}(\nabla^{k-1}\chi_n)), D_S(f^2 s_{k-1}(\nabla^{k-1}\chi_n)) \rangle d\text{vol}_g \\
&= \int_M f^2 |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g \\
&\quad + \int_M k \langle D_S(s_{k-1}(\nabla^{k-1}\chi_n)), 2f s_k(df \otimes s_{k-1}(\nabla^{k-1}\chi_n)) \rangle d\text{vol}_g \\
&\geq (1-\delta) \int_M f^2 |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g - \frac{k^2}{\delta} \int_M |\nabla f|^2 |\nabla^{k-1}\chi_n|^2 d\text{vol}_g,
\end{aligned}$$

for any $\delta > 0$. Substituting (45) in (43), then this latter, (50) and (51) in (46), we hence get that

$$\begin{aligned}
&(-\alpha)C \int_M f^2 |s_{k-1}(\nabla^{k-1}\chi_n)|^2 d\text{vol}_g + (1+\eta) \int_M f^2 |\nabla(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g \\
&+ \frac{1}{\eta} \int_M |\nabla f|^2 |s_{k-1}(\nabla^{k-1}\chi_n)|^2 d\text{vol}_g \\
&\geq (1-\delta) \int_M f^2 |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 d\text{vol}_g - \frac{k^2}{\delta} \int_M |\nabla f|^2 |\nabla^{k-1}\chi_n|^2 d\text{vol}_g \\
&\quad - \int_M C(m,k) f^2 \left(|\Delta \nabla^{k-2}\chi_n|^2 + |\text{Riem}|^2 |\nabla^{k-2}\chi_n|^2 + \dots + |\nabla^{k-3}\text{Riem}|^2 |\nabla\chi_n|^2 \right) d\text{vol}_g \\
&\quad - \int_M |\nabla f|^2 |\nabla^{k-1}\chi_n|^2 d\text{vol}_g.
\end{aligned}$$

Using our assumptions, the properties of the rough Laplacian cut-off functions χ_n , Proposition 5.3, the dominated convergence theorem, and the fact that $f \in W^{k,2}(M)$, this yields that

$$(52) \quad \limsup_{n \rightarrow \infty} \int_M f^2 \left[(1-\delta) |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 - (1+\eta) |\nabla s_{k-1}(\nabla^{k-1}\chi_n)|^2 \right] d\text{vol}_g \leq 0.$$

Let us first study the term $|\nabla s_{k-1}(\nabla^{k-1}\chi_n)|^2$. Since, by definition,

$$s_{k-1}(\nabla^{k-1}\chi_n)(E_{i_1}, \dots, E_{i_{k-1}}) = \frac{1}{(k-1)!} \left(\sum_{\pi \in \Pi_{k-1}} \nabla_{i_{\pi(1)} \dots i_{\pi(k-1)}}^{k-1} \chi_n \right),$$

using (49), we have that

$$\begin{aligned}
&((k-1)!)^2 |\nabla s_{k-1}(\nabla^{k-1}\chi_n)|^2 \\
&= \sum_{i_1, \dots, i_k} \left[\sum_{\pi \in \Pi_{k-1}} \nabla_{i_k i_{\pi(1)} \dots i_{\pi(k-1)}}^k \chi_n \right]^2 \\
&= (k-1)! |\nabla^k \chi_n|^2 + \sum_{i_1, \dots, i_k} \sum_{\pi \in \Pi_{k-1}} (\nabla_{i_k i_{\pi(1)} \dots i_{\pi(k-1)}}^k \chi_n \cdot \sum_{\sigma \in \Pi_{k-1} \setminus \{\pi\}} \nabla_{i_k i_{\sigma(1)} \dots i_{\sigma(k-1)}}^k \chi_n) \\
&= ((k-1)! + ((k-1)! - 1)(k-1)!) |\nabla^k \chi_n|^2 \\
&\quad + \left(\nabla^k \chi_n * \left(\text{Riem} * \nabla^{k-2} \chi_n + \nabla \text{Riem} * \nabla^{k-3} \chi_n + \dots \nabla^{k-3} \text{Riem} * \nabla \chi_n \right) \right) \\
&= ((k-1)!)^2 |\nabla^k \chi_n|^2 + \left(\nabla^k \chi_n * \left(\text{Riem} * \nabla^{k-2} \chi_n + \nabla \text{Riem} * \nabla^{k-3} \chi_n + \dots \nabla^{k-3} \text{Riem} * \nabla \chi_n \right) \right).
\end{aligned}$$

Hence, using Young's inequality,

(53)

$$|\nabla s_{k-1}(\nabla^{k-1}\chi_n)|^2 \leq |\nabla^k \chi_n|^2 + C(m, k) \left[\varepsilon |\nabla^k \chi_n|^2 + \frac{1}{\varepsilon} \left(|\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + |\nabla \text{Riem}|^2 |\nabla^{k-3} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right) \right]$$

for any $\varepsilon > 0$. On the other hand, concerning the other term in (52), we have that

$$|D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 = k^2 |s_k(\nabla s_{k-1}(\nabla^{k-1}\chi_n))|^2.$$

Since $s_k(\nabla s_{k-1}(\nabla^{k-1}\chi_n)) = s_k(\nabla^k \chi_n)$, we thus have that

$$\begin{aligned} |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 &= k^2 |s_k(\nabla^k \chi_n)|^2 \\ &= \frac{k^2}{(k!)^2} \sum_{i_1, \dots, i_k} \left[\sum_{\pi \in \Pi_k} \nabla_{i_{\pi(1)} \dots i_{\pi(k)}}^k \chi_n \right]^2 \\ &= \frac{k^2}{(k!)^2} \left[k! |\nabla^k \chi_n|^2 + \sum_{i_1, \dots, i_k} \sum_{\pi \in \Pi_k} (\nabla_{i_{\pi(1)} \dots i_{\pi(k)}}^k \chi_n \cdot \sum_{\sigma \in \Pi_k \setminus \{\pi\}} \nabla_{i_{\sigma(1)} \dots i_{\sigma(k)}}^k \chi_n) \right] \\ &= \frac{k^2}{(k!)^2} \left[(k! + (k! - 1)k!) |\nabla^k \chi_n|^2 + \left(\nabla^k \chi_n * \left(\text{Riem} * \nabla^{k-2} \chi_n + \nabla \text{Riem} * \nabla^{k-3} \chi_n + \dots + \nabla^{k-3} \text{Riem} * \nabla \chi_n \right) \right) \right] \\ &= \frac{k^2}{(k!)^2} \left[(k!)^2 |\nabla^k \chi_n|^2 + \left(\nabla^k \chi_n * \left(\text{Riem} * \nabla^{k-2} \chi_n + \nabla \text{Riem} * \nabla^{k-3} \chi_n + \dots + \nabla^{k-3} \text{Riem} * \nabla \chi_n \right) \right) \right]. \end{aligned}$$

Using again Young's inequality we get

$$(54) \quad |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 \geq k^2 |\nabla^k \chi_n|^2 + C(m, k) \left[-\varepsilon |\nabla^k \chi_n|^2 - \frac{1}{\varepsilon} \left(|\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + |\nabla \text{Riem}|^2 |\nabla^{k-3} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right) \right],$$

for any $\varepsilon > 0$. Using (53) and (54), we get that

$$\begin{aligned} &\int_M f^2 [(1 - \delta)(k^2 - C(m, k)\varepsilon) - (1 + \eta)(1 + C(m, k)\varepsilon)] |\nabla^k \chi_n|^2 d\text{vol}_g \\ &\leq \int_M f^2 \left[(1 - \delta) |D_S(s_{k-1}(\nabla^{k-1}\chi_n))|^2 - (1 + \eta) |\nabla s_{k-1}(\nabla^{k-1}\chi_n)|^2 \right] d\text{vol}_g \\ &\quad + \int_M f^2 \left[(1 - \delta) C(m, k) \frac{1}{\varepsilon} \left(|\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right) \right] d\text{vol}_g \\ &\quad + \int_M f^2 \left[(1 + \eta) C(m, k) \frac{1}{\varepsilon} \left(|\text{Riem}|^2 |\nabla^{k-2} \chi_n|^2 + \dots + |\nabla^{k-3} \text{Riem}|^2 |\nabla \chi_n|^2 \right) \right] d\text{vol}_g. \end{aligned}$$

By (52) and reasoning as above we know that the RHS converge to 0 as $n \rightarrow \infty$. Hence, suitably choosing $\varepsilon, \eta, \delta \ll 1$ we finally obtain the desired conclusion:

$$\int_M f^2 |\nabla^k \chi_n|^2 d\text{vol}_g \rightarrow 0,$$

as $n \rightarrow \infty$.

6. SOME SHARP APPLICATIONS

6.1. Disturbed Sobolev inequalities. First, we point out the following

Theorem 6.1. *Let (M^m, g) be a smooth, complete non-compact Riemannian manifold without boundary. Let $o \in M$, $r(x) \doteq \text{dist}_g(x, o)$ and suppose that for some $\eta > 0$, $D > 0$ and some $i_0 > 0$,*

$$|\text{Ric}_g|(x) \leq D^2(1 + r(x)^2)^\eta, \quad \text{inj}_g(x) \geq \frac{i_0}{D(1 + r(x))^\eta}.$$

Let $p \in [1, m)$ and $q \in [p, mp/(m - p)]$. Then there exist constants $A_1 > 0$, $A_2 > 0$, depending on m, p, q and the constant C from Theorem 1.2, such that for all $\varphi \in C_c^\infty(M)$ it holds

$$(55) \quad \left(\int_M |\varphi|^q d\text{vol}_g \right)^{\frac{1}{q}} \leq A_1 \left(\int_M |\nabla \varphi|^p d\text{vol}_g \right)^{1/p} + A_2 \left(\int_M H^{2\eta} |\varphi|^p d\text{vol}_g \right)^{1/p},$$

where $H \in C^\infty(M)$ is the distance-like function given by Theorem 1.2.

Remark 6.2. Theorem 6.1 was proved in [20] with $\eta \leq 1$. However the proof therein works also when $\eta > 1$ up to replace [20, Theorem 1.5] with the $k = 2$ case of Theorem 1.2 above, with $\lambda(t) = D^2(1 + t^2)^{\frac{\eta}{2}}$. Note that one could also state the theorem for more general growth functions $\lambda(r)$ as in Theorem 1.2.

Since, under a C^k -control on the curvature, there exists distance-like functions with controlled higher order derivatives, one naturally expects that some sort of improved higher order Sobolev inequality should be obtained exploiting the control on the higher derivatives of the curvature. However, for the moment, this possible phenomenon remains unclear to us. Indeed, generalizing a fact remarked in [4, Proposition 2.11] for the standard (i.e. non-disturbed) Sobolev inequalities, higher order disturbed Sobolev inequalities hold true under exactly the same assumptions as Theorem 6.1.

Proposition 6.3. *Let (M^m, g) be a smooth, complete non-compact Riemannian manifold without boundary. Let $o \in M$, $r(x) \doteq \text{dist}_g(x, o)$ and suppose that for some $\eta > 0$, $D > 0$ and some $i_0 > 0$,*

$$|\text{Ric}_g|(x) \leq D^2(1 + r(x)^2)^\eta, \quad \text{inj}_g(x) \geq \frac{i_0}{D(1 + r(x))^\eta}.$$

Let $p \in [1, m)$ and let k be an integer in $[1, \frac{m}{p})$. Then there exists a constant $A > 0$ depending on m, p, k and the constant C from Theorem 1.2, such that for all $\varphi \in C_c^\infty(M)$ it holds

$$(56) \quad \|\varphi\|_{L^{\frac{pm}{m-kp}}(M)} \leq A \sum_{r=-1}^{k-1} \left\| H^{\frac{\eta}{mp}(r+1)(2m-rp)} |\nabla^{k-r-1} \varphi| \right\|_{L^p(M)},$$

where $H \in C^\infty(M)$ is the distance-like function given by Theorem 1.2.

Proof. For the ease of notation we will write $\|\cdot\|_p$ for $\|\cdot\|_{L^p(M)}$. For $j = 1, \dots, k$, define $q_j \doteq pm/(m - jp)$. We prove that for every $s = 0, \dots, k - 1$ it holds

$$(57) \quad \|\varphi\|_{q_k} \leq C \sum_{r=s-1}^{k-1} \left\| H^{2\eta \sum_{j=s}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_s},$$

with the convention that $\sum_{j=0}^{-1} q_j^{-1} = 0$. Since $\sum_{j=0}^r q_j^{-1} = (2m - rp)(r + 1)/(2mp)$, the inequality (56) is equivalent to (57) when $s = 0$.

Note that $p = q_0 < q_1 < \dots < q_k$, and that $q_j = nq_{j-1}/(n - q_{j-1})$, so that Theorem 6.1 applies with $p = q_{j-1}$ and $q = q_j$. In particular, (57) holds true when $s = k - 1$. For general s , we prove

now the validity of (57) by backward induction. Namely, suppose that for some integer $t \in [1, k-1]$, (57) holds true with $s = t$, i.e.

$$\|\varphi\|_{q_k} \leq C \sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_t}.$$

Applying (55) with $p = q_{t-1}$ and $q = q_t$ at each terms of RHS gives

$$\begin{aligned} \|\varphi\|_{q_k} &\leq C \left[\sum_{r=t-1}^{k-1} \left\| H^{\frac{2\eta}{q_{t-1}}} H^{2\eta \sum_{j=t}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} + \sum_{r=t-1}^{k-1} \left\| \nabla \left(H^{2\eta \sum_{j=t}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right) \right\|_{q_{t-1}} \right] \\ &\leq C \left[\sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t-1}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} + \sum_{r=t-1}^{k-1} \left\| \left(H^{2\eta \sum_{j=t}^r q_j^{-1-1}} |\nabla H| |\nabla^{k-r-1} \varphi| \right) \right\|_{q_{t-1}} \right. \\ &\quad \left. + \sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t}^r q_j^{-1}} \nabla |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} \right]. \end{aligned}$$

Recalling that $|\nabla H| \leq 1$, and that $|\nabla |\nabla^{k-r-1} \varphi|| \leq |\nabla^{k-r} \varphi|$ (see [4, p. 36, (1)]), we obtain

$$\begin{aligned} \|\varphi\|_{q_k} &\leq C \left[\sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t-1}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} + \sum_{r=t-1}^{k-1} \left\| \left(H^{2\eta \sum_{j=t}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right) \right\|_{q_{t-1}} \right. \\ &\quad \left. + \sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t}^r q_j^{-1}} |\nabla^{k-r} \varphi| \right\|_{q_{t-1}} \right] \\ &\leq C \left[\sum_{r=t-1}^{k-1} \left\| H^{2\eta \sum_{j=t-1}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} + \sum_{r=t-2}^{k-2} \left\| H^{2\eta \sum_{j=t}^{r+1} q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}} \right]. \end{aligned}$$

Since $\sum_{j=t}^{r+1} q_j^{-1} \leq \sum_{j=t-1}^r q_j^{-1}$, we finally get

$$\|\varphi\|_{q_k} \leq C \sum_{r=t-2}^{k-1} \left\| H^{2\eta \sum_{j=t-1}^r q_j^{-1}} |\nabla^{k-r-1} \varphi| \right\|_{q_{t-1}},$$

i.e. (57) holds true for $s = t-1$ as desired. \square

6.2. Calderón-Zygmund inequalities. Calderón-Zygmund inequalities are a powerful tool in Euclidean analysis which permits to control the L^p -norm of the Hessian of a function u in terms of the L^p -norms of the Laplacian of u and of u itself. On a complete non-compact Riemannian manifolds (M, g) , it was proved by B. Güneysu and S. Pigola in [16] that the global Calderón-Zygmund inequality

$$(CZ(p)) \quad \forall u \in C_c^\infty(M), \quad \|\text{Hess } u\|_g^p \leq A_1 \|u\|_{L^p}^p + A_2 \|\Delta u\|_{L^p}^p$$

holds for $p \in [1, +\infty)$ if either

- $p = 2$ and $\text{Ric}_g \geq -C$ for some $C > 0$, or
- $|\text{Ric}_g| \leq C$ for some $C > 0$ and $\text{inj}_g(M) > i_0 > 0$.

In general there is no hope to get (CZ(p)) on an arbitrary complete non-compact manifold, due to counterexamples, [16, 21]. However one can weaken the assumptions in [16] obtaining a weaker version of (CZ(p)) in which an unbounded weight function appears in the $\|u\|_{L^p}$ term of (CZ(p)). This approach was considered for instance in [20], where the authors proved (a slightly weaker version of) the following

Theorem 6.4. *Let (M^m, g) be a smooth, complete non-compact Riemannian manifold without boundary. Let $o \in M$, $r(x) \doteq \text{dist}_g(x, o)$ and suppose that one of the following curvature assumptions holds*

(a) *for some $\eta > 0$, some $D > 0$ and some $i_0 > 0$,*

$$|\text{Ric}_g|(x) \leq D^2(1 + r(x)^2)^\eta, \quad \text{inj}_g(x) \geq \frac{i_0}{D(1 + r(x))^\eta} > 0 \quad \text{on } M.$$

(b) *for some $\eta > 0$ and some $D > 0$,*

$$|\text{Sect}_g|(x) \leq D^2(1 + r(x)^2)^\eta.$$

Then there exist constants $A_1 > 0$, $A_2 > 0$, depending on m , η , D and the constant C from Theorem 1.2, such that for all $\varphi \in C_c^\infty(M)$ it holds

$$(58) \quad \|\text{Hess } \varphi|_g\|_{L^2}^2 \leq A_1 \|H^{2\eta} \varphi\|_{L^2}^2 + A_2 \|\Delta \varphi\|_{L^2}^2,$$

where $H \in C^\infty(M)$ is the distance-like function given by Theorem 1.2.

Remark 6.5. The same observation as in Remark 6.2 applies also to Theorem 6.4.

As commented in [20], obtaining a weighted L^p Calderón-Zygmund inequality under the same assumptions of Theorem 6.4 is a non-trivial problem. The main issue is to keep a control on the injectivity radius under the conformal deformation. However, it turns out that this can be done at least under slightly stronger assumptions, i.e. if we assume both a control on the sectional curvatures and on the injectivity radius. Accordingly, we can obtain Theorem 1.8 stated in Section 1, which we state here again for readers' convenience.

Theorem 6.6. *Let (M^m, g) be a smooth, complete non-compact Riemannian manifold without boundary. Let $o \in M$, $r(x) \doteq \text{dist}_g(x, o)$ and suppose that for some $\eta > 0$, some $D > 0$ and some $i_0 > 0$,*

$$|\text{Sect}_g|(x) \leq D^2(1 + r(x)^2)^\eta, \quad \text{inj}_g(x) \geq \frac{i_0}{D(1 + r(x))^\eta} > 0 \quad \text{on } M.$$

Then there exist constants $A > 0$ depending on m , η , D , i_0 and the constant C from Theorem 1.2, such that for all $\varphi \in C_c^\infty(M)$ it holds

$$\|\text{Hess } \varphi|_g\|_{L^p}^p \leq A [\|H^{2\eta} \varphi\|_{L^p}^p + \|\Delta \varphi\|_{L^p}^p],$$

where $H \in C^\infty(M)$ is the distance-like function given by Theorem 1.2.

Proof. We can write the geometric assumption in the following form: for some $\eta > 0$, some $D' > 0$ and some $i'_0 > 0$,

$$|\text{Sect}_g|(x) \leq \bar{D}^2 r(x)^{2\eta}, \quad \text{inj}_g(x) \geq \frac{\bar{i}_0}{r(x)^\eta} > 0 \quad \text{on } M \setminus B_1(o).$$

As in the proof of [20, Theorem 1.7], define the new complete conformal metric $\tilde{g} = H^{2s}g$, which satisfies in particular

$$(59) \quad |\text{Sect}_{\tilde{g}}| \leq K^2$$

on M for some constant $K > 0$. Here H is the second order distance-like function whose existence is guaranteed in our assumptions by Theorem 1.2. For later purposes, observe that reasoning as in the proof of [20, Lemmas 7.3 and 7.4] one can prove the existence of a constant \tilde{c} such that for all j large enough and all $\rho > 0$,

$$B_{\tilde{c}^{-1}\rho r(x_j) - \eta}^g(x_j) \subset B_{\tilde{c}\rho}^{\tilde{g}}(x_j) \subset B_{\tilde{c}\rho r - \eta(r_j)}^g(x_j).$$

We claim that

$$(60) \quad \exists \tilde{i}_0 > 0, \quad \forall x \in M, \quad \text{inj}_{\tilde{g}}(x) \geq \tilde{i}_0 > 0.$$

Suppose it is not the case. Then there exists a sequence of points $\{x_j\}_{j=1}^\infty \subset M \setminus B_1^g(0)$ such that $\text{inj}_{\tilde{g}}(x_j) < \frac{1}{j}$ and $\text{dist}_{\tilde{g}}(x_j, o) \rightarrow \infty$ as $j \rightarrow \infty$.

We call $F_j : \mathbb{B}_{\frac{\pi}{2K}}(0) \subset T_{x_j}M \rightarrow M$ the exponential map \exp_{x_j} of the metric \tilde{g} restricted to $\mathbb{B}_{\frac{\pi}{2K}}(0) \subset T_{x_j}M$. Since $\text{Sect}_{\tilde{g}} \leq K^2$, by Rauch comparison theorem in $B_{\frac{\tilde{g}}{2K}}(x_j)$ there are no points conjugated to x_j for the metric \tilde{g} . In particular F_j is a smooth local diffeomorphism onto $B_{\frac{\tilde{g}}{2K}}(x_j)$ and a smooth surjective local isometry once $T_{x_j}M$ is endowed with the pulled-back metric $\exp_{x_j}^* \tilde{g}$. Hence, $\text{inj}_{\tilde{g}}(x_j) < \frac{1}{j}$ implies that for $j > \frac{2K}{\pi}$, there exists a point $y_j \in B_{\frac{\tilde{g}}{j-1}}(x_j)$ such that $y_j = F(Y_{j,1}) = F(Y_{j,2})$ for two different points $Y_{j,1}$ and $Y_{j,2}$ in $\mathbb{B}_{j-1}(0) \subset T_{x_j}M$. Also, there are two different constant speed minimizing geodesics $\tilde{\gamma}_{j,1}(t)$ and $\tilde{\gamma}_{j,2}(t)$ wrt \tilde{g} which connect x_j to y_j and lifts to $\tilde{\Gamma}_{j,1}(t) = tY_{j,1}$ and $\tilde{\Gamma}_{j,2}(t) = tY_{j,2}$ respectively.

Let $j > \frac{2\tilde{c}^2 K}{\pi}$, so that

$$(61) \quad B_{\frac{\tilde{g}}{j-1}}(x_j) \subset B_{\frac{g}{\tilde{c}j^{-1}r^{-\eta}(r_j)}}(x_j) \subset B_{\frac{\tilde{g}}{\tilde{c}2j-1}}(x_j) \subset B_{\frac{\tilde{g}}{2K}}(x_j).$$

Since F is a local isometry, $F^{-1}(y_j)$ is a discrete set, so that $\tilde{\gamma}_{j,1}$ and $\tilde{\gamma}_{j,2}$ belong necessarily to two different relative homotopy classes $[\tilde{\gamma}_{j,1}]_{\{x_j, y_j\}}$ and $[\tilde{\gamma}_{j,2}]_{\{x_j, y_j\}}$ of paths in $B_{\frac{\tilde{g}}{\tilde{c}2j-1}}(x_j)$ with fixed boundary points x_j and y_j . Because of (61) we have also that $\gamma_{j,1}$ and $\gamma_{j,2}$ are curves contained in $B_{\frac{g}{\tilde{c}j^{-1}r^{-\eta}(r_j)}}(x_j)$ which belong to two different relative homotopy classes $[\tilde{\gamma}_{j,1}]_{\{x_j, y_j\}}$ and $[\tilde{\gamma}_{j,2}]_{\{x_j, y_j\}}$ of paths in $B_{\frac{g}{\tilde{c}j^{-1}r^{-\eta}(r_j)}}(x_j)$ with fixed boundary points x_j and y_j . However,

$$\text{inj}_g(x_j) \geq \frac{\bar{i}_0}{r(x_j)^\eta} > \tilde{c}j^{-1}r^{-\eta}(x_j),$$

as soon as $j > \tilde{c}/i_0$. Hence, for j large enough, $B_{\frac{g}{\tilde{c}j^{-1}r^{-\eta}(r_j)}}(x_j)$ is contractible, which contradicts the existence of two different homotopy classes of curves. The claim (60) is thus proved.

Recalling also (59), we have thus the validity on (M, \tilde{g}) of an L^p Calderón-Zygmund inequality, [16, Theorem C], i.e.,

$$(62) \quad \forall u \in C_c^\infty(M), \quad \|\text{Hess}_{\tilde{g}} u\|_{\tilde{g}}^p \leq A_1 \|u\|_{L^p}^p + A_2 \|\Delta_{\tilde{g}} u\|_{L^p}^p.$$

Here and on $\|\cdot\|_{L^p}$ is the L^p norm of a function computed with respect to the conformally deformed metric \tilde{g} on M . Given $\varphi \in C_c^\infty(M)$, define $u \in C_c^\infty(M)$ by $H^\eta(\frac{m}{p}-2)u = \varphi$. Setting $e^\phi = H^\eta$ and computing as in [20, Section 7], we get that

$$\begin{aligned} e^{4\phi} |\text{Hess}_{\tilde{g}} u|_{\tilde{g}}^2 &= |\text{Hess} u|_g^2 + 2|\nabla u|_g^2 |\nabla \phi|_g^2 + (m-2)g(\nabla u, \nabla \phi)^2 - 2g(\nabla u, \nabla \phi) \Delta_g u - 4\text{Hess} u(\nabla u, \nabla \phi) \\ &\geq |\text{Hess} u|_g^2 + 2|\nabla u|_g^2 |\nabla \phi|_g^2 - (m-2)|\nabla u|_g^2 |\nabla \phi|_g^2 - |\nabla u|_g^2 |\nabla \phi|_g^2 - |\Delta_g u|^2 \\ &\quad - \frac{1}{2} |\text{Hess} u|_g^2 - 8|\nabla u|_g^2 |\nabla \phi|_g^2 \\ &\geq \frac{1}{2} |\text{Hess} u|_g^2 - (m+5)|\nabla u|_g^2 |\nabla \phi|_g^2 - |\Delta_g u|^2. \end{aligned}$$

Accordingly

$$|\text{Hess} u|_g^2 \leq C [H^{4\eta} |\text{Hess}_{\tilde{g}} u|_{\tilde{g}}^2 + \eta^2 |\nabla u|_g^2 |\nabla \log H|_g^2 + |\Delta_g u|^2],$$

and

$$(63) \quad H^{(m-2p)\eta} |\text{Hess} u|_g^p d\text{vol}_g \leq C \left[|\text{Hess}_{\tilde{g}} u|_{\tilde{g}}^p d\text{vol}_{\tilde{g}} + H^{(m-2p)\eta} |\nabla u|_g^p |\nabla \log H|_g^p d\text{vol}_g + H^{(m-2p)\eta} |\Delta_g u|^p d\text{vol}_g \right].$$

From (62) we get

$$\begin{aligned}
\int_M |\text{Hess}_{\bar{g}} u|_{\bar{g}}^p d\text{vol}_{\bar{g}} &\leq A_1 \int_M |u|^p d\text{vol}_{\bar{g}} + A_2 \int_M |\Delta_{\bar{g}} u|^p d\text{vol}_{\bar{g}} \\
&\leq C \left[\int_M |u|^p d\text{vol}_{\bar{g}} + \int_M H^{-2\eta p} |\Delta_g u + (m-2)g(\nabla u, \eta \nabla \log H)|^p d\text{vol}_{\bar{g}} \right] \\
&\leq C \left[\int_M H^{m\eta} |u|^p d\text{vol}_g + \int_M H^{(m-2p)\eta} |\Delta_g u|^p d\text{vol}_g \right. \\
&\quad \left. + \int_M H^{(m-2p)\eta} |\nabla u|_g^p |\nabla \log H|_g^p d\text{vol}_g \right].
\end{aligned}$$

Inserting in (63) gives

$$\begin{aligned}
(64) \quad \int_M H^{(m-2p)\eta} |\text{Hess } u|_g^p d\text{vol}_g &\leq C \left[\int_M H^{m\eta} |u|^p d\text{vol}_g + \int_M H^{(m-2p)\eta} |\Delta_g u|^p d\text{vol}_g \right. \\
&\quad \left. + \int_M H^{(m-2p)\eta} |\nabla u|_g^p |\nabla \log H|_g^p d\text{vol}_g \right].
\end{aligned}$$

Recall that $u = H^{-\eta(\frac{m}{p}-2)}\varphi$. Then

$$\begin{aligned}
H^{\eta(\frac{m}{p}-2)} |\nabla_g u| &\leq C[|\nabla_g \varphi| + H^{-1}\varphi|\nabla_g H|], \\
H^{\eta(\frac{m}{p}-2)} |\Delta_g u| &\leq |\Delta_g \varphi| + C[H^{-1}|\nabla_g \varphi||\nabla_g H| + H^{-2}\varphi|\Delta_g H|],
\end{aligned}$$

and

$$H^{\eta(\frac{m}{p}-2)} |\text{Hess } u| \geq |\text{Hess } \varphi| - C[H^{-1}|\nabla_g \varphi||\nabla_g H| + H^{-2}\varphi|\text{Hess } H|].$$

Combining these latter with (64) we obtain

$$\begin{aligned}
\int_M |\text{Hess } \varphi|_g^p d\text{vol}_g &\leq C \left[\int_M H^{2p\eta} |\varphi|^p d\text{vol}_g + \int_M |\Delta_g \varphi|^p d\text{vol}_g \right. \\
&\quad \left. + \int_M H^{-p} |\nabla \varphi|_g^p |\nabla H|_g^p d\text{vol}_g + \int_M H^{-2p} |\varphi|^p |\text{Hess } H|_g^p d\text{vol}_g \right] \\
&\leq C \left[\int_M H^{2p\eta} |\varphi|^p d\text{vol}_g + \int_M |\Delta_g \varphi|^p d\text{vol}_g + \int_M |\nabla \varphi|_g^p d\text{vol}_g \right].
\end{aligned}$$

where we used the fact that H is a strictly positive exhaustion function, $|\nabla H|$ is uniformly bounded and $|\text{Hess } H| \leq CH^\eta$. Applying [16, Proposition 3.10 a)] with ε small enough, we finally get

$$\int_M |\text{Hess } \varphi|_g^p d\text{vol}_g \leq C \left[\int_M H^{2p\eta} |\varphi|^p d\text{vol}_g + \int_M |\Delta_g \varphi|^p d\text{vol}_g \right]$$

as desired. \square

6.3. Omori-Yau maximum principle. We end this section with a remark on the Omori-Yau maximum principle. Recall that a Riemannian manifold (M, g) is said to satisfy the full Omori-Yau maximum principle for the Hessian if for any function $u \in C^2(M)$ with $u^* = \sup_M u < +\infty$, there exists a sequence $\{x_n\}_n \subset M$ with the properties

$$\text{(i) } u(x_n) > u^* - \frac{1}{n}, \quad \text{(ii) } |\nabla u(x_n)| < \frac{1}{n}, \quad \text{(iii) } \text{Hess}(u)(x_n) < \frac{1}{n}g,$$

for each $n \in \mathbb{N}$. In [24] it was proved that the full Omori-Yau maximum principle for the Hessian holds e.g. if the radial sectional curvature of M (i.e. the sectional curvature of 2-planes containing ∇r), satisfies

$$\text{Sect}_{\text{rad}} \geq -C^2(1+r^2) \prod_{j=1}^{\bar{j}} \left(\ln^{[j]}(r) \right)^2,$$

where $\ln^{[j]}$ stands for the j -th iterated logarithm. In [20] we proved that the same is true if

$$|\text{Ric}_g|(x) \leq D^2(1+r(x)^2), \quad \text{inj}_g(x) \geq \frac{i_0}{D(1+r(x))} > 0 \quad \text{on } M.$$

Using Theorem 1.2 and reasoning as in [20], we get that assuming

$$|\text{Ric}_g|(x) \leq C^2 r^2 \prod_{j=1}^{\bar{j}} \left(\ln^{[j]}(r) \right)^2, \quad \text{inj}_g(x) \geq \frac{i_0}{r \prod_{j=1}^{\bar{j}} \ln^{[j]}(r)}$$

outside a compact set of M is enough.

APPENDIX A. SOME COMMUTATION FORMULAS

Using again the “*” notation defined in Section 2, we have the validity of the following

Lemma A.1. *Let $u \in C^\infty(M)$. Then*

$$\Delta \nabla^{q-2} u - \nabla^{q-2} \Delta u = \text{Riem} * \nabla^{q-2} u + \nabla \text{Ric} * \nabla^{q-3} u + \dots \nabla^{q-3} \text{Ric} * \nabla u$$

Proof. (Sketch). We work in a normal frame $\{E_i\}$ orthonormal at $p \in M$, and in frame computations we will use the convention of lowering all indices, summing over repeated indices. Recall that we are adopting the following sign conventions for curvatures:

$$\text{R}(X, Y) = \nabla_X \nabla_Y - \nabla_Y \nabla_X - \nabla_{[X, Y]};$$

$$\text{Riem}(X, Y, Z, W) = g(\text{R}(X, Y)Z, W)$$

$$\text{Ric}(X, Y) = \text{tr} \text{Riem}(X, \cdot, \cdot, Y).$$

Moreover, for the ease of notation we will write in coordinates R_{ijkl} for Riem_{ijkl} . Letting $u \in C^\infty(M)$, recall the following commutation rules

$$(65) \quad (\nabla_i \nabla_j - \nabla_j \nabla_i) u = 0,$$

$$(66) \quad (\nabla_k \nabla_j - \nabla_j \nabla_k) \nabla_i u = -\nabla_t u R_{tijk}.$$

More generally one can compute that, for any $l \geq 2$,

$$(s) \quad (\nabla_{i_l} \nabla_{i_{l-1}} - \nabla_{i_{l-1}} \nabla_{i_l}) \nabla_{i_{l-2}} \dots \nabla_{i_1} u = -\nabla_{i_{l-2}} \dots \nabla_{i_2} \nabla_t u R_{ti_1 i_{l-1} i_l} - \nabla_{i_{l-2}} \dots \nabla_{i_3} \nabla_t \nabla_{i_1} u R_{ti_2 i_{l-1} i_l} \\ - \dots - \nabla_t \nabla_{i_{l-3}} \nabla_{i_1} u R_{ti_{l-2} i_{l-1} i_l}.$$

If one is interested in the commutation rule for the $(s-1)$ -th and the s -th derivative of u of a total of q derivatives, it suffices to take ∇^{q-s} of formula (s). When studying a term like $\Delta \nabla^{q-2} u - \nabla^{q-2} \Delta u$, one hence realizes that the higher order derivatives of curvature terms arise from commutators of the 2-nd and the 3-rd (of the total q) derivatives. There are two such terms in the telescopic development of $\Delta \nabla^{q-2} u - \nabla^{q-2} \Delta u$, namely

$$\nabla_{i_q \dots i_6}^{q-5} \nabla_{i_5} \nabla_p [(\nabla_p \nabla_{i_2} \nabla_{i_1} - \nabla_{i_2} \nabla_p \nabla_{i_1}) u] \\ = \nabla_{i_q \dots i_6}^{q-5} \nabla_{i_5} \nabla_p [-\nabla_t u R_{ti_1 i_2 p}] \\ = -\nabla_{i_q \dots i_6}^{q-5} [\nabla_{i_5} \nabla_p \nabla_t u R_{ti_1 i_2 p} + \nabla_p \nabla_t u \nabla_{i_5} R_{ti_1 i_2 p} + \nabla_{i_5} \nabla_t u \nabla_{i_5} \nabla_p R_{ti_1 i_2 p} + \nabla_t u \nabla_{i_5} \nabla_p R_{ti_1 i_2 p}],$$

and

$$\nabla_{i_q \dots i_6}^{q-5} \nabla_{i_5} \nabla_{i_2} [(\nabla_p \nabla_{i_1} \nabla_p - \nabla_{i_1} \nabla_p \nabla_p) u] = \nabla_{i_q \dots i_6}^{q-5} \nabla_{i_5} \nabla_{i_2} [-\nabla_t u R_{tpi_1 p}]$$

Using Bianchi’s identities each of these terms can be written in the form

$$\text{Riem} * \nabla^{q-2} u + \nabla \text{Ric} * \nabla^{q-3} u + \dots + \nabla^{q-3} \text{Ric} * \nabla u.$$

□

APPENDIX B. WEITZENBÖCK FORMULA FOR Δ_{Sym}

Given h a totally symmetric $(0, k-1)$ tensor, $\{E_i\}$ a local orthonormal frame on M , recall that

$$(D_S h)(X_1, X_2, \dots, X_k) = k s_k(\nabla h)(X_1, X_2, \dots, X_k) = \frac{1}{(k-1)!} \left(\sum_{\sigma \in \Pi_k} (\nabla h)(X_{\sigma(1)}, \dots, X_{\sigma(k)}) \right)$$

$$(D_S^* h)(X_1, \dots, X_{k-2}) = - \sum_i (\nabla_{E_i} h)(E_i, X_1, \dots, X_{k-2}).$$

For the sake of completeness we provide here a proof of the following

Lemma B.1 ([26]). *Letting $\Delta_{\text{Sym}} = D_S^* D_S - D_S D_S^*$, we have that*

$$(67) \quad \begin{aligned} \Delta_{\text{Sym}} h &= \Delta_B h - \tilde{\mathfrak{R}}^{(k-1)}(h) + \tilde{\mathfrak{G}}^{(k-1)}(h) \\ &= \Delta_B h - \mathfrak{Ric}(h), \end{aligned}$$

where

$$\begin{aligned} \tilde{\mathfrak{R}}^{(k-1)}(h)(X_1, \dots, X_{k-1}) &\doteq \sum_{i=1}^{k-1} \sum_j h(X_1, \dots, E_j, \dots, X_{k-1}) \text{Ric}(X_i, E_j); \\ \tilde{\mathfrak{G}}^{(k-1)}(h)(X_1, \dots, X_{k-1}) &\doteq \sum_{p < i} \sum_{j, l} h(X_1, \dots, E_l, \dots, E_j, \dots, X_{k-1}) \\ &\quad \cdot (\text{Riem}(E_l, X_p, X_i, E_j) + \text{Riem}(E_l, X_i, X_p, E_j)), \end{aligned}$$

the definition of \mathfrak{Ric} has been recalled in (41), and $\Delta_B = -\text{tr}_{12}(\nabla^2) = \nabla^* \nabla$, with ∇^* the formal L^2 -adjoint of ∇ .

Proof. Assume that we perform computation at a point $p \in M$ in a normal frame, so that all covariant derivatives of vector fields vanish at p . We have that

$$\begin{aligned} (D_S^* D_S h)(X_1, \dots, X_{k-1}) &= - \sum_i (\nabla_{E_i} D_S h)(E_i, X_1, \dots, X_{k-1}) \\ &= - \frac{1}{(k-1)!} \sum_i \nabla_{E_i} ((k-1)! \nabla_{E_i} h(X_1, \dots, X_{k-1}) + (k-1)! \nabla_{X_1} h(E_i, X_2, \dots, X_{k-1}) + \dots \\ &\quad + (k-1)! \nabla_{X_{k-1}} h(E_i, X_1, \dots, X_{k-2})) \\ &= - \sum_i (\nabla_{E_i} \nabla_{E_i} h(X_1, \dots, X_{k-1}) + \nabla_{E_i} \nabla_{X_1} h(E_i, X_2, \dots, X_{k-1}) \\ &\quad + \dots + \nabla_{E_i} \nabla_{X_{k-1}} h(E_i, X_1, \dots, X_{k-2})). \end{aligned}$$

While,

$$\begin{aligned} (D_S D_S^* h)(X_1, \dots, X_{k-1}) &= (k-1) s_{k-1}(\nabla D_S^* h)(X_1, \dots, X_{k-1}) \\ &= \frac{1}{(k-2)!} ((k-2)! \nabla_{X_1} D_S^* h(X_2, \dots, X_{k-1}) + \dots + (k-2)! \nabla_{X_{k-1}} D_S^* h(X_1, \dots, X_{k-2})) \\ &= - \sum_i (\nabla_{X_1} \nabla_{E_i} h(E_i, X_2, \dots, X_{k-1}) + \dots + \nabla_{X_{k-1}} \nabla_{E_i} h(E_i, X_1, \dots, X_{k-2})). \end{aligned}$$

Hence

$$\begin{aligned} \Delta_{\text{Sym}} h &= (D_S^* D_S - D_S D_S^*) h \\ &= \Delta_B h + \sum_i (\nabla_{X_1} \nabla_{E_i} h(E_i, X_2, \dots, X_{k-1}) - \nabla_{E_i} \nabla_{X_1} h(E_i, X_2, \dots, X_{k-1})) \\ &\quad + \nabla_{X_2} \nabla_{E_i} h(E_i, X_1, X_3, \dots, X_{k-1}) - \nabla_{E_i} \nabla_{X_2} h(E_i, X_1, X_3, \dots, X_{k-1}) + \dots \\ &\quad + \nabla_{X_{k-1}} \nabla_{E_i} h(E_i, X_1, \dots, X_{k-2}) - \nabla_{E_i} \nabla_{X_{k-1}} h(E_i, X_1, \dots, X_{k-2}) \end{aligned}$$

Since h is symmetric, a standard computation gives that

$$\begin{aligned} (R(X, Y)h)(X_1, \dots, X_{k-1}) &= - \sum_l (h(E_l, X_2, \dots, X_{k-1}) \text{Riem}(X, Y, X_1, E_l)) \\ &\quad + \dots + h(X_1, \dots, E_l) \text{Riem}(X, Y, X_{k-1}, E_l). \end{aligned}$$

Hence,

$$\begin{aligned} &\sum_i (\nabla_{X_1} \nabla_{E_i} h(E_i, X_2, \dots, X_{k-1}) - \nabla_{E_i} \nabla_{X_1} h(E_i, X_2, \dots, X_{k-1})) \\ &= \sum_i (R(X_1, E_i)h)(E_i, X_2, \dots, X_{k-1}) \\ &= - \sum_{i,l} (h(E_l, X_2, \dots, X_{k-1}) \text{Riem}(X_1, E_i, E_l, E_l) + h(E_i, E_l, X_3, \dots, X_{k-1}) \text{Riem}(X_1, E_i, X_2, E_l)) \\ &\quad + \dots + h(E_i, X_2, \dots, X_{k-2}, E_l) \text{Riem}(X_1, E_i, X_{k-1}, E_l), \end{aligned}$$

and analogously for the other terms. We thus get the validity of the first equality in (67). The second equality is a direct computation, using the expression (41) for \mathfrak{Ric} . \square

Acknowledgement. *The first and the second authors gratefully acknowledge the partial support of the PRIN project 2017 “Real and Complex Manifolds: Topology, Geometry and holomorphic dynamics” and of the MIUR project “Dipartimenti di Eccellenza 2018-2022”- CUP: E11G18000350001. The first author is partially supported by INdAM-GNSAGA. The second and the third authors are partially supported by INdAM-GNAMPA. The authors would like to thank the anonymous referees for the careful reading of the paper and for useful suggestions.*

REFERENCES

- [1] R. A. Adams, *Sobolev spaces*, Pure and Applied Mathematics, vol. 65, Academic Press [A subsidiary of Harcourt Brace Jovanovich, Publishers], New York-London, 1975.
- [2] M. T. Anderson, *Convergence and rigidity of manifolds under Ricci curvature bounds*, *Invent. Math.* **102** (1990), no. 2, 429–445. MR 1074481
- [3] T. Aubin, *Espaces de Sobolev sur les variétés riemanniennes*, *Bull. Sci. Math. (2)* **100** (1976), no. 2, 149–173. MR 0488125
- [4] ———, *Some nonlinear problems in Riemannian geometry*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 1998.
- [5] L. Bandara, *Density problems on vector bundles and manifolds*, *Proc. Amer. Math. Soc.* **142** (2014), no. 8, 2683–2695.
- [6] D. Bianchi and A. G. Setti, *Laplacian cut-offs, porous and fast diffusion on manifolds and other applications*, *Calc. Var. Partial Differential Equations* **57** (2018), no. 1, Art. 4, 33. MR 3735744
- [7] J. Cheeger and M. Gromov, *Bounds on the von Neumann dimension of L^2 -cohomology and the Gauss-Bonnet theorem for open manifolds*, *J. Differential Geom.* **21** (1985), no. 1, 1–34. MR 806699
- [8] B. Chow, S.-C. Chu, D. Glickenstein, C. Guenther, J. Isenberg, T. Ivey, D. Knopf, P. Lu, F. Luo, and L. Ni, *The Ricci flow: techniques and applications. Part III. Geometric-analytic aspects*, Mathematical Surveys and Monographs, vol. 163, American Mathematical Society, Providence, RI, 2010. MR 2604955
- [9] M. P. Gaffney, *The conservation property of the heat equation on Riemannian manifolds*, *Comm. Pure Appl. Math.* **12** (1959), 1–11. MR 0102097

- [10] D. Gilbarg and N. S. Trudinger, *Elliptic partial differential equations of second order*, Classics in Mathematics, Springer-Verlag, Berlin, 2001, Reprint of the 1998 edition. MR 1814364
- [11] R. E. Greene and H. Wu, C^∞ approximations of convex, subharmonic, and plurisubharmonic functions, *Ann. Sci. École Norm. Sup. (4)* **12** (1979), no. 1, 47–84. MR 532376
- [12] D. Guidetti, B. Güneysu, and D. Pallara, L^1 -elliptic regularity and $H = W$ on the whole L^p -scale on arbitrary manifolds, *Ann. Acad. Sci. Fenn. Math.*, **42** (2017), no. 1, 497–521 MR 3558546
- [13] B. Güneysu, *Sequences of Laplacian cut-off functions*, *J. Geom. Anal.* **26** (2016), no. 1, 171–184. MR 3441508
- [14] ———, *Covariant Schrödinger semigroups on Riemannian manifolds*, *Operator Theory: Advances and Applications*, vol. 264, Birkhäuser/Springer, Cham, 2017. MR 3751359
- [15] B. Güneysu and S. Pigola, *Quantitative C^1 -estimates on manifolds*, *Int. Math. Res. Not. IMRN* 2018, no. 13, 4103–4119.
- [16] ———, *The Calderón-Zygmund inequality and Sobolev spaces on noncompact Riemannian manifolds*, *Adv. Math.* **281** (2015), 353–393. MR 3366843
- [17] Q. Han and F. Lin, *Elliptic partial differential equations*, Courant Lecture Notes in Mathematics, vol. 1, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1997. MR 1669352
- [18] E. Hebey, *Nonlinear analysis on manifolds: Sobolev spaces and inequalities*, Courant Lecture Notes in Mathematics, vol. 5, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 1999.
- [19] S. Huang, *A note on existence of exhaustion functions and its applications*, *J. Geom. Anal.* **29** (2019), no. 2, 1649–1659.
- [20] D. Impera, M. Rimoldi, and G. Veronelli, *Density problems for second order Sobolev spaces and cut-off functions on manifolds with unbounded geometry*, *Int. Math. Res. Not. IMRN* 2021, no. 14, 10521–10558. MR 4285729.
- [21] S. Li, *Counterexamples to the L^p -Calderón-Zygmund estimate on open manifolds*, *Ann. Global Anal. Geom.* **57** (2020), no. 1, 61–70. MR 4057451
- [22] A. Lichnerowicz, *Propagateurs et commutateurs en relativité générale*, *Inst. Hautes Études Sci. Publ. Math.* (1961), no. 10, 56. MR 0157736
- [23] P. Petersen, *Riemannian geometry*, third ed., Graduate Texts in Mathematics, vol. 171, Springer, Cham, 2016. MR 3469435
- [24] S. Pigola, M. Rigoli, and A. G. Setti, *Maximum principles on Riemannian manifolds and applications*, *Mem. Amer. Math. Soc.* **174** (2005), no. 822, x+99. MR 2116555
- [25] M. Rimoldi and G. Veronelli, *Extremals of log Sobolev inequality on non-compact manifolds and Ricci soliton structures*, *Calc. Var. Partial Differential Equations* **58** (2019), no. 2, Art. 66, 26.
- [26] J. H. Sampson, *On a theorem of Chern*, *Trans. Amer. Math. Soc.* **177** (1973), 141–153. MR 0317221
- [27] R. Schoen and S.-T. Yau, *Lectures on differential geometry*, Conference Proceedings and Lecture Notes in Geometry and Topology, I, International Press, Cambridge, MA, 1994, Lecture notes prepared by Wei Yue Ding, Kung Ching Chang [Gong Qing Zhang], Jia Qing Zhong and Yi Chao Xu, Translated from the Chinese by Ding and S. Y. Cheng, With a preface translated from the Chinese by Kaising Tso. MR 1333601
- [28] L.-F. Tam, *Exhaustion functions on complete manifolds*, Recent advances in geometric analysis, *Adv. Lect. Math. (ALM)*, vol. 11, Int. Press, Somerville, MA, 2010, pp. 211–215.
- [29] P. Topping, *Lectures on the Ricci flow*, London Mathematical Society Lecture Note Series, vol. 325, Cambridge University Press, Cambridge, 2006.

(Debora Impera) DIPARTIMENTO DI SCIENZE MATEMATICHE "GIUSEPPE LUIGI LAGRANGE", POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI, 24, TORINO, ITALY, I-10129

E-mail address: debora.impera@polito.it

(Michele Rimoldi) DIPARTIMENTO DI SCIENZE MATEMATICHE "GIUSEPPE LUIGI LAGRANGE", POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI, 24, TORINO, ITALY, I-10129

E-mail address: michele.rimoldi@polito.it

(Giona Veronelli) DIPARTIMENTO DI MATEMATICA E APPLICAZIONI, UNIVERSITÀ DI MILANO BICOCCA, VIA R. COZZI 53, I-20125 MILANO, ITALY

E-mail address: giona.veronelli@unimib.it