

Transoral robotic surgery for bilateral parenchymal submandibular stones: the Flex Robotic System

Original

Transoral robotic surgery for bilateral parenchymal submandibular stones: the Flex Robotic System / Capaccio, Pasquale; Cammarota, Raffaella; Riva, Giuseppe; Albera, Andrea; Albera, Roberto; Pecorari, Giancarlo. - In: JOURNAL OF OTOLARYNGOLOGY-HEAD AND NECK SURGERY. - ISSN 1916-0216. - 17:1(2021), pp. 45-48. [10.5152/B-ENT.2021.20190]

Availability:

This version is available at: 11583/2946592 since: 2021-12-20T08:54:50Z

Publisher:

AVES

Published

DOI:10.5152/B-ENT.2021.20190

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Transoral robotic surgery for bilateral parenchymal submandibular stones: the Flex Robotic System

Pasquale Capaccio¹ , Raffaella Cammarota² , Giuseppe Riva² , Andrea Albera² , Roberto Albera² , Giancarlo Pecorari² 

¹Otorhinolaryngology Unit, Department of Biomedical, Surgical and Dental Sciences, University of Milan, Fondazione IRCCS Ca' Granda Ospedale Maggiore Policlinico, Milan, Italy

²Otorhinolaryngology Unit, Department of Surgical Sciences, University of Turin, Turin, Italy

Cite this article as: Capaccio P, Cammarota R, Riva G, Albera A, Albera R, Pecorari G. Transoral robotic surgery for bilateral parenchymal submandibular stones: the Flex Robotic System. *B-ENT* 2021; 17(1): 45-8.

ABSTRACT

Traditional management of the submandibular gland stones is based on transcervical sialadenectomy. Recently, conservative treatments emerged. In particular, transoral robotic surgery (TORS) has been used for the removal of submandibular sialoliths. Previous case series of TORS reported the use of the Da Vinci system with a high success rate. We describe an uncommon case of bilateral hilo-parenchymal submandibular stone removed via transoral approach with the Flex Robotic System. We report the step-by-step procedure for the transoral robotic approach to hilo-parenchymal submandibular sialoliths through the oral floor. The procedure was successful without complications. The total procedure time was 130 minutes, including 20 minutes for the robotic setting. The Flex Robotic System appeared to be a safe conservative surgical tool for hilo-parenchymal submandibular sialoliths by simultaneously preserving submandibular glands and avoiding severe complications on anatomical structures of the oral floor.

Keywords: Conservative salivary surgery, oral surgical procedures, robotic surgical procedures, salivary calculi, salivary glands

Introduction

Conventional treatment for submandibular gland stones consists of transcervical sialadenectomy with known complications (1). The traditional transoral removal was introduced by Downtown and Quist in 1960 but was recently described using robot-assisted glandular surgery (2-4). Conservative salivary techniques, such as sialendoscopy-assisted transoral surgery, ensured the ability to treat obstructing symptoms with stones removal (5, 6).

First cases of transoral robotic submandibular stones removal were performed with the Da Vinci Si HD (Intuitive Surgical, Sunnyvale, CA, USA), reporting 100% procedural success (7, 8). Another minimally invasive robotic approach in head and neck surgery is the Flex Robotic System (Medrobotics Inc., Raynham, MA, USA). It consists of a new combination of 2 instruments manually controlled by the surgeon with a fully robotic snake arm monocular robotic camera (9). The Flex Robotic System was successfully used for head and neck cancer surgery and was validated in multicenter studies (10).

By combining our experience with transoral robotic surgery (TORS) for head and neck cancer and conservative transoral

salivary surgery (4, 5), we present the use of Flex Robotic System in a case of bilateral submandibular stones. The aim of this paper was to present a step-by-step description of the transoral robotic approach to hilo-parenchymal submandibular sialoliths through the oral floor with the Flex Robotic System.

Case Presentation

A female patient aged 68 years reported recurrent bilateral submandibular swelling with mild pain lasting for 6 months. During oral examination, purulent discharge from the orifice of Wharton's ducts flushed out. Ultrasonography showed bilateral salivary stones with dilation of the intraparenchymal duct system; moreover, submandibular glands had a dishomogeneous appearance owing to recurrent inflammation. Antibiotic and steroid therapy was administered with mild clinical improvement. Computed tomography showed the presence of large bilateral salivary stones located in the parenchyma near the glandular hilum (Figure 1). The patient's consent was obtained.

TORS was scheduled to remove stones through the oral floor with preservation of the submandibular glands. General anesthesia with a nasotracheal tube was necessary to achieve

Corresponding Author: Pasquale Capaccio, pasquale.capaccio@unimi.it

Received: November 27, 2020 **Accepted:** March 14, 2021

Available online at www.b-ent.be



CC BY 4.0: Copyright@Author(s), "Content of this journal is licensed under a Creative Commons Attribution 4.0 International License."



Figure 1. Axial computed tomography that shows the presence of large bilateral salivary stones in the Wharton ducts (15 × 11 mm at the right side and 11 × 8 mm at the left side) near the submandibular hilum

broad exposure and access to the floor of the mouth, bilaterally. The tongue was retracted to the contralateral side, and the oral floor was flattened by positioning the Flex retractor. The Flex Robotic System was placed behind the head of the patient. The Flex scope was moved by the first surgeon using a 3-dimensional (3D) high definition (3D-HD) monitor through a controller on the Flex console. Suction and external pressure over the submandibular gland were provided by the first assistant. The surgeon marked the site of the left salivary stone on the mucosal surface through palpation (Figure 2). Mucosal incision was performed using monopolar cautery of the robot along the line demarcated previously (Figure 3). The Wharton duct was exposed through a blunt dissection using the Maryland dissector. The lingual nerve was identified and was gently mobilized from the duct and retracted medially to visualize the gland hilum (Figure 4). Then, external finger pressure of the assistant surgeon was mandatory for the next steps. A duct

Main Points:

- Transoral robotic submandibular stone removal is emerging, and the Da Vinci Si HD is the most used robotic system.
- The Flex Robotic System showed great potential as a surgical tool in the surgery of the oral floor, providing the following: flexible instruments, excellent visualization, maneuverability, and tactile feedback.
- Our case report showed that Flex Robotic system represents a viable and safe conservative surgical tool for hilo-parenchymal submandibular stone removal.



Figure 2. The location of the left salivary stone was marked on the mucosal surface through palpation



Figure 3. Transoral Robotic Surgery with Flex Robotic System. The mucosal incision was performed using the monopolar cautery of the robot along the line demarcated previously



Figure 4. The lingual nerve was identified and gently dissected to separate it from the Wharton duct



Figure 5. A duct incision was performed over the sialolith, and the stone was carefully dissected from its attachment to the hilum of the submandibular gland

incision over the sialolith and a careful dissection of the stone from its attachment to the gland hilum were performed (Figure 5). The sialolith was then removed. The surgical incisional area was then irrigated with saline solution to remove any debris. A hemostatic and antimicrobial surgical net (Tabotamp, Johnson & Johnson Medical Ltd, Skipton, United Kingdom) was placed over the incisional area, and closure of the oral mucosa was done using absorbable 3-0 Vycril stitches. The same procedure was repeated for the right salivary stone.

The stones were successfully removed using the Flex Robotic System. Visualization and exposure of the lingual nerve, the Wharton's duct, and the parenchyma harboring the stone were excellent. The total procedure time was 130 minutes, including 20 minutes for the robotic setting. No intraoperative complications occurred. The patient was discharged on the second day after the surgery. Moderate pain in the oral floor was described by the patient for 1 week after the surgery; the patient also referred to bilateral lingual hypoesthesia lasting for 1 month. Perioperative and postoperative antibiotic (amoxicillin + clavulanic acid) and steroid therapy was administered during the hospital stay. Then, antibiotic was further prescribed for 5 days.

Medical examinations were performed 7, 30, and 90 days after the surgery. An ultrasonographic examination was done 3 months after the procedure to check the glandular parenchyma echogenicity and to assess any ductal system dilation or residual stones. No persistent gland swelling, ranula, or ductal stenosis were observed. Ultrasonography did not reveal any residual stones or ductal dilation.

Discussion

About 80%-90% of salivary stones involve the submandibular gland, being usually located at the distal tract of the Wharton duct, hilum, and hilo-parenchymal area (11, 12). Stone recurrence after the transoral approach is possible, probably because of the deep and narrow surgical area. Indeed, an appropriate view of oral floor anatomical landmarks after incision of glandular parenchyma is difficult (5, 6). The introduction of the

transoral robotic approach made it possible to overcome some of these limitations (13). In particular, because the robotic approach guarantees a 3D and enhanced depth perception of the oral floor, it allows to follow the anatomical course of the lingual nerve and to maintain healthy submandibular duct until incision to the parenchyma.

TORS with the Da Vinci system has some limitations, such as a limited number of cutting instruments, rigid and relatively bulky robotic arms, and high costs (14). Recently, the Flex Robotic System combined the benefit of a robotic system with a flexible endoscope that supports cutting instruments and flexible devices for TORS (9). Furthermore, because the devices are controlled by the surgeon's hands rather than by a remote robotic system, the Flex Robotic System ensures tactile sensation and manual haptic feedback (9).

Despite the increasing literature showing successful results with the Flex Robotic System for head and neck lesions, it has never been applied to the anterior oral floor (15). In our case, bilateral large hilo-parenchymal submandibular sialoliths were removed using the Flex Robotic System. The procedure was safe because no intraoperative or postoperative and persistent unfavorable effects, such as lingual nerve injury, tongue tingling, ranula, or hilar stenosis with recurrent sialadenitis, were reported. Moreover, this was the first reported case of bilateral submandibular stone removal using TORS.

An adequate exposure of the oral floor is necessary through correct positioning of the Flex retractor. Moreover, a blunt dissection of the Wharton duct and the stones using the Maryland dissector guarantees a clean surgical field with a small amount of blood. This ensures a better visualization of the deep surgical area, which is enhanced by the 3D-HD camera. The role of the assistant surgeon is important while performing suction, traction of the tissue, and external gland push up for a better exposure of the gland parenchyma in the oral floor. Finally, the procedure lasted for 130 minutes for bilateral stones removal, but the surgical time can be further decreased by increasing the experience of the surgeon.

In conclusion, the Flex Robotic System seems to be a safe conservative surgical tool for hilo-parenchymal submandibular stones removal.

Informed Consent: Verbal informed consent was obtained from the patients who agreed to take part in the study.

Peer-review: Externally peer-reviewed.

Author Contributions: Supervision – P.C., R.A., G.P.; Design – P.C., G.P.; Resources – P.C., R.A., G.P.; Materials – R.C., G.R., A.A.; Data Collection and/or Processing – R.C., G.R., A.A.; Analysis and/or Interpretation – P.C., G.R.; Literature Search – G.R., A.A.; Writing Manuscript – P.C., G.R.; Critical Review – P.C., R.A., G.P.

Conflict of Interest: The authors have no conflict of interest to declare.

Financial Disclosure: The authors declared that this study has received no financial support.

References

1. Papaspyrou G, Werner JA, Sesterhenn AM. Transcervical extirpation of the submandibular gland: the University of Marburg experience. *Eur Arch Otorhinolaryngol* 2014; 271: 2009-12. [\[Crossref\]](#)
2. Prosser JD, Bush CM, Solares CA, et al. Trans-oral robotic submandibular gland removal. *J Robot Surg* 2013; 7: 87-90. [\[Crossref\]](#)
3. Lin X, Liang L, Shao X, et al. Trans-oral robotic surgery of submandibular gland removal with preservation of sublingual gland and Wharton's duct. *J Craniofac Surg* 2019; 30:237-8. [\[Crossref\]](#)
4. Capaccio P, Montevecchi F, Meccariello G, et al. Transoral robotic surgery for hilo-parenchymal submandibular stones: step-by-step description and reasoned approach. *Int J Oral Maxillofac Surg* 2019; 14:1520-4. [\[Crossref\]](#)
5. Capaccio P, Clemente IA, McGurk M, et al. Transoral removal of hiloparenchymal submandibular calculi: a long-term clinical experience. *Eur Arch Otorhinolaryngol* 2011; 268: 1081-6. [\[Crossref\]](#)
6. Capaccio P, Gaffuri M, Rossi V, et al. Sialendoscope-assisted transoral removal of hilo-parenchymal submandibular stones: surgical results and subjective scores. *Acta Otorhinolaryngol Ital* 2017; 37:122-7. [\[Crossref\]](#)
7. Walvekar RR, Tayler PD, Tammareddi N, Peters G. Robotic-Assisted transoral removal of a submandibular megalith. *Laryngoscope* 2011; 121: 534-7. [\[Crossref\]](#)
8. Razavi C, Pascheles C, Samara G, Marzouk M. Robot-Assisted sialolithotomy with sialendoscopy for the management of large submandibular gland stones. *Laryngoscope* 2016; 126:345-51. [\[Crossref\]](#)
9. Mattheis S, Hasskamp P, Holtmann L, et al. Flex Robotic System in transoral robotic surgery: The first 40 patients. *Head Neck* 2017; 39: 471-5. [\[Crossref\]](#)
10. Lang S, Mattheis S, Hasskamp P, et al. A european multicenter study evaluating the flex robotic system in transoral robotic surgery. *Laryngoscope* 2017; 127: 391-5. [\[Crossref\]](#)
11. Capaccio P, Torretta S, Ottavian F, Sambataro G, Pignataro L. Modern management of obstructive salivary diseases. *Acta Otorhinolaryngol Ital* 2007; 27: 161-72.
12. Capaccio P, Bottero A, Pompilio M, Ottaviani F. Conservative transoral removal of hilar submandibular salivary calculi. *Laryngoscope* 2005; 115: 750-2. [\[Crossref\]](#)
13. de Almeida JR, Li R, Magnuson JS, et al. Oncologic outcomes after transoral robotic surgery: a multi-institutional study. *JAMA Otolaryngol Head Neck Surg* 2015; 141: 1043-51. [\[Crossref\]](#)
14. Hoffmann TK, Schuler PJ, Bankfalvi A, et al. Comparative analysis of resection tools suited for transoral robot-assisted surgery. *Eur Arch Otorhinolaryngol* 2014; 271:1207-13. [\[Crossref\]](#)
15. Cammaroto G, Stringa LM, Zhang H, et al. Alternative Applications of Trans-Oral Robotic Surgery (TORS): A Systematic Review. *J Clin Med* 2020; 9. [\[Crossref\]](#)