

Asymptotic analysis of Poisson shot noise processes, and applications

Original

Asymptotic analysis of Poisson shot noise processes, and applications / Torrisi, G. L.; Leonardi, E.. - In: STOCHASTIC PROCESSES AND THEIR APPLICATIONS. - ISSN 0304-4149. - ELETTRONICO. - 144:(2022), pp. 229-270. [10.1016/j.spa.2021.11.008]

Availability:

This version is available at: 11583/2948532 since: 2022-01-09T09:46:35Z

Publisher:

Elsevier B.V.

Published

DOI:10.1016/j.spa.2021.11.008

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

Elsevier postprint/Author's Accepted Manuscript

© 2022. This manuscript version is made available under the CC-BY-NC-ND 4.0 license
<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:
<http://dx.doi.org/10.1016/j.spa.2021.11.008>

(Article begins on next page)

Asymptotic analysis of Poisson shot noise processes, and applications

Giovanni Luca Torrisi* and Emilio Leonardi†

Abstract

Poisson shot noise processes are natural generalizations of compound Poisson processes that have been widely applied in insurance, neuroscience, seismology, computer science and epidemiology. In this paper we study sharp deviations, fluctuations and the stable probability approximation of Poisson shot noise processes. Our achievements extend, improve and complement existing results in the literature. We apply the theoretical results to Poisson cluster point processes, including generalized linear Hawkes processes, and risk processes with delayed claims. Many examples are discussed in detail.

Keywords: Central limit theorem; Hawkes processes; Poisson cluster processes; Poisson shot noise processes; Ruin probabilities; Sharp deviations; Stable laws.

1 Introduction

We consider Poisson shot noise processes $\{S_t\}_{t \geq 0}$ of the form

$$S_t := \sum_{n \geq 1} H(t - T_n, M_n) \mathbf{1}_{(0, t]}(T_n), \quad (1)$$

where $\{T_n\}_{n \geq 1}$ is a homogeneous Poisson process on $(0, \infty)$ with intensity $\lambda > 0$, $\{M_n\}_{n \geq 1}$ is a sequence of random variables with values on some measurable space (M, \mathcal{M}) , independent of $\{T_n\}_{n \geq 1}$, $H : [0, \infty) \times M \rightarrow \mathbb{R}$ is a measurable function and $\mathbf{1}_A(\cdot)$ is the indicator function of a set A . We suppose that the random variables $\{M_n\}_{n \geq 1}$ are independent and identically distributed.

Poisson shot noise processes are natural generalizations of compound Poisson processes, which have found applications in different fields, due to their versatility and mathematical tractability. In insurance mathematics, Poisson shot noise processes arise as models of Incurred But Not Reported Claims [21, 22]. In this context, T_n represents the instant at which the n th claim arrives, for any $m \in M$, the function $H(\cdot, m)$ is non-negative and non-decreasing, the random quantity $H(\infty, M_n)$ models the total pay-off caused by the n th claim and the random function $H(\cdot - T_n, M_n)$ models the evolution of the pay-off process for the n th claim. We refer the reader to [5, 14, 17, 21, 22, 29, 30, 37] (and the literature cited therein) for specific applications in insurance mathematics of Poisson shot noise processes, such as estimates of ruin probabilities of risk processes with delay in claim settlement. In neurophysiology, Poisson shot noise processes appear as models of synaptic input. In this context, T_n describes the n th presynaptic event happened in the time interval $(0, t]$, $H(\cdot, \cdot)$ is the impulse-response function and M_n models a possible synaptic inhomogeneity

*Consiglio Nazionale delle Ricerche, Via dei Taurini 19, 00185 Roma, Italy. e-mail: giovanniluca.torrisi@cnr.it

†Dipartimento di Elettronica e Telecomunicazioni, Politecnico di Torino, Italy. e-mail: emilio.leonardi@polito.it

[6, 33]. In computer science, Poisson shot noise processes are used e.g. as traffic, queues or caches models [3, 16, 24, 25, 28, 37]. Poisson shot noise processes are also exploited to model earthquake aftershocks [38] and epidemics [31]. Spatial versions of $\{S_t\}_{t>0}$ are proposed in [1] to model the interference in wireless communication networks, see also [2, 15, 34, 36]. We study sharp estimates, fluctuations and stable probability approximation of spatial Poisson shot noise processes, with applications to communication networks, in a companion paper.

Over the years, the mathematics of Poisson shot noise processes have been investigated by many authors. The central limit theorem and the Berry-Esseen bound have been proved in [26, 27]. The law of the large numbers and functional central limit theorems have been studied in [22]. Scalar and sample path large deviations are investigated in [5, 16, 29, 35]. In [23], the authors study the weak convergence to a multivariate infinite-variance stable distribution of the finite-dimensional distributions of a properly normalized and centred Poisson shot noise process.

In words, the main theoretical contributions of this paper are: (i) The sharp deviations, at scales $O(t)$, of $\{S_t\}_{t>0}$ from its asymptotic mean; this result improves the tail estimates based on large deviations given in [5, 29]; (ii) The fluctuations, at scales $o(t)$, of $\{S_t\}_{t>0}$ from its asymptotic mean; these results improve the central limit theorem in [26]; (iii) In the case of a multiplicative noise, we provide quantitative limit theorems for the weak convergence, as $t \rightarrow \infty$, of S_t (properly re-scaled) to a random variable S with a stable law; these results complement the research started in [23]. We emphasize that the results about stable approximations of Poisson shot noise processes cover only stable laws with stability parameter α and skewness parameter β such that either $\alpha \neq 1$ or $\alpha = 1$ and $\beta = 0$ (see Subsection 2.3). We remark that, although stable laws have been defined also for $\alpha = 1$ and $\beta \neq 0$, the most relevant and well-known stable laws (i.e., the Gaussian, the Cauchy and the Lévy) are all encompassed by our study.

From the point of view of the applications, our main achievements concern: (i) The sharp deviations and fluctuations of Poisson cluster processes and, in particular, of generalized linear Hawkes processes, extending in this way the results in [18]; (ii) Estimates of ruin probabilities of risk processes with delayed claims, refining the large deviation approximation provided in [5].

Most of our theoretical contributions are achieved by means of the recently developed mod- ϕ convergence theory [12, 13]. Roughly speaking, provided that a natural normalization of the characteristic function of a stochastic process converges to some non-trivial limiting function, mod- ϕ convergence theory allows us to obtain precise deviations and fluctuations of the process from its asymptotic mean, improving classical results stemming from large deviations and central limit theorems.

The paper is structured as follows. In Section 2 we give some preliminaries on mod- ϕ convergence theory, compound Poisson and stable laws. Moreover, we state an elementary inequality between complex numbers and the Faà di Bruno formula which will be exploited a lot of times throughout the paper. In Section 3 we present the results on sharp deviations and fluctuations from the asymptotic mean of Poisson shot noise processes. Applications to Poisson cluster processes and ruin probabilities are described in Sections 4 and 5, respectively. In particular, in Section 4 we shall consider extensions of the classical linear Hawkes process (see [19]), where the number of offspring of any parent is not necessarily Poisson distributed and the law of the birth times is not necessarily absolutely continuous with respect to the Lebesgue measure. In Section 6 we state the results concerning the stable approximation of Poisson shot noise processes (with a multiplicative noise), which extend well-known results for the compound Poisson process (which are indeed recovered considering a constant shot shape). All the proofs are given in Section 7.

2 Preliminaries

2.1 Mod- ϕ convergence

We preliminary recall that a real-valued random variable X (or its law) is said infinitely divisible if, for any $n \in \mathbb{N} := \{1, 2, \dots\}$,

$$X \stackrel{d}{=} X_1 + \dots + X_n,$$

for some independent and identically distributed random variables X_1, \dots, X_n . Here the symbol $\stackrel{d}{=}$ denotes the equality in law.

We proceed providing the definition of mod- ϕ convergence, see [12].

Definition 2.1 *Let ϕ be a non-constant infinitely divisible law on \mathbb{R} and let $D \subseteq \mathbb{C}$ be a subset of the complex plane which contains 0. We assume that the Laplace transform of ϕ is defined on D , i.e.,*

$$\left| \int_{\mathbb{R}} e^{zx} \phi(dx) \right| < \infty, \quad \text{for all } z \in D,$$

and it has Lévy exponent $\eta(\cdot)$ on D , i.e.,

$$\int_{\mathbb{R}} e^{zx} \phi(dx) = e^{\eta(z)}, \quad z \in D.$$

Let $\{X_t\}_{t>0}$ be a real-valued stochastic process with Laplace transform defined on D , i.e., such that $|\mathbb{E}[e^{zX_t}]| < \infty$, for all $t > 0$ and $z \in D$, and let $x(t)$, $t > 0$, be a positive function such that $x(t) \rightarrow +\infty$, as $t \rightarrow +\infty$.

We say that $\{X_t\}_{t>0}$ converges mod- ϕ on D , with parameter function $x(\cdot)$ and limiting function $\psi(\cdot)$, if $\psi : D \rightarrow \mathbb{C}$ is analytic, it does not vanish on $\text{Re}D := \{\text{Re}z : z \in D\}$ and

$$\text{for any compact } K \subseteq D, \quad \lim_{t \rightarrow \infty} \sup_{z \in K} |\psi_t(z) - \psi(z)| = 0,$$

where

$$\psi_t(z) := \mathbb{E}[e^{zX_t}] e^{-x(t)\eta(z)}.$$

We say that $\{X_t\}_{t>0}$ converges mod- ϕ on D with speed $O(x(t)^{-\sigma})$, for some $\sigma \in \mathbb{N}$, and limiting function $\psi(\cdot)$, if $\psi : D \rightarrow \mathbb{C}$ is analytic, it does not vanish on $\text{Re}D$ and

$$\text{for any compact } K \subseteq D \text{ there exists } C_K > 0: \quad \sup_{z \in K} |\psi_t(z) - \psi(z)| \leq C_K x(t)^{-\sigma}.$$

In this paper we consider two different classes of infinitely divisible (reference) laws: the compound Poisson law and the stable law.

2.2 Compound Poisson laws

Let X be a non-negative random variable such that

$$a_X \in (0, \infty], \quad \text{where } a_X := \sup\{\gamma : \mathbb{E}[e^{\gamma X}] < \infty\}. \quad (2)$$

Throughout this paper, we denote by $\phi_{\lambda, X}$ the compound Poisson law with Lévy exponent

$$\eta_{\lambda, X}(z) := \lambda(\mathbb{E}[e^{zX}] - 1), \quad z \in D_{cp}(a_X) := \{z \in \mathbb{C} : \text{Re}z < a_X\}, \quad (3)$$

i.e., a random variable Y has law $\phi_{\lambda, X}$ if and only if

$$\mathbb{E}[e^{zY}] = e^{\eta_{\lambda, X}(z)}, \quad z \in D_{cp}(a_X).$$

2.3 Stable laws

Let $c > 0$ (the scale parameter), $\alpha \in (0, 2]$ (the stability parameter) and $\beta \in [-1, 1]$ (the skewness parameter) be fixed. We consider the stable law $\phi_{c,\alpha,\beta}$ with parameters (c, α, β) , whose Fourier transform (or characteristic function) has Lévy exponent

$$\eta_{c,\alpha,\beta}(\mathbf{i}\xi) := -|c\xi|^\alpha(1 - \mathbf{i}\beta h(\alpha, \xi)\text{sgn}(\xi)), \quad \xi \in \mathbb{R} \setminus \{0\}, \quad \eta_{c,\alpha,\beta}(0) := 0,$$

i.e., a random variable S has law $\phi_{c,\alpha,\beta}$ if and only if $\mathbb{E}[e^{\mathbf{i}\xi S}] = e^{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}$, $\xi \in \mathbb{R}$. Here

$$h(\alpha, \xi) := \mathbf{1}\{\alpha \neq 1\} \tan\left(\frac{\pi\alpha}{2}\right) - \mathbf{1}\{\alpha = 1\} \frac{2}{\pi} \log|\xi| \quad \text{and} \quad \text{sgn}(\xi) \text{ denotes the sign of } \xi.$$

Since

$$|e^{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}| = e^{-|c\xi|^\alpha}, \quad \xi \in \mathbb{R}, \quad (4)$$

the characteristic function of $\phi_{c,\alpha,\beta}$ is integrable. Therefore, the stable law with parameters (c, α, β) has a density with respect to the Lebesgue measure. A standard computation shows that such a density is bounded above by $\frac{\Gamma(\frac{1}{\alpha})}{\alpha\pi c}$, where $\Gamma(\cdot)$ denotes the Euler gamma function.

For later purposes, we recall the following scaling property of the Lévy exponent $\eta_{c,\alpha,\beta}(\cdot)$. For any $t > 0$ and $\xi \in \mathbb{R}$, it holds:

$$t\eta_{c,\alpha,\beta}\left(\frac{\mathbf{i}\xi}{t^{1/\alpha}}\right) = \eta_{c,\alpha,\beta}(\mathbf{i}\xi) \quad \text{if either } \alpha \neq 1 \text{ or } \alpha = 1 \text{ and } \beta = 0. \quad (5)$$

Finally, we recall some famous stable laws: the standard Gaussian distribution corresponds to the stable law with parameters $(2^{-1/2}, 2, 0)$, the standard Cauchy distribution corresponds to the stable law with parameters $(1, 1, 0)$ and the standard Lévy distribution corresponds to the stable law with parameters $(1, 2^{-1}, 1)$.

2.4 An elementary inequality and the Faà di Bruno formula

Throughout this paper we exploit the following elementary inequality between complex numbers:

Lemma 2.2 *It holds:*

$$|e^{z_1} - e^{z_2}| \leq |z_1 - z_2| e^{\max\{\text{Re}z_1, \text{Re}z_2\}}, \quad z_1, z_2 \in \mathbb{C}.$$

Since we have not found a proof of this inequality in standard textbooks of complex analysis, we show it in Section 7.

Hereafter, for a sufficiently smooth function f we denote by $f^{(n)}$ its derivative of order $n \in \mathbb{N}$.

Lemma 2.3 (*Faà di Bruno formula*) *For any sufficiently smooth functions g and h ,*

$$(g \circ h)^{(j)}(x) = j! \sum_{i=1}^j \frac{g^{(i)}(h(x))}{i!} \sum_{m_1+m_2+\dots+m_i=j} \frac{h^{(m_1)}(x)}{m_1!} \dots \frac{h^{(m_i)}(x)}{m_i!}, \quad j \in \mathbb{N},$$

where the sum is taken over all the $m_1, \dots, m_i \in \mathbb{N}$ such that $m_1 + \dots + m_i = j$.

3 Sharp deviations and fluctuations of Poisson shot noise processes

As already mentioned in the Introduction, our analysis relies on the mod- ϕ convergence theory. Specifically, we obtain different sharp deviations estimates (at scale $O(t)$) of Poisson shot noise processes depending on whether the reference measure ϕ is non-lattice or lattice, as it can be realized by comparing the formulas (15) and (18). The distinction “ ϕ non-lattice” and “ ϕ lattice”, instead, has no impact on the results about the fluctuations (at scales $o(t)$) of Poisson shot noise processes. For the sake of completeness, we recall that a probability law is lattice if its support is included in a set of the form $\gamma_1 + \gamma_2\mathbb{Z}$, for some parameters $\gamma_1 \in \mathbb{R}$ and $\gamma_2 > 0$.

In both cases (non-lattice and lattice), the shot shape $H(\cdot, \cdot)$ is supposed to be a non-negative function. Moreover we assume that

$$Z := \sup_{t \geq 0} H(t, M_1) \text{ is such that } a := a_Z \in (0, \infty]. \quad (6)$$

Although in the non-lattice case, i.e., under the assumption

$$\phi_{\lambda, Z} \text{ is non-lattice,} \quad (7)$$

we simply suppose that:

$$\text{function } H : [0, \infty) \times M \rightarrow [0, \infty) \text{ is non-negative,} \quad (8)$$

in the lattice case, we naturally assume that:

$$\text{function } H : [0, \infty) \times M \rightarrow \mathbb{N} \cup \{0\} \text{ takes values in } \mathbb{N} \cup \{0\}. \quad (9)$$

A crucial hypothesis to prove the mod- $\phi_{\lambda, Z}$ convergence of Poisson shot noise processes is

$$\int_0^\infty (Z - H(s, M_1)) \, ds \in L^q(\mathbb{P}), \text{ for any } q > 1. \quad (10)$$

However, when sharp deviations are concerned and $\phi_{\lambda, Z}$ is lattice we need to strengthen (10) by assuming

$$\exists \sigma \in \mathbb{N} \text{ and } \{\kappa_q\}_{q>1} \subset (0, \infty): \quad (11)$$

$$\sup_{q>1} \kappa_q^{-1} \left\| \int_t^\infty (Z - H(s, M_1)) \, ds \right\|_{L^q(\mathbb{P})} \leq t^{-\sigma}, \text{ for all } t \text{ large enough.}$$

Indeed, this condition guarantees the mod- $\phi_{\lambda, Z}$ convergence of Poisson shot noise processes with speed $O(t^{-\sigma})$. Finally, we mention that to prove the fluctuations of Poisson shot noise processes when $\phi_{\lambda, Z}$ is non-lattice, we need to assume that $\phi_{\lambda, Z}$ has a density, i.e.,

$$\phi_{\lambda, Z} \text{ is absolutely continuous with respect to the Lebesgue measure.} \quad (12)$$

Hereon, under (6) (and either (8) or (9))

$\forall x \in (0, \lambda \mathbb{E}[Ze^{aZ}])$, we denote by $\theta_x \in (-\infty, a)$ the unique solution of the equation $\lambda \mathbb{E}[Ze^{\theta_x Z}] = x$.

Note that $0 < x < \lambda \mathbb{E}[Z]$ if and only if $\theta_x < 0$; $x = \lambda \mathbb{E}[Z]$ if and only if $\theta_x = 0$; $x \in (\lambda \mathbb{E}[Z], \lambda \mathbb{E}[Ze^{aZ}])$ if and only if $\theta_x \in (0, a)$.

Throughout this article, we denote by $\eta_{\lambda,Z}^*$ the Fenchel-Legendre transform of $\eta_{\lambda,Z}$ (see (3)), i.e.,

$$\eta_{\lambda,Z}^*(x) := \sup_{\theta \in \mathbb{R}} (\theta x - \eta_{\lambda,Z}(\theta)), \quad x > 0.$$

Under (6) (and either (8) or (9)), standard computations show that

$$\eta_{\lambda,Z}^*(x) = x\theta_x - \lambda(\mathbb{E}[e^{\theta_x Z}] - 1), \quad x > 0. \quad (13)$$

Hereafter, for ease of notation, we also set

$$\varphi(z) := \int_0^\infty (\mathbb{E}[e^{zH(s, M_1)}] - \mathbb{E}[e^{zZ}]) \, ds, \quad z \in \mathbb{C}. \quad (14)$$

3.1 Sharp deviations at scales $O(t)$

The following theorems provide the exact asymptotic behavior of the tail of the Poisson shot noise when the reference compound Poisson law $\phi_{\lambda,Z}$ is either non-lattice or lattice, respectively.

Theorem 3.1 (Non-lattice case) *Assume (6), (7), (8) and (10). Then, for any $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$, we have*

$$\mathbb{P}(S_t \geq tx) = \frac{\exp(-t\eta_{\lambda,Z}^*(x) + \lambda\varphi(\theta_x))}{\theta_x \sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} (1 + o(1)), \quad \text{as } t \rightarrow +\infty. \quad (15)$$

Theorem 3.2 (Lattice case). *Assume (6), (9) and (11). Then:*

(i) *For any $x \in (0, \lambda\mathbb{E}[Ze^{aZ}])$ such that $tx \in \mathbb{N}$, we have*

$$\mathbb{P}(S_t = tx) = \frac{\exp(-t\eta_{\lambda,Z}^*(x) + \lambda\varphi(\theta_x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} (1 + o(1)), \quad \text{as } t \rightarrow +\infty. \quad (16)$$

More in general, for any $x \in (0, \lambda\mathbb{E}[Ze^{aZ}])$ such that $tx \in \mathbb{N}$, we have

$$\mathbb{P}(S_t = tx) = \frac{\exp(-t\eta_{\lambda,Z}^*(x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} \left(e^{\lambda\varphi(\theta_x)} + \sum_{k=1}^{\sigma-1} \frac{a_k(\theta_x)}{t^k} + O\left(\frac{1}{t^\sigma}\right) \right), \quad \text{as } t \rightarrow +\infty, \quad (17)$$

where the quantities $a_k(\theta_x)$ are computed in Proposition 7.3.

(ii) *For any $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$ such that $tx \in \mathbb{N}$, we have*

$$\mathbb{P}(S_t \geq tx) = \frac{\exp(-t\eta_{\lambda,Z}^*(x) + \lambda\varphi(\theta_x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} \frac{1}{1 - e^{-\theta_x}} (1 + o(1)), \quad \text{as } t \rightarrow +\infty. \quad (18)$$

More in general, for any $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$ such that $tx \in \mathbb{N}$, we have

$$\mathbb{P}(S_t \geq tx) = \frac{\exp(-t\eta_{\lambda,Z}^*(x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} \frac{1}{1 - e^{-\theta_x}} \left(e^{\lambda\varphi(\theta_x)} + \sum_{k=1}^{\sigma-1} \frac{b_k(\theta_x)}{t^k} + O\left(\frac{1}{t^\sigma}\right) \right), \quad \text{as } t \rightarrow +\infty, \quad (19)$$

where the quantities $b_k(\theta_x)$ are computed in Proposition 7.3.

We emphasize that Theorem 3.1 and Theorem 3.2(ii) refine the tail estimates provided by the corresponding large deviations principle in [16] (see also [29]).

We shall give many examples where the quantities θ_x , $\mathbb{E}[e^{\theta_x Z}]$ and $\mathbb{E}[Z^2 e^{\theta_x Z}]$ can be computed. We shall give also some examples where the integral $\varphi(\theta_x)$ can be computed. When this is not possible, our starting point to provide estimates of the integral is the following elementary proposition.

Proposition 3.3 *Under the foregoing assumptions and notation, we have:*

(i) *If $\theta_x < 0$ (i.e., $x \in (0, \lambda\mathbb{E}[Z])$), then*

$$0 \leq \varphi(\theta_x) \leq -\theta_x \int_0^\infty \mathbb{E}[Z - H(s, M_1)] ds.$$

(ii) *If $\theta_x > 0$ (i.e., $x \in (\lambda\mathbb{E}[Z], \mathbb{E}[Ze^{aZ}])$), then for any $q, q' > 1$ such that $q^{-1} + q'^{-1} = 1$, we have*

$$-\theta_x \|e^{\theta_x Z}\|_{L^q(\mathbb{P})} \left\| \int_0^\infty (Z - H(s, M_1)) ds \right\|_{L^{q'}(\mathbb{P})} \leq \varphi(\theta_x) \leq 0.$$

We remark that this proposition is exploited e.g. in the application to Poisson cluster processes to exhibit explicit estimates of the integral $\varphi(\theta_x)$, see Proposition 4.1(iii) and Proposition 4.2(vi).

3.2 Fluctuations at scales $o(t)$

The fluctuations of the Poisson shot noise process from its asymptotic mean, at any scale which is $o(t)$ as $t \rightarrow \infty$, are provided by the following theorem, where $N(0, 1)$ denotes a random variable distributed according to the standard Gaussian law. We refer the reader to Remark 3.5 for a brief discussion on the range of the scaling functions for the process $\{S_t - \lambda\mathbb{E}[Z]t\}_{t>0}$.

Theorem 3.4 *Assume either (6), (8), (10) and (12) or (6), (9) and (10). Then:*

(i) *For any function $y(\cdot)$ such that $y(t) = o(t^{1/6})$, it holds*

$$\mathbb{P} \left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} > y(t) \right) = \mathbb{P}(N(0, 1) > y(t))(1 + o(1)), \quad \text{as } t \rightarrow +\infty. \quad (20)$$

In particular, if $y(t) \rightarrow \infty$ as $t \rightarrow +\infty$, then (by the asymptotics of the tail of $N(0, 1)$)

$$\mathbb{P} \left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} > y(t) \right) = \frac{e^{-\frac{y(t)^2}{2}}}{y(t)\sqrt{2\pi}}(1 + o(1)), \quad \text{as } t \rightarrow +\infty.$$

(ii) *For any function $y(\cdot)$ such that $y(t) \rightarrow +\infty$, as $t \rightarrow \infty$, and $y(t) = o(t^{1/2})$, it holds*

$$\begin{aligned} \mathbb{P} \left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} > y(t) \right) &= \frac{e^{-t\eta_{\lambda, Z}^*(v(t))}}{\theta_{v(t)} \sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_{v(t)} Z}]}} (1 + o(1)) \\ &= \frac{e^{-t\eta_{\lambda, Z}^*(v(t))}}{y(t)\sqrt{2\pi}} (1 + o(1)), \quad \text{as } t \rightarrow +\infty, \end{aligned} \quad (21)$$

and

$$t\eta_{\lambda, Z}^*(v(t)) = \frac{y(t)^2}{2}(1 + o(1)). \quad (22)$$

Here

$$v(t) := \lambda\mathbb{E}[Z] + \frac{y(t)}{\sqrt{t}} \sqrt{\lambda\mathbb{E}[Z^2]}.$$

(iii) *For any function $y(\cdot)$ such that $y(t) \rightarrow +\infty$, as $t \rightarrow \infty$, and $y(t) = o\left(t^{\frac{1}{2} - \frac{1}{m}}\right)$, $m \geq 3$ integer, it holds*

$$\mathbb{P} \left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} > y(t) \right) = \frac{\exp \left(-\frac{y(t)^2}{2} \left(1 + 2 \sum_{j=1}^{m-2} (\lambda\mathbb{E}[Z^2])^{(j+2)/2} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(j+1)}}{(j+2)!} \left(\frac{y(t)}{\sqrt{t}} \right)^j \right) \right)}{y(t)\sqrt{2\pi}} (1 + o(1)), \quad (23)$$

as $t \rightarrow +\infty$. Here, the derivatives of θ_x evaluated at $\lambda\mathbb{E}[Z]$, denoted by $\theta_{\lambda\mathbb{E}[Z]}^{(j+1)}$, can be recursively computed by the formula:

$$\theta_{\lambda\mathbb{E}[Z]}^{(j+1)} = j! \sum_{i=1}^j \frac{g^{(i)}(0)}{i!} \sum_{m_1+m_2+\dots+m_i=j} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(m_1)}}{m_1!} \dots \frac{\theta_{\lambda\mathbb{E}[Z]}^{(m_i)}}{m_i!}, \quad 1 \leq j \leq m-2, \quad (24)$$

where the sum is taken over all the $m_1, \dots, m_i \in \mathbb{N}$ such that $m_1 + \dots + m_i = j$,

$$\theta'_{\lambda\mathbb{E}[Z]} = \frac{1}{\lambda\mathbb{E}[Z^2]} \quad \text{and} \quad g(x) := \frac{1}{\lambda\mathbb{E}[Z^2 e^{xZ}]}.$$

3.3 Some remarks

We conclude this section with the following three remarks.

Remark 3.5 (On the range of the scaling functions) Theorems 3.1 and 3.4 cover the whole range of scalings for the process $\{S_t - \lambda\mathbb{E}[Z]t\}_{t>0}$ up to the order of t . Indeed, Theorem 3.4(i) covers scalings, say $s(t)$, of $S_t - \lambda\mathbb{E}[Z]t$ such that $s(t) \ll t^{1/6}\sqrt{t} = t^{2/3}$. Theorem 3.4(ii) covers scalings of $S_t - \lambda\mathbb{E}[Z]t$ such that either $s(t) \sim t^{1/6}\sqrt{t} = t^{2/3}$ or $t^{2/3} \ll s(t) \ll t$. Finally, Theorem 3.1 refers to scalings of $S_t - \lambda\mathbb{E}[Z]t$ of order t .

Remark 3.6 (Central Limit Theorem and Extended Central Limit Theorem) Let the assumptions of Theorem 3.4 prevail. Then it is well-known that the classical Central Limit Theorem for the Poisson shot noise holds, i.e.,

$$\frac{S_t - \mathbb{E}[S_t]}{\sqrt{\text{Var}(S_t)}} \rightarrow N(0, 1) \quad \text{in law, as } t \rightarrow \infty.$$

(see e.g. Theorem 2.3 in [22]). This Central Limit Theorem can be retrieved by using Theorem 3.4(i). Indeed, as we shall check later on (see the comment after the statement of Lemma 7.1), the conditions (6), (8) and (10) guarantee

$$\mathbb{E}[S_t] = \lambda\mathbb{E}[Z]t + O(1) \quad \text{and} \quad \text{Var}(S_t) = \lambda\mathbb{E}[Z^2]t + O(1). \quad (25)$$

It follows

$$\frac{S_t - \mathbb{E}[S_t]}{\sqrt{\text{Var}(S_t)}} = \left(1 + \frac{O(1)}{\sqrt{t}}\right)^{-1/2} \left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} - \frac{O(1)}{\sqrt{t}}\right).$$

Therefore, for any $x \in \mathbb{R}$,

$$\mathbb{P}\left(\frac{S_t - \mathbb{E}[S_t]}{\sqrt{\text{Var}(S_t)}} > x\right) = \mathbb{P}\left(\frac{S_t - \lambda\mathbb{E}[Z]t}{\sqrt{\lambda\mathbb{E}[Z^2]t}} > x(t)\right) = \mathbb{P}(N(0, 1) > x(t))(1 + o(1)), \quad (26)$$

where the latter relation is a consequence of (20) since

$$x(t) := x \left(1 + \frac{O(1)}{\sqrt{t}}\right)^{1/2} + \frac{O(1)}{\sqrt{t}} = o(t^{1/6}). \quad (27)$$

The Central Limit Theorem for the Poisson shot noise process stated at the beginning follows combining (26) with the trivial relation

$$\mathbb{P}(N(0, 1) > x(t)) = \mathbb{P}(N(0, 1) > x)(1 + o(1)).$$

In fact, Theorem 3.4 is a big improvement of the classical Central Limit Theorem for the Poisson shot noise process, yielding the Extended Central Limit Theorem for the Poisson shot noise process.

Remark 3.7 As it will be clear from the proofs, in all the theorems above, conditions (10) and (11) can be replaced respectively by

$$Z \in L^\infty(\mathbb{P}) \text{ and } \int_0^\infty (Z - H(s, M_1)) \, ds \in L^1(\mathbb{P}),$$

and

$$Z \in L^\infty(\mathbb{P}) \text{ and } \exists \sigma \in \mathbb{N} \text{ and } \kappa > 0:$$

$$\left\| \int_t^\infty (Z - H(s, M_1)) \, ds \right\|_{L^1(\mathbb{P})} \leq \kappa t^{-\sigma} \text{ for all } t \text{ large enough.}$$

Note also that if $Z \in L^\infty(\mathbb{P})$, then Z satisfies (6) with $a := a_Z = +\infty$.

4 Application to Poisson cluster processes

Let $N_n(\cdot)$, $n \geq 1$, be independent and identically distributed finite and simple point processes on $[0, \infty)$, independent on $\{T_n\}_{n \geq 1}$. We assume $N_n(\{0\}) = 1$ for any $n \geq 1$. Denoting by $\{S_{n,k}\}_{k \geq 0}$, $S_{n,0} := 0$ the (ordered) points of N_n , and interpreting T_n as the ancestor of the point process $\theta_{T_n} N_n \equiv \{S_{n,k} + T_n\}_{k \geq 0}$ of offspring, we have that, at time $t > 0$, the total number of offspring generated by the ancestors (and including the ancestors) is equal to

$$\sum_{n \geq 1} \mathbf{1}_{(0,t]}(T_n) \theta_{T_n} N_n([T_n, t]) = \sum_{n \geq 1} \mathbf{1}_{(0,t]}(T_n) N_n([0, t - T_n]) = \sum_{n \geq 1} \mathbf{1}_{(0,t]}(T_n) H(t - T_n, M_n), \quad (28)$$

where $M_n := N_n$, i.e., M_n is a random variable with values on $\mathcal{M} := \mathcal{N}$, i.e., the space of finite and simple counting measures on $[0, \infty)$ (endowed with the usual vague topology), and $H : [0, \infty) \times \mathcal{M} \rightarrow \mathbb{N} \cup \{0\}$ is defined by

$$H(t, m) = H(t, \mu) := \mu([0, t]).$$

Thus, assuming that $\{T_n\}_{n \geq 1}$ is a homogeneous Poisson process with intensity λ , at time $t > 0$, the total number of offspring generated by the ancestors (including the ancestors) is equal to the Poisson shot noise process S_t defined by (28).

We put $L_n := \sup_{k \geq 1} S_{n,k}$, $n \geq 1$, and note that L_n is the “length” of the cluster point process $N_n(\cdot)$. Clearly, the random variables $\{L_n\}_{n \geq 1}$ are independent and identically distributed, and we set $L := L_1$. The following proposition holds.

Proposition 4.1 *Let $\{S_t\}_{t \geq 0}$ be the Poisson shot noise process (28), i.e., S_t denotes the number of offspring generated by the Poissonian ancestors $\{T_n\}_{n \geq 1}$ in the time interval $(0, t]$ (including the ancestors). Assume that $Z := N_1([0, \infty))$ satisfies (6), and that*

$$\mathbb{E}[L^k] < \infty, \forall k \in \mathbb{N}. \quad (29)$$

Then, setting $H(s, M_1) := N_1([0, s])$, $s \in [0, t]$, we have that:

- (i) *The formulas (20), (21) and (23) of Theorem 3.4 hold.*
- (ii) *For any arbitrarily fixed $\sigma \in \mathbb{N}$, the formulas (17) and (19) of Theorem 3.2 hold.*
- (iii) *The following estimates hold:*
 - (1) *For any $x \in (0, \lambda \mathbb{E}[Z])$ and any $q, q' > 1$ such that $q^{-1} + q'^{-1} = 1$, we have*

$$0 \leq \varphi(\theta_x) \leq -\theta_x \|L\|_{L^q(\mathbb{P})} \|Z\|_{L^{q'}(\mathbb{P})}. \quad (30)$$

(2) For any $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$ and any $q, q', q_1, q_2 > 1$ such that $q^{-1} + q'^{-1} = 1$ and $q_1^{-1} + q_2^{-1} = 1$, we have

$$-\theta_x \|e^{\theta_x Z}\|_{L^q(\mathbb{P})} \|L\|_{L^{q'q_1}(\mathbb{P})} \|Z\|_{L^{q'q_2}(\mathbb{P})} \leq \varphi(\theta_x) \leq 0. \quad (31)$$

Here the function $\varphi(\cdot)$ is given by (14) with Z and $H(\cdot, M_1)$ defined at the beginning of the statement.

We note that the Part (ii) of this proposition refines the tail estimates provided by the large deviations principle in [4].

4.1 Poisson cluster processes with a Galton-Watson branching structure or generalized linear Hawkes processes

We consider Poisson cluster processes whose clusters have a Galton-Watson branching structure. Clearly, to define such processes it suffices to describe the structure of the cluster point process N_1 .

The points in the cluster C_0 , generated by a common ancestor placed in the origin, are partitioned in generations $g \in \mathbb{N} \cup \{0\}$. Every point belonging to the $(g-1)$ th generation will give rise to a random number of “children” points in the g th generation. By definition, N_1 is the point process with support C_0 . Let

- K_h be the number of points in the h th generation,
- W_h be the total number of points in the cluster C_0 up the h th generation,
- $\{B_1^h, \dots, B_{K_h}^h\}$ be the birth times (i.e., the times at which points are placed) in the h th generation.

We assume that the birth times are arranged in increasing order, i.e.,

$$B_1^h < \dots < B_{K_h}^h.$$

Let $\{P_{i,j}\}_{(i,j) \in \mathbb{N}^2}$ be a sequence of $\mathbb{N} \cup \{0\}$ -valued independent random variables with law $\{p_k\}_{k \geq 0}$, where P_{ij} represents the number of children generated by the j th individual of the i th generation. Let $\{B_{i,j,k}\}_{(i,j,k) \in \mathbb{N}^3}$ be a sequence of non-negative independent and identically distributed random variables, independent of $\{P_{i,j}\}_{(i,j) \in \mathbb{N}^2}$, where B_{ijk} represents the time lag between the birth time of the j th individual of the i th generation and the birth time of its k th child.

The point $\{0\}$ constitutes the 0th generation. By construction we have $K_0 = 1$ and $W_0 = 1$. The birth time of the unique individual in this generation is given by $B_1^0 := 0$. The points in the g th generation are generated by the points in the $(g-1)$ th generation according to the following rule:

(i) If $K_{g-1} = 0$, then the $(g-1)$ th generation is empty and then the g th generation will be empty as well. We set $K_g := 0$ and $W_g := W_{g-1}$.

(ii) If $K_{g-1} > 0$, then:

- $K_g := P_{g,1} + \dots + P_{g,K_{g-1}}$ is the number of births in the g th generation.
- $W_g := W_{g-1} + K_g$ is the total number of points in the cluster C_0 until the g th generation.
- The birth times in the g th generation are given by the union of the following sets:

$$\{B_1^{g-1} + B_{g,1,1}, \dots, B_1^{g-1} + B_{g,1,P_{g,1}}\},$$

which are the birth times of the children of the parent born at time B_1^{g-1} ,

$$\{B_2^{g-1} + B_{g,2,1}, \dots, B_2^{g-1} + B_{g,2,P_{g,2}}\},$$

which are the birth times of the children of the parent born at time B_2^{g-1}

...

$$\{B_{K_{g-1}}^{g-1} + B_{g,K_{g-1},1}, \dots, B_{K_{g-1}}^{g-1} + B_{g,K_{g-1},P_{g,K_{g-1}}}\},$$

which are the birth times of the children of the parent born at time $B_{K_{g-1}}^{g-1}$.

The points in the g th generation are then arranged in increasing order as

$$\{B_1^g, \dots, B_{K_g}^g\}.$$

If the law of $B_{1,1,1}$ has a probability density $h(\cdot)/\int_0^\infty h(t) dt$, where $h : (0, \infty) \rightarrow [0, \infty)$ is an integrable function, and the law of $P_{1,1}$ is Poisson with mean $\int_0^\infty h(t) dt$, then the corresponding Poisson shot noise process $\{S_t\}_{t>0}$, where S_t denotes the number of points generated by the Poissonian ancestors in the time interval $(0, t]$, is a classical Hawkes process (or linear Hawkes process) [19], for which large deviations were studied in [4] and sharp deviations and fluctuations have been recently investigated in [18].

The sequence $\{K_n\}_{n \in \mathbb{N} \cup \{0\}}$ is a Galton-Watson process from an initial population of one individual and with offspring law $\{p_k\}_{k \geq 0}$. Hereon, we assume that

$$0 < \mathbb{E}[P_{1,1}] < 1, \tag{32}$$

so that the Galton-Watson process is subcritical and the total progeny

$$Z := N_1([0, \infty)) = \sum_{n \geq 0} K_n$$

has mean equal to $(1 - \mathbb{E}[P_{1,1}])^{-1}$. For later purposes, we recall that the distribution of the total progeny $Z := N_1([0, \infty))$ is related to the offspring distribution by the formula

$$\mathbb{P}(Z = k) = \frac{1}{k} \mathbb{P}(P_{1,1} + \dots + P_{1,k} = k - 1), \quad k \in \mathbb{N} \tag{33}$$

(the reader is referred to e.g. [20] for an introduction to branching processes). Hereafter, we also suppose that

$$b := a_{P_{1,1}} \in (0, \infty], \tag{34}$$

where $a_{P_{1,1}}$ is defined by (2). So (being the branching process subcritical) by Theorem 2.1 in [32] we have that the total progeny Z satisfies (6).

The “length” of N_1 is given by

$$L := \sup_{g \in \mathbb{N}} \sup_{j \leq K_g} B_j^g.$$

Define

$$V_g := \sum_{k=1}^{K_{g-1}} \sum_{j=1}^{P_{g,k}} B_{g,k,j}.$$

Clearly, V_1 is an upper bound of the latest birth in the first generation; $V_1 + V_2$ is an upper bound of the latest birth until the second generation and by induction

$$V := \sum_{g \geq 1} V_g \quad (35)$$

is clearly an upper bound for L . Hereon, we suppose

$$\mathbb{E}[B_{1,1,1}^k] < \infty, \text{ for any } k \in \mathbb{N}, \quad (36)$$

and denote by $G_{P_{1,1}}$ the probability generating function of $P_{1,1}$. In the next proposition, we provide the fluctuations and the sharp deviations of the number of points S_t up to time t .

Proposition 4.2 *Assume (32), (34) and (36). Then, setting $H(s, M_1) := N_1([0, s])$, $s \in [0, t]$, we have that:*

- (i) *The formulas (20), (21) and (23) of Theorem 3.4 hold.*
- (ii) *For any arbitrarily fixed $\sigma \in \mathbb{N}$, the formulas (17) and (19) of Theorem 3.2 hold.*
- (iii) *Setting*

$$a_c := \sup_{\theta \geq 0} (\theta - \log \mathbb{E}[e^{\theta P_{1,1}}]),$$

we have $a_c \in (0, a_Z]$.

- (iv) *Set $b_c := \sup \Theta$, where*

$$\Theta := \{\theta : \theta < a_c \text{ and } \mathbb{E}[e^{\theta Z}] < e^b\},$$

and b is defined by (34) (note that $b_c > 0$). We have

$$\mathbb{E}[e^{\theta Z}] = e^{\theta} G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) < \infty, \quad \text{for any } \theta \in (-\infty, b_c),$$

and the moments of Z satisfy the formula:

$$\begin{aligned} \mathbb{E}[Z^n] = 1 + \sum_{k=1}^n k! \binom{n}{k} \sum_{i=1}^k \frac{\mathbb{E}[P_{1,1}(P_{1,1}-1) \dots (P_{1,1}-(i-1))]}{i!} \\ \sum_{m_1+m_2+\dots+m_i=k} \frac{\mathbb{E}[Z^{m_1}]}{m_1!} \dots \frac{\mathbb{E}[Z^{m_i}]}{m_i!}, \quad \text{for any } n \geq 1, \end{aligned} \quad (37)$$

where the third sum is taken over all the $m_1, \dots, m_i \in \mathbb{N}$ such that $m_1 + \dots + m_i = k$.

- (v) *For any $x \in (0, \lambda \mathbb{E}[Z e^{b_c Z}])$, let $\varrho_x \in (0, \mathbb{E}[e^{b_c Z}])$ be the unique solution of the equation in ϱ :*

$$\frac{\lambda \varrho}{1 - \varrho \frac{G'_{P_{1,1}}(\varrho)}{G_{P_{1,1}}(\varrho)}} = x. \quad (38)$$

Then

$$\theta_x = \log \frac{\varrho_x}{G_{P_{1,1}}(\varrho_x)}, \quad \varrho_x = \mathbb{E}[e^{\theta_x Z}],$$

and

$$\mathbb{E}[Z^2 e^{\theta_x Z}] = \varrho_x \frac{1 + e^{\theta_x} (G'_{P_{1,1}}(\varrho_x) + G''_{P_{1,1}}(\varrho_x))}{(1 - e^{\theta_x} G'_{P_{1,1}}(\varrho_x))^2}.$$

- (vi) *The following estimates hold:*

- (1) *For any $x \in (0, \lambda(1 - \mathbb{E}[P_{1,1}])^{-1})$ and any $q, q' > 1$ such that $q^{-1} + q'^{-1} = 1$, we have*

$$p_1 \mathbb{E}[B_{1,1,1}] e^{\theta_x} (1 - \mathbb{E}[e^{\theta_x Z}]) \leq \varphi(\theta_x) \leq -\theta_x \|B_{1,1,1}\|_{L^q(\mathbb{P})} \|Z\|_{L^q(\mathbb{P})} \|Z\|_{L^{q'}(\mathbb{P})}. \quad (39)$$

Note that these bounds are non trivial for any choice of the conjugate exponents.

(2) For any $x \in (\lambda(1 - \mathbb{E}[P_{1,1}])^{-1}, \lambda \mathbb{E}[Ze^{b_c Z}])$ and any $q, q', q_1, q_2 > 1$ such that $q^{-1} + q'^{-1} = 1$ and $q_1^{-1} + q_2^{-1} = 1$, we have

$$-\theta_x \|e^{\theta_x Z}\|_{L^q(\mathbb{P})} \|B_{1,1,1}\|_{L^{q'q_1}(\mathbb{P})} \|Z\|_{L^{q'q_1}(\mathbb{P})} \|Z\|_{L^{q'q_2}(\mathbb{P})} \leq \varphi(\theta_x) \leq p_1 \mathbb{E}[B_{1,1,1}] e^{\theta_x} (1 - \mathbb{E}[e^{\theta_x Z}]). \quad (40)$$

Note that the upper bound is always non trivial. The lower bound is non trivial for any $q \in (1, b_c/\theta_x)$.

Here, the function $\varphi(\cdot)$ involved in the relations (39) and (40) is defined as in the statement of Proposition 4.1.

4.2 Example 1: Binomial offspring distribution

Suppose that $P_{1,1}$ has a binomial distribution with parameters (m, p) such that $\mathbb{E}[P_{1,1}] = mp < 1$, $m \geq 1$, $p \in (0, 1)$. Then condition (32) is satisfied, and the assumption (34) holds with $b = +\infty$, indeed

$$G_{P_{1,1}}(\varrho) = (\varrho p + (1 - p))^m, \quad \varrho \in \mathbb{R}.$$

A standard computation gives

$$a_c = b_c = \log \left(\frac{1}{mp} \left(\frac{m-1}{m(1-p)} \right)^{m-1} \right)$$

(since $0^0 := 1$, if $m = 1$ then $a_c = b_c = -\log p$). Using the formula (33), we have that Z is distributed according to the Consul distribution, i.e.,

$$\mathbb{P}(Z = k) = \frac{1}{k} \binom{km}{k-1} p^{k-1} (1-p)^{km-k+1}, \quad k \in \mathbb{N}$$

(see e.g. [7]). Therefore

$$\begin{aligned} \mathbb{E}[e^{b_c Z}] &= \sum_{k \geq 1} \left(\frac{1}{mp} \left(\frac{m-1}{m(1-p)} \right)^{m-1} \right)^k \frac{1}{k} \binom{km}{k-1} p^{k-1} (1-p)^{km-k+1} \\ &= \frac{1-p}{p} \sum_{k \geq 1} \frac{1}{k} \binom{mk}{k-1} \left(\frac{(m-1)^{m-1}}{m^m} \right)^k = +\infty, \end{aligned} \quad (41)$$

and so $\mathbb{E}[Ze^{b_c Z}] = +\infty$. To check that the infinite sum in (41) diverges, we note that it is trivially equal to $+\infty$ if $m = 1$ and that, by construction, the total progeny of a Galton-Watson process with offspring distribution the binomial law with parameters (m, p) is bigger than or equal to the total progeny of a Galton-Watson process with offspring distribution the Bernoulli law with parameter p . Note that the equation (38) reads

$$\frac{\lambda p \varrho^2 + \lambda(1-p)\varrho}{p(1-m)\varrho + (1-p)} = x,$$

i.e.,

$$\lambda p \varrho^2 + (\lambda(1-p) + (m-1)px)\varrho - (1-p)x = 0, \quad x \in (0, \infty),$$

which gives

$$\varrho_x = \frac{-(\lambda(1-p) + (m-1)px) + \sqrt{(\lambda(1-p) + (m-1)px)^2 + 4xp(1-p)\lambda}}{2\lambda p}.$$

Therefore, we know explicitly the quantities θ_x , $\mathbb{E}[e^{\theta_x Z}]$ and $\mathbb{E}[Z^2 e^{\theta_x Z}]$.

Note that the formulas (16) and (18) involve also the integral $\varphi(\theta_x)$ whose estimates (39) and (40) involve, in turn, the moments of Z and of $e^{\theta_x Z}$. Hereafter, for the sake of completeness we briefly describe how those estimates can be computed. For any $x \in (0, \lambda(1 - mp)^{-1})$, choose, for instance, $q = q' = 2$ in the upper bound of (39). Then both the lower and the upper bounds on the integral are explicit (once the law of $B_{1,1,1}$ is fixed). Indeed, $p_1 = mp(1 - p)^{m-1}$, we already computed the quantities θ_x and $\mathbb{E}[e^{\theta_x Z}]$ and by (37) one has $\mathbb{E}[Z] = (1 - \mathbb{E}[P_{1,1}])^{-1}$, and

$$\mathbb{E}[Z^2] = 1 + 2\mathbb{E}[P_{1,1}]\mathbb{E}[Z] + \mathbb{E}[P_{1,1}]\mathbb{E}[Z^2] + (\mathbb{E}[P_{1,1}^2] - \mathbb{E}[P_{1,1}]) (\mathbb{E}[Z])^2,$$

from which $\mathbb{E}[Z^2]$ is readily calculated. For any $x \in (\lambda(1 - mp)^{-1}, \infty)$, we note that (once the law of $B_{1,1,1}$ is fixed) the upper bound in (40) is always explicit. As far as the lower bound in (40) is concerned, we note that, fixed $q \in (1, b_c/\theta_x)$, the finite quantity $\|e^{\theta_x Z}\|_{L^q(\mathbb{P})}$ can be computed e.g. numerically (the law of Z is explicitly known). Taking e.g. $q_1 = q_2 = 2$ and letting q' be the conjugate exponent of q , we have

$$\|Z\|_{2q'} \leq \mathbb{E}[Z^{\lceil 2q' \rceil}]^{1/(2q')}$$

(where $\lceil \cdot \rceil$ denotes the ceiling function), which yields a computable lower bound (recall that $-\theta_x < 0$), using the formula (37).

4.3 Example 2: Geometric offspring distribution

Suppose that $P_{1,1}$ has a geometric distribution with parameter $p \in (1/2, 1)$. Then condition (32) is satisfied, and the assumption (34) holds with $b = -\log(1 - p)$, indeed

$$G_{P_{1,1}}(\varrho) = \frac{p}{1 - \varrho(1 - p)}, \quad |\varrho| < (1 - p)^{-1}.$$

A standard computation gives

$$a_c = -\log(4p(1 - p)).$$

Using the formula (33), we have

$$\mathbb{P}(Z = k) = \frac{1}{k} \binom{2(k-1)}{k-1} (1 - p)^{k-1} p^k, \quad k \in \mathbb{N}.$$

For all $\theta \leq a_c = -\log(4p(1 - p))$, it holds

$$\begin{aligned} \sum_{k \geq 1} \frac{1}{k} \binom{2(k-1)}{k-1} [(1 - p)pe^\theta]^k &= (1 - p)pe^\theta \sum_{h \geq 0} \frac{1}{h+1} \binom{2h}{h} [(1 - p)pe^\theta]^h \\ &= \frac{1 - \sqrt{1 - 4(1 - p)pe^\theta}}{2} < 1, \end{aligned}$$

where the latter equality follows recognizing the generating function of the Catalan numbers. Therefore, for all $\theta \leq a_c$ we have $\mathbb{E}[e^{\theta Z}] < 1/(1 - p)$, which implies $b_c = a_c$. Moreover,

$$\mathbb{E}[e^{b_c Z}] = \frac{1}{4(1 - p)} \sum_{k \geq 1} \frac{1}{k} \binom{2(k-1)}{k-1} \left(\frac{1}{4}\right)^{k-1} = \frac{1}{8(1 - p)},$$

and

$$\mathbb{E}[Ze^{b_c Z}] = \frac{1}{4(1 - p)} \sum_{h \geq 0} \binom{2h}{h} \left(\frac{1}{4}\right)^h = +\infty.$$

Note that the equation (38) reads

$$\frac{\lambda \varrho(1 - (1 - p)\varrho)}{1 - 2(1 - p)\varrho} = x,$$

i.e.,

$$-\lambda(1 - p)\varrho^2 + (2(1 - p)x + \lambda)\varrho - x = 0, \quad x > 0,$$

which gives

$$\varrho_x = \frac{(2(1 - p)x + \lambda) - \sqrt{(2(1 - p)x + \lambda)^2 - 4\lambda x(1 - p)}}{2\lambda(1 - p)}.$$

Therefore, we know explicitly the quantities θ_x , $\mathbb{E}[e^{\theta_x Z}]$ and $\mathbb{E}[Z^2 e^{\theta_x Z}]$. Similar considerations as in the case of a binomial offspring distribution yield estimates of the integral $\varphi(\theta_x)$, which also appears in the formulas (16) and (18).

4.4 Example 3: Poisson offspring distribution, i.e., linear Hawkes processes with a general displacement distribution

As already mentioned, precise deviations and fluctuations of classical Hawkes processes (or linear Hawkes processes) have been studied in [18]. Here, for the sake of completeness, we briefly explain how to apply Proposition 4.2 even to the linear Hawkes process.

We emphasize that to apply Proposition 4.2 we need to assume that the displacement distribution, i.e., the law of $B_{1,1,1}$, has all moments finite. The results in [18], instead, require only the existence of some moments for the displacement distribution, which, however, must have a density with respect to the Lebesgue measure. Hereon, instead, we do not require the law of $B_{1,1,1}$ to be absolutely continuous with respect to the Lebesgue measure.

So, suppose that $P_{1,1}$ has a Poisson distribution with mean $\mu \in (0, 1)$. Then condition (32) is satisfied, and the assumption (34) holds with $b = +\infty$, indeed

$$G_{P_{1,1}}(\varrho) = e^{\mu(\varrho-1)}, \quad y \in \mathbb{R}.$$

A standard computation gives

$$a_c = \mu - 1 - \log \mu.$$

Using the formula (33), we have that Z is distributed according to the Borel distribution, i.e.,

$$\mathbb{P}(Z = k) = \frac{(\mu k)^{k-1}}{k!} e^{-\mu k}, \quad k \in \mathbb{N}$$

(see e.g. [7]). Clearly, $b_c = a_c$. Moreover, standard computations yield

$$\mathbb{E}[e^{b_c Z}] = \mu^{-1} \quad \text{and} \quad \mathbb{E}[Z e^{b_c Z}] = +\infty.$$

Note that the equation (38) reads

$$\frac{\lambda \varrho}{1 - \mu \varrho} = x,$$

which gives

$$\varrho_x = \frac{x}{\lambda + \mu x}.$$

Therefore, we know explicitly the quantities θ_x , $\mathbb{E}[e^{\theta_x Z}]$ and $\mathbb{E}[Z^2 e^{\theta_x Z}]$. Similar considerations as in the case of a binomial offspring distribution yield estimates of the integral $\varphi(\theta_x)$, which also appears in the formulas (16) and (18).

5 Further applications

5.1 Ruin probabilities of risk processes with delayed claims

Consider an insurance company with initial capital $u > 0$ and premium rate $c > 0$. As already mentioned in the Introduction, the total claim amount, up to time $t > 0$, due to Incurred But Not Reported Claims is often modeled by a Poisson shot noise process S_t of the form (1), see [21, 22]. We recall that, in this context, it is assumed

$$\text{For any } m \in \mathbb{M}, \text{ the function } H(\cdot, m) \text{ is non-negative and non-decreasing.} \quad (42)$$

The ruin probability of the insurance company is clearly given by

$$\psi_{IBNR}(u) := \mathbb{P} \left(\sup_{t>0} (S_t - ct) \geq u \right), \quad u > 0.$$

Letting

$$\psi_{CL}(u) := \mathbb{P} \left(\sup_{t>0} (C_t - ct) \geq u \right), \quad u > 0,$$

denote the ruin probability of the “associated” Cramér-Lundberg risk process, where

$$C_t := \sum_{n \geq 1} H(\infty, M_n) \mathbf{1}_{(0, t]}(T_n), \quad t > 0,$$

it is well-known that (see e.g. Theorem 1.2.2 in [11]) if

$$(2) \text{ holds with } Z := H(\infty, M_1) \text{ in place of } X, \lambda(\mathbb{E}[e^{aZ}] - 1) - ca > 0, \quad (43)$$

$$\int_0^\infty x e^{wx} \mathbb{P}(Z > x) dx < \infty, \quad (44)$$

and the “net profit” condition

$$c > \lambda \mathbb{E}[Z] \quad (45)$$

is satisfied, then

$$\psi_{CL}(u) = \left(\frac{\lambda w}{c - \lambda \mathbb{E}[Z]} \int_0^\infty x e^{wx} \mathbb{P}(Z > x) dx \right)^{-1} e^{-wu} (1 + o(1)), \quad \text{as } u \rightarrow \infty. \quad (46)$$

Here $w > 0$ is the unique positive solution of the equation in γ :

$$\lambda(\mathbb{E}[e^{\gamma Z}] - 1) - c\gamma = 0. \quad (47)$$

Using large deviations techniques, it was proved in [5] that, under the foregoing assumptions,

$$\lim_{u \rightarrow \infty} \frac{1}{u} \log \psi_{IBNR}(u) = -w.$$

The next proposition improves such ruin probability estimate.

Proposition 5.1 *Assume (42), (43), (44) and (45). Then:*

(i) *If (7) and (10) hold, then*

$$\begin{aligned} \frac{\exp(\lambda \varphi(w))}{w \sqrt{2\lambda\pi(\lambda \mathbb{E}[Z e^{wZ}] - c)^{-1} \mathbb{E}[Z^2 e^{wZ}]}} \frac{e^{-wu}}{\sqrt{u}} (1 + o(1)) &\leq \psi_{IBNR}(u) \\ &\leq \left(\frac{\lambda w}{c - \lambda \mathbb{E}[Z]} \int_0^\infty x e^{wx} \mathbb{P}(Z > x) dx \right)^{-1} e^{-wu} (1 + o(1)), \quad \text{as } u \rightarrow \infty. \end{aligned}$$

(ii) If (9) and (11) hold, then

$$\begin{aligned} \frac{\exp(\lambda\varphi(w))}{w(1-e^{-w})\sqrt{2\lambda\pi(\lambda\mathbb{E}[Ze^{wZ}] - c)^{-1}\mathbb{E}[Z^2e^{wZ}]}} \frac{e^{-wu}}{\sqrt{u}}(1+o(1)) &\leq \psi_{IBNR}(u) \\ &\leq \left(\frac{\lambda w}{c - \lambda\mathbb{E}[Z]} \int_0^\infty xe^{wx}\mathbb{P}(Z > x) dx \right)^{-1} e^{-wu}(1+o(1)), \quad \text{as } u \rightarrow \infty. \end{aligned}$$

5.1.1 Example 4: a risk process with delayed claims

We consider a risk process with delay in claim settlement and total claim amount at time $t > 0$ given by the Poisson shot noise process

$$S_t := \sum_{n \geq 1} F(t - T_n) M_n \mathbf{1}_{(0, t]}(T_n),$$

where F is the distribution function of a law on $(0, \infty)$ with a finite mean. We suppose that the random variable M_1 is non-negative, it has a density with respect to the Lebesgue measure and it is such that $a_{M_1} \in (0, \infty]$ (see (2)). Then, setting $\bar{F} := 1 - F$, we have

$$M_1 \int_0^\infty \bar{F}(s) ds \in L^q(\mathbb{P}), \quad \text{for any } q > 1,$$

and the compound Poisson law ϕ_{λ, M_1} has a density with respect to the Lebesgue measure. Note that Z is distributed as M_1 and the assumptions of Theorems 3.1 and 3.4 are satisfied.

Exponentially distributed claims : sharp deviations at scales $O(t)$

Suppose that M_1 is exponentially distributed with mean $1/a$, $a \in (0, \infty)$. Then, a straightforward computation shows

$$\theta_x = a - \sqrt{\frac{\lambda a}{x}}, \quad x > \lambda/a, \quad (48)$$

$$\mathbb{E}[e^{\theta_x M_1}] = \frac{a}{a - \theta_x} = \sqrt{\frac{ax}{\lambda}}, \quad (49)$$

$$\mathbb{E}[M_1^2 e^{\theta_x M_1}] = \frac{2a}{(a - \theta_x)^3} = 2\frac{x}{\lambda} \sqrt{\frac{x}{\lambda a}}, \quad (50)$$

and

$$\begin{aligned} \varphi(\theta_x) &= \int_0^\infty (\mathbb{E}[e^{\theta_x F(s) M_1}] - \mathbb{E}[e^{\theta_x M_1}]) ds = \int_0^\infty \left(\frac{a}{a - \theta_x F(s)} - \frac{a}{a - \theta_x} \right) ds \\ &= -a \left(\sqrt{\frac{ax}{\lambda}} - 1 \right) \Phi(\lambda, a, x, F), \end{aligned}$$

where

$$\Phi(\lambda, a, x, F) := \int_0^\infty \frac{\bar{F}(s)}{a - \left(a - \sqrt{\frac{\lambda a}{x}} \right) F(s)} ds.$$

So by Theorem 3.1 we have that, for any $x > \lambda/a$, as $t \rightarrow \infty$,

$$\mathbb{P}(S_t \geq tx) = \frac{(\lambda/ax)^{1/4}}{2\sqrt{\pi}(\sqrt{ax} - \sqrt{\lambda})} \exp\left(-a(\sqrt{\lambda ax} - \lambda)\Phi(\lambda, a, x, F)\right) \frac{1}{\sqrt{t}} e^{-(xa + \lambda - 2\sqrt{\lambda ax})t}(1 + o(1)).$$

For specific choices of $F(\cdot)$ the quantity Φ can be computed. For instance, if $F(\cdot)$ is the distribution function of the uniform law on $(0, 1)$, then

$$\int_0^\infty \frac{\bar{F}(s)}{a - \left(a - \sqrt{\frac{\lambda a}{x}}\right) F(s)} ds = -\frac{\sqrt{\lambda x/a}}{(\sqrt{ax} - \sqrt{\lambda})^2} \log \sqrt{\frac{ax}{\lambda}} + \frac{1}{a - \sqrt{\frac{\lambda a}{x}}}.$$

For this model, one may easily compute the ruin probabilities estimates provided by Proposition 5.1(i). We omit the details.

Fluctuations at scales $o(t)$

Hereon, by applying Theorem 3.4(iii) with $y(t) := o(t^{1/4})$, $y(t) \rightarrow \infty$, we study the fluctuations of S_t around its asymptotic mean at a scale of order $o(t^{3/4})$. To perform the computation, we do not need to know the distribution of M_1 (as it happens to study the sharp deviations of S_t from its asymptotic mean, where we need to compute the function θ), but only the first four moments of M_1 . Indeed, under the foregoing assumptions, the formula (23) holds with $Z = M_1$ and $m = 4$ and, assuming the knowledge of the quantities $\mathbb{E}[M_1^i]$, $i = 1, 2, 3, 4$, we only need to compute $\theta_{\lambda\mathbb{E}[M_1]}^{(j)}$ for $j = 2, 3$. To this aim, by the recursive formula (24) we have

$$\theta_{\lambda\mathbb{E}[M_1]}^{(2)} = g'(0)\theta_{\lambda\mathbb{E}[M_1]}^{(1)} = -\frac{\lambda\mathbb{E}[M_1^3]}{(\lambda\mathbb{E}[M_1^2])^3}$$

and

$$\begin{aligned} \theta_{\lambda\mathbb{E}[M_1]}^{(3)} &= g'(0)\theta_{\lambda\mathbb{E}[M_1]}^{(2)} + g''(0)(\theta_{\lambda\mathbb{E}[M_1]}^{(1)})^2 \\ &= \frac{(\lambda\mathbb{E}[M_1^3])^2}{(\lambda\mathbb{E}[M_1^2])^5} - \frac{\lambda}{(\lambda\mathbb{E}[M_1^2])^2} \left(\mathbb{E}[M_1^4] - 2\lambda \frac{(\mathbb{E}[M_1^3])^2}{\mathbb{E}[M_1^2]} \right). \end{aligned}$$

5.2 A teletraffic model

In this subsection we briefly discuss the application of Theorems 3.1 and 3.4 to a teletraffic model proposed in [24].

We consider an infinite servers queuing system, handling in parallel active jobs (connections). New jobs arrive at the system according to a homogeneous Poisson process $\{T_n\}_{n \geq 1}$. Let $\{M_n\}_{n \geq 1}$ denote the sequence of the processing times of the jobs, which are assumed to be independent and identically distributed. Every job en-queued is served by the system at unit rate. At time t , the number of active jobs in the system is given by

$$X_t := \sum_{n \geq 1} \mathbf{1}_{(0, M_n]}(t - T_n) \mathbf{1}_{(0, t]}(T_n).$$

Then, the total workload processed by the system up to time t is given by the Poisson shot noise process

$$S_t := \int_0^t X_s ds = \sum_{n \geq 1} [(t - T_n) \wedge M_n] \mathbf{1}_{(0, t]}(T_n),$$

where, for $a, b \in \mathbb{R}$, the symbol $a \wedge b$ denotes the minimum between a and b . Hereon, we suppose that the random variable M_1 has a density with respect to the Lebesgue measure and that $a_{M_1} \in (0, \infty]$ (see (2)). Then

$$\int_0^\infty (M_1 - s \wedge M_1) ds = \frac{M_1^2}{2} \in L^q(\mathbb{P}), \quad \text{for any } q > 1,$$

and the compound Poisson law ϕ_{λ, M_1} is absolutely continuous with respect to the Lebesgue measure. Here again, Z is distributed as M_1 and the assumptions of Theorems 3.1 and 3.4 are satisfied. Assuming the knowledge of only the first four moments of M_1 , by the same application of Theorem 3.4(iii) with $y(t) := o(t^{1/4})$, $y(t) \rightarrow \infty$, discussed in the Example 4, we can quantify the fluctuations of S_t around its asymptotic mean at a scale of order $o(t^{3/4})$. If we assume that M_1 is exponentially distributed with mean $1/a$, $a \in (0, \infty)$, then the function θ is given by (48) and the quantities $\mathbb{E}[e^{\theta_x M_1}]$ and $\mathbb{E}[M_1^2 e^{\theta_x M_1}]$ are provided by (49) and (50), respectively. Moreover

$$\begin{aligned} \varphi(\theta_x) &= \int_0^\infty (\mathbb{E}[e^{\theta_x(s \wedge M_1)}] - \mathbb{E}[e^{\theta_x M_1}]) \, ds = \mathbb{E} \left[\int_0^{M_1} (e^{\theta_x s} - e^{\theta_x M_1}) \, ds \right] \\ &= -\frac{1}{\theta_x} + \frac{a}{\theta_x(a - \theta_x)} - \frac{a}{(a - \theta_x)^2} \\ &= \sqrt{\frac{x}{\lambda a}} - \frac{x}{\lambda}. \end{aligned}$$

So by Theorem 3.1 we have the explicit asymptotic expression of the tail of the processed workload.

6 Stable probability approximation of Poisson shot noise processes

We recall that throughout this paper we denote by $\phi_{c, \alpha, \beta}$ the stable law with scale parameter $c > 0$, stability parameter $\alpha \in (0, 2]$ and skewness parameter $\beta \in [-1, 1]$ (see Subsection 2.3 for the definition of stable distribution and its first properties). In the context of the stable probability approximation of the Poisson shot noise, we consider the following assumptions on the model (1):

The random variable M_1 is distributed according to $\phi_{c, \alpha, \beta}$ with either $\alpha \neq 1$ or $\alpha = 1$ and $\beta = 0$. (51)

For any $t > 0$ and $m \in \mathbb{R}$, $H(t, m) := mF(t)$, for some distribution function $F : [0, \infty) \rightarrow [0, 1]$. (52)

The following theorem holds.

Theorem 6.1 *Assume (51) and (52) and let S be a random variable with law $\phi_{c\lambda^{1/\alpha}, \alpha, \beta}$. Then $S_t/t^{1/\alpha}$ converges in law to S , as $t \rightarrow \infty$.*

The next theorem quantifies the speed of such weak convergence. For the sake of readability, we state this result by giving estimates of the Kolmogorov distance between $S_t/t^{1/\alpha}$ and S in terms of big O functions. However, in the proof we compute explicitly the constants involved in the bounds; anyway, such constants have a complicated expression.

Letting X, Y denote two real-valued random variables, we recall that the Kolmogorov distance between X and Y is defined by

$$d_{\text{Kol}}(X, Y) := \sup_{x \in \mathbb{R}} |\mathbb{P}(X \leq x) - \mathbb{P}(Y \leq x)|.$$

Theorem 6.2 *Let the assumptions and notation of Theorem 6.1 prevail. Then:*

(i) *Define*

$$A := \left\{ \eta \in [\max\{0, (\alpha - 1)/\alpha\}, 1/2] : \limsup_{t \rightarrow \infty} \frac{\int_0^t (1 - F(s)^\alpha) \, ds}{t^\eta} < \infty \right\}$$

and $\eta_0 := \inf A$.

If

$$\eta_0 \in A \tag{53}$$

then

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) = O\left(\frac{1}{t^{1-\eta_0}}\right), \quad \text{as } t \rightarrow \infty.$$

Instead, if

$$A \neq \emptyset \quad \text{and} \quad \eta_0 \notin A,$$

then for any $\tilde{\eta}_0 \in (\eta_0, 1)$ it holds

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) = O\left(\frac{1}{t^{1-\tilde{\eta}_0}}\right), \quad \text{as } t \rightarrow \infty.$$

(ii) If

$$\int_0^\infty (1 - F(s)^\alpha) ds \in [0, \infty) \tag{54}$$

then

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) = O\left(\frac{1}{t^{1/\alpha}}\right) \mathbf{1}\{\alpha \in (1, 2]\} + O\left(\frac{1}{t}\right) \mathbf{1}\{\alpha \in (0, 1) \cup \{\alpha = 1, \beta = 0\}\}, \quad \text{as } t \rightarrow \infty.$$

Remark 6.3 In [23] the authors study the weak convergence to a multivariate α -stable distribution of the finite-dimensional distributions of a properly centred and scaled Poisson shot noise process. In particular, Section 3.3 of [23] considers Poisson shot noise processes of the form

$$X_t := \sum_{n \geq 1} M_n h(t - T_n) \mathbf{1}_{(0, t]}(T_n), \quad t > 0,$$

where $\{T_n\}_{n \geq 1}$ is a homogeneous Poisson process on $(0, \infty)$, independent of the sequence of independent and identically distributed random variables $\{M_n\}_{n \geq 1}$, and $h(\cdot)$ is a measurable and positive function. By the theory developed in [23] it follows that if the law of M_1 is regularly varying at infinity with a constant sign, and the shot shape function $h(\cdot)$ is bounded on the compacts and regularly varying too, then X_t , properly centred and scaled, converges weakly to a stable law. Although the Poisson shot noise model S_t considered in Theorem 6.1 has a multiplicative noise, we consider different hypotheses on the mark M_1 and on the shot shape function $F(\cdot)$. For instance, we do not assume that both the distribution of M_1 and the shot shape function are regularly varying. Furthermore, Theorem 6.2 provides the speed of the weak convergence to the stable law, which is far from the achievements in [23].

6.1 Example 5: shot shapes for the stable probability approximation

Assuming (51) and (52), hereon we discuss some possible choices of the shot shape $F(\cdot)$ which allows us to apply Theorem 6.2.

We start noticing that if the law with distribution function $F(\cdot)$ has a finite mean, then by Theorem 6.2(ii) we have

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) = O\left(\frac{1}{t^{1/\alpha}}\right) \mathbf{1}\{\alpha \in (1, 2]\} + O\left(\frac{1}{t}\right) \mathbf{1}\{\alpha \in (0, 1) \cup \{\alpha = 1, \beta = 0\}\}.$$

Indeed, since $\alpha \in (0, 2]$ we have $F(t)^2 \leq F(t)^\alpha$ for any $t > 0$. Therefore

$$\int_0^\infty (1 - F(s)^\alpha) ds \leq \int_0^\infty (1 - F(s)^2) ds \leq 2 \int_0^\infty (1 - F(s)) ds < \infty.$$

If either $\alpha \in (1, 2]$ or $\alpha = 1$ and $\beta = 0$, and the law with distribution function $F(\cdot)$ has an infinite mean, then Theorem 6.2(ii) can not be applied, indeed $F(t) \geq F(t)^\alpha$ for any $t > 0$. In such a case we can hope to apply Theorem 6.2(i). This is the case, for instance, when $F(\cdot)$ is the distribution function of a Pareto law with parameter 1, i.e.,

$$F(t) := \frac{t}{1+t} \mathbf{1}_{[0, \infty)}(t).$$

Indeed, we have

$$\int_0^\infty (1 - F(t)) dt = \infty,$$

and so Theorem 6.2(ii) does not apply (condition (54) is not satisfied). However, by applying twice de l'Hopital's theorem, for any $\eta \in (0, 1/2]$ we have

$$\begin{aligned} \lim_{t \rightarrow \infty} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t^\eta} &= \lim_{t \rightarrow \infty} \frac{1 - F(t)^\alpha}{\eta t^{\eta-1}} \\ &= \frac{\alpha}{\eta(1-\eta)} \lim_{t \rightarrow \infty} \frac{F(t)^{\alpha-1} \frac{t^2}{(t+1)^2}}{t^\eta} = 0. \end{aligned} \quad (55)$$

If $\alpha \in (1, 2]$, then $\eta_0 = (\alpha - 1)/\alpha \in A$ (the set A is defined in the statement of Theorem 6.2) and so by Theorem 6.2(i) we have $d_{\text{Kol}}(S_t/t^{1/\alpha}, S) = O(1/t^{1/\alpha})$. If $\alpha = 1$ and $\beta = 0$, then $\eta_0 = 0 \notin A$ and so by Theorem 6.2(i) we have $d_{\text{Kol}}(S_t/t, S) = O(1/t^{1-\eta})$, for any $\eta \in (0, 1)$. Of interest for applications in insurance it is the case when the total pay-off M_1 has the heavy tail Lévy distribution, which corresponds to the stable law with parameters $c = 1$, $\alpha = 1/2$ and $\beta = 1$. In such a case if the delay in claim settlement is modeled by a distribution function $F(\cdot)$ with a finite mean, then we fall in the first case discussed above and so

$$d_{\text{Kol}}(S_t/t^2, S) = O\left(\frac{1}{t}\right).$$

If, for instance, $F(\cdot)$ is the distribution function of a Pareto law with parameter 1, then (by the same computation as in (55))

$$\lim_{t \rightarrow \infty} \frac{\int_0^t (1 - F(s)^{1/2}) ds}{t^\eta} = 0, \quad \text{for any } \eta \in (0, 1/2],$$

and so $\eta_0 = 0 \notin A$. Therefore by Theorem 6.2(i) we have $d_{\text{Kol}}(S_t/t^2, S) = O(1/t^{1-\eta})$, for any $\eta \in (0, 1)$.

7 Proofs

7.1 Proof of Theorem 3.1

As already mentioned in the Introduction, the proof is based on the mod- ϕ convergence theory. So, first we state a lemma which concerns the mod-compound Poisson convergence of Poisson shot noise processes (i.e., the mod- ϕ convergence of Poisson shot noise processes when ϕ is the compound Poisson law), then we prove Theorem 3.1 applying the results in [12].

The following lemma, whose proof is postponed at the end of the subsection, holds.

Lemma 7.1 Assume (6), (8) and (10). Then $\{S_t\}_{t>0}$ converges mod- $\phi_{\lambda,Z}$ on $D_{cp} := D_{cp}(a)$ with parameter function $x(t) := t$ and limiting function

$$\psi(z) := e^{\lambda\varphi(z)}, \quad (56)$$

where $\varphi(\cdot)$ is defined by (14).

Before proving Theorem 3.1, we note that, under the assumptions and notation of Lemma 7.1, by the theory of mod- ϕ convergence (see p. 20 of [12]) we have the asymptotic expressions of the mean and the variance of S_t anticipated in (25).

Proof of Theorem 3.1. Let $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$ be fixed. Reasoning by contradiction, suppose

$$\limsup_{t \rightarrow \infty} \frac{\mathbb{P}(S_t \geq tx)}{\frac{\exp(-t\eta_{\lambda,Z}^*(x) + \lambda\varphi(\theta_x))}{\theta_x \sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}}} \neq 1, \quad (57)$$

and let $\{t_k\}_{k \geq 1} \subset (0, \infty)$ be a divergent sequence that realizes the limsup. By Lemma 7.1 the stochastic process $\{S_{t_k}\}_{k \geq 1}$ converges mod- $\phi_{\lambda,Z}$ on D_{cp} with parameter sequence $x(t_k) := t_k$ and limiting function (56). So, by Theorem 4.2.1 in [12] easily follows that the relation (15) holds with t_k in place of t . Indeed, in our framework, the functions $F(\cdot)$, h , and $\eta(\cdot)$ in the Theorem 4.2.1 of [12] are given, respectively, by the functions $\eta_{\lambda,Z}^*(\cdot)$ (in (13)), θ , and $\eta_{\lambda,Z}(\cdot)$. Moreover, the function $\psi(\cdot)$ and the quantities c and d of Theorem 4.2.1 in [12] are given, respectively, by the function $\psi(\cdot)$ defined by (56), and $c = -\infty$ and $d = a$. Therefore we reached a contradiction and in (57) we have the equality. The same arguments hold if in (57) we replace the limsup with the liminf. The proof is completed.

□

Proof of Lemma 7.1. Note that $\{(T_n, M_n)\}_{n \geq 1}$ is a Poisson process on $(0, \infty) \times M$ with mean measure $\lambda dt \mathbb{P}_{M_1}(dm)$. So, by the expression of the Laplace functional of a Poisson process (see e.g. [8]), we have

$$\mathbb{E}[e^{zS_t}] = \exp\left(\lambda \int_0^t (\mathbb{E}[e^{zH(s, M_1)}] - 1) ds\right), \quad \text{for any } t > 0 \text{ and } z \in D_{cp}.$$

By the elementary relation $|e^z| = e^{\operatorname{Re} z}$, $z \in \mathbb{C}$, we have

$$\begin{aligned} |\mathbb{E}[e^{zS_t}]| &\leq \mathbb{E}[e^{(\operatorname{Re} z)S_t}] \\ &= \exp\left(\lambda \int_0^t (\mathbb{E}[e^{(\operatorname{Re} z)H(s, M_1)}] - 1) \mathbf{1}\{\operatorname{Re} z \leq 0\} ds + \lambda \int_0^t (\mathbb{E}[e^{(\operatorname{Re} z)H(s, M_1)}] - 1) \mathbf{1}\{\operatorname{Re} z > 0\} ds\right) \\ &\leq \exp(\lambda t (\mathbb{E}[e^{(\operatorname{Re} z)Z}] - 1) \mathbf{1}\{\operatorname{Re} z > 0\}) < \infty, \quad \text{for any } t > 0 \text{ and } z \in D_{cp}. \end{aligned}$$

Therefore, the Laplace transform of S_t , $t > 0$, is defined on D_{cp} . We now check that $\psi(\cdot)$, defined by (56), does not vanish on $\operatorname{Re} D_{cp} = (-\infty, a)$ and it is analytic on D_{cp} . By the Lemma 2.2, for any $z \in D_{cp}$ and $s > 0$, we have

$$\begin{aligned} |e^{zH(s, M_1)} - e^{zZ}| &\leq |z|(Z - H(s, M_1))e^{\max\{(\operatorname{Re} z)H(s, M_1), (\operatorname{Re} z)Z\}} \\ &= |z|(Z - H(s, M_1))e^{\max\{(\operatorname{Re} z)H(s, M_1), (\operatorname{Re} z)Z\}} \mathbf{1}\{\operatorname{Re} z \leq 0\} \\ &\quad + |z|(Z - H(s, M_1))e^{\max\{(\operatorname{Re} z)H(s, M_1), (\operatorname{Re} z)Z\}} \mathbf{1}\{\operatorname{Re} z > 0\} \\ &\leq |z|(Z - H(s, M_1))e^{\max\{0, \operatorname{Re} z\}Z} \quad \mathbb{P}\text{-a.s..} \end{aligned}$$

Therefore, for any $z \in D_{cp}$ and $s > 0$,

$$\mathbb{E}[|e^{zH(s, M_1)} - e^{zZ}|] \leq |z| \mathbb{E}[(Z - H(s, M_1))e^{\max\{0, \operatorname{Re} z\}Z}]. \quad (58)$$

So, applying Hölder's inequality with $p \in (1, a/\max\{0, \operatorname{Re} z\})$, where we conventionally set $a/0 := \infty$, and $q > 1$ such that $p^{-1} + q^{-1} = 1$ (note that the conjugate exponents depend on z), we have

$$|\varphi(z)| \leq |z| \left\| \int_0^\infty (Z - H(s, M_1)) ds \right\|_{L^q(\mathbb{P})} \left\| e^{\max\{0, \operatorname{Re} z\}Z} \right\|_{L^p(\mathbb{P})} < \infty, \quad \text{for any } z \in D_{cp},$$

where the latter term is finite due to the assumptions (6) and (10). So

$$\psi(x) = e^{\lambda\varphi(x)} > 0, \quad \text{for any } x \in (-\infty, a).$$

To prove that $\psi(\cdot)$ is analytic on D_{cp} we show that $\varphi(\cdot)$ is such, and to this aim we prove that

$$\sup_{z \in K} |\varphi_t(z) - \varphi(z)| \rightarrow 0, \quad \text{as } t \rightarrow \infty, \text{ for any compact } K \subset D_{cp}, \quad (59)$$

where

$$\varphi_t(z) := \int_0^t (\mathbb{E}[e^{zH(s, M_1)}] - \mathbb{E}[e^{zZ}]) ds, \quad z \in D_{cp}$$

(then the analiticity of $\varphi(\cdot)$ on D_{cp} it follows by the analiticity of φ_t on D_{cp} for any $t > 0$). Let $K \subset D_{cp}$ be an arbitrarily fixed compact. Without loss of generality, we suppose that there exists $a' \in (0, a)$ such that $K \cap \{z : 0 < \operatorname{Re} z \leq a'\} \neq \emptyset$. Indeed, if $K \subset \{z : \operatorname{Re} z \leq 0\}$, then there exists $K' \subset D_{cp}$ compact such that $K \subset K'$ and $K' \cap \{z : 0 < \operatorname{Re} z \leq a'\} \neq \emptyset$. Hereon, we set

$$m_K := \max(\operatorname{Re} K).$$

By construction, $0 < m_K < a$, and, setting $\overline{M}_K := \sup_{z \in K} |z|$, by (58) it follows

$$\mathbb{E}[|e^{zH(s, M_1)} - e^{zZ}|] \leq \overline{M}_K \mathbb{E}[(Z - H(s, M_1))e^{m_K Z}], \quad \forall s > 0 \text{ and } z \in K.$$

By this relation and the definition of φ_t and φ , for any $t > 0$ and $z \in K$, we have

$$|\varphi_t(z) - \varphi(z)| \leq \overline{M}_K \mathbb{E} \left[e^{m_K Z} \int_t^\infty (Z - H(s, M_1)) ds \right].$$

Applying Hölder's inequality with $p \in (1, a/m_K)$ and $q > 1$ such that $p^{-1} + q^{-1} = 1$ (note that in this case the conjugate exponents depend on K , but not on z) for any $t > 0$ and $z \in K$, we have

$$|\varphi_t(z) - \varphi(z)| \leq \mathcal{M}_K(t) := \overline{M}_K \left(\mathbb{E}[e^{p m_K Z}]^{1/p} \right) \left\| \int_t^\infty (Z - H(s, M_1)) ds \right\|_{L^q(\mathbb{P})}. \quad (60)$$

The claim (59) follows taking first the supremum over K on this latter relation, and then letting t tend to infinity. Indeed, by the assumption (10) and the dominated convergence theorem we immediately have

$$\left\| \int_t^\infty (Z - H(s, M_1)) ds \right\|_{L^q(\mathbb{P})}^q \rightarrow 0, \quad \text{as } t \rightarrow \infty.$$

Finally we prove that

$$\psi_t(z) := e^{-t\eta_{\lambda, z}(z)} \mathbb{E}[e^{zS_t}], \quad t > 0, z \in D_{cp},$$

converges to $\psi(\cdot)$ uniformly on the compacts of D_{cp} . Let $K \subset D_{cp}$ be an arbitrarily fixed compact. For any $z \in K$, by first applying Lemma 2.2, then using the elementary inequality $\operatorname{Re} w \leq |w|$, $w \in \mathbb{C}$, and finally exploiting (60), we have

$$\begin{aligned} |\psi_t(z) - \psi(z)| &= \left| \exp \left(\lambda \int_0^t (\mathbb{E}[e^{zH(s, M_1)}] - \mathbb{E}[e^{zZ}]) \, ds \right) - \exp(\lambda \varphi(z)) \right| \\ &\leq \lambda \int_t^\infty \mathbb{E}[|e^{zH(s, M_1)} - e^{zZ}|] \, ds \times \exp \left(\lambda \int_0^\infty \mathbb{E}[|e^{zH(s, M_1)} - e^{zZ}|] \, ds \right) \leq \left(\lambda e^{\lambda \mathcal{M}_K(0)} \right) \mathcal{M}_K(t). \end{aligned} \quad (61)$$

The claimed local uniform convergence follows by first taking the supremum over K on this relation and then letting t tend to infinity.

□

7.2 Proof of Theorem 3.2

Here again, the proof is based on the mod- ϕ convergence theory. However, we need to refine Lemma 7.1 since we need the mod-compound Poisson convergence with a polynomial decay (see Definition 2.1). So, first we state such a refinement in a lemma and then we prove Theorem 3.2 applying the results in [12].

The following lemma, whose proof is postponed at the end of the subsection, holds.

Lemma 7.2 *Assume (6), (9) and (11). Then $\{S_t\}_{t>0}$ converges mod- $\phi_{\lambda, Z}$ on $D_{cp} := D_{cp}(a)$ with speed $O(t^{-\sigma})$ and limiting function (56).*

Proof of Theorem 3.2. Although the proof is conceptually similar to the proof of Theorem 3.1, since it exploits different results of the mod- ϕ convergence theory, we provide some details.

Proof of Part (i).

Let $x \in (0, \lambda \mathbb{E}[Ze^{aZ}])$ be such that $tx \in \mathbb{N}$. Reasoning by contradiction, suppose

$$\limsup_{t \rightarrow \infty} t^{-\sigma} \frac{\left| \mathbb{P}(S_t = tx) - \frac{\exp(-t\eta_{\lambda, Z}^*(x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}} \left(\psi(\theta_x) + \sum_{k=1}^{\sigma-1} \frac{a_k(\theta_x)}{t^k} \right) \right|}{\frac{\exp(-t\eta_{\lambda, Z}^*(x))}{\sqrt{2\lambda\pi t \mathbb{E}[Z^2 e^{\theta_x Z}]}}} = +\infty, \quad (62)$$

and let $\{t_k\}_{k \geq 1} \subset (0, \infty)$ be a divergent sequence that realizes the limsup. By Lemma 7.2 the stochastic process $\{S_{t_k}\}_{k \geq 1}$ converges mod- $\phi_{\lambda, Z}$ on D_{cp} with speed $O(1/(t_k)^\sigma)$ and limiting function (56). So, by Theorem 3.2.2(1) in [12] easily follows that the relation (17) holds with t_k in place of t . Indeed, in our framework, the functions $F(\cdot)$, h , and $\eta(\cdot)$ in the Theorem 3.2.2 of [12] are given, respectively, by the functions $\eta_{\lambda, Z}^*(\cdot)$ (in (13)), θ , and $\eta_{\lambda, Z}(\cdot)$. Moreover, the function $\psi(\cdot)$ and the quantities c and d of Theorem 3.2.2 in [12] are given, respectively, by the function $\psi(\cdot)$ defined by (56), and $c = -\infty$, $d = a$. Therefore we reached a contradiction and the limsup in (62) is finite. The proof is completed.

Proof of Part (ii).

Similar to the proof of Part (i). One has to apply Theorem 3.2.2(2) in [12] in place of Theorem 3.2.2(1) in [12].

□

Proof of Lemma 7.2. Note that the assumptions imply (10). Indeed, for any $t > 0$ and $q > 1$,

$$\mathbb{E} \left[\left(\int_0^t (Z - H(s, M_1)) \, ds \right)^q \right] \leq t^q \mathbb{E}[Z^q] < \infty, \quad (63)$$

where the latter quantity is finite by (6). So (10) follows by (63), (11) and Minkowski's inequality. Consequently, all the claims in the proof of Lemma 7.1 hold, and using the same notation of the lemma, we only have to prove that for any compact $K \subset D_{cp}$, there exists $C_K > 0$ such that

$$\sup_{z \in K} |\psi_t(z) - \psi(z)| \leq C_K t^{-\sigma}, \quad \text{for any } t > 0.$$

To this aim, we note that by the relations (60) and (61), for any $t > 0$ and $K \subset D_{cp}$, we have

$$|\psi_t(z) - \psi(z)| \leq \lambda e^{\lambda \mathcal{M}_K(0)} \overline{M}_K \left(\mathbb{E}[e^{p m_K Z}]^{1/p} \right) \left\| \int_t^\infty (Z - H(s, M_1)) ds \right\|_{L^q(\mathbb{P})}.$$

The claim follows by the assumption (11) (note that the conjugate exponents p and q in this latter inequality depend on K , but not on z).

□

7.3 Computation of the quantities $a_k(\theta_x)$ and $b_k(\theta_x)$, $1 \leq k \leq \sigma - 1$, and proof of Proposition 3.3

The following proposition provides the expression of the functions $a_k(\cdot)$ and $b_k(\cdot)$ in Theorem 3.2.

Proposition 7.3 *Let the assumptions and notation of Theorem 3.2 prevail. Then, for any $x \in (0, \lambda \mathbb{E}[Z e^{aZ}])$ and $k = 1, \dots, \sigma - 1$, we have*

$$\begin{aligned} a_k(\theta_x) &= \sum_{\ell=0}^{2k} \frac{\psi^{(2k-\ell)}(\theta_x)}{(2k-\ell)!} \sum_{(m_1, \dots, m_\ell) \in \mathcal{S}_\ell} \frac{(-1)^{m_1 + \dots + m_\ell}}{m_1! 1!^{m_1} m_2! 2!^{m_2} \dots m_\ell! \ell!^{m_\ell}} \\ &\quad \times \prod_{j=1}^{\ell} \left(\frac{\mathbb{E}[Z^{j+2} e^{\theta_x Z}]}{(j+1)(j+2) \mathbb{E}[Z^2 e^{\theta_x Z}]} \right)^{m_j} \frac{(-1)^k (2(k + m_1 + \dots + m_\ell) - 1)!!}{(\lambda \mathbb{E}[Z^2 e^{\theta_x Z}])^k} \end{aligned}$$

and

$$\begin{aligned} b_k(\theta_x) &= \sum_{n=0}^{2k} \sum_{(m_1, \dots, m_n) \in \mathcal{S}_n} \frac{e^{(m_1 + \dots + m_n) \theta_x} (m_1 + \dots + m_n)! (1 - e^{-\theta_x})^{-(m_1 + \dots + m_n) - 1}}{m_1! 1!^{m_1} m_2! 2!^{m_2} \dots m_n! n!^{m_n}} \prod_{j=1}^n (-1)^{j m_j} \\ &\quad \times \sum_{\ell=0}^{2k-n} \frac{\psi^{(2k-n-\ell)}(\theta_x)}{(2k-n-\ell)!} \sum_{\mathcal{S}_\ell} \frac{(-1)^{m_1 + \dots + m_\ell}}{m_1! 1!^{m_1} m_2! 2!^{m_2} \dots m_\ell! \ell!^{m_\ell}} \\ &\quad \times \prod_{j=1}^{\ell} \left(\frac{\mathbb{E}[Z^{j+2} e^{\theta_x Z}]}{(j+1)(j+2) \mathbb{E}[Z^2 e^{\theta_x Z}]} \right)^{m_j} \frac{(-1)^k (2(k + m_1 + \dots + m_\ell) - 1)!!}{(\lambda \mathbb{E}[Z^2 e^{\theta_x Z}])^k}. \end{aligned}$$

Here \mathcal{S}_n denotes the set of n -tuples of non-negative integers (m_1, \dots, m_n) such that $1 \cdot m_1 + 2 \cdot m_2 + \dots + n \cdot m_n = n$, $f^{(k)}$ denotes the derivative of order $k \in \mathbb{N}$ of a sufficiently smooth function f and $f^{(0)} = f$. The derivatives of the function $\psi(\cdot)$, defined by (56), can be computed by using the recursive formula:

$$\psi^{(n)}(\theta_x) = n! \sum_{i=1}^n \lambda^i \frac{e^{\lambda \varphi(\theta_x)}}{i!} \sum_{m_1 + m_2 + \dots + m_i = n} \frac{\varphi^{(m_1)}(\theta_x)}{m_1!} \dots \frac{\varphi^{(m_i)}(\theta_x)}{m_i!}, \quad n \in \mathbb{N},$$

where the sum is taken over all the $m_1, \dots, m_i \in \mathbb{N}$ such that $m_1 + \dots + m_i = n$, and

$$\varphi^{(n)}(\theta_x) = \int_0^\infty (\mathbb{E}[H(s, M_1)^n e^{\theta_x H(s, M_1)}] - \mathbb{E}[Z^n e^{\theta_x Z}]) ds.$$

Proof. The general expression of the functions $a_k(\cdot)$ and $b_k(\cdot)$ can be computed following the suggestions of Remark 3.2.5 in [12]. See Proposition 1 in [18] for the details. The recursive formula for the derivatives of ψ is easily obtained by applying the Faà di Bruno formula (see Lemma 2.3). \square

Proof of Proposition 3.3.

Proof of Part (i).

The lower bound is trivial. The upper bound follows immediately by applying Lemma 2.2.

Proof of Part (ii).

The upper bound is trivial. For the lower bound, note that by Lemma 2.2 and Hölder's inequality it follows

$$\int_0^\infty (\mathbb{E}[e^{\theta_x Z}] - \mathbb{E}[e^{\theta_x H(s, M_1)}]) \, ds \leq \theta_x \|e^{\theta_x Z}\|_{L^p(\mathbb{P})} \left\| \int_0^\infty (Z - H(s, M_1)) \, ds \right\|_{L^q(\mathbb{P})}.$$

\square

7.4 Proof of Theorem 3.4

Proof of Part (i).

Reasoning by contradiction, suppose

$$\limsup_{t \rightarrow \infty} \frac{\mathbb{P}\left(\frac{S_t - \lambda \mathbb{E}[Z]t}{\sqrt{\lambda \mathbb{E}[Z^2]t}} > y(t)\right)}{\mathbb{P}(N(0, 1) > y(t))} \neq 1, \quad (64)$$

and let $\{t_k\}_{k \geq 1} \subset (0, \infty)$ be a divergent sequence that realizes the \limsup . The proof of (20) then proceeds similarly to the proof of (15) in Theorem 3.1 [i.e., applying Lemma 7.1 and then Theorem 4.3.1 in [12] (in place of Theorem 4.2.1 in [12]) to reach a contradiction]. By a similar argument we reach a contradiction if we replace the \limsup with the \liminf in (64).

Proof of Part (ii).

From now on, for ease of notation, we denote by f the Fenchel-Legendre transform $\eta_{\lambda, Z}^*$ in (13). For technical reasons (related to the application of Dini's implicit function theorem), we assume f to be defined in a neighborhood of $\lambda \mathbb{E}[Z]$. The relations in (21) follow (reasoning again by contradiction as in the Part (i)) by Lemma 7.1 and Theorem 4.3.1 in [12]. As far as (22) is concerned, note that by the definition of the function θ and Dini's implicit function theorem we have that θ is infinitely differentiable on the domain of f and

$$\theta'_x = \frac{1}{\lambda \mathbb{E}[Z^2 e^{\theta_x Z}]}.$$

In particular, f is infinitely differentiable on its domain and, by Taylor's formula, in a neighborhood of $\lambda \mathbb{E}[Z]$, we have

$$f(x) = f(\lambda \mathbb{E}[Z]) + f'(\lambda \mathbb{E}[Z])(x - \lambda \mathbb{E}[Z]) + \frac{f''(\lambda \mathbb{E}[Z])}{2}(x - \lambda \mathbb{E}[Z])^2 + o((x - \lambda \mathbb{E}[Z])^2).$$

Note that $\theta_{\lambda \mathbb{E}[Z]} = 0$ and so $f(\lambda \mathbb{E}[Z]) = 0$. A straightforward computation yields

$$f'(x) = \theta_x + x\theta'_x - \lambda\theta'_x \mathbb{E}[Z e^{\theta_x Z}] = \theta_x \quad \text{and} \quad f''(x) = \theta'_x.$$

Therefore,

$$f'(\lambda \mathbb{E}[Z]) = \theta_{\lambda \mathbb{E}[Z]} = 0 \quad \text{and} \quad f''(\lambda \mathbb{E}[Z]) = \theta'_{\lambda \mathbb{E}[Z]} = \frac{1}{\lambda \mathbb{E}[Z^2]}.$$

So, as $x \rightarrow \lambda\mathbb{E}[Z]$, we have

$$f(x) = \frac{1}{2\lambda\mathbb{E}[Z^2]}(x - \lambda\mathbb{E}[Z])^2 + o((x - \lambda\mathbb{E}[Z])^2).$$

In particular, as $t \rightarrow \infty$, by the expression of $v(\cdot)$ we have

$$f(v(t)) = \frac{y(t)^2}{2t} + o\left(\frac{y(t)^2}{t}\right),$$

which gives the claim.

Proof of Part (iii).

Note that, in particular, $y(t) = o(t^{1/2})$, and therefore by the previous Part (ii) we have that the relations in (21) hold. In the previous Part (ii) we noticed that f is infinitely differentiable on its domain with $f''(\cdot) = \theta'$. Therefore, the k th derivative of f is given by $f^{(k)}(x) = \theta_x^{(k-1)}$, $k \geq 1$. By Taylor's formula, as $x \rightarrow \lambda\mathbb{E}[Z]$, we have

$$f(x) = \sum_{k=2}^m \frac{\theta_{\lambda\mathbb{E}[Z]}^{(k-1)}}{k!} (x - \lambda\mathbb{E}[Z])^k + o((x - \lambda\mathbb{E}[Z])^m).$$

In particular, as $t \rightarrow \infty$, by the expression of $v(\cdot)$ we have

$$tf(v(t)) = \sum_{k=2}^m (\lambda\mathbb{E}[Z^2])^{k/2} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(k-1)}}{k!} \frac{y(t)^k}{t^{k/2-1}} + t \cdot o\left(\frac{y(t)^m}{t^{m/2}}\right). \quad (65)$$

Note that

$$\lim_{t \rightarrow \infty} t \cdot o\left(\frac{y(t)^m}{t^{m/2}}\right) = \lim_{t \rightarrow \infty} \frac{o\left(\frac{y(t)^m}{t^{m/2}}\right)}{\frac{y(t)^m}{t^{m/2}}} \lim_{t \rightarrow \infty} \left(\frac{y(t)}{t^{\frac{1}{2} - \frac{1}{m}}}\right)^m = 0, \quad (66)$$

where we used the assumption on $y(\cdot)$. Moreover,

$$\begin{aligned} \sum_{k=2}^m (\lambda\mathbb{E}[Z^2])^{k/2} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(k-1)}}{k!} \frac{y(t)^k}{t^{k/2-1}} &= y(t)^2 \left(\frac{1}{2} + \sum_{k=3}^m (\lambda\mathbb{E}[Z^2])^{k/2} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(k-1)}}{k!} \frac{y(t)^{k-2}}{t^{k/2-1}} \right) \\ &= y(t)^2 \left(\frac{1}{2} + \sum_{j=1}^{m-2} (\lambda\mathbb{E}[Z^2])^{(j+2)/2} \frac{\theta_{\lambda\mathbb{E}[Z]}^{(j+1)}}{(j+2)!} \left(\frac{y(t)}{\sqrt{t}}\right)^j \right) \rightarrow \infty, \quad \text{as } t \rightarrow \infty. \end{aligned} \quad (67)$$

The claim follows combining (21) with (65), (66) and (67). Finally, the recursive formula for $\theta_{\lambda\mathbb{E}[Z]}^{(j)}$, $2 \leq j \leq m-1$, easily follows by the Faà di Bruno formula (see Lemma 2.3). The proof is completed.

7.5 Proof of Propositions 4.1 and 4.2

Proof of Proposition 4.1.

Proof of Parts (i) and (ii).

The claim follows by applying Theorems 3.2 and 3.4. To this aim, we show that, under the foregoing

assumptions, one has (11) (then, due to the fact that $Z := N_1([0, \infty))$ satisfies (6), one also has that condition (10) holds). We start noticing that, for $t \geq 0$, we have

$$\begin{aligned} \int_t^\infty (N_1([0, \infty)) - N_1([0, s])) ds &= \int_t^\infty (N_1([0, L]) - N_1([0, s])) ds \\ &= \mathbf{1}(L > t) \int_t^\infty (N_1([0, L]) - N_1([0, s])) \mathbf{1}(L > s) ds \\ &\leq (L - t)^+ N_1((t, L]), \end{aligned} \quad (68)$$

where $x^+ := \max\{x, 0\}$. For an arbitrary $q > 1$, set $\tilde{p}_1 := \frac{m}{q}$, $m > q$, $m \in \mathbb{N}$, and let $\tilde{p}_2 := m/(m - q)$. Note that $\tilde{p}_1, \tilde{p}_2 > 1$ and $(\tilde{p}_1)^{-1} + (\tilde{p}_2)^{-1} = 1$. Therefore by Hölder's inequality we have

$$\left\| \int_t^\infty (N_1([0, L]) - N_1([0, s])) ds \right\|_{L^q(\mathbb{P})} \leq \mathbb{E}[(L - t)^+]^{1/m} \mathbb{E}[N_1((t, L])^{q\tilde{p}_2}]^{1/(q\tilde{p}_2)}.$$

For $\theta^* \in (0, a)$ fixed, we have

$$\mathbb{E}[N_1((t, L])^{q\tilde{p}_2}]^{1/(q\tilde{p}_2)} \leq \mathbb{E}[(N_1([0, \infty))]^{q\tilde{p}_2}] \leq \frac{(\lceil q\tilde{p}_2 \rceil)!}{(\theta^*)^{\lceil q\tilde{p}_2 \rceil}} \mathbb{E}[e^{\theta^* N_1([0, \infty))}] =: \kappa_q^{(1)},$$

where $x \mapsto \lceil x \rceil$ denotes the ceiling function. So

$$\left\| \int_t^\infty (N_1([0, L]) - N_1([0, s])) ds \right\|_{L^q(\mathbb{P})} \leq \kappa_q^{(1)} \mathbb{E}[(L - t)^+]^{1/m}. \quad (69)$$

Let $\sigma \in \mathbb{N}$ be arbitrarily fixed and set $\theta := \sigma m + 1$. Since $(L - t)^+$ is a non-negative random variable, for any $t \geq 1$ by Markov's inequality we have

$$\begin{aligned} \mathbb{E}[(L - t)^+]^m &= m \int_0^\infty u^{m-1} \mathbb{P}((L - t)^+ > u) du \\ &= m \int_0^\infty u^{m-1} \mathbb{P}(L^{\theta m} > (u + t)^{\theta m}) du \\ &\leq m \mathbb{E}[L^{\theta m}] \int_0^\infty \frac{u^{m-1}}{(u + t)^{\theta m}} du \\ &= \mathbb{E}[L^{\theta m}] \int_0^\infty \frac{1}{(z^{1/m} + t)^{\theta m}} dz \\ &\leq \mathbb{E}[L^{\theta m}] \int_0^\infty \frac{1}{z^\theta + t^\theta} dz \\ &= \mathbb{E}[L^{\theta m}] t^{-(\theta-1)} \int_0^\infty \frac{1}{z^\theta + 1} dz \\ &\leq \mathbb{E}[L^{\theta m}] t^{-(\theta-1)} \left(1 + \int_1^\infty \frac{1}{z^2 + 1} dz \right) := \kappa_q^{(2)} t^{-(\theta-1)}, \end{aligned}$$

where the quantity which multiplies $t^{-(\theta-1)}$ depends on q since $m = m(q)$. So

$$\mathbb{E}[(L - t)^+]^{1/m} \leq (\kappa_q^{(2)})^{1/m} t^{-(\theta-1)/m} \leq (\kappa_q^{(2)})^{1/m} t^{-\sigma},$$

where we used the definition of θ and the fact that $t \geq 1$. Combining this latter inequality with (69), for any $\sigma \in \mathbb{N}$ fixed and $t \geq 1$, we have

$$\left\| \int_t^\infty (N_1([0, L]) - N_1([0, s])) ds \right\|_{L^q(\mathbb{P})} \leq \kappa_q t^{-\sigma},$$

where $\kappa_q := \kappa_q^{(1)}(\kappa_q^{(2)})^{1/m}$. Therefore, for any $\sigma \in \mathbb{N}$ fixed we have

$$\sup_{q>1} \kappa_q^{-1} \left\| \int_t^\infty (N_1([0, L]) - N_1([0, s])) ds \right\|_{L^q(\mathbb{P})} \leq t^{-\sigma}, \quad \text{for all } t \geq 1.$$

Thus condition (11) holds, and the proof of (i) and (ii) is completed.

Proof of Part (iii).

The claim easily follows by Proposition 3.3. Here, we limit ourselves to note that by (68) we have

$$\int_0^\infty (Z - N_1([0, s])) ds \leq LZ,$$

and so the upper bound in (30) and the lower bound in (31) follow, respectively, by the upper bound in the Part (i) of Proposition 3.3 and the lower bound in the Part (ii) of Proposition 3.3, and Hölder's inequality.

□

Proof of Proposition 4.2.

Proof of Parts (i) and (ii).

We are going to apply Proposition 4.1, and so we verify the assumptions therein. We already noticed that $Z := N_1([0, \infty))$ satisfies (6). So we only need to check (29). Let $\{U_n\}_{n \geq 1}$ be a sequence of independent random variables with the same law as $B_{1,1,1}$, independent of Z . Since V , defined by (35), has the same law as

$$\sum_{n=1}^{Z-1} U_n,$$

and V is an upper bound of L , we have

$$L \leq L' := \sum_{n=1}^Z U_n. \quad (70)$$

Such random variable L' has all the moments finite, indeed for any $k \in \mathbb{N}$, using Minkowski's inequality we have

$$\mathbb{E} \left[\left(\sum_{n=1}^Z U_n \right)^k \right] \leq \mathbb{E}[Z^k] \mathbb{E}[U_1^k] < \infty, \quad (71)$$

where the latter term is finite due to (36) and the fact that Z satisfies (6). The proof is completed.

Proof of Part (iii).

By the random walk perspective to branching processes (see e.g. [20] p. 90), we have that Z has the same law of the hitting time to zero of the random walk $\{Z'_n\}_{n \geq 0}$ defined by

$$Z'_0 := 1 \quad Z'_n := P_{1,1} + \dots + P_{1,n} - (n-1), \quad n \in \mathbb{N},$$

i.e.

$$Z \stackrel{d}{=} \inf\{n \in \mathbb{N} : Z'_n = 0\}$$

(we define the right-hand side equal to $+\infty$ if $\{\dots\} = \emptyset$). So

$$\mathbb{P}(Z > n) \leq \mathbb{P}(Z'_n > 0) = \mathbb{P} \left(\frac{1}{n} \sum_{k=1}^n P_{1,k} \geq 1 \right) \leq e^{-nac}, \quad n \in \mathbb{N}, \quad (72)$$

where the latter inequality follows by the Chernoff's bound (see e.g. [20] Theorem 2.19 p. 68, and recall that $\mathbb{E}[P_{1,1}] < 1$). Standard convexity arguments (combined with the subcritical assumption) guarantee $a_c \in (0, \infty)$. A straightforward computation shows that (72) implies $\mathbb{E}[e^{\theta Z}] < \infty \forall \theta < a_c$, which completes the proof.

Proof of Part (iv).

Let $\{Z_j\}_{j \geq 1}$ be independent copies of Z and let K_1 be the random variable defined at the beginning of Subsection 4.1. For any $\theta \in (-\infty, b_c)$, by standard computations we have

$$\begin{aligned} \mathbb{E}[e^{\theta Z}] &= e^\theta \sum_{k \geq 0} \mathbb{E}[e^{\theta \sum_{j=1}^k Z_j} | K_1 = k] p_k \\ &= e^\theta \sum_{k \geq 0} \mathbb{E}[e^{\theta Z}]^k p_k = e^\theta G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) < \infty. \end{aligned} \quad (73)$$

As far as the moments of Z are concerned, we combine the Faà di Bruno formula with the elementary relation:

$$\frac{d^n}{dx^n}(f(x)g(x)) = \sum_{k=0}^n \binom{n}{k} f^{(n-k)}(x) g^{(k)}(x), \quad n \in \mathbb{N}, \quad (74)$$

for sufficiently smooth functions f and g . For any $\theta \in (-\infty, b_c)$, by (73) and (74) we have

$$\frac{d^n}{d\theta^n} \mathbb{E}[e^{\theta Z}] = e^\theta \sum_{k=0}^n \binom{n}{k} \frac{d^k}{d\theta^k} G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]).$$

By the Faà di Bruno formula we have

$$\frac{d^k}{d\theta^k} G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) = k! \sum_{i=1}^k \frac{G_{P_{1,1}}^{(i)}(\mathbb{E}[e^{\theta Z}])}{i!} \sum_{m_1+m_2+\dots+m_i=k} \frac{\frac{d^{m_1}}{d\theta^{m_1}} \mathbb{E}[e^{\theta Z}]}{m_1!} \dots \frac{\frac{d^{m_i}}{d\theta^{m_i}} \mathbb{E}[e^{\theta Z}]}{m_i!}, \quad k \in \mathbb{N},$$

where the sum is taken over all the $m_1, \dots, m_i \in \mathbb{N}$ such that $m_1 + \dots + m_i = k$. The claim follows recalling that

$$\mathbb{E}[Z^n] = \frac{d^n}{d\theta^n} \mathbb{E}[e^{\theta Z}] \Big|_{\theta=0}$$

and that

$$G_{P_{1,1}}^{(i)}(1) = \mathbb{E}[P_{1,1}(P_{1,1}-1) \dots (P_{1,1}-(i-1))]$$

is the i th factorial moment of $P_{1,1}$.

Proof of Part (v).

Differentiating (73) with respect to θ , for any $\theta \in (-\infty, b_c)$ we have

$$\mathbb{E}[Z e^{\theta Z}] = e^\theta G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) + e^\theta \mathbb{E}[Z e^{\theta Z}] G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) = \mathbb{E}[e^{\theta Z}] + e^\theta \mathbb{E}[Z e^{\theta Z}] G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]),$$

where the latter equality follows by (73). Therefore

$$\mathbb{E}[Z e^{\theta Z}] = \frac{\mathbb{E}[e^{\theta Z}]}{1 - e^\theta G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}])}. \quad (75)$$

Differentiating again we have

$$\mathbb{E}[Z^2 e^{\theta Z}] = \mathbb{E}[e^{\theta Z}] \frac{1 + e^\theta (G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) + G''_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]))}{(1 - e^\theta G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]))^2}. \quad (76)$$

Recall that, for any $x \in (0, \lambda \mathbb{E}[Ze^{b_c Z}])$ we denote by $\theta_x \in (-\infty, b_c)$ the unique solution to $\lambda \mathbb{E}[Ze^{\theta_x Z}] = x$. Using (73), we rewrite the relation (75) as

$$\mathbb{E}[Ze^{\theta Z}] = \frac{\mathbb{E}[e^{\theta Z}]}{1 - \mathbb{E}[e^{\theta Z}] \frac{G'_{P_{1,1}}(\mathbb{E}[e^{\theta Z}])}{G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}])}}.$$

Taking $\theta = \theta_x$ in this relation, we have

$$\frac{x}{\lambda} = \frac{\mathbb{E}[e^{\theta_x Z}]}{1 - \mathbb{E}[e^{\theta_x Z}] \frac{G'_{P_{1,1}}(\mathbb{E}[e^{\theta_x Z}])}{G_{P_{1,1}}(\mathbb{E}[e^{\theta_x Z}])}},$$

and so $\varrho_x = \mathbb{E}[e^{\theta_x Z}]$ is solution to (38). The expressions of θ_x and $\mathbb{E}[Z^2 e^{\theta_x Z}]$ are readily obtained by (73) and (76), respectively. The proof is completed.

Proof of Part (vi).

For a fixed $r > 0$, let $\{N_r^{(j)}(\cdot)\}_{j \geq 1}$ be independent and identically distributed point processes on $[r, \infty)$ with the same branching structure as N_1 , but ancestor in r (when $r = 0$ they are independent copies of N_1). Hereon, we denote by $N_r^{(j)}(s)$ the number of points of $N_r^{(j)}$ on the interval $[r, s]$ and by Q the law of $B_{1,1,1}$. For any $\theta \in (-\infty, b_c)$ and $s > 0$, similarly to (73), we have

$$\begin{aligned} \mathbb{E}[e^{\theta N_1([0,s])}] &= \sum_{k \geq 0} \mathbb{E}[e^{\theta N_1([0,s])} \mid K_1 = k] p_k \\ &= \sum_{k \geq 0} \mathbb{E} \left[e^{\theta \left(1 + \sum_{j=1}^k N_{B_{1,0,j}}^{(j)}(s) \right)} \mid K_1 = k \right] p_k \\ &= e^\theta \sum_{k \geq 0} p_k \left(\int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr) \right)^k \\ &= e^\theta G_{P_{1,1}} \left(\int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr) \right). \end{aligned}$$

By this relation, (73) and the mean value theorem, we have (since $N_r^{(1)}(s) = 0$ for $r > s$)

$$\begin{aligned} \mathbb{E}[e^{\theta N_1([0,s])}] - \mathbb{E}[e^{\theta Z}] &= e^\theta \left(G_{P_{1,1}} \left(\int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr) \right) - G_{P_{1,1}}(\mathbb{E}[e^{\theta Z}]) \right) \\ &= e^\theta \left(\int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr) - \mathbb{E}[e^{\theta Z}] \right) G'_{P_{1,1}}(\xi) \\ &= e^\theta \left(\int_{[0,s]} (\mathbb{E}[e^{\theta N_r^{(1)}(s)}] - \mathbb{E}[e^{\theta Z}]) Q(dr) + (1 - \mathbb{E}[e^{\theta Z}]) Q((s, \infty)) \right) G'_{P_{1,1}}(\xi), \end{aligned}$$

for some

$$\xi \in \left(\min \left\{ \int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr), \mathbb{E}[e^{\theta Z}] \right\}, \max \left\{ \int_{[0,\infty)} \mathbb{E}[e^{\theta N_r^{(1)}(s)}] Q(dr), \mathbb{E}[e^{\theta Z}] \right\} \right).$$

Since $\xi \geq 0$, a simple computation shows that $G'_{P_{1,1}}(\xi) \geq p_1$. Therefore

$$\begin{aligned} \mathbb{E}[e^{\theta N_1([0,s])}] - \mathbb{E}[e^{\theta Z}] &\leq e^\theta \left(\int_{[0,s]} (\mathbb{E}[e^{\theta N_r^{(1)}(s)}] - \mathbb{E}[e^{\theta Z}]) Q(dr) + (1 - \mathbb{E}[e^{\theta Z}]) Q((s, \infty)) \right) p_1 \\ &\leq p_1 e^\theta (1 - \mathbb{E}[e^{\theta Z}]) Q((s, \infty)), \quad \text{for any } \theta \in (0, b_c), \end{aligned}$$

and

$$\mathbb{E}[e^{\theta N_1([0,s])}] - \mathbb{E}[e^{\theta Z}] \geq p_1 e^\theta (1 - \mathbb{E}[e^{\theta Z}]) Q((s, \infty)), \quad \text{for any } \theta < 0.$$

The claim follows combining these inequalities with (30), (31), (70) and (71).

□

7.6 Proof of Proposition 5.1

We only prove Part (i). Indeed, mutatis mutandis (i.e. applying Theorem 3.2(ii) in place of Theorem 3.1) the proof of Part (ii) is similar. We start noticing that the upper bound is a simple consequence of the inequality $\psi_{IBNR}(u) \leq \psi_{CL}(u)$, $u > 0$, and (46). As far as the lower bound is concerned, note that, for any $u, d > 0$, we have

$$\psi_{IBNR}(u) \geq \mathbb{P}(S_{ud} - cud \geq u) = \mathbb{P}\left(\frac{S_{ud}}{ud} \geq c + \frac{1}{d}\right). \quad (77)$$

By the strict convexity of the function

$$(0, a) \ni \gamma \mapsto \lambda(\mathbb{E}[e^{\gamma Z}] - 1) - c\gamma,$$

we have $\lambda\mathbb{E}[Ze^{wZ}] - c > 0$. Set $d := (\lambda\mathbb{E}[Ze^{wZ}] - c)^{-1} > 0$ and $x := c + d^{-1} = \lambda\mathbb{E}[Ze^{wZ}]$. By the “net profit” condition and the fact that $\mathbb{E}[Ze^{aZ}] > \mathbb{E}[Ze^{wZ}] > 0$, we have $x \in (\lambda\mathbb{E}[Z], \lambda\mathbb{E}[Ze^{aZ}])$. Moreover, a straightforward computation gives

$$\theta_x = w \quad \text{and} \quad x\theta_x - \lambda(\mathbb{E}[e^{\theta_x Z}] - 1) = w(\lambda\mathbb{E}[Ze^{wZ}] - c).$$

The lower bound follows by these relations, (77) and Theorem 3.1.

7.7 Proof of Theorems 6.1 and 6.2

In this section we prove Theorems 6.1 and 6.2. In particular, we emphasize that the proof of Theorem 6.2 exploits the ideas and the techniques developed in [13].

Proof of Theorem 6.1. We divide the proof in two steps. In the first step we prove the inequality

$$|\psi_t(\mathbf{i}\xi) - 1| \leq (H_1(t)|\xi|^\alpha + H_2(t)|\xi|^{2\alpha}) \exp(H_1(t)|\xi|^\alpha + H_2(t)|\xi|^{2\alpha}), \quad \forall (t, \xi) \in (0, \infty) \times \mathbb{R}, \quad (78)$$

where

$$\psi_t(\mathbf{i}\xi) := \mathbb{E}[e^{\mathbf{i}\xi(S_t/t^{1/(2\alpha)})}] e^{-\sqrt{t}\eta_{c,\lambda^{1/\alpha},\alpha,\beta}(\mathbf{i}\xi)}, \quad t > 0, \xi \in \mathbb{R}, \quad (79)$$

$$H_1(t) := \lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{\int_0^t (1 - F(s)^\alpha) ds}{\sqrt{t}}, \quad t > 0, \quad (80)$$

and

$$H_2(t) := \frac{\lambda c^{2\alpha}}{2} [1 + (\beta \tan(\pi\alpha/2))^2], \quad t > 0. \quad (81)$$

In the second step we conclude the proof.

Step 1: Proof of (78).

For any $t > 0$ and $\xi \in \mathbb{R}$, by Lemma 2.2 and the expression of the Laplace functional of a Poisson process (see e.g. [8]), we have

$$\mathbb{E}[e^{\mathbf{i}\xi S_t/t^{1/\alpha}}] = \exp\left(\lambda \int_0^t (e^{\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s)/t^{1/\alpha})} - 1) ds\right) = \exp\left(\lambda \int_0^t (e^{\frac{1}{t}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - 1) ds\right). \quad (82)$$

Again, by the scaling property of the Lévy exponent $\eta_{c,\alpha,\beta}(\cdot)$ and its definition, we have

$$\eta_{c,\alpha,\beta}\left(\mathbf{i}\xi t^{\frac{1}{2\alpha}}\right) = \eta_{c,\alpha,\beta}\left(\frac{\mathbf{i}\xi}{(t^{-1/2})^{1/\alpha}}\right) = \sqrt{t}\eta_{c,\alpha,\beta}(\mathbf{i}\xi), \quad t > 0, \xi \in \mathbb{R}, \quad (83)$$

and $\eta_{c\lambda^{1/\alpha},\alpha,\beta}(\mathbf{i}\xi) = \lambda\eta_{c,\alpha,\beta}(\mathbf{i}\xi)$, $\xi \in \mathbb{R}$. By these relations and (82), for any $t > 0$ and $\xi \in \mathbb{R}$, it follows

$$\begin{aligned} \psi_t(\mathbf{i}\xi) &= \exp\left(\lambda \int_0^t \left(e^{\frac{1}{t}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - 1\right) ds - \lambda\sqrt{t}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)\right) \\ &= \exp\left(\lambda \int_0^t \left(e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - 1 - \frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)\right) ds\right) \\ &= \exp\left(\lambda \int_0^t \left(e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}\right) ds + \lambda t \left(e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)} - 1 - \frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)\right)\right). \end{aligned} \quad (84)$$

By Lemma 2.2 and the definition of the Lévy exponent $\eta_{c,\alpha,\beta}(\cdot)$, for any $t > 0$, $s \in (0, t)$ and $\xi \in \mathbb{R}$, we have

$$\begin{aligned} \left|e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}\right| &\leq \frac{1}{\sqrt{t}} \left|\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s)) - \eta_{c,\alpha,\beta}(\mathbf{i}\xi)\right| e^{\frac{1}{\sqrt{t}} \max\{\operatorname{Re}(\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))), \operatorname{Re}(\eta_{c,\alpha,\beta}(\mathbf{i}\xi))\}} \\ &= \frac{1}{\sqrt{t}} |\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s)) - \eta_{c,\alpha,\beta}(\mathbf{i}\xi)| e^{-\frac{1}{\sqrt{t}}|c\xi F(s)|^\alpha} \\ &= \frac{1}{\sqrt{t}} |1 - \mathbf{i}\tan(\pi\alpha/2)\operatorname{sgn}(\xi)| |c\xi|^\alpha - |c\xi F(s)|^\alpha e^{-\frac{1}{\sqrt{t}}|c\xi F(s)|^\alpha} \\ &= c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{(1 - F(s)^\alpha) e^{-\frac{1}{\sqrt{t}}|c\xi F(s)|^\alpha}}{\sqrt{t}} |\xi|^\alpha. \end{aligned}$$

Therefore, for any $t > 0$ and $\xi \in \mathbb{R}$,

$$\left|\lambda \int_0^t \left(e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi F(s))} - e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}\right) ds\right| \leq H_1(t) |\xi|^\alpha. \quad (85)$$

Applying Taylor's formula with integral remainder to the function

$$u \mapsto e^{u \frac{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}{\sqrt{t}}}, \quad u \in [0, 1],$$

for any $t > 0$ and $\xi \in \mathbb{R}$, we have

$$e^{\frac{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}{\sqrt{t}}} - 1 = \frac{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}{\sqrt{t}} + \frac{(\eta_{c,\alpha,\beta}(\mathbf{i}\xi))^2}{t} \int_0^1 (1-u) e^{u \frac{\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}{\sqrt{t}}} du.$$

Therefore, for any $t > 0$ and $\xi \in \mathbb{R}$,

$$\left|\lambda t \left(e^{\frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)} - 1 - \frac{1}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)\right)\right| \leq \frac{\lambda}{2} |\eta_{c,\alpha,\beta}(\mathbf{i}\xi)|^2 = H_2(t) |\xi|^{2\alpha}, \quad (86)$$

where $H_2(\cdot)$ is the constant function defined in (81). Note that in (86) we used the relation

$$\left|e^{\frac{u}{\sqrt{t}}\eta_{c,\alpha,\beta}(\mathbf{i}\xi)}\right| = \left|e^{\eta_{c,\alpha,\beta}\left(\frac{\mathbf{i}\xi}{\left(\frac{\sqrt{t}}{u}\right)^{1/\alpha}}\right)}\right| = e^{-\frac{u}{\sqrt{t}}|c\xi|^\alpha} \leq 1, \quad u \in [0, 1],$$

which follows by (4) and Lemma 2.2. By Lemma 2.2 and $\max\{\operatorname{Re} z, 0\} \leq |z|$, $z \in \mathbb{C}$, we have

$$|e^z - 1| \leq |z|e^{|z|}, \quad z \in \mathbb{C}.$$

The claim follows combining this elementary inequality with (84), (85) and (86).

Step 2: Conclusion of the proof.

By (78), for any $t > 0$ and $\xi \in \mathbb{R}$, we have

$$\begin{aligned} \left| \psi_t \left(\mathbf{i} \frac{\xi}{t^{1/(2\alpha)}} \right) - 1 \right| &\leq \left(\lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t} |\xi|^\alpha + \frac{\lambda c^{2\alpha}}{2t} [1 + (\beta \tan(\pi\alpha/2))^2] |\xi|^{2\alpha} \right) \\ &\quad \times \exp \left(\lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t} |\xi|^\alpha + \frac{\lambda c^{2\alpha}}{2t} [1 + (\beta \tan(\pi\alpha/2))^2] |\xi|^{2\alpha} \right). \end{aligned} \quad (87)$$

The right-hand side of this latter inequality goes to zero as $t \rightarrow \infty$, indeed by de l'Hopital's theorem and the fact that $F(\cdot)$ is a distribution function, we have

$$\lim_{t \rightarrow \infty} \frac{1}{t} \int_0^t (1 - F(s)^\alpha) ds = 0.$$

The claim then follows by Lévy's continuity theorem noticing that, for any $t > 0$ and $\xi \in \mathbb{R}$,

$$\psi_t \left(\mathbf{i} \frac{\xi}{t^{1/(2\alpha)}} \right) = \frac{\mathbb{E}[e^{\mathbf{i}(\xi/t^{1/(2\alpha)})t^{-1/(2\alpha)}S_t}]}{e^{\sqrt{t}\eta_{c\lambda^{1/\alpha}, \alpha, \beta}(\mathbf{i}(\xi/t^{1/(2\alpha)}))}}} = \frac{\mathbb{E}[e^{\mathbf{i}\xi(S_t/t^{1/\alpha})}]}{e^{\eta_{c\lambda^{1/\alpha}, \alpha, \beta}(\mathbf{i}\xi)}}, \quad (88)$$

where the latter equality follows by (83).

□

Proof of Theorem 6.2. By (78), for any $t > 0$ and $\xi \in \mathbb{R}$, we have (87), which, for $\eta \in (0, 1]$, we rewrite as

$$\left| \psi_t \left(\mathbf{i} \frac{\xi}{t^{1/(2\alpha)}} \right) - 1 \right| \leq \left(\frac{H_{1,\eta}(t)}{t^{1-\eta}} |\xi|^\alpha + \frac{H_2}{t} |\xi|^{2\alpha} \right) \exp \left(\frac{H_{1,\eta}(t)}{t^{1-\eta}} |\xi|^\alpha + \frac{H_2}{t} |\xi|^{2\alpha} \right), \quad (89)$$

where

$$H_{1,\eta}(t) := \lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t^\eta}, \quad t > 0,$$

and $H_2(\cdot) \equiv H_2$ is defined by (81); note that $H_{1,1/2}(\cdot) \equiv H_1(\cdot)$, where $H_1(\cdot)$ is defined by (80). Letting \mathbb{P}_X denote the law of a random variable X , we consider the signed measure

$$\mu_t(dx) := \mathbb{P}_{t^{-1/\alpha}S_t}(dx) - \mathbb{P}_S(dx), \quad t > 0,$$

and note that

$$\widehat{\mu}_t(\xi) := \mathbb{E}[e^{\mathbf{i}\xi(S_t/t^{1/\alpha})}] - \mathbb{E}[e^{\mathbf{i}\xi S}] = \mathbb{E}[e^{\mathbf{i}\xi S}] \left(\psi_t \left(\mathbf{i} \frac{\xi}{t^{1/(2\alpha)}} \right) - 1 \right), \quad t > 0, \xi \in \mathbb{R},$$

where the latter equality follows by (88). Therefore, by (89) and (4), for any $t > 0$ and $\xi \in \mathbb{R} \setminus \{0\}$, we have

$$\begin{aligned} \left| \frac{\widehat{\mu}_t(\xi)}{\xi} \right| &= e^{-\lambda|c\xi|^\alpha} \frac{1}{|\xi|} \left| \psi_t \left(\mathbf{i} \frac{\xi}{t^{1/(2\alpha)}} \right) - 1 \right| \\ &\leq \left(\frac{H_{1,\eta}(t)}{t^{1-\eta}} |\xi|^{\alpha-1} + \frac{H_2}{t} |\xi|^{2\alpha-1} \right) \exp \left(-\lambda|c\xi|^\alpha + \frac{H_{1,\eta}(t)}{t^{1-\eta}} |\xi|^\alpha + \frac{H_2}{t} |\xi|^{2\alpha} \right). \end{aligned} \quad (90)$$

Let $\rho(\cdot)$ be the kernel function provided by Lemma 2.13 in [13], and for $\varepsilon > 0$ and $x, a \in \mathbb{R}$, put $\rho_\varepsilon(x) := \varepsilon^{-1}\rho(x/\varepsilon)$, $f_\varepsilon(x) := \mathbf{1}_{(-\infty, 0]} * \rho_\varepsilon(x)$, and $f_{a,\varepsilon}(x) := f_\varepsilon(x - a)$. Here the symbol $*$ denotes the convolution product. We continue by first proving the Part (i) of the theorem, and then the Part (ii). We present a proof which is a little bit more technical than the necessary since we provide a constant involved in the big O notation.

Proof of Part (i).

We only prove the claim under the assumption (53). Indeed, if $\eta_0 \notin A$, then $\eta_0 + \varepsilon \in A$ for any $\varepsilon > 0$ sufficiently small. So let $\eta_0 \in A$ and let $t^* > 0$ be such that

$$\sup_{t \geq t^*} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t^{\eta_0}} \in [0, \infty).$$

For an arbitrarily fixed

$$\gamma \in \left(-\frac{\eta_0}{\alpha(1-\eta_0)}, 1 - \frac{\eta_0}{\alpha(1-\eta_0)} \right], \quad (91)$$

we define

$$\varepsilon(t) := \frac{1}{K_0 t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}}, \quad t > 0, \quad K_0 := \left(\frac{\lambda}{4H_2} \right)^{\frac{1}{\alpha}} c, \quad (92)$$

and consider the family of functions $\{f_{a,\varepsilon(t)}\}_{t>0}$. By Proposition 2.14 in [13], we have that the functions $f_{a,\varepsilon(t)}$ are smooth test distributions in $\mathcal{T}_1(\mathbb{R})$ (we refer the reader to Definition 2.9 on p. 11 of [13] for the rigorous definition of such space) with Fourier transform compactly supported on the interval

$$I_t := [-K_0 t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}, K_0 t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}] \quad (93)$$

and $\|\partial f_{a,\varepsilon(t)}\|_{L^1(\mathbb{R}, dx)} = 1$. Note that

$$\alpha\gamma - 1 \leq 0. \quad (94)$$

Indeed, by (91) and the fact that $\alpha \leq 1/(1-\eta_0)$ (since $\eta_0 \geq (\alpha-1)/\alpha$), we have

$$\alpha\gamma - 1 \leq \alpha - \frac{\eta_0}{1-\eta_0} - 1 = \alpha - \frac{1}{1-\eta_0} \leq 0.$$

For $t \geq 1$ and $\xi \in I_t$, we have

$$\begin{aligned} \frac{|\xi|^{2\alpha}}{t} &= |\xi|^\alpha \frac{|\xi|^\alpha}{t} \leq K_0^\alpha t^{(1-\eta_0)(\alpha\gamma + \frac{\eta_0}{1-\eta_0})-1} |\xi|^\alpha \\ &= K_0^\alpha t^{(1-\eta_0)(\alpha\gamma-1)} |\xi|^\alpha \leq \frac{\lambda}{4H_2} |c\xi|^\alpha, \end{aligned} \quad (95)$$

where for the latter inequality we used that $t^{(1-\eta_0)(\gamma\alpha-1)} \leq 1$ since $t \geq 1$ and $\gamma\alpha - 1 \leq 0$ by (94). Let $t' \geq 1$ be such that

$$\frac{\int_0^t (1 - F(s)^\alpha) ds}{t} \leq \frac{\lambda c^\alpha}{4\sqrt{2\lambda H_2}}, \quad \text{for any } t \geq t'. \quad (96)$$

Note that such a t' exists since the left-hand side of (96) tends to zero as $t \rightarrow \infty$ by de l'Hopital's theorem. By (96) we have

$$\frac{H_{1,\eta_0}(t)}{t^{1-\eta_0}} \leq \lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \frac{\lambda c^\alpha}{4\sqrt{2\lambda H_2}} = \frac{\lambda c^\alpha}{4}, \quad \text{for any } t \geq t'. \quad (97)$$

By (90), (95) and (97), for any $t \geq t'' := \max\{t', t^*\}$ and $\xi \in I_t \setminus \{0\}$, we have

$$\left| \frac{\widehat{\mu}_t(\xi)}{\xi} \right| \leq \left(K'_1 \frac{|\xi|^{\alpha-1}}{t^{1-\eta_0}} + H_2 \frac{|\xi|^{2\alpha-1}}{t} \right) \exp \left(-\frac{\lambda|c\xi|^\alpha}{2} \right), \quad (98)$$

where

$$K'_1 := \lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \sup_{t \geq t^*} \frac{\int_0^t (1 - F(s)^\alpha) ds}{t^{\eta_0}}.$$

We note that, for any $\nu, \kappa, \rho > 0$,

$$\int_{\mathbb{R}} |\xi|^{\nu-1} e^{-\kappa|\xi|^\rho} d\xi = 2 \int_0^\infty \xi^{\nu-1} e^{-\kappa\xi^\rho} d\xi = 2 \int_0^\infty x^{(\nu-1)/\rho} e^{-\kappa x} dx^{1/\rho} = \frac{2\Gamma\left(\frac{\nu}{\rho}\right)}{\rho\kappa^{\nu/\rho}}, \quad (99)$$

where $\Gamma(\cdot)$ is the Euler gamma function. In particular, for any $t \geq t''$, the function $\xi \mapsto \frac{\widehat{\mu}_t(\xi)}{\xi}$ is integrable. Therefore by Remark 2.11 in [13] we have that the extended Parseval formula applies and, letting $\widehat{f}(\xi) := \int_{\mathbb{R}} e^{i\xi x} f(x) dx$ denote the Fourier transform of $f \in L^1(\mathbb{R}, dx)$, we have

$$\begin{aligned} |\mathbb{E}[f_{a,\varepsilon(t)}(S_t/t^{1/\alpha})] - \mathbb{E}[f_{a,\varepsilon(t)}(S)]| &= \frac{1}{2\pi} \left| \int_{I_t} \widehat{\partial f_{a,\varepsilon(t)}}(\xi) \frac{\widehat{\mu}_t(-\xi)}{\xi} d\xi \right| \\ &\leq \frac{\|\partial f_{a,\varepsilon(t)}\|_{L^1(\mathbb{R}, dx)}}{2\pi} \int_{I_t} \left| \frac{\widehat{\mu}_t(-\xi)}{\xi} \right| d\xi \end{aligned} \quad (100)$$

$$= \frac{1}{2\pi} \int_{I_t} \left| \frac{\widehat{\mu}_t(-\xi)}{\xi} \right| d\xi, \quad \text{for any } t \geq t'', \quad (101)$$

where in (100) we used the elementary inequality $|\widehat{f}(\xi)| \leq \|f\|_{L^1(\mathbb{R}, dx)}$, for any $\xi \in \mathbb{R}$. By (98) and (99), for any $t \geq t''$, we have

$$\begin{aligned} \int_{I_t} \left| \frac{\widehat{\mu}_t(-\xi)}{\xi} \right| d\xi &\leq \frac{K'_1}{t^{1-\eta_0}} \int_{I_t} |\xi|^{\alpha-1} e^{-\frac{\lambda|c|^\alpha}{2}|\xi|^\alpha} d\xi + \frac{H_2}{t} \int_{I_t} |\xi|^{2\alpha-1} e^{-\frac{\lambda|c|^\alpha}{2}|\xi|^\alpha} d\xi \\ &\leq \frac{K'_1}{t^{1-\eta_0}} \frac{4}{\lambda\alpha c^\alpha} + \frac{H_2}{t} \frac{8}{\alpha\lambda^2 c^{2\alpha}} \leq \frac{4}{\lambda\alpha c^\alpha} \left(K'_1 + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t^{1-\eta_0}}, \end{aligned}$$

where the latter inequality follows noticing that $t'' \geq 1$. Combining this with (101), for any $t \geq t''$, we have

$$\begin{aligned} |\mathbb{E}[f_{a,\varepsilon(t)}(S_t/t^{1/\alpha})] - \mathbb{E}[f_{a,\varepsilon(t)}(S)]| &\leq \frac{2}{\pi\lambda\alpha c^\alpha} \left(K'_1 + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t^{1-\eta_0}} \\ &\leq \frac{2}{\pi\lambda\alpha c^\alpha} \left(K'_1 + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}} = B\varepsilon(t), \end{aligned} \quad (102)$$

where

$$B := \frac{2K_0}{\pi\lambda\alpha c^\alpha} \left(K'_1 + \frac{2H_2}{\lambda c^\alpha} \right)$$

and the inequality (102) is a consequence of $\gamma \leq 1 - \frac{\eta_0}{\alpha(1-\eta_0)}$ and $t'' \geq 1$. By Theorem 2.15 in [13], noticing that S has a density with respect to the Lebesgue measure bounded above by $\frac{1}{\alpha\pi c\lambda^{1/\alpha}}\Gamma\left(\frac{1}{\alpha}\right)$ (see Subsection 2.3), we have

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) \leq \frac{C}{t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}}, \quad \text{for any } t \geq t'',$$

where

$$C := \inf_{\zeta > 0} \left\{ (1 + \zeta) \left(\frac{2}{\pi \lambda \alpha c^\alpha} \left(K_1' + \frac{2H_2}{\lambda c^\alpha} \right) + \frac{(4H_2)^{\frac{1}{\alpha}} \Gamma\left(\frac{1}{\alpha}\right)}{\pi^{4/3} \alpha c^2 \lambda^{2/\alpha}} (4(1 + \zeta^{-1})^{1/3} + 3^{4/3}) \right) \right\}. \quad (103)$$

The claim follows noticing that

$$\inf_{-\frac{\eta_0}{\alpha(1-\eta_0)} < \gamma \leq 1 - \frac{\eta_0}{\alpha(1-\eta_0)}} \frac{1}{t^{(1-\eta_0)(\gamma + \frac{\eta_0}{\alpha(1-\eta_0)})}} = \frac{1}{t^{1-\eta_0}}, \quad t \geq t''.$$

Proof of Part (ii).

We take

$$\gamma \in \left(-\frac{1}{\alpha}, \min \left\{ 2 - \frac{1}{\alpha}, \frac{1}{\alpha} \right\} \right] \quad (104)$$

and define $\varepsilon(t)$ and I_t as in (92) and (93) with $1/2$ in place of η_0 , i.e.,

$$\varepsilon(t) := \frac{1}{K_0 t^{\frac{1}{2}(\gamma + \frac{1}{\alpha})}}, \quad t > 0, \quad (105)$$

and $I_t = [-\varepsilon(t)^{-1}, \varepsilon(t)^{-1}]$. Then we consider the family of functions $\{f_{a,\varepsilon(t)}\}_{t>0}$ (the function $f_{a,\varepsilon}$ is defined at the beginning of the proof) which, by Proposition 2.14 in [13], are smooth test distributions in $\mathcal{T}_1(\mathbb{R})$ with Fourier transform compactly supported on I_t . Note that the relations (95) and (97) hold with $1/2$ in place of η_0 . By the assumption (54), it follows

$$\frac{H_1(t)}{\sqrt{t}} \leq \frac{K_1''}{t}, \quad \text{for any } t > 0,$$

where $H_1(\cdot)$ is given by (80) and

$$K_1'' := \lambda c^\alpha \sqrt{1 + (\beta \tan(\pi\alpha/2))^2} \int_0^\infty (1 - F(s)^\alpha) ds \in [0, \infty).$$

Combining this with (90), (95) and (97) (again with $1/2$ in place of η_0), for any $t \geq t'$ and $\xi \in I_t \setminus \{0\}$, we have

$$\left| \frac{\widehat{\mu}_t(\xi)}{\xi} \right| \leq \left(K_1'' \frac{|\xi|^{\alpha-1}}{t} + H_2 \frac{|\xi|^{2\alpha-1}}{t} \right) \exp \left(-\frac{\lambda |c\xi|^\alpha}{2} \right). \quad (106)$$

As in the proof of Part (i), we have that, for any $t \geq t'$, the function $\xi \mapsto \frac{\widehat{\mu}_t(\xi)}{\xi}$ is integrable, and one can apply the extended Parseval formula to get (101), with $\varepsilon(t)$ defined by (105) and t' in place of t'' . By (106) and (99), for any $t \geq t'$, we have

$$\begin{aligned} \int_{I_t} \left| \frac{\widehat{\mu}_t(-\xi)}{\xi} \right| d\xi &\leq \frac{K_1''}{t} \int_{I_t} |\xi|^{\alpha-1} e^{-\frac{\lambda |c\xi|^\alpha}{2}} d\xi + \frac{H_2}{t} \int_{I_t} |\xi|^{2\alpha-1} e^{-\frac{\lambda |c\xi|^\alpha}{2}} d\xi \\ &\leq \frac{K_1''}{t} \frac{4}{\alpha \lambda c^\alpha} + \frac{H_2}{t} \frac{8}{\alpha \lambda^2 c^{2\alpha}} = \frac{4}{\lambda \alpha c^\alpha} \left(K_1'' + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t}. \end{aligned}$$

Combining this with (101) (with $\varepsilon(t)$ given by (105) and t' in place of t''), for any $t \geq t'$, we have

$$\begin{aligned} |\mathbb{E}[f_{a,\varepsilon(t)}(S_t/t^{1/\alpha})] - \mathbb{E}[f_{a,\varepsilon(t)}(S)]| &\leq \frac{2}{\pi \lambda \alpha c^\alpha} \left(K_1'' + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t} \\ &\leq \frac{2}{\pi \lambda \alpha c^\alpha} \left(K_1'' + \frac{2H_2}{\lambda c^\alpha} \right) \frac{1}{t^{\frac{1}{2}(\gamma + \frac{1}{\alpha})}} = B' \varepsilon(t), \end{aligned} \quad (107)$$

where

$$B' := \frac{2K_0}{\pi\lambda\alpha c^\alpha} \left(K_1'' + \frac{2H_2}{\lambda c^\alpha} \right)$$

and the inequality (107) is a consequence of $\gamma \leq 2 - \frac{1}{\alpha}$ and $t' \geq 1$. By Theorem 2.15 in [13], noticing that S has a density (with respect to the Lebesgue measure) bounded above by $\frac{1}{\alpha\pi c\lambda^{1/\alpha}}\Gamma\left(\frac{1}{\alpha}\right)$, we have

$$d_{\text{Kol}}(S_t/t^{1/\alpha}, S) \leq \frac{C'}{t^{\frac{1}{2}(\gamma+\frac{1}{\alpha})}}, \quad \text{for any } t \geq t',$$

where the constant C' is defined as C in (103), but with K_1'' in place of K_1' . The claim follows taking the infimum over γ , which satisfies the constraint (104). Indeed, if $\alpha \in (1, 2]$, then $1/\alpha < 2 - 1/\alpha$, and so

$$\inf_{-1/\alpha < \gamma \leq \min\{1/\alpha, 2-1/\alpha\}} \frac{1}{t^{\frac{1}{2}(\gamma+\frac{1}{\alpha})}} = t^{-\frac{1}{\alpha}}, \quad t \geq t';$$

if, instead, $\alpha \in (0, 1]$, then $1/\alpha \geq 2 - 1/\alpha$, and so

$$\inf_{-1/\alpha < \gamma \leq \min\{1/\alpha, 2-1/\alpha\}} \frac{1}{t^{\frac{1}{2}(\gamma+\frac{1}{\alpha})}} = t^{-1}, \quad t \geq t'.$$

□

7.8 Proof of Lemma 2.2

For ease of notation, let $a_i := \text{Re} z_i$ and $b_i := \text{Im} z_i$, $i = 1, 2$. Hereon, without loss of generality, we assume $a_1 \geq a_2$. We have:

$$|e^{z_1} - e^{z_2}| = |e^{a_1+ib_1} - e^{a_2+ib_1} + e^{a_2+ib_1} - e^{a_2+ib_2}| = |e^{a_1} - e^{a_2} + e^{a_2}(1 - e^{i(b_2-b_1)})|.$$

By the mean value theorem we have $e^{a_1} - e^{a_2} = (a_1 - a_2)e^c$, for some $c = c(a_1, a_2) \in [a_2, a_1]$. Therefore

$$|e^{z_1} - e^{z_2}| = |(a_1 - a_2)e^c + e^{a_2}(1 - e^{i(b_2-b_1)})| \leq |(a_1 - a_2)e^{a_1} + e^{a_2}(1 - e^{i(b_2-b_1)})|.$$

Setting $\alpha := a_1 - a_2 \geq 0$ and $\beta := b_2 - b_1$, we have

$$|(a_1 - a_2)e^{a_1} + e^{a_2}(1 - e^{i(b_2-b_1)})| = e^{a_1}|\alpha + e^{-\alpha}(1 - e^{i\beta})| = e^{a_1}|\alpha + e^{-\alpha}(1 - \cos \beta - i \sin \beta)|.$$

The claim follows if we prove the inequality

$$|\alpha + e^{-\alpha}(1 - \cos \beta) - i e^{-\alpha} \sin \beta| \leq \sqrt{\alpha^2 + \beta^2}. \quad (108)$$

Indeed $|z_1 - z_2| = \sqrt{\alpha^2 + \beta^2}$ and

$$|\alpha + e^{-\alpha}(1 - \cos \beta - i \sin \beta)| = |\alpha + e^{-\alpha}(1 - \cos \beta) - i e^{-\alpha} \sin \beta|.$$

Note that (108) is equivalent to

$$f_1(\alpha, \beta) := [\alpha + e^{-\alpha}(1 - \cos \beta)]^2 + e^{-2\alpha}(\sin \beta)^2 \leq \alpha^2 + \beta^2 =: f_2(\alpha, \beta). \quad (109)$$

To prove this latter inequality, we start noticing that $f_1(\alpha, 0) = f_2(\alpha, 0) = \alpha^2$ and

$$f_1(\alpha, \beta) = \alpha^2 + 2\alpha e^{-\alpha}(1 - \cos \beta) + 2e^{-2\alpha}(1 - \cos \beta) = \alpha^2 + 2(\alpha + e^{-\alpha})e^{-\alpha}(1 - \cos \beta).$$

Then we distinguish two cases: $\beta > 0$ and $\beta < 0$. Assume first $\beta > 0$. For any $\gamma > 0$, we have

$$\frac{\partial f_1(\alpha, \gamma)}{\partial \gamma} = 2(\alpha + e^{-\alpha})e^{-\alpha} \sin \gamma \leq 2\gamma = \frac{\partial f_2(\alpha, \gamma)}{\partial \gamma},$$

where we used the elementary inequalities $\sin \gamma \leq |\sin \gamma| \leq |\gamma| = \gamma$ and $0 < (\alpha + e^{-\alpha})e^{-\alpha} \leq 1$ (the latter follows by the elementary relations $e^\alpha \geq 1 + \alpha \geq e^{-\alpha} + \alpha$; recall that $\alpha \geq 0$). Therefore

$$f_1(\alpha, \beta) = f_1(\alpha, 0) + \int_0^\beta \frac{\partial f_1(\alpha, \gamma)}{\partial \gamma} d\gamma \leq f_2(\alpha, 0) + \int_0^\beta \frac{\partial f_2(\alpha, \gamma)}{\partial \gamma} d\gamma = f_2(\alpha, \beta), \quad (110)$$

and the claim (109) for $\beta > 0$ is proved. Now, assume $\beta < 0$. By (110) we have $f_1(\alpha, -\beta) \leq f_2(\alpha, -\beta)$ and the claim (109) for $\beta < 0$ follows noticing that $f_i(\alpha, -\beta) = f_i(\alpha, \beta)$, $i = 1, 2$.

Acknowledgments We would like to thank the Editor and two anonymous Reviewers for a careful reading of the paper. We would like to acknowledge support for the project titled “Epidemics and Counting Structures in Erdős–Rényi Random Graphs” from the Istituto Nazionale di Alta Matematica “Francesco Severi”.

References

- [1] F. Baccelli and B. Błaszczyszyn. On a coverage process ranging from the Boolean model to the Poisson-Voronoi tessellation with applications to wireless communications. *Advances in Applied Probability*, 33, pp. 293–323, 2001.
- [2] F. Baccelli and B. Błaszczyszyn. *Stochastic Geometry and Wireless Networks. Volume I: Theory*. Now Publishers, London, 2009.
- [3] C. Barakat, P. Thiran, G. Iannaccone, C. Diot and P. Owezarski. A flow-based model for internet backbone traffic *IMW '02: Proceedings of the 2nd ACM SIGCOMM Workshop on Internet measurement*, pp. 35–47, November 2002.
- [4] C. Bordenave and G.L. Torrisi. Large deviations of Poisson cluster processes. *Stochastic Models*, 23, pp. 593–625, 2007.
- [5] P. Brémaud. An insensitivity property of Lundberg’s estimate for delayed claims. *Journal of Applied Probability*, 37, pp. 914–917, 2000.
- [6] M. Brigham and A. Destexhe. Nonstationary filtered shot-noise processes and applications to neuronal membranes. *Physical Review E*, 91: 062102, 2015.
- [7] P.C. Consul and F. Famoye. *Lagrangian Probability Distributions*. Birkhäuser, Berlin, 2006.
- [8] D. J. Daley and D. Vere-Jones. *An Introduction to the Theory of Point Processes*. Springer, New York, 2003.
- [9] M. Draief and L. Massoulié. *Epidemics and Rumours in Complex Networks*. Springer, Berlin, 2016.
- [10] K. Duffy and G.L. Torrisi. Sample path large deviations of Poisson shot noise with heavy tail semi-exponential distributions. *Journal of Applied Probability*, 48, pp. 688–698, 2011.

- [11] P. Embrechts, C. Klüppelberg and T. Mikosch. *Modelling Extremal Events*. Springer, Berlin, 2003.
- [12] V. Féray, P.L. Méliot and A. Nikeghbali. *Mod- ϕ Convergence. Normality Zones and Precise Deviations*. Springer, Berlin, 2016.
- [13] V. Féray, P.L. Méliot and A. Nikeghbali. *Mod- ϕ Convergence II: Estimates on the Speed of Convergence*. In: Donati-Martin C., Lejay A., Rouault A. (eds) Séminaire de Probabilités L. Lecture Notes in Mathematics, vol 2252. Springer, Berlin, 2019.
- [14] A. Ganesh and G.L. Torrisi. A class of risk processes with delayed claims: ruin probability estimates under heavy tail conditions. *Journal of Applied Probability*, 43, pp. 916–926, 2006.
- [15] A. Ganesh and G.L. Torrisi. Large deviations of the interference in a wireless communication model. *IEEE Transactions on Information Theory*, 54, pp. 3505–3517, 2008.
- [16] A. Ganesh, C. Macci and G.L. Torrisi. Sample path large deviations for Poisson shot noise processes, and applications. *Electronic Journal of Probability*, 10, pp. 1026–1043, 2005.
- [17] A. Ganesh, C. Macci and G.L. Torrisi. A class of risk processes with reserve-dependent premium rate: sample path large deviations and importance sampling. *Queueing Systems*, 55, pp. 83–94, 2007.
- [18] F. Gao and L. Zhu. Precise deviations of Hawkes processes. *Bernoulli*, to appear, 2021.
- [19] A.G. Hawkes. Spectra of some self-exciting and mutually exciting point processes. *Biometrika*, 58, pp. 83–90, 1971.
- [20] R. van der Hofstad. *Random Graphs and Complex Networks. Volume 1*. Cambridge Series in Statistical and Probabilistic Mathematics, 2017.
- [21] C. Klüppelberg and T. Mikosch. Modelling delay in claim settlement. *Scandinavian Actuarial Journal*, pp. 154 – 168, 1995
- [22] C. Klüppelberg and T. Mikosch. Explosive Poisson shot noise processes with applications to risk reserves. *Bernoulli*, 1, pp. 125–147, 1995.
- [23] C. Klüppelberg, T. Mikosch and A. Schärf Regular variation in the mean and stable limits for Poisson shot noise. *Bernoulli*, 9, pp. 467–496, 2003.
- [24] T. Konstantopoulos and S.J. Lin. Macroscopic models for long-range dependent network traffic. *Queueing Systems*, 28, pp. 214–243, 1998.
- [25] T.G. Kurtz. Limit theorems for workload input models. In: F.P. Kelly, S. Zachary and I. Ziedins (eds), *Stochastic Networks: Theory and Applications*, Oxford University Press, Oxford, pp. 119–139, 1997.
- [26] J.A. Lane. The central limit theorem for the Poisson shot-noise process. *Journal of Applied Probability*, 21, pp. 287 – 301, 1984.
- [27] J.A. Lane. The Berry–Esseen bound for the Poisson shot-noise. *Advances in Applied Probability*, 19, pp. 512 – 514, 1987.
- [28] E. Leonardi and G.L. Torrisi. Modeling LRU caches with shot noise request processes. *SIAM Journal on Applied Mathematics*, 77, pp. 361–383, 2017.

- [29] C. Macci and G.L. Torrisi. Asymptotic results for perturbed risk processes with delayed claims. *Insurance: Mathematics and Economics*, 34, pp. 307–320, 2004.
- [30] C. Macci, G. Stabile and G.L. Torrisi. Lundberg parameters for non standard risk processes. *Scandinavian Actuarial Journal*, 6, pp. 417-432, 2005.
- [31] S. Mehra, J.M. McCaw, M.B. Flegg, P.G. Taylor and J.A. Flegg. Antibody dynamics for plasmodium vivax malaria: a mathematical model *Bulletin of Mathematical Biology*, 83(1): 6, 2021.
- [32] M.K. Nakayama, P. Shahabuddin and K. Sigman. On finite exponential models for branching processes and busy periods for queues. *Journal of Applied Probability*, 41, pp. 273-280, 2004.
- [33] N. Privault. Nonstationary shot-noise modeling of neuron membrane potentials by closed-form moments and Gram-Charlier expansions. *Biological Cybernetics*, to appear, 2021.
- [34] N. Privault and G.L. Torrisi. Density estimation of functionals of spatial point processes with application to wireless networks. *SIAM Journal on Mathematical Analysis*, 43, pp. 1311-1344, 2011.
- [35] G. Stabile and G.L. Torrisi. Large deviations of Poisson shot noise processes, under heavy tail semi-exponential conditions. *Statistics and Probability Letters*, 80, pp. 1200-1209, 2010.
- [36] G.L. Torrisi and E. Leonardi. Simulating the tail of the interference in a Poisson network model. *IEEE Transactions on Information Theory*, 59, 1773–1787, 2013.
- [37] G.L. Torrisi. Simulating the ruin probability of risk processes with delay in claim settlement. *Stochastic Processes and their Applications*, 112, pp. 225-244, 2004.
- [38] D. Vere-Jones. Stochastic models for earthquake occurrences. *Journal of the Royal Statistical Society Ser. B*, 32, pp. 1–42, 1970.