

Flame Retardant Coatings for Plastics and Textiles

Original

Flame Retardant Coatings for Plastics and Textiles / Malucelli, Giulio. - In: COATINGS. - ISSN 2079-6412. - ELETTRONICO. - 15:11(2025), pp. 1-3. [10.3390/coatings15111342]

Availability:

This version is available at: 11583/3005289 since: 2025-11-20T08:09:39Z

Publisher:

MDPI

Published

DOI:10.3390/coatings15111342

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)




ELSEVIER

Contents lists available at ScienceDirect

Journal of Symbolic Computation

www.elsevier.com/locate/jsc



Degrees of Kalman varieties of tensors

Zahra Shahidi ^a, Luca Sodomaco ^b, Emanuele Ventura ^c

^a Università di Firenze, Dipartimento di Matematica e Informatica, Viale Morgagni 67/A, 50134 Firenze, Italy

^b Department of Mathematics and Systems Analysis, Aalto University, Espoo, Finland

^c Politecnico di Torino, Dipartimento di Scienze Matematiche "G. L. Lagrange", Corso Duca degli Abruzzi 24, 10129 Torino, Italy



ARTICLE INFO

Article history:

Available online 22 April 2022

MSC:

14N07

14N05

14N10

15A69

15A18

Keywords:

Tensors

Singular vector tuples

Kalman variety

Generating function

Asymptotics

ABSTRACT

Kalman varieties of tensors are algebraic varieties consisting of tensors whose singular vector k -tuples lay on prescribed subvarieties. They were first studied by Ottaviani and Sturmfels in the context of matrices. We extend recent results of Ottaviani and the first author to the partially symmetric setting. We describe a generating function whose coefficients are the degrees of these varieties and we analyze its asymptotics, providing analytic results à la Zeilberger and Pantone. We emphasize the special role of isotropic vectors in the spectral theory of tensors and describe the totally isotropic Kalman variety as a dual variety.

© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license

(<http://creativecommons.org/licenses/by/4.0/>).

1. Introduction

Singular vector k -tuples are the partially symmetric tensor analog of singular vector pairs of rectangular matrices. Their definition is recalled in §2. We refer to Qi and Luo (2017) for background and applications.

In this article, we tackle the problem of describing the set of partially symmetric tensors admitting a singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$, where (the equivalence class of) each entry \mathbf{x}_i lies on a fixed irreducible subvariety $Z_i \subset \mathbb{P}(V_i)$ of the i -th factor. This set is an algebraic variety in the projective

E-mail addresses: z.shahidi80@gmail.com (Z. Shahidi), luca.sodomaco@aalto.fi (L. Sodomaco), emanuele.ventura@polito.it, emanueleventura.sw@gmail.com (E. Ventura).

<https://doi.org/10.1016/j.jsc.2022.04.016>

0747-7171/© 2022 The Author(s). Published by Elsevier Ltd. This is an open access article under the CC BY license (<http://creativecommons.org/licenses/by/4.0/>).

space $\mathbb{P}(S^{\omega_1} V_1 \otimes \cdots \otimes S^{\omega_k} V_k)$ of partially symmetric tensors with k factors called *generalized Kalman variety* of tensors.

Recently, the spectral theory of tensors along with its connections to pure and applied algebraic geometry and combinatorics is witnessing several interesting results; see e.g. Beorchia et al. (2021), Teixeira Turatti (2021) for recent geometric results. Despite this progress, algebraic varieties providing a clean geometric picture of spectral properties of tensors await to be described. Kalman varieties are therefore central objects in this context. This paper is a contribution to this circle of ideas from the perspective of degrees of varieties, their generating functions and asymptotics.

The name *Kalman variety* was first introduced by Ottaviani and Sturmfels in Ottaviani and Sturmfels (2013) to indicate the variety of square matrices possessing at least one eigenvector on a fixed linear subspace. They determined its codimension, degree and studied its singular locus. Thereafter, Sam (2012) and Huang (2020) determined their defining equations. More recently, Ottaviani and the first author Ottaviani and Shahidi (2021) rephrased the original setting for singular vector pairs, extending it to the case of singular vector k -tuples. Their Kalman variety is the variety of tensors having a singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$, where (the equivalence class of) the first entry \mathbf{x}_1 lies on a fixed linear subspace $L \subset \mathbb{P}(V_1)$.

Our point of departure is (Ottaviani and Shahidi, 2021, Theorem 1.2), where the authors showed that the codimension of this Kalman variety is equal to the codimension of $L \subset \mathbb{P}(V_1)$. Moreover, they established an elegant formula for the degrees of the Kalman variety for symmetric and non-symmetric tensors. In the latter case, letting $n_i = \dim(V_i)$ and $\delta = \text{codim}(L)$, the desired degree is the coefficient of the monomial $h^\delta t_1^{n_1 - \delta - 1} \prod_{i \geq 2} t_i^{n_i - 1}$ in the polynomial

$$\prod_{i=1}^k \frac{(\widehat{t}_i + h)^{n_i} - t_i^{n_i}}{(\widehat{t}_i + h) - t_i}, \quad \widehat{t}_i := \left(\sum_{j=1}^k t_j \right) - t_i.$$

This expression is similar to the formula for the number of singular vector k -tuples of a general tensor. The latter quantity coincides with another well-known metric invariant of an algebraic variety, called the *ED degree* of the Segre variety $\mathbb{P}(V_1) \times \cdots \times \mathbb{P}(V_k)$. For more details on ED degrees of algebraic varieties, we refer to Draisma et al. (2016). Ottaviani and Friedland (Friedland and Ottaviani, 2014, Theorem 1) computed the above ED degree using Chern classes of a suitable vector bundle on the Segre variety.

The aforementioned results establish the *enumerative* nature of the subject. When enumerative structures appear, it is a natural problem to determine a generating function whose coefficients are the counted quantities. Generating functions are tremendously useful tools to have a global picture of the enumerated objects. See MacMahon (1915), Stanley (2012), Wilf (1990) for fascinating introductions to this topic along with its applications to combinatorics and analysis. Ekhad and Zeilberger (2016) found a generating function for the Friedland-Ottaviani’s formula and hence for the ED degrees. The asymptotic behavior was then analyzed by Pantone (2017). Zeilberger’s generating function is strikingly similar to the generating function of degrees of hyperdeterminants. An asymptotic analysis of the latter was performed in (Ottaviani et al., 2021, Theorem 3.8).

Our first contribution is a generalization of (Ottaviani and Shahidi, 2021, Theorem 1.2) and (Friedland and Ottaviani, 2014, Theorem 12). More specifically, we determine the degrees of generalized Kalman varieties of partially symmetric tensors.

Theorem 1. *For every $i \in [k]$, let $Z_i \subset \mathbb{P}(V_i)$ be an irreducible projective variety of codimension δ_i and let $Z = \prod_{i=1}^k Z_i$. We assume that each Z_i is not contained in the isotropic quadric $Q_i \subset \mathbb{P}(V_i)$. Let $\delta = (\delta_1, \dots, \delta_k)$. The generalized Kalman variety*

$$\kappa_{\mathbf{n}, \omega}(Z) := \{T \in \mathbb{P}(S^\omega V) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z\}$$

is connected of codimension $\delta := \sum_{i=1}^k \delta_i$. The degree of $\kappa_{\mathbf{n}, \omega}(Z)$ is $d(\mathbf{n}, \delta, \omega) \prod_{i=1}^k \deg(Z_i)$, where $d(\mathbf{n}, \delta, \omega)$ is the coefficient of the monomial $h^\delta \prod_{i=1}^k t_i^{n_i - \delta_i - 1}$ in the polynomial

$$\prod_{i=1}^k \frac{(\widehat{t}_i + h)^{n_i} - t_i^{n_i}}{(\widehat{t}_i + h) - t_i}, \quad \widehat{t}_i := \left(\sum_{j=1}^k \omega_j t_j \right) - t_i.$$

We study separately the case when each subvariety Z_i coincides with the isotropic quadric Q_i . The corresponding Kalman variety is called *totally isotropic Kalman variety*. In Theorem 25, we describe the totally isotropic Kalman variety as a dual variety of a specific Segre-Veronese variety. This specializes to Theorem 27 in the symmetric case; this last result is related to (Beorchia et al., 2021, Proposition 2.10).

Our second main result furnishes a Zeilberger-type (Ekhad and Zeilberger, 2016, Proposition 1) generating function for the degrees of generalized Kalman varieties in the case $\delta = (\delta, 0, \dots, 0)$.

Theorem 2. *Keep the assumptions of Theorem 1. If $\delta = (\delta, 0, \dots, 0)$ for some $\delta \geq 0$, the generating function for the coefficients $d(\mathbf{n}, \delta, \omega)$ is*

$$\sum_{\mathbf{n} \in \mathbb{N}^k} \sum_{\delta=0}^{\infty} d(\mathbf{n}, \delta, \omega) \mathbf{x}^{\mathbf{n}} y^{\delta} = \frac{1}{H_{\omega}(\mathbf{x}, y)} \prod_{i=1}^k \frac{x_i}{1 - x_i} \tag{1.1}$$

where

$$H_{\omega}(\mathbf{x}, y) := -y x_1 \prod_{i=2}^k (1 + x_i) + \prod_{i=1}^k (1 + x_i) - \sum_{j=1}^k \omega_j x_j \prod_{i \neq j} (1 + x_i). \tag{1.2}$$

Our third result offers an asymptotic study of the degrees $d(\mathbf{n}, \delta, \omega)$ in three different regimes: first, when $n_i \rightarrow \infty$ for a single index i . Secondly, in the binary format $n_1 = \dots = n_k = 2$ as $k \rightarrow \infty$. Finally, in the hypercubical format $n_1 = \dots = n_k = n$ for $n \rightarrow \infty$. The last asymptotic study is more involved and leads to the following result, which agrees with (Pantone, 2017, Theorem 1.3) for $\delta = 0$ and $\omega = 1$. We refer to the first two paragraphs of Section 2 for the preliminary notations and definitions used.

Theorem 3. *Consider the factor $d(n, \delta, \omega)$ in the degree of the generalized Kalman variety $\kappa_{n, \omega 1}(Z)$. Assume that either $k = 2$ and $w \geq 2$ or $k \geq 3$. Then asymptotically, for $n \rightarrow +\infty$,*

$$d(n, \delta, \omega) = \frac{(\omega k - 1)^{k-1}}{(2\pi)^{\frac{k-1}{2}} (\omega k)^{\frac{k-2}{2}} (\omega k - 2)^{\frac{3k-1}{2}}} \left(\frac{\omega k}{\omega k - 1} \right)^{\delta} \frac{(\omega k - 1)^{kn}}{n^{\frac{k-1}{2} - \delta}} \left[1 + O\left(\frac{1}{n}\right) \right].$$

In Definition 14, we introduce the *Kalman strata*. These are, roughly speaking, the building blocks of generalized Kalman varieties, and are meant to take into account the presence of *isotropic* vectors in the singular vector tuple. The proof of Theorem 1 relies on determining the linear space of tensors having a singular vector k -tuple over a specific Kalman stratum. This is achieved in Theorem 13. This result is interesting on its own right, giving an intrinsic description of the fibers in terms of orthogonal spaces. Another consequence of Theorem 13 is Proposition 19, where we show the irreducibility of the *spectral variety* (see Definition 11), which is a natural incidence correspondence in this setting.

In §6, we introduce Kalman varieties of symmetric singular vector k -tuples and derive their codimensions. The degrees of these new interesting varieties seem more challenging even in the case of matrices. We leave their determination as an open question.

Acknowledgments

The idea of this project was conceived during Shahidi’s postdoc at Università di Firenze. We thank Giorgio Ottaviani for very useful discussions and encouragement. We thank Jan Draisma for explaining to us a way of deriving the equations in Example 41. The first author thanks Dr. Alireza Firoozfar and

Dr. Mohsen Afsharchi for their support. The second author is partially supported by the Academy of Finland Grant 323416. During most of the preparation of the manuscript, the third author was a postdoc at Universität Bern, supported by Vici Grant 639.033.514 of Jan Draisma from the Netherlands Organisation for Scientific Research. We thank two anonymous referees for their useful comments and questions that also helped to improve the presentation.

2. Kalman varieties of tensors

Notation. Throughout the paper, if not specified we denote by \mathbf{j} a vector (j_1, \dots, j_k) of k variables, while $\mathbf{x}^{\mathbf{m}}$ stands for the monomial $x_1^{m_1} \dots x_k^{m_k}$. We often use the shorthand $[k]$ to denote the set of indices $\{1, \dots, k\}$. Define $\mathbf{1} = (1, \dots, 1) \in \mathbb{N}^k$ and, for $m \in \mathbb{N}$, let $m\mathbf{1} = (m, \dots, m) \in \mathbb{N}^k$.

For every $i \in [k]$ we consider an n_i -dimensional vector space V_i over the field $\mathbb{F} = \mathbb{R}$ or $\mathbb{F} = \mathbb{C}$. If $\mathbb{F} = \mathbb{R}$, then we prefer the notation $V_i^{\mathbb{R}}$. Let $\boldsymbol{\omega} = (\omega_1, \dots, \omega_k)$ be a vector of nonnegative integers. For each $i \in [k]$, let $S^{\omega_i} V_i$ be the ω_i -th symmetric power of V_i , as a subspace of the tensor product $V_i^{\otimes \omega_i}$. Moreover, we denote by $S^{\boldsymbol{\omega}} V$ the tensor product $\bigotimes_{i=1}^k S^{\omega_i} V_i$. This is the space of *partially symmetric tensors* of format $n_1^{\times \omega_1} \times \dots \times n_k^{\times \omega_k}$. Every element of $S^{\boldsymbol{\omega}} V$ is a linear combination of *decomposable partially symmetric tensors*, that is, tensors of the form $T = \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}$ for some vectors $\mathbf{x}_i \in V_i$. On each projective space $\mathbb{P}(V_i)$ we fix a smooth projective quadric hypersurface $Q_i = \mathcal{V}(q_i)$, where q_i is the homogeneous polynomial in $\mathbb{C}[x_{i,1}, \dots, x_{i,n_i}]_2$ associated to a positive definite real quadratic form $q_i^{\mathbb{R}} : V_i^{\mathbb{R}} \rightarrow \mathbb{R}$. We refer to Q_i as the *isotropic quadric* in the i -th factor $\mathbb{P}(V_i)$. Finally, we denote by \mathbb{P} the product $\prod_{i=1}^k \mathbb{P}(V_i)$ and by $\Pi_{\mathbf{n}, \boldsymbol{\omega}}$ the product $\mathbb{P}(S^{\boldsymbol{\omega}} V) \times \mathbb{P}$.

Definition 4. The *Frobenius (or Bombieri-Weyl) inner product* of two complex decomposable tensors $T = \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}$ and $T' = \mathbf{y}_1^{\omega_1} \otimes \dots \otimes \mathbf{y}_k^{\omega_k}$ of $S^{\boldsymbol{\omega}} V$ is

$$q_F(T, T') := q_1(\mathbf{x}_1, \mathbf{y}_1)^{\omega_1} \dots q_k(\mathbf{x}_k, \mathbf{y}_k)^{\omega_k}, \tag{2.1}$$

and it is naturally extended to every vector in $S^{\boldsymbol{\omega}} V$. We identify all the vector spaces with their duals using the Frobenius inner product.

Definition 5. Let $T \in S^{\boldsymbol{\omega}} V$. A *singular vector k -tuple* of T is a k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ of nonzero vectors $\mathbf{x}_i \in V_i$ such that

$$\text{rank} \left(\begin{matrix} T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) \\ \mathbf{x}_i \end{matrix} \right) \leq 1 \quad \forall i \in [k], \tag{2.2}$$

where $T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k})$ is the tensor contraction of T with respect to $\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}$. If $\omega_i = 1$ for some $i \in [k]$, then $\mathbf{x}_i^{\omega_i-1} = \mathbf{x}_i^0 = 1$ as an element of $S^0 V_i = \mathbb{F}$. If we interpret T as a multi-homogeneous polynomial in the coordinates of each vector \mathbf{x}_i , then the previous tensor contraction corresponds the gradient $\nabla_i T$ with respect to the coordinates of \mathbf{x}_i .

A singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is *normalized* if $q_i(\mathbf{x}_i) = 1$ for all $i \in [k]$. A singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is *isotropic* if $q_i(\mathbf{x}_i) = 0$ for some $i \in [k]$. When $k = 1$, $\mathbf{n} = (n)$ and $\boldsymbol{\omega} = (\omega)$, a vector $\mathbf{x} \in V$ satisfying (2.2) is called an *eigenvector* of the symmetric tensor $T \in S^{\boldsymbol{\omega}} V$.

Definition 6. Let $T \in S^{\boldsymbol{\omega}} V$ and let $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ be a singular vector k -tuple of T . For every $i \in [k]$, the value $\sigma_i \in \mathbb{C}$ such that

$$T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) = \sigma_i \mathbf{x}_i \tag{2.3}$$

is called the *i -th singular value* of $(\mathbf{x}_1, \dots, \mathbf{x}_k)$.

Remark 7. Let $T \in S^\omega V$ and let $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ be a singular vector k -tuple of T with the associated k -tuple $(\sigma_1, \dots, \sigma_k)$ of singular values. For every $i \in [k]$, the identity (2.3) can be interpreted as the identity in $(V_i)^*$

$$q_F(T, \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot _ \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) = \sigma_i q_i(\mathbf{x}_i, _).$$

Evaluating the left-hand side of the previous identity at the vector \mathbf{x}_i yields the number $q_F(T, \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k})$ which does not depend on the specific index i . Thus we have $\sigma_1 q_1(\mathbf{x}_1, \mathbf{x}_1) = \dots = \sigma_k q_k(\mathbf{x}_k, \mathbf{x}_k)$. Therefore

- (i) $\sigma_1 = \dots = \sigma_k$ if $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is normalized. The common value $\sigma := \sigma_1 = \dots = \sigma_k$ is often called the *singular value* of the normalized singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$.
- (ii) $\sigma_j = 0$ for all $j \in [k]$ such that $q_j(\mathbf{x}_j, \mathbf{x}_j) \neq 0$ if $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is isotropic.

If $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is a singular vector k -tuple of $T \in S^\omega V$, then it is immediate to check that, for every tuple $(\lambda_1, \dots, \lambda_k)$ of nonzero complex numbers, the k -tuple $(\lambda_1 \mathbf{x}_1, \dots, \lambda_k \mathbf{x}_k)$ is also a singular vector k -tuple of T . For this reason, we consider $\mathbb{P} = \prod_{i=1}^k \mathbb{P}(V_i)$ and say that $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$ is a singular vector k -tuple of T if its representative $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is.

Definition 8. For every $i \in [k]$, we denote by $SO(V_i)$ the (complex) *special orthogonal group* of automorphisms of V_i with determinant 1 that preserve the bilinear product in V_i associated to q_i .

Remark 9. The notion of singular vector k -tuple is $SO(V_1) \times \dots \times SO(V_k)$ -equivariant. More explicitly, given a singular vector k -tuple $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ of T and an element $g_i \in SO(V_i)$ for all $i \in [k]$, then $(g_1(\mathbf{x}_1), \dots, g_k(\mathbf{x}_k))$ is a singular vector k -tuple of $(g_1, \dots, g_k) \cdot T$. This is because the contraction (or scalar product) in the i -th factor is preserved by the action of $SO(V_i)$.

Lemma 10. Consider a tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$. For all $i \in [k]$, define the subspace

$$\begin{aligned} W_i &:= \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \langle \mathbf{x}_i \rangle^\perp \otimes \dots \otimes \mathbf{x}_k^{\omega_k} \\ &= \{ \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{w}_i \otimes \dots \otimes \mathbf{x}_k^{\omega_k} \mid \mathbf{w}_i \in \langle \mathbf{x}_i \rangle^\perp \} \subset S^\omega V, \end{aligned} \tag{2.4}$$

where $\langle \mathbf{x}_i \rangle^\perp$ is the orthogonal complement of $\langle \mathbf{x}_i \rangle$ in V_i with respect to the fixed quadratic form q_i . Given a tensor $T \in S^\omega V$, we have that $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$ is a singular vector k -tuple of T if and only if $T \in (W_1 + \dots + W_k)^\perp$, where in this case the sign \perp denotes the orthogonal complement with respect to the Frobenius inner product in $S^\omega V$.

Proof. By definition, the tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$ is a singular vector k -tuple of T if and only if the relations in (2.2) hold. For a fixed $i \in [k]$, the corresponding relation in (2.2) is equivalent to $T \in W_i^\perp$. Hence $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$ is a singular vector k -tuple of T if and only if $T \in W_1^\perp \cap \dots \cap W_k^\perp$, and one has the equality $W_1^\perp \cap \dots \cap W_k^\perp = (W_1 + \dots + W_k)^\perp$. \square

We introduce an incidence correspondence that will play a fundamental role in the upcoming proofs.

Definition 11. Let $\Pi_{\mathbf{n}, \omega} = \mathbb{P}(S^\omega V) \times \mathbb{P}$. The *spectral variety of type (\mathbf{n}, ω)* is

$$\Sigma_{\mathbf{n}, \omega} := \{ ([T], [\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \Pi_{\mathbf{n}, \omega} \mid ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \text{ is a singular vector } k\text{-tuple for } T \}. \tag{2.5}$$

Remark 12. Note that $\Sigma_{\mathbf{n}, \omega}$, as described set-theoretically above, is closed in the Zariski topology: equipped with its reduced structure, it is the subvariety whose ideal is the radical ideal of

$$(J_1 + \dots + J_k) : \left(\prod_{i=1}^k \langle \mathbf{x}_i \rangle \right)^\infty \subset \mathbb{C}[\mathbf{x}_1, \dots, \mathbf{x}_k],$$

where $\langle \mathbf{x}_i \rangle$ is the ideal generated by the coordinates of \mathbf{x}_i and

$$J_i = \left\langle 2 \times 2 \text{ minors of } \begin{pmatrix} T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) \\ \mathbf{x}_i \end{pmatrix} \right\rangle \quad \forall i \in [k].$$

In the following, we will often consider the diagram

$$\begin{array}{ccc} & \Sigma_{\mathbf{n}, \omega} & \\ \alpha \swarrow & & \searrow \beta \\ \mathbb{P}(S^\omega V) & & \mathbb{P}, \end{array} \tag{2.6}$$

where α and β denote the two projections along the two factors of $\Pi_{\mathbf{n}, \omega}$.

Theorem 13. *Let $\Sigma_{\mathbf{n}, \omega} \subset \Pi_{\mathbf{n}, \omega}$ be the spectral variety of type (\mathbf{n}, ω) . The projection $\beta : \Sigma_{\mathbf{n}, \omega} \rightarrow \mathbb{P}$ is surjective and every fiber of β over a k -tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k])$ with r isotropic components is a projective subspace in $\mathbb{P}(S^\omega V)$ of codimension*

$$\text{codim}(\beta^{-1}([\mathbf{x}_1], \dots, [\mathbf{x}_k])) = \sum_{i=1}^k (n_i - 1) - \max\{0, r - 1\}. \tag{2.7}$$

Proof. Consider a tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in \mathbb{P}$. The fiber $\beta^{-1}([\mathbf{x}_1], \dots, [\mathbf{x}_k])$ is isomorphic to the projective subspace of classes $[T] \in \mathbb{P}(S^\omega V)$ such that $([\mathbf{x}_1], \dots, [\mathbf{x}_k])$ is a singular vector k -tuple of T . By Lemma 10, we have that

$$\beta^{-1}([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \cong \mathbb{P}((W_1 + \dots + W_k)^\perp),$$

where the subspaces W_i have been defined in (2.4). Hence

$$\text{codim}(\beta^{-1}([\mathbf{x}_1], \dots, [\mathbf{x}_k])) = \dim(W_1 + \dots + W_k).$$

The proof goes by induction on k . First, we assume $k = 2$. We have

$$\dim(W_1 + W_2) = \dim(W_1) + \dim(W_2) - \dim(W_1 \cap W_2) = \sum_{i=1}^2 (n_i - 1) - \dim(W_1 \cap W_2),$$

so it remains to compute $\dim(W_1 \cap W_2)$ depending on the number of isotropic vectors in the pair $([\mathbf{x}_1], [\mathbf{x}_2])$. Consider nonzero elements $F_1 \in W_1$ and $F_2 \in W_2$, written explicitly as $F_1 = \mathbf{x}_1^{\omega_1-1} \cdot \mathbf{w}_1 \otimes \mathbf{x}_2^{\omega_2}$ and $F_2 = \mathbf{x}_1^{\omega_1} \otimes \mathbf{x}_2^{\omega_2-1} \cdot \mathbf{w}_2$ for some vectors $\mathbf{w}_1 \in (\mathbf{x}_1)^\perp$ and $\mathbf{w}_2 \in (\mathbf{x}_2)^\perp$. Then necessarily $F_1 = F_2$ only if $q_1(\mathbf{x}_1) = q_2(\mathbf{x}_2) = 0$, in which case the equality is attained for $\mathbf{w}_i \in \langle \mathbf{x}_i \rangle^\perp$, $i \in [2]$. In particular $\dim(W_1 \cap W_2) = 1$ and the identity (2.7) is satisfied. Otherwise $W_1 \cap W_2 = \{0\}$ if at least one among $\mathbf{x}_1, \mathbf{x}_2$ is not isotropic, and (2.7) is satisfied as well.

Now we assume that the identity (2.7) is satisfied for $k - 1$ factors among the given k . Without loss of generality, we apply the induction step to the first $k - 1$ factors. In particular

$$\dim(W_1 + \dots + W_k) = \dim(W_1 + \dots + W_{k-1}) + \dim(W_k) - \dim[(W_1 + \dots + W_{k-1}) \cap W_k],$$

so it remains to compute $\dim[(W_1 + \dots + W_{k-1}) \cap W_k]$ depending on the number of isotropic vectors in the tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k])$. Pick nonzero tensors $F_1 \in W_1 + \dots + W_{k-1}$ and $F_2 \in W_k$, which can be written explicitly as

$$F_1 = \left(\sum_{i=1}^{k-1} \mathbf{x}_1^{\omega_1} \otimes \cdots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{w}_i \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}} \right) \otimes \mathbf{x}_k^{\omega_k}, \mathbf{w}_i \in \langle \mathbf{x}_i \rangle^\perp$$

$$F_2 = \mathbf{x}_1^{\omega_1} \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}} \otimes \mathbf{x}_k^{\omega_k-1} \cdot \mathbf{w}_k, \mathbf{w}_k \in \langle \mathbf{x}_k \rangle^\perp.$$

If $q_k(\mathbf{x}_k) \neq 0$, then necessarily $F_1 \neq F_2$, so $(W_1 + \cdots + W_{k-1}) \cap W_k = \{0\}$. In this case the identity (2.7) follows applying the induction step. Otherwise $q_k(\mathbf{x}_k) = 0$ and in this case $F_1 = F_2$ only if $\mathbf{w}_k \in \langle \mathbf{x}_k \rangle \subset \langle \mathbf{x}_k \rangle^\perp$ and

$$\sum_{i=1}^{k-1} \mathbf{x}_1^{\omega_1} \otimes \cdots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{w}_i \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}} = \mathbf{x}_1^{\omega_1} \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}}. \tag{2.8}$$

Now suppose that $q_i(\mathbf{x}_i) \neq 0$ for some $i \in [k - 1]$, for simplicity $i = 1$. Since $\langle \mathbf{x}_1 \rangle^\perp \oplus \langle \mathbf{x}_1 \rangle = \mathbb{C}^{n_1}$, we have the decomposition

$$\left(\mathbf{x}_1^{\omega_1-1} \langle \mathbf{x}_1 \rangle^\perp \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i \right) \oplus \left(\mathbf{x}_1^{\omega_1-1} \langle \mathbf{x}_1 \rangle \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i \right) = \mathbf{x}_1^{\omega_1-1} \mathbb{C}^{n_1} \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i.$$

On one hand, the left-hand side of (2.8) can be rewritten as

$$\mathbf{x}_1^{\omega_1-1} \cdot \mathbf{w}_1 \otimes \mathbf{x}_2^{\omega_2} \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}} + \sum_{i=2}^{k-1} \mathbf{x}_1^{\omega_1} \otimes \cdots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{w}_i \otimes \cdots \otimes \mathbf{x}_{k-1}^{\omega_{k-1}}.$$

The first summand is a nonzero element of $\mathbf{x}_1^{\omega_1-1} \langle \mathbf{x}_1 \rangle^\perp \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i$, whereas the second summand is a nonzero element of $\mathbf{x}_1^{\omega_1-1} \langle \mathbf{x}_1 \rangle \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i$. On the other hand, the right-hand side of (2.8) lives only in $\mathbf{x}_1^{\omega_1-1} \langle \mathbf{x}_1 \rangle \otimes \bigotimes_{i=2}^{k-1} S^{\omega_i} V_i$. Hence $F_1 \neq F_2$ and $(W_1 + \cdots + W_{k-1}) \cap W_k = \{0\}$. The identity (2.7) again follows applying the induction step.

We conclude that, if $(W_1 + \cdots + W_{k-1}) \cap W_k \neq \{0\}$, then $q_i(\mathbf{x}_i) = 0$ for all $i \in [k]$. Under this assumption, we have that $\dim[(W_1 + \cdots + W_{k-1}) \cap W_k] = 1$: indeed, the only possibility to have $F_1 = F_2$ is that $\mathbf{w}_i \in \langle \mathbf{x}_i \rangle$ for all $i \in [k]$. Hence, by induction,

$$\begin{aligned} \dim(W_1 + \cdots + W_k) &= \dim(W_1 + \cdots + W_{k-1}) + \dim(W_k) - \dim[(W_1 + \cdots + W_{k-1}) \cap W_k] \\ &= \sum_{i=1}^k (n_i - 1) - (k - 1), \end{aligned}$$

which agrees with (2.7) when $r = k$. \square

Definition 14 (Kalman strata). For every $i \in [k]$, let $Z_i \subset \mathbb{P}(V_i)$ be a projective variety and consider the product $Z = \prod_{i=1}^k Z_i$. Given a subset $J \subset [k]$, we define the product

$$Q_J := \prod_{j \in J} Q_j \times \prod_{j \notin J} [\mathbb{P}(V_j) \setminus Q_j] \subset \mathbb{P}.$$

The partially isotropic Kalman strata with respect to Z are

$$\kappa_{\mathbf{n}, \omega}^J(Z) := \overline{\{T \in \mathbb{P}(S^\omega V) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z \cap Q_J\}}.$$

In particular,

- for $J = \emptyset$, we call $\kappa_{\mathbf{n}, \omega}^{nor}(Z) := \kappa_{\mathbf{n}, \omega}^\emptyset(Z)$ the normalized Kalman variety with respect to Z because we may assume that all components \mathbf{x}_i are normalized with respect to the inner product q_i . We also denote by Q_{nor} the product $Q_\emptyset = \prod_{j=1}^k [\mathbb{P}(V_j) \setminus Q_j]$.

- for $J = [k]$, we call $\kappa_{\mathbf{n},\omega}^{iso}(Z) := \kappa_{\mathbf{n},\omega}^{[k]}(Z)$ the *totally isotropic Kalman variety* with respect to Z because all components \mathbf{x}_i are isotropic. In this case

$$\kappa_{\mathbf{n},\omega}^{iso}(Z) = \{T \in \mathbb{P}(S^\omega V) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z \cap Q\},$$

namely the right-hand side is already closed. Here we use the shorthand $Q := Q_{[k]} = \prod_{i=1}^k Q_i$. If $Z = Q$, we indicate this variety simply with $\kappa_{\mathbf{n},\omega}^{iso}$.

Note that all the loci $\kappa_{\mathbf{n},\omega}^J(Z)$ are closed by definition, and they may be reducible. We generally refer to all of them (to their irreducible components) as *Kalman strata*. If $\omega = \mathbf{1}$, we use the shorthand $\kappa_{\mathbf{n}}^J(Z) = \kappa_{\mathbf{n},\mathbf{1}}^J(Z)$.

Definition 15. For every $i \in [k]$, let $Z_i \subset \mathbb{P}(V_i)$ be a projective variety. The *generalized Kalman variety* with respect to $Z = \prod_{i=1}^k Z_i$ is

$$\kappa_{\mathbf{n},\omega}(Z) := \{T \in \mathbb{P}(S^\omega V) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z\}. \tag{2.9}$$

The Kalman strata $\kappa_{\mathbf{n},\omega}^J(Z)$ of Definition 14 are closed subvarieties of the generalized Kalman variety. Moreover, if $Z = Q = \prod_{i=1}^k Q_i$, then $\kappa_{\mathbf{n},\omega}(Q) = \kappa_{\mathbf{n},\omega}^{iso}$.

Similarly as in Remark 12, the right-hand side of (2.9) is closed in the Zariski topology.

Definition 16. Consider the spectral variety $\Sigma_{\mathbf{n},\omega} \subset \Pi_{\mathbf{n},\omega}$. For every $i \in [k]$, let $Z_i \subset \mathbb{P}(V_i)$ be a projective variety and consider the product $Z = \prod_{i=1}^k Z_i \subset \mathbb{P}$. The *spectral variety of type (\mathbf{n}, ω) restricted to Z* is the incidence variety $\Sigma_{\mathbf{n},\omega}(Z) := \Sigma_{\mathbf{n},\omega} \cap [\mathbb{P}(S^\omega V) \times Z]$. In particular,

$$\Sigma_{\mathbf{n},\omega}(Z) = \{(T, [\mathbf{x}_1], \dots, [\mathbf{x}_k]) \mid ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z \text{ is a singular vector } k\text{-tuple for } T\}.$$

Similarly as in (2.6), in the following proofs we will consider the diagram of projections

$$\begin{array}{ccc} & \Sigma_{\mathbf{n},\omega}(Z) & \\ \alpha_Z \swarrow & & \searrow \beta_Z \\ \mathbb{P}(S^\omega V) & & Z. \end{array} \tag{2.10}$$

Before we proceed, we recall a standard lemma needed in the next proofs. The (omitted) proof relies on a direct application of (Hartshorne, 1977, Proposition 7.10) and (Vakil, 2017, Exercise 11.4.C).

Lemma 17. Let $\varphi: M \rightarrow N$ be a surjective morphism of projective varieties and assume that N is connected. Suppose there exists a finite collection $\{S_i\}_{i \in I}$ of irreducible quasi-projective subvarieties of N such that $N = \bigcup_{i \in I} S_i$ with the property that, for each $i \in I$, the restriction $\varphi|_{\varphi^{-1}(S_i)}: \varphi^{-1}(S_i) \rightarrow S_i$ has equidimensional linear projective fibers. Then:

- (i) Each closed subvariety $\overline{\varphi^{-1}(S_i)} \subset M$ is irreducible.
- (ii) M is connected and $M = \bigcup_{i \in I} \overline{\varphi^{-1}(S_i)}$. In particular, the irreducible components of M have the form $\overline{\varphi^{-1}(S_i)}$ for some $i \in I$.
- (iii) One has $\varphi^{-1}(S_i) \setminus \overline{\varphi^{-1}(S_i)} \subset \bigcup_{j \in I \setminus \{i\}} \varphi^{-1}(S_j)$.

Proposition 18. For every $i \in [k]$, let $Z_i \subset \mathbb{P}(V_i)$ be an irreducible projective variety of codimension δ_i and consider the product $Z = \prod_{i=1}^k Z_i \subset \mathbb{P}$. We assume that $Z_i \not\subset Q_i$ for all $i \in [k]$ and let $\delta := \sum_{i=1}^k \delta_i$. Then:

- (i) The variety $\kappa_{\mathbf{n},\omega}^{nor}(Z)$ is irreducible of dimension $\dim(\kappa_{\mathbf{n},\omega}^{nor}(Z)) = \dim(\mathbb{P}(S^\omega V)) - \delta$.

(ii) For any $J \neq \emptyset$, we have $\dim(\kappa_{\mathbf{n},\omega}^J(Z)) < \dim(\kappa_{\mathbf{n},\omega}^{nor}(Z))$.

Proof. (i). Consider the restricted spectral variety $\Sigma_{\mathbf{n},\omega}(Z)$ along with the two projections α_Z and β_Z introduced in diagram (2.10). The morphism β_Z is surjective and the fiber $\beta_Z^{-1}([\mathbf{x}_1], \dots, [\mathbf{x}_k])$ is set-theoretically the collection of tensors $T \in \mathbb{P}(S^\omega V)$ that possess $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z$ as a singular vector k -tuple.

Let $Q = \prod_{i=1}^k Q_i \subset \mathbb{P}$ and $Y = \overline{\beta_Z^{-1}(Z \setminus Q)} \subset \Sigma_{\mathbf{n},\omega}(Z)$. Now consider the morphism

$$\psi := \beta_Z|_{\beta_Z^{-1}(Z \setminus Q)} : \beta_Z^{-1}(Z \setminus Q) \rightarrow Z \setminus Q.$$

By Theorem 13, ψ is a surjective morphism with equidimensional linear projective fibers of codimension $\sum_{i=1}^k (n_i - 1)$ in $\mathbb{P}(S^\omega V)$. By Lemma 17(i), Y is irreducible.

Now consider the first projection $\alpha_Z : \Sigma_{\mathbf{n},\omega}(Z) \rightarrow \mathbb{P}(S^\omega V)$. The morphism α_Z is projective and hence closed. One has $\kappa_{\mathbf{n},\omega}^{nor}(Z) = \alpha_Z(Y)$. To see this, notice that by definition we have:

$$\alpha_Z(\beta_Z^{-1}(Z \setminus Q)) = \{T \in \mathbb{P}(S^\omega V) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Z \setminus Q\}.$$

One finds $\kappa_{\mathbf{n},\omega}^{nor}(Z) = \overline{\alpha_Z(\beta_Z^{-1}(Z \setminus Q))} = \overline{\alpha_Z(\beta_Z^{-1}(Z \setminus Q))} = \overline{\alpha_Z(Y)} = \alpha_Z(Y)$. Here the first equality follows by Definition 14, the second by continuity of α_Z , and the last follows from α_Z being closed. Moreover, since Y is irreducible, then so is $\kappa_{\mathbf{n},\omega}^{nor}(Z)$.

In order to calculate the codimension of $\kappa_{\mathbf{n},\omega}^{nor}(Z)$, note that $\dim(Y) = \dim(\mathbb{P}(S^\omega V)) + \dim(Z) - \sum_{i=1}^k (n_i - 1) = \dim(\mathbb{P}(S^\omega V)) - \delta$. Observe that the general fiber of the restriction of α_Z to Y is finite. Indeed, a point of the product $Z \setminus Q$ is the equivalence class of a tuple of non-isotropic vectors. The group $SO(V_1) \times \dots \times SO(V_k)$ acts transitively on equivalence classes of non-isotropic vectors. Thus a general tensor in the fiber under β_Z of a point in $Z \setminus Q$ has a finite number of singular vector k -tuples. This implies the finiteness of the restriction of α_Z to Y . Then $\dim(\kappa_{\mathbf{n},\omega}^{nor}(Z)) = \dim(Y)$. So $\text{codim}(\kappa_{\mathbf{n},\omega}^{nor}(Z)) = \delta$.

(ii). Fix a nonempty subset $J \subset [k]$ and consider the quasi-projective subvariety $Z \cap Q_J$ of Z . By Theorem 13, the fiber of β_Z at every point of $Z \cap Q_J$ has dimension $\dim(\mathbb{P}(S^\omega V)) - \sum_{i=1}^k (n_i - 1) + (|J| - 1)$. Let $Y_J = \beta_Z^{-1}(Z \cap Q_J)$. Thus

$$\begin{aligned} \dim(Y_J) &= \sum_{j \notin J} \dim(Z_j) + \sum_{j \in J} [\dim(Z_j) - 1] + \dim(\mathbb{P}(S^\omega V)) - \sum_{i=1}^k (n_i - 1) + (|J| - 1) \\ &= \dim(\mathbb{P}(S^\omega V)) - \delta - 1. \end{aligned}$$

As before, we find that $\kappa_{\mathbf{n},\omega}^J(Z) = \alpha_Z(Y_J)$. Thus the dimension of $\kappa_{\mathbf{n},\omega}^J(Z)$ is at most that of Y_J , which finishes the proof. \square

The last step before proving Theorem 1 is establishing the irreducibility of the spectral variety $\Sigma_{\mathbf{n},\omega}$.

Proposition 19. *The spectral variety $\Sigma_{\mathbf{n},\omega} \subset \Pi_{\mathbf{n},\omega}$ is irreducible and $\Sigma_{\mathbf{n},\omega} = \overline{\beta^{-1}(\mathbb{P} \setminus Q)}$.*

Proof. The product $\mathbb{P} = \prod_{i=1}^k \mathbb{P}(V_i)$ comes equipped with the i -th projection map $\pi_i : \mathbb{P} \rightarrow \mathbb{P}(V_i)$. On the product \mathbb{P} , consider the pull-back vector bundles

$$\varepsilon_i := \pi_i^* \mathcal{Q}_{\mathbb{P}(V_i)}(\omega_1, \dots, \omega_{i-1}, \omega_i - 1, \omega_{i+1}, \dots, \omega_k) \quad \forall i \in [k].$$

The fiber of each vector bundle ε_i at $[\mathbf{x}] = [\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}]$ is isomorphic to

$$\text{Hom}(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i - 1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}, V_i / \langle \mathbf{x}_i \rangle).$$

Let $\omega_i = (\omega_1, \dots, \omega_i - 1, \dots, \omega_k)$. Every tensor $T \in S^{\omega_i} V$ induces a section s_T of ε_i which corresponds to the composition

$$(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) \xrightarrow{i} S^{\omega_i} V \xrightarrow{T} V_i \xrightarrow{\pi} \frac{V_i}{\langle \mathbf{x}_i \rangle}.$$

This section vanishes on $[\mathbf{x}]$ if and only if

$$\text{rank} \left(T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) \right) \leq 1.$$

Define the vector bundle $\varepsilon := \bigoplus_{i=1}^k \varepsilon_i$ on \mathbb{P} . For every $T \in S^{\omega} V$, the diagonal section $(s_T, \dots, s_T) \in H^0(\varepsilon)$ vanishes on \mathbf{x} if and only if $(\mathbf{x}_1, \dots, \mathbf{x}_k)$ is a partially symmetric singular k -tuple of T .

Using the vector bundle ε and the natural projections $\tilde{\alpha}: \Pi_{\mathbf{n},\omega} \rightarrow \mathbb{P}(S^{\omega} V)$ and $\tilde{\beta}: \Pi_{\mathbf{n},\omega} \rightarrow \mathbb{P}$, we introduce the *Friedland-Ottaviani vector bundle* on $\Pi_{\mathbf{n},\omega}$:

$$E := \tilde{\alpha}^*(\mathcal{O}_{\mathbb{P}(S^{\omega} V)}(1)) \otimes \tilde{\beta}^*(\varepsilon).$$

The vector bundle E has rank $\sum_i (n_i - 1)$ and was first introduced by Friedland and Ottaviani for studying singular vector k -tuples; see (Friedland and Ottaviani, 2014, Theorem 12). The vanishing locus $Z(s)$ of the section $s \in H^0(E)$ given by the map $T \mapsto (s_T, \dots, s_T)$ equals the spectral variety $\Sigma_{\mathbf{n},\omega} \subset \Pi_{\mathbf{n},\omega}$ (Friedland and Ottaviani, 2014, Lemma 9).

The variety $\beta^{-1}(\mathbb{P} \setminus Q)$ is an irreducible component of $\Sigma_{\mathbf{n},\omega}$ by Lemma 17(i). By Theorem 13 and Lemma 17(ii), if some another irreducible component of $\Sigma_{\mathbf{n},\omega}$ exists, its codimension must be strictly higher than $\text{codim}(\beta^{-1}(\mathbb{P} \setminus Q)) = \sum_i (n_i - 1) = \text{rank}(E)$. In any affine open $U \subset \Pi_{\mathbf{n},\omega}$ trivializing the vector bundle E , $Z(s)|_U = \Sigma_{\mathbf{n},\omega}|_U$ is defined by the vanishing of $\text{rank}(E)$ equations. Therefore, by Krull’s principal ideal theorem, on the affine chart $U \subset X$ there cannot be any irreducible component of $\Sigma_{\mathbf{n},\omega}|_U$ whose codimension is strictly higher than $\text{rank}(E)$. Gluing an affine open cover of $\Pi_{\mathbf{n},\omega}$ trivializing E , then one finds that there cannot be other irreducible components of $\Sigma_{\mathbf{n},\omega}$ besides $\beta^{-1}(\mathbb{P} \setminus Q)$. This proves the equality and irreducibility. \square

We are ready to prove Theorem 1.

Proof of Theorem 1. Consider the product $\Pi_{\mathbf{n},\omega}$ and the spectral variety $\Sigma_{\mathbf{n},\omega}(Z)$ restricted to Z . Moreover, we consider the projections β_Z and α_Z of diagram (2.10). By Theorem 13 and Lemma 17(ii), the irreducible components of $\Sigma_{\mathbf{n},\omega}(Z)$ are, up to permuting factors, of the form $Y'_j = \beta_Z^{-1}(Z'_j)$, where Z'_j is an irreducible component of $Z \cap Q_j$ for some $j \subset [k]$, as $Z \cap Q_j$ might be reducible. Note that the generalized Kalman variety is $\kappa_{\mathbf{n},\omega}(Z) = \alpha_Z(\Sigma_{\mathbf{n},\omega}(Z))$. Therefore the irreducible components of the generalized Kalman variety are of the form $\alpha_Z(Y'_j)$, each one of those is a Kalman stratum.

From the dimensions calculated in Proposition 18, it follows that the unique highest-dimensional irreducible component of $\kappa_{\mathbf{n},\omega}(Z)$ is the normalized Kalman variety $\kappa_{\mathbf{n},\omega}^{nor}(Z)$. Therefore codimension and degree of the generalized Kalman variety coincide with the ones of $\kappa_{\mathbf{n},\omega}^{nor}(Z)$.

Again by Lemma 17(ii), $\Sigma_{\mathbf{n},\omega}(Z)$ is connected. Since α_Z is continuous, $\kappa_{\mathbf{n},\omega}(Z)$ is connected.

We are left with the computation of $\text{deg}(\kappa_{\mathbf{n},\omega}(Z)) = \text{deg}(\kappa_{\mathbf{n},\omega}^{nor}(Z))$. As in the proof of Proposition 19, we work with the Friedland-Ottaviani vector bundle E of rank $\sum_i (n_i - 1)$ on $\Pi_{\mathbf{n},\omega}$.

Denote by $D = \prod_{i=1}^k \binom{n_i + \omega_i - 1}{\omega_i} - 1$ the dimension of $\mathbb{P}(S^{\omega} V)$. We recall that α and β appearing in diagram (2.6) are the restrictions of $\tilde{\alpha}$ and $\tilde{\beta}$ to $\Sigma_{\mathbf{n},\omega}$. By Proposition 19, $\Sigma_{\mathbf{n},\omega} = Z(s) = \beta^{-1}(\mathbb{P} \setminus Q)$ and has codimension $\sum_i (n_i - 1)$ in $\Pi_{\mathbf{n},\omega}$. Therefore the top Chern class $c_{\sum_i (n_i - 1)}(E)$ of E satisfies $c_{\sum_i (n_i - 1)}(E) = [Z(s)] \in A^*(\Pi_{\mathbf{n},\omega})$, where $A^*(\Pi_{\mathbf{n},\omega}) = \mathbb{C}[h, t_1, \dots, t_k]/(h^{D+1}, t_1^{n_1}, \dots, t_k^{n_k})$ ($h = c_1(\mathcal{O}_{\mathbb{P}(S^{\omega} V)}(1))$ and $t_i = c_1(\mathcal{O}_{\mathbb{P}(V_i)}(1))$) is the Chow ring of $\Pi_{\mathbf{n},\omega}$.

We now look at the intersection $Z(s) \cap \beta^{-1}(Z) \subset \Pi_{\mathbf{n},\omega}$, which may be reducible. Note that this intersection contains $\beta^{-1}(Z \setminus Q)$, which has codimension $\sum_i (n_i - 1) + \text{codim}(Z) = \sum_i (n_i - 1) + d$. We now check whether the intersection contains other components whose codimension is smaller

than or equal to the codimension of $\overline{\beta^{-1}(Z \setminus Q)}$. By Lemma 17(iii), $Z(s) \setminus \beta^{-1}(\mathbb{P} \setminus Q)$ is a union of quasi-projective varieties of the form $\beta^{-1}(\prod_{j \in J} Q_j \times \prod_{j \notin J} [\mathbb{P}(V_j) \setminus Q_j])$. Therefore $(Z(s) \setminus \beta^{-1}(\mathbb{P} \setminus Q)) \cap \beta^{-1}(Z)$ is a union of quasi-projective varieties of the form $\beta^{-1}(\prod_{j \in J} Q_j \cap Z_j \times \prod_{j \notin J} [Z_j \setminus Q_j])$. As in the proof of Proposition 18, all these quasi-projective varieties have dimensions strictly smaller than the dimension of $\overline{\beta^{-1}(Z \setminus Q)}$. In conclusion, the intersection $Z(s) \cap \beta^{-1}(Z) \subset \Pi_{\mathbf{n}, \omega}$ contains a unique irreducible component in codimension $\sum_i (n_i - 1) + \delta$; the other potential components are in strictly higher codimensions.

The previous paragraph shows that the element $c_{\sum_i (n_i - 1)}(E) \prod_{i=1}^k \text{deg}(Z_i) \cdot \beta^*(\pi_i^*(\mathcal{O}_{Z_i}(1)^{\delta_i}))$ of the Chow ring is nonzero in codimension $\sum_i (n_i - 1) + \delta$ and zero in smaller codimension. Thus the intersection number

$$\alpha^*(c_1(\mathcal{O}_{\mathbb{P}(S \otimes V)}(1))^{D-\delta} \cdot c_{\sum_i (n_i - 1)}(E) \prod_{i=1}^k \text{deg}(Z_i) \cdot \beta^*(\pi_i^*(\mathcal{O}_{Z_i}(1)^{\delta_i}))$$

in the Chow ring $A^*(\Pi_{\mathbf{n}, \omega})$ is the degree of $\alpha(\overline{\beta^{-1}(Z \setminus Q)}) = \kappa_{\mathbf{n}, \omega}^{\text{nor}}(Z)$.

The Friedland-Ottaviani vector bundle E is the direct sum of k summands. By Euler’s exact sequence, each of these summands has Chern polynomial

$$\frac{(1 + \hat{t}_i + h)^{n_i}}{1 + \hat{t}_i + h - t_i} = \frac{(1 + \hat{t}_i + h)^{n_i - 1}}{1 - \frac{t_i}{1 + \hat{t}_i + h}} = \sum_{j=0}^{\infty} (1 + \hat{t}_i + h)^{n_i - 1 - j} t_i^j = \sum_{j=0}^{n_i - 1} (1 + \hat{t}_i + h)^{n_i - 1 - j} t_i^j,$$

where $\hat{t}_i := (\sum_{j=1}^k \omega_j t_j) - t_i$. The statement then follows as in the proof of (Ottaviani and Shahidi, 2021, Theorem 3.4), noting that we carry as extra factor the product of the degrees $\text{deg}(Z_i)$. \square

Remark 20. We do not know whether the generalized Kalman variety $\kappa_{\mathbf{n}, \omega}(Z)$, under the assumptions of Theorem 1, is always irreducible. The issue is that we do not know in general how the Kalman strata intersect.

The next examples suggest that these varieties might be subtle (even for matrices) and show irreducibility is not true when the assumptions $Z_i \not\subset Q_i$ of Theorem 1 are weakened.

Example 21. Let $k = 2$, $\mathbf{n} = (2, 3)$, and $\omega = (1, 1)$. We assume that $\mathbb{P}(V_1) \cong \mathbb{P}_{\mathbb{C}}^1$, $\mathbb{P}(V_2) \cong \mathbb{P}_{\mathbb{C}}^2$, $Q_1 = \mathcal{V}(x_{1,1}^2 + x_{1,2}^2) \subset \mathbb{P}_{\mathbb{C}}^1$ and $Q_2 = \mathcal{V}(x_{2,1}^2 + x_{2,2}^2 + x_{2,3}^2) \subset \mathbb{P}_{\mathbb{C}}^2$. On one hand, the product $\mathbb{P}_{\mathbb{C}}^1 \times Q_2$ is irreducible in $\mathbb{P}_{\mathbb{C}}^1 \times \mathbb{P}_{\mathbb{C}}^2$. On the other hand, the generalized Kalman variety $\kappa_{\mathbf{n}}(\mathbb{P}_{\mathbb{C}}^1 \times Q_2)$ is a degree 8 hypersurface with three irreducible components:

- (i) the totally isotropic Kalman variety $\kappa_{\mathbf{n}}^{\text{iso}}$, that is a degree 4 hypersurface with two irreducible components. Using coordinates a_{ij} for the space $\mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^3) \cong \mathbb{P}_{\mathbb{C}}^5$, its equation is

$$4(a_{11}a_{21} + a_{12}a_{22} + a_{13}a_{23})^2 + (a_{11}^2 + a_{12}^2 + a_{13}^2 - a_{21}^2 - a_{22}^2 - a_{23}^2)^2.$$

The first summand corresponds to the Euclidean inner product of the two rows of $A = (a_{ij}) \in \mathbb{C}^2 \otimes \mathbb{C}^3$. The second summand is the difference between the Euclidean norms of the two rows of A .

- (ii) the Kalman stratum $\kappa_{\mathbf{n}}^{[2]}(\mathbb{P}_{\mathbb{C}}^1 \times Q_2)$, that is a degree 4 irreducible hypersurface of equation

$$(a_{11}a_{22} - a_{12}a_{21})^2 + (a_{11}a_{23} - a_{13}a_{21})^2 + (a_{12}a_{23} - a_{13}a_{22})^2,$$

namely the sum of squares of the three maximal minors of A .

If $\mathbf{n} = (2, 4)$, then the generalized Kalman variety $\kappa_{\mathbf{n}}(\mathbb{P}_{\mathbb{C}}^1 \times Q_2)$, where $Q_2 = \mathcal{V}(x_{2,1}^2 + \dots + x_{2,4}^2) \subset \mathbb{P}_{\mathbb{C}}^3$, is a degree 8 hypersurface in $\mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^4) \cong \mathbb{P}_{\mathbb{C}}^7$ with three analogous irreducible components. For $\mathbf{n} = (3, 3)$, one finds $\kappa_{\mathbf{n}}(\mathbb{P}_{\mathbb{C}}^2 \times Q_2) = \kappa_{\mathbf{n}}^{\text{iso}} = \kappa_{\mathbf{n}}(Q_1 \times \mathbb{P}_{\mathbb{C}}^2)$.

Example 22. Let us consider the matrix case $k = 2$, $\mathbf{n} = (4, 4)$, $\delta = (2, 1)$ and $\omega = (1, 1)$. Consider the twisted cubic curve $Z_1 \subset \mathbb{P}_{\mathbb{C}}^3$ and a general quadric surface $Z_2 \subset \mathbb{P}(V_2) \cong \mathbb{P}_{\mathbb{C}}^3$. Applying Theorem 1, we have

$$\deg(\kappa_{\mathbf{n}}^{nor}(Z)) = \deg(Z_1) \deg(Z_2) d((4, 4), (2, 1), (1, 1)),$$

where $d((4, 4), (2, 1), (1, 1))$ is the coefficient of the monomial $h^3 t_1 t_2^2$ in the polynomial

$$\sum_{i,j=0}^3 (t_1 + h)^{3-j} (t_2 + h)^{3-1} t_1^i t_2^j = \dots + 20h^3 t_1 t_2^2 + \dots,$$

therefore $\deg(\kappa_{\mathbf{n}}^{nor}(Z)) = 3 \cdot 2 \cdot 20 = 120$. We provide a Macaulay2 code Grayson and Stillman (1997) to verify symbolically this degree computation. We speed up our degree computation by working over the finite field $\mathbb{Z}/\mathbb{Z}_{101}$ and restricting to a 3-dimensional subspace $H \subset \mathbb{P}(\mathbb{C}^4 \otimes \mathbb{C}^4)$, since $\kappa_{\mathbf{n}}^{nor}(Z)$ has codimension $2 + 1 = 3$:

```
K = ZZ/101; R = K[u_0..u_3, x_(1,1)..x_(2,4)];
coeffs = toList((1,1)..(4,4));
for f in coeffs do a_f = sum(m+1, i-> random(K)*u_i);
M = sum(4, i-> sum(4, j-> a_(i+1,j+1)*x_(1,i+1)*x_(2,j+1)));
xx1 = matrix{{x_(1,1)..x_(1,4)}}, xx2 = matrix{{x_(2,1)..x_(2,4)}};
I = minors(2, contract(xx1,M) | |xx1) + minors(2, contract(xx2,M) | |xx2) +
    ideal(sum(4, i-> x_(1,i+1)^2)-1, sum(4, i-> x_(2,i+1)^2)-1);
```

Here I is the ideal of relations among the singular vector pairs in the symbolic coordinates of M . We impose the existence of a singular vector pair on $Z = Z_1 \times Z_2$ as follows:

```
Z_1 = minors(2, matrix{{x_(1,1)..x_(1,3)}, {x_(1,2)..x_(1,4)}});
Z_2 = ideal((symmetricPower(2,xx2)*random(K^10,K^1))_(0,0));
J = I + Z_1 + Z_2;
```

Finally, the ideal of $\kappa_{\mathbf{n}}^{nor}(Z)$ is computed with

```
Kalman = eliminate(toList(x_(1,1)..x_(2,4)), J);
degree Kalman -- this confirms that the variety has degree 120
```

Remark 23. Note that Theorem 1 includes the symmetric case. Consider an irreducible variety $Z \subset \mathbb{P}(V)$ not contained in the isotropic quadric $Q \subset \mathbb{P}(V)$ and the generalized Kalman variety $\kappa_{\mathbf{n},\omega}(Z) := \kappa_{\mathbf{n},\omega}(Z)$, where $\delta = (\delta)$, $\delta = \text{codim}(Z)$, $\mathbf{n} = (n)$ and $\omega = (\omega)$. Its degree is $d(n, \delta, \omega) \deg(Z)$, where $d(n, \delta, \omega)$ is the coefficient of the monomial $h^\delta t^{n-\delta-1}$ in the polynomial

$$\begin{aligned} \frac{[(\omega - 1)t + h]^n - t^n}{(\omega - 1)t + h - t} &= \sum_{i=0}^{n-1} [(\omega - 1)t + h]^{n-1-i} t^i \\ &= \sum_{i=0}^{n-1} \sum_{j=0}^{n-1-i} \binom{n-1-i}{j} (\omega - 1)^{n-1-i-j} t^{n-1-j} h^j. \end{aligned}$$

The coefficient of the monomial $h^\delta t^{n-\delta-1}$ in the last polynomial is

$$\sum_{i=0}^{n-1} \binom{n-1-i}{\delta} (\omega - 1)^{n-\delta-1-i} = \sum_{i=0}^{n-\delta-1} \binom{n-1-i}{\delta} (\omega - 1)^{n-\delta-1-i} = \sum_{j=0}^{n-\delta-1} \binom{\delta+j}{j} (\omega - 1)^j,$$

which recovers (Ottaviani and Shahidi, 2021, Theorem 2.3).

3. The totally isotropic Kalman variety

In Theorem 1 we assumed that each Z_j is not contained in the corresponding isotropic quadric Q_j . On the opposite side, we exhibit a description of totally isotropic Kalman varieties as dual varieties. Before we prove the precise statement, we need the following lemma.

Lemma 24. Consider the product $Q = \prod_{i=1}^k Q_i$ of isotropic quadrics $Q_i \subset \mathbb{P}(V_i)$. Let $v_{\mathbf{n},\omega}: \mathbb{P} \rightarrow \mathbb{P}(S^\omega V)$ be the degree- ω Segre-Veronese embedding of $\mathbb{P} = \prod_{i=1}^k \mathbb{P}(V_i)$. Then the dual variety $[v_{\mathbf{n},\omega}(Q)]^\vee$ is a hypersurface for every choice of \mathbf{n} and ω .

Proof. Considering a slight modification of (Gel'fand et al., 1994, Chapter 1, Corollary 5.10), the variety $[v_{\mathbf{n},\omega}(Q)]^\vee$ is a hypersurface if and only if

$$\dim(v_{n_j,\omega_j}(Q_j)) + \text{codim}([v_{n_j,\omega_j}(Q_j)]^\vee) - 1 \leq \dim(v_{\mathbf{n},\omega}(Q)) \quad \forall j \in [k], \tag{3.1}$$

where each variety $v_{n_j,\omega_j}(Q_j)$, which is the degree- ω_j Veronese embedding of Q_j . The variety $[v_{n_j,\omega_j}(Q_j)]^\vee$ is a hypersurface for all $\omega_j \geq 1$ by (Sodomaco, 2018, Corollary 4.8). Therefore, for all $j \in [k]$ the corresponding inequality in (3.1) becomes $n_j - 2 \leq \dim(v_{\mathbf{n},\omega}(Q)) = n_1 + \dots + n_k - 2k$, which is clearly satisfied. \square

Theorem 25. Assume $n_j \geq 3$ for all $j \in [k]$. Consider the product $Q = \prod_{i=1}^k Q_i$ of isotropic quadrics $Q_i \subset \mathbb{P}(V_i)$. Let $\kappa_{\mathbf{n},\omega}^{\text{iso}}$ be the totally isotropic Kalman variety, i.e., the variety of partially symmetric tensors having a singular vector k -tuple in Q . Let $v_{\mathbf{n},\omega}: \mathbb{P} \rightarrow \mathbb{P}(S^\omega V)$ be the degree- ω Segre-Veronese embedding of $\mathbb{P} = \prod_{i=1}^k \mathbb{P}(V_i)$. Then

$$\kappa_{\mathbf{n},\omega}^{\text{iso}} = [v_{\mathbf{n},\omega}(Q)]^\vee. \tag{3.2}$$

In particular, $\kappa_{\mathbf{n},\omega}^{\text{iso}}$ is an irreducible hypersurface of $\mathbb{P}(S^\omega V)$. Its degree is equal to the degree of $[v_{\mathbf{n},\omega}(Q)]^\vee$, that is

$$2^k \sum_{j=0}^N (-1)^j (N+1-j)! \sum_{|\alpha|=j} \left[\prod_{l=1}^k \frac{\omega_l^{n_l-2-\alpha_l}}{(n_l-2-\alpha_l)!} \sum_{\beta_l=0}^{\alpha_l} \prod_{l \in [k]} \binom{n_l}{\beta_l} (-2)^{\alpha_l-\beta_l} \right] \tag{3.3}$$

where $N = \dim(v_{\mathbf{n},\omega}(Q)) = n_1 + \dots + n_k - 2k$.

If $J := \{j \in [k] \mid n_j = 2\} \neq \emptyset$, then identity (3.2) still holds. In such a case, $\kappa_{\mathbf{n},\omega}^{\text{iso}}(Q)$ has $2^{|J|}$ irreducible components, where each one of them is isomorphic to the Kalman variety of the irreducible product $\prod_{j \notin J} Q_j$.

Proof. A tensor $T \in \mathbb{P}(S^\omega V)$ is a hyperplane $H_T \subset \mathbb{P}(S^\omega V)^*$. Recall that we identify all the vector spaces with their duals using the Frobenius inner product.

Suppose that, for some k -tuple $([\mathbf{x}_1], \dots, [\mathbf{x}_k]) \in Q$, the affine tangent space $\widehat{T}_{[\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}]} v_{\mathbf{n},\omega}(Q)$ at $[\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_k^{\omega_k}]$ to $v_{\mathbf{n},\omega}(Q)$ is a subspace of the affine cone \widehat{H}_T of H_T . Any vector \mathbf{v} of this affine tangent space may be written as

$$\mathbf{v} = \sum_{i=1}^k \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{y}_i \otimes \dots \otimes \mathbf{x}_k^{\omega_k}, \quad \mathbf{y}_i \in \widehat{T}_{[\mathbf{x}_i]} Q_i \quad \forall i \in [k].$$

By linearity, we may suppose that $\mathbf{v} = \mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{y}_i \otimes \dots \otimes \mathbf{x}_k^{\omega_k}$ for some $i \in [k]$. Then $\mathbf{v} \in \widehat{H}_T$ for every choice of $\mathbf{y}_i \in \widehat{T}_{[\mathbf{x}_i]} Q_i$. This implies that the contraction $T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{y}_i \otimes \dots \otimes \mathbf{x}_k^{\omega_k})$ vanishes for every $\mathbf{y}_i \in \widehat{T}_{[\mathbf{x}_i]} Q_i$. Equivalently, the vector $\mathbf{w}_i := T(\mathbf{x}_1^{\omega_1} \otimes \dots \otimes \mathbf{x}_i^{\omega_i-1} \cdot \mathbf{e}_i \otimes \dots \otimes \mathbf{x}_k^{\omega_k}) \in V_i$ is contained in the affine conormal space $\widehat{N}_{[\mathbf{x}_i]} Q_i = (\widehat{T}_{[\mathbf{x}_i]} Q_i)^\perp$. Observing that

$$\widehat{N}_{[\mathbf{x}_i]}Q_i = \left\{ \mathbf{y}_i \in V_i \mid \text{rank} \begin{pmatrix} \nabla q_i(\mathbf{x}_i) \\ \mathbf{y}_i \end{pmatrix} \leq 1 \right\} = \left\{ \mathbf{y}_i \in V_i \mid \text{rank} \begin{pmatrix} \mathbf{x}_i \\ \mathbf{y}_i \end{pmatrix} \leq 1 \right\} = \langle \mathbf{x}_i \rangle,$$

we conclude that the vector \mathbf{w}_i defined above is proportional to the vector \mathbf{x}_i . Applying this argument for each $i \in [k]$, we find that $([\mathbf{x}_1], \dots, [\mathbf{x}_k])$ is a singular vector k -tuple for the tensor T . This means that $T \in \kappa_{\mathbf{n},\omega}^{iso}$. Recall that

$$[v_{\mathbf{n},\omega}(Q)]^\vee = \overline{\{T \in \mathbb{P}(S^\omega V) \mid \widehat{T}_{[\mathbf{x}]}v_{\mathbf{n},\omega}(Q) \subset H_T \text{ for some } [\mathbf{x}] \in v_{\mathbf{n},\omega}(Q)\}}.$$

Upon taking closures, we have proven the inclusion $[v_{\mathbf{n},\omega}(Q)]^\vee \subset \kappa_{\mathbf{n},\omega}^{iso}$. The variety $[v_{\mathbf{n},\omega}(Q)]^\vee$ is a hypersurface for every choice of \mathbf{n} and ω by Lemma 24. On the other hand, the totally isotropic Kalman variety $\kappa_{\mathbf{n},\omega}^{iso}$ is an irreducible hypersurface if $n_j \geq 3$ for all $j \in [k]$. Hence equality $\kappa_{\mathbf{n},\omega}^{iso} = [v_{\mathbf{n},\omega}(Q)]^\vee$ follows.

Since $[v_{\mathbf{n},\omega}(Q)]^\vee$ is a hypersurface, its degree is equal to the polar class $\delta_0(v_{\mathbf{n},\omega}(Q))$ by (Holme, 1988, Theorem 3.4). This invariant can be computed using the Chern classes of $v_{\mathbf{n},\omega}(Q)$ by the relation

$$\delta_0(v_{\mathbf{n},\omega}(Q)) = \sum_{j=0}^N (-1)^j (N+1-j) c_j(v_{\mathbf{n},\omega}(Q)) \cdot h^{N-j},$$

where $h = c_1(\mathcal{O}_{v_{\mathbf{n},\omega}(Q)}(1))$ is the hyperplane class. This computation was done in (Sodomaco, 2020, Proposition 5.3.1) in a slightly more general setting and leads to the degree (3.3).

Suppose $J \neq \emptyset$. For $1 \leq j \leq 2^{|J|}$, define Y_j to be the j th irreducible component of Q . Notice that each Y_j is isomorphic to a product of quadrics. We now describe $[v_{\mathbf{n},\omega}(Q)]^\vee$ in terms of the dual varieties $[v_{\mathbf{n},\omega}(Y_j)]^\vee$.

By definition, $v_{\mathbf{n},\omega}(Q) = \bigcup_{j \in J} v_{\mathbf{n},\omega}(Y_j)$. Consider the conormal variety $\mathcal{N}_{v_{\mathbf{n},\omega}(Q)}$ and the diagram

$$\begin{array}{ccc}
 & \mathcal{N}_{v_{\mathbf{n},\omega}(Q)} & \\
 p_1 \swarrow & & \searrow p_2 \\
 v_{\mathbf{n},\omega}(Q) & & \mathbb{P}(S^\omega V).
 \end{array} \tag{3.4}$$

The preimage under p_1 of each irreducible component $v_{\mathbf{n},\omega}(Y_j)$ is a projective bundle and therefore irreducible (Vakil, 2017, Exercise 11.4.C). The image of such a component is the irreducible hypersurface $[v_{\mathbf{n},\omega}(Y_j)]^\vee$. This shows that $[v_{\mathbf{n},\omega}(Q)]^\vee = \bigcup_{j \in J} [v_{\mathbf{n},\omega}(Y_j)]^\vee$. To see that these irreducible components are all distinct, assume $[v_{\mathbf{n},\omega}(Y_j)]^\vee = [v_{\mathbf{n},\omega}(Y_i)]^\vee$ for some $i \neq j \in J$. Applying the dual construction again and the fact that every projective irreducible variety is reflexive in characteristic zero, we find that $v_{\mathbf{n},\omega}(Y_j) = [v_{\mathbf{n},\omega}(Y_j)]^{\vee\vee} = [v_{\mathbf{n},\omega}(Y_i)]^{\vee\vee} = v_{\mathbf{n},\omega}(Y_i)$, a contradiction, because Y_i and Y_j are distinct.

Now, the totally isotropic Kalman variety is $\kappa_{\mathbf{n},\omega}^{iso} = \bigcup_{j \in J} \kappa_{\mathbf{n},\omega}^{iso}(Y_j)$, where we do not know a priori that all the irreducible varieties in the union are distinct. However, by the first part of this proof, we find that $\kappa_{\mathbf{n},\omega}^{iso}(Y_j) = [v_{\mathbf{n},\omega}(Y_j)]^\vee$. Moreover, as we have checked, all the irreducible components $[v_{\mathbf{n},\omega}(Y_j)]^\vee$ are distinct. Therefore $\kappa_{\mathbf{n},\omega}^{iso} = \bigcup_{j \in J} [v_{\mathbf{n},\omega}(Y_j)]^\vee = [v_{\mathbf{n},\omega}(Q)]^\vee$ has $2^{|J|}$ components, where each of them is a dual variety of the corresponding irreducible quadric. \square

Example 26. Let $k = 2$ and $\mathbf{n} = (2, 2)$. We assume that $\mathbb{P}(V_1) = \mathbb{P}(V_2) \cong \mathbb{P}_\mathbb{C}^1$ and $Q_1 = Q_2 = \mathcal{V}(x_1^2 + x_2^2) \subset \mathbb{P}_\mathbb{C}^1$. One has

$$\kappa_{\mathbf{n}}(\mathbb{P}_\mathbb{C}^1 \times Q_2) = \kappa_{\mathbf{n}}^{iso} = \kappa_{\mathbf{n}}(Q_1 \times \mathbb{P}_\mathbb{C}^1).$$

This is a reducible hypersurface of degree 4 whose irreducible components are four planes in $\mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2) \cong \mathbb{P}_\mathbb{C}^3$, dual to the four points of $Q_1 \times Q_2$ as predicted by Theorem 25. Denoting a_{ij} the homogeneous coordinates of $\mathbb{P}_\mathbb{C}^3$, it is defined by the vanishing of the polynomial $[(a_{11} - a_{22})^2 + (a_{12} + a_{21})^2][(a_{11} + a_{22})^2 + (a_{12} - a_{21})^2]$. There are four lines in $\mathbb{P}_\mathbb{C}^3$ obtained as the

intersection of the four pairs of non-conjugate planes. Each of the Kalman strata $\kappa_n^{(1)}(Q_1 \times \mathbb{P}_C^1) = \kappa_n^{(1)}(\mathbb{P}_C^1 \times \mathbb{P}_C^1)$ and $\kappa_n^{(2)}(\mathbb{P}_C^1 \times Q_2) = \kappa_n^{(2)}(\mathbb{P}_C^1 \times \mathbb{P}_C^1)$ consists of two such lines. Therefore, both these Kalman strata have degree 2 and codimension 2.

In the symmetric case, Theorem 25 specializes to the following result, which is related to (Beorchia et al., 2021, Proposition 2.10).

Theorem 27. *Let $\kappa_{n,\omega}^{iso}$ be the Kalman variety of symmetric tensors having an isotropic eigenvector. Let $v_{n,\omega}: \mathbb{P}(V) \rightarrow \mathbb{P}(S^\omega V)$ be the degree- ω Veronese embedding of $\mathbb{P}(V)$. Then*

$$\kappa_{n,\omega}^{iso} = [v_{n,\omega}(Q)]^\vee.$$

In particular, $\kappa_{n,\omega}^{iso}$ is an irreducible hypersurface of $\mathbb{P}(S^\omega V)$. Its degree is

$$\deg(\kappa_{n,\omega}^{iso}) = 2 \sum_{j=0}^{n-2} (j+1)(\omega-1)^j. \tag{3.5}$$

Example 28. Consider the totally isotropic Kalman variety $\kappa_{3,2}^{iso}$ of ternary quadrics possessing an isotropic eigenvector. By Theorem 27, this is an irreducible hypersurface of degree 6 in $\mathbb{P}(S^\omega V) \cong \mathbb{P}^5$. We include a Macaulay2 code to verify this, using both descriptions according to Theorem 27. Below we denote the homogeneous coordinates of $\mathbb{P}(S^\omega V)$ by a_0, \dots, a_5 .

As Kalman variety:

```
R = QQ[x_1..x_3, a_0..a_5, c_0..c_5];
MX = matrix{{x_1^2, 2*x_1*x_2, 2*x_1*x_3, x_2^2, 2*x_2*x_3, x_3^2}};
aa = matrix{{a_0..a_5}}; xx = matrix{{x_1..x_3}};
f = (MX*transpose(aa))_(0,0);
Ivect = minors(2, diff(xx, f) || xx);
IQ = ideal(sum(3, i-> x_(i+1)^2));
sat = saturate(Ivect+IQ, ideal xx);
Kalman = eliminate(first entries xx, sat);
```

As dual variety $[v_{3,2}(Q)]^\vee$:

```
cc = matrix{{c_0..c_5}};
IVQ = eliminate(first entries xx,
  saturate(ideal(first entries (MX-aa))+IQ, ideal xx));
jacIVQ = diff(aa, transpose gens IVQ);
norIVQ = saturate(IVQ + minors(codim(IVQ)+1, jacIVQ | cc), ideal aa);
dualIVQ = eliminate(first entries aa, norIVQ);
eq = sub(dualIVQ_0, apply(6, i-> c_i=>a_i));
eq == Kalman_0 -- the two equations coincide
```

We point that the defining polynomial eq of $[v_{n,\omega}(Q)]^\vee$ has a role in the theory of characteristic polynomials of symmetric tensors. The second author showed that the leading coefficient of the characteristic polynomial of a symmetric tensor in $S^\omega V$ is equal, up to scaling, to the defining polynomial of $[v_{n,\omega}(Q)]^\vee$ with multiplicity $\omega - 2$ Sodomaco (2018).

4. Generating functions

A generating function is a device somewhat similar to a bag. Instead of carrying many little objects detachedly, which could be embarrassing, we put them all in a bag, and then we have only one object to carry, the bag.

[George Pólya (Pólya, 1954, Chapter VI)]

In the previous section, we determined the degrees of the generalized Kalman varieties $\kappa_{\mathbf{n},\omega}(Z)$ by computing the coefficients $d(\mathbf{n}, \delta, \omega)$. In the spirit of Pólya’s quote, Theorem 2 furnishes a generating function for the coefficients $d(\mathbf{n}, \delta, \omega)$ when $\delta = (\delta, 0, \dots, 0)$, or equivalently when we consider only one subvariety $Z_1 \subset \mathbb{P}(\mathbb{C}^{n_1})$ of codimension δ . For this particular choice of δ , we use the notation $d(\mathbf{n}, \delta, \omega)$ in place of $d(\mathbf{n}, \delta, \omega)$. Before proceeding to the proof of Theorem 2, we start with a lemma.

Lemma 29. Let $d(\mathbf{m}) = \sum_{j_1=0}^{m_1-1} \dots \sum_{j_k=0}^{m_k-1} f(j_1, \dots, j_k)$. Then

$$\sum_{\mathbf{m} \in \mathbb{N}^k} d(\mathbf{m}) \mathbf{x}^{\mathbf{m}} = \left(\prod_{i=1}^k \frac{x_i}{1-x_i} \right) \sum_{\mathbf{j} \in \mathbb{N}^k} f(\mathbf{j}) \mathbf{x}^{\mathbf{j}}.$$

Proof. The proof is an induction on k . For $k = 1$, we have

$$\begin{aligned} \sum_{m=0}^{\infty} \left(\sum_{j=0}^{m-1} f(j) \right) x^m &= f(0)x + (f(0) + f(1))x^2 + (f(0) + f(1) + f(2))x^3 + \dots \\ &= (x + x^2 + x^3 + \dots) \left(f(0) + f(1)x + f(2)x^2 + \dots \right) \\ &= \frac{x}{1-x} \sum_{j=0}^{\infty} f(j)x^j. \end{aligned}$$

The induction step is similar. \square

As in Zeilberger’s approach Ekhad and Zeilberger (2016), we shall employ a classical and powerful theorem of MacMahon (MacMahon, 1915, §3, Chapter 2, 66).

Theorem 30 (MacMahon Master Theorem). Let $A = (a_{ij})$ be an $m \times m$ complex matrix, and let $\mathbf{z} = (z_1, \dots, z_m)$ be a vector of formal variables. Let $f(\mathbf{p})$ be the coefficient of the monomial $\mathbf{z}^{\mathbf{p}}$ in the product $\prod_{i=1}^m (a_{i1}z_1 + \dots + a_{im}z_m)^{p_i}$. Let $\mathbf{w} = (w_1, \dots, w_m)$ be another vector of formal variables, $T = \text{diag}(\mathbf{w})$ and denote by I_m the identity matrix of size m . Then

$$\sum_{\mathbf{p} \in \mathbb{N}^m} f(\mathbf{p}) \mathbf{w}^{\mathbf{p}} = \frac{1}{\det(I_m - TA)}.$$

We are now ready to prove Theorem 2.

Proof of Theorem 2. Recall that, for any vector of codimensions δ , the integer $d(\mathbf{n}, \delta, \omega)$ is the coefficient of the monomial $h^\delta \prod_{i=1}^k t_i^{n_i - \delta_i - 1}$ in the polynomial

$$\prod_{i=1}^k \frac{(\widehat{t}_i + h)^{n_i} - t_i^{n_i}}{(\widehat{t}_i + h) - t_i} = \prod_{i=1}^k \sum_{j_i=0}^{n_i-1} (\widehat{t}_i + h)^{j_i} t_i^{n_i-1-j_i},$$

where $\delta = \sum_{i=1}^k \delta_i$ and $\widehat{t}_i := \left(\sum_{j=1}^k \omega_j t_j \right) - t_i$. Equivalently, $d(\mathbf{n}, \delta, \omega)$ is the constant term of

$$\sum_{j_1=0}^{n_1-1} \dots \sum_{j_k=0}^{n_k-1} h^{-\delta} \prod_{i=1}^k (\widehat{t}_i + h)^{j_i} t_i^{\delta_i - j_i},$$

namely the sum of the constant terms in the products $h^{-\delta} \prod_{i=1}^k (\widehat{t}_i + h)^{j_i} t_i^{\delta_i - j_i}$. Observe that the constant term of each product $h^{-\delta} \prod_{i=1}^k (\widehat{t}_i + h)^{j_i} t_i^{\delta_i - j_i}$ is the coefficient of $h^\delta \prod_{i=1}^k t_i^{j_i}$ in $\prod_{i=1}^k (\widehat{t}_i + h)^{j_i} t_i^{\delta_i}$, which we may call $f(\mathbf{j}, \delta)$. Therefore

$$d(\delta, \mathbf{n}, \omega) = \sum_{j_1=0}^{n_1-1} \cdots \sum_{j_k=0}^{n_k-1} f(\mathbf{j}, \delta).$$

Applying Lemma 29 with respect to \mathbf{n} , we have

$$\sum_{\mathbf{n}, \delta \in \mathbb{N}^k} d(\delta, \mathbf{n}, \omega) \mathbf{x}^{\mathbf{n}} \mathbf{y}^{\delta} = \prod_{i=1}^k \frac{x_i}{1-x_i} \sum_{\mathbf{j}, \delta \in \mathbb{N}^k} f(\mathbf{j}, \delta) \mathbf{x}^{\mathbf{j}} \mathbf{y}^{\delta}.$$

From now on, we assume that $\delta_1 = \delta$ and $\delta_i = 0$ for all $i \geq 2$ and we call $f(\mathbf{j}, \delta)$ the coefficient of $h^{\delta} \prod_{i=1}^k t_i^{j_i}$ in $t_1^{\delta} \prod_{i=1}^k (\widehat{t}_i + h)^{j_i}$. In this special case, we can apply MacMahon's Theorem 30 with $m = k + 1$, $f(\mathbf{j}, \delta)$ in place of $f(\mathbf{p})$, and considering the product

$$t_1^{\delta} \prod_{i=1}^k (\widehat{t}_i + h)^{j_i} = t_1^{\delta} \prod_{i=1}^k (a_{i,1}t_1 + \cdots + a_{i,k}t_k + a_{i,k+1}h)^{j_i}.$$

Here $A = (a_{ij})$ is the $(k + 1) \times (k + 1)$ matrix defined as

$$A := \left(\begin{array}{cccc|c} & & & & 1 \\ & B & & & \vdots \\ & & & & 1 \\ \hline 1 & 0 & \cdots & 0 & 0 \end{array} \right),$$

where B is the $k \times k$ matrix whose (i, j) -th entry is $\omega_j - \delta_{ij}$ ($\delta_{ij} = 1$ if $i = j$, and 0 otherwise). Then

$$\sum_{\mathbf{j} \in \mathbb{N}^k} \sum_{\delta=0}^{\infty} f(\mathbf{j}, \delta) \mathbf{x}^{\mathbf{j}} \mathbf{y}^{\delta} = \frac{1}{\det(I_{k+1} - TA)},$$

where $T = \text{diag}(\mathbf{x}, y)$. Summing up, we have obtained the formula

$$\sum_{\mathbf{n} \in \mathbb{N}^k} \sum_{\delta=0}^{\infty} d(\mathbf{n}, \delta, \omega) \mathbf{x}^{\mathbf{n}} \mathbf{y}^{\delta} = \prod_{i=1}^k \frac{x_i}{1-x_i} \frac{1}{\det(I_{k+1} - TA)}. \tag{4.1}$$

It remains to compute $\det(I_{k+1} - TA)$. Define $M := I_{k+1} - TA$. More explicitly

$$M = \left(\begin{array}{cccc|cccc} 1 - (\omega_1 - 1)x_1 & -\omega_2x_1 & -\omega_3x_1 & \cdots & -\omega_{k-1}x_1 & -\omega_kx_1 & -x_1 \\ -\omega_1x_2 & 1 - (\omega_2 - 1)x_2 & -\omega_3x_2 & \cdots & -\omega_{k-1}x_2 & -\omega_kx_2 & -x_2 \\ \vdots & \vdots & \vdots & & \vdots & \vdots & \vdots \\ -\omega_1x_k & -\omega_2x_k & -\omega_3x_k & \cdots & -\omega_{k-1}x_k & 1 - (\omega_k - 1)x_k & -x_k \\ \hline -y & 0 & 0 & \cdots & 0 & 0 & 1 \end{array} \right).$$

By expanding along the last row, we derive two $k \times k$ submatrices

$$M' := \begin{pmatrix} -\omega_2x_1 & -\omega_3x_1 & \cdots & -\omega_kx_1 & -x_1 \\ 1 - (\omega_2 - 1)x_2 & -\omega_3x_2 & \cdots & -\omega_kx_2 & -x_2 \\ \vdots & \vdots & & \vdots & \vdots \\ -\omega_2x_k & -\omega_3x_k & \cdots & 1 - (\omega_k - 1)x_k & -x_k \end{pmatrix},$$

$$M'' := \begin{pmatrix} 1 - (\omega_1 - 1)x_1 & -\omega_2x_1 & \cdots & -\omega_kx_1 \\ -\omega_1x_2 & 1 - (\omega_2 - 1)x_2 & \cdots & -\omega_kx_2 \\ \vdots & \vdots & & \vdots \\ -\omega_1x_k & -\omega_2x_k & \cdots & 1 - (\omega_k - 1)x_k \end{pmatrix}.$$

The next step is to compute the determinants of M' and M'' .

Claim 1: We have

$$\det(M') = (-1)^k x_1 \prod_{i=2}^k (1 + x_i). \tag{4.2}$$

First of all, the polynomial $\det(M')$ has degree $\leq k$ in x_1, \dots, x_k . The first row of M' is a multiple of x_1 , hence x_1 divides $\det(M')$. Note that, if $x_i = -1$ then the first and the i -th row of M' are proportional and so $\det(M')$ vanishes. By Euclidean division, we may write $\det(M') = (1 + x_i)q + R$, where R is the remainder, which then depends only on the variables x_j with $j \neq i$. Hence $R(x_1, \dots, \widehat{x_i}, \dots, x_k) = 0$ for every value of the variables. Since we are in characteristic zero, $R = 0$. The left-hand side is divisible by $x_1 \prod_{i \geq 2} (1 + x_i)$ of degree k , and hence there is a scalar γ such that $\det(M') = \gamma x_1 \prod_{i \geq 2} (1 + x_i)$. In order to determine that $\gamma = (-1)^k$, we substitute $x_1 = 1$ and $x_2 = \dots = x_k = 0$ in M' and observe that the determinant of the resulting matrix is $(-1)^k$.

Claim 2: We have

$$\det(M'') = \prod_{i=1}^k (1 + x_i) - \sum_{j=1}^k \omega_j x_j \prod_{i \neq j} (1 + x_i). \tag{4.3}$$

Write $M'' = D - \mathbf{x}^T \boldsymbol{\omega}$, where $\mathbf{x} = (x_1, \dots, x_k)$ and $\boldsymbol{\omega} = (\omega_1, \dots, \omega_k)$ are row vectors and $D = \text{diag}(\mathbf{1} + \mathbf{x})$ is the diagonal matrix whose i -th diagonal entry is $1 + x_i$. Since D is invertible over the fraction field $\mathbb{C}(\mathbf{x})$, we have $M'' = D[I_k - (D^{-1} \mathbf{x}^T) \boldsymbol{\omega}]$. By the matrix determinant lemma (Hogben, 2007, §4.1, Fact 22) over the fraction field $\mathbb{C}(\mathbf{x})$, one has

$$\det(I_k - (D^{-1} \mathbf{x}^T) \boldsymbol{\omega}) = 1 - \boldsymbol{\omega} D^{-1} \mathbf{x}^T.$$

Thus

$$\det(M'') = \det(D - \mathbf{x}^T \boldsymbol{\omega}) = (1 - \boldsymbol{\omega} D^{-1} \mathbf{x}^T) \det(D) = \prod_{i=1}^k (1 + x_i) - \sum_{j=1}^k \omega_j x_j \prod_{i \neq j} (1 + x_i).$$

From the formulas (4.2) and (4.3), we conclude that

$$\begin{aligned} \det(I_{k+1} - TA) &= \det(M) = (-1)^{k+1} y \det(M') + \det(M'') = \\ &= -y x_1 \prod_{i=2}^k (1 + x_i) + \prod_{i=1}^k (1 + x_i) - \sum_{j=1}^k \omega_j x_j \prod_{i \neq j} (1 + x_i), \end{aligned}$$

which establishes the result. \square

Example 31. For $k = 2$ and $\boldsymbol{\omega} = (1, 1)$ we have

$$\begin{aligned} \sum_{\mathbf{n} \in \mathbb{N}^k} \sum_{\delta=0}^{\infty} d(\mathbf{n}, \delta, \boldsymbol{\omega}) \mathbf{x}^{\mathbf{n}} y^{\delta} &= \frac{x_1}{1-x_1} \cdot \frac{x_2}{1-x_2} \cdot \begin{vmatrix} 1 & -x_1 & -x_1 \\ -x_2 & 1 & -x_2 \\ -y & 0 & 1 \end{vmatrix}^{-1} \\ &= \frac{x_1}{1-x_1} \cdot \frac{x_2}{1-x_2} \cdot \frac{1}{1-x_1 y - x_1 x_2 - x_1 x_2 y} \\ &= 2yx_1^2 x_2^2 + 3y^2 x_1^3 x_2^2 + \dots \end{aligned}$$

This fits with (Ottaviani and Shahidi, 2021, Table 1). Note that for $y = 0$, when substituted into the first expression of the second line, we recover the formula in (Friedland and Ottaviani, 2014, Theorem 1).

Remark 32. If $\omega_1 = \dots = \omega_k = \omega$, the polynomial $H_\omega(\mathbf{x}, y)$ may be rewritten more symmetrically as

$$\begin{aligned} H_\omega(\mathbf{x}, y) &= -y x_1 \prod_{i=2}^k (1 + x_i) + \prod_{i=1}^k (1 + x_i) - \omega \sum_{j=1}^k x_j \prod_{i \neq j} (1 + x_i) \\ &= -y x_1 \sum_{i=0}^{k-1} e_i(\widehat{\mathbf{x}}_1) + \sum_{i=0}^k e_i(\mathbf{x}) - \omega \sum_{j=0}^{k-1} \sum_{i=1}^k x_i e_j(\widehat{\mathbf{x}}_i) \\ &= -y x_1 \sum_{i=0}^{k-1} e_i(\widehat{\mathbf{x}}_1) + \sum_{i=0}^k (1 - \omega i) e_i(\mathbf{x}), \end{aligned}$$

where $\widehat{\mathbf{x}}_i$ denotes the vector $(x_1, \dots, x_{i-1}, x_{i+1}, \dots, x_k)$ and for a vector $\mathbf{x} = (z_1, \dots, z_k)$, we define $e_0(\mathbf{x}) := 1$ and $e_i(\mathbf{x}) := \sum_{1 \leq j_1 < \dots < j_i \leq k} z_{j_1} \cdots z_{j_i} \quad \forall i \in [k]$.

5. Asymptotics

5.1. Asymptotic behavior of $d(\mathbf{n}, \delta, \omega)$ for $n_i \rightarrow \infty$

Fix an index $i \in [k]$. We study the asymptotic behavior of $d(\mathbf{n}, \delta, \omega)$ when the dimension n_i goes to infinity. If $\delta_i = 0$ and $\omega = (1, \dots, 1)$, we recover (Ottaviani and Shahidi, 2021, Corollary 3.5). If $\delta_i \geq 1$ and $\omega_i = 1$, regardless of the other entries of δ and ω , we obtain a new stabilization property.

Proposition 33. Fix $i \in [k]$ and assume that $\omega_i = 1$. Let $n_i - 1 = \sum_{j \neq i} (n_j - 1) + \delta_i$. Then

$$d(n_1, \dots, n_{i-1}, m, n_{i+1}, \dots, n_k, \delta, \omega) = d(n_1, \dots, n_{i-1}, n_i, n_{i+1}, \dots, n_k, \delta, \omega) \quad \forall m \geq n_i.$$

Proof. Recall that $\delta = \sum_{i=1}^k \delta_i$. Let $T_i = \prod_{j \neq i} t_j^{n_j - \delta_j - 1}$. We need to compare the coefficients of

$$\begin{aligned} (1) & h^\delta t_i^{m - \delta_i - 1} T_i \quad \text{in} \\ & \frac{(\widehat{t}_i + h)^m - t_i^m}{\widehat{t}_i + h - t_i} \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j} = \left[\sum_{l=0}^{m-1} (\widehat{t}_i + h)^{m-1-l} t_i^l \right] \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j} \\ (2) & h^\delta t_i^{n_i - \delta_i - 1} T_i \quad \text{in} \\ & \frac{(\widehat{t}_i + h)^{n_i} - t_i^{n_i}}{\widehat{t}_i + h - t_i} \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j} = \left[\sum_{l=0}^{n_i-1} (\widehat{t}_i + h)^{n_i-1-l} t_i^l \right] \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j}. \end{aligned}$$

Observe that we can replace (2) with the coefficient of $t_i^{m-n_i} h^\delta t_i^{n_i - \delta_i - 1} T_i = h^\delta t_i^{m - \delta_i - 1} T_i$ in

$$\begin{aligned} & \left[\sum_{l=0}^{n_i-1} (\widehat{t}_i + h)^{n_i-1-l} t_i^{l+m-n_i} \right] \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j} \\ &= \left[\sum_{s=m-n_i}^{m-1} (\widehat{t}_i + h)^{m-1-s} t_i^s \right] \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j}. \end{aligned}$$

Now consider the product

$$\left[\sum_{s=0}^{m-n_i-1} (\widehat{t}_i + h)^{m-1-s} t_i^s \right] \prod_{j \neq i} \frac{(\widehat{t}_j + h)^{n_j} - t_j^{n_j}}{\widehat{t}_j + h - t_j}. \tag{5.1}$$

Recall that $\widehat{t}_i := (\sum_{j=1}^k \omega_j t_j) - t_i$. Since $\omega_i = 1$, the maximum degree of t_i in the first factor of (5.1) is $m - n_i - 1$. Moreover, the maximum degree of t_i in the second factor of (5.1) is $\sum_{j \neq i} (n_j - 1)$. Summing up, the maximum total degree of t_i in (5.1) is

$$m - n_i - 1 + \sum_{j \neq i} (n_j - 1) = m - \delta_i - 1 + \left(\delta_i + \sum_{j \neq i} (n_j - 1) - n_i \right) < m - \delta_i - 1$$

because $\delta_i + \sum_{j \neq i} (n_j - 1) - n_i < 0$ by hypothesis. Therefore, the product (5.1) gives no contribution to the coefficient of $h^\delta t_i^{m-\delta_i-1} T_i$. This implies the stabilization. \square

Remark 34. Proposition 33 does not hold if we study the asymptotic behavior of $d(\mathbf{n}, \delta, \omega)$ when $n_i \rightarrow \infty$ for $\omega_i > 1$. An immediate counterexample can be found in the symmetric case $k = 1$, see Remark 23.

5.2. Asymptotic behavior of $d(\mathbf{n}, \delta, \omega)$ in the binary format for $k \rightarrow \infty$

Assume $n_1 = \dots = n_k = 2$. Here $\delta_i \in \{0, 1\}$ for all $i \in [k]$. By (Ottaviani and Shahidi, 2021, Theorem 1.2), the integer $d(\mathbf{n}, \delta, \omega)$ is the coefficient of $h^\delta \prod_{i=1}^k t_i^{1-\delta_i}$ in

$$(\omega_1 t_1 + \dots + \omega_k t_k + h)^k = \sum_{j_1 + \dots + j_k + l = k} \binom{k}{j_1, \dots, j_k, l} (\omega_1 t_1)^{j_1} \dots (\omega_k t_k)^{j_k} h^l,$$

where $\binom{k}{j_1, \dots, j_k, l}$ is the multinomial coefficient. Setting $\mathcal{P}_0 := \{i \in [k] \mid \delta_i = 0\}$, we have that $|\mathcal{P}_0| = k - \delta$ and $d(\mathbf{n}, \delta, \omega) = \binom{k}{\delta} \prod_{i \in \mathcal{P}_0} \omega_i$. In particular, the growth of $d(\mathbf{n}, \delta, \omega)$ is factorial in k .

5.3. Asymptotic behavior of $d(\mathbf{n}, \delta, \omega)$ in the hypercubical format n^k for $n \rightarrow \infty$

In Theorem 2 we derived a generating function for the integers $d(\mathbf{n}, \delta, \omega)$, hence for $\delta = (\delta, 0, \dots, 0)$. Observe that the polynomial $H_\omega(\mathbf{x}, y)$ in (1.2) is of the form $H_\omega(\mathbf{x}, y) = -H_{1,\omega}(\mathbf{x})y + H_{2,\omega}(\mathbf{x})$. Therefore, we can rewrite the generating function of the degrees $d(\mathbf{n}, \delta, \omega)$ as

$$\sum_{\mathbf{n} \in \mathbb{N}^k} \sum_{\delta=0}^\infty d(\mathbf{n}, \delta, \omega) \mathbf{x}^{\mathbf{n}} y^\delta = \frac{1}{-H_{1,\omega}(\mathbf{x})y + H_{2,\omega}(\mathbf{x})} \prod_{i=1}^k \frac{x_i}{1 - x_i} = \sum_{\delta=0}^\infty F(\mathbf{x}) y^\delta, \tag{5.2}$$

where

$$F_\omega(\mathbf{x}) := \frac{H_{1,\omega}(\mathbf{x})^\delta}{H_{2,\omega}(\mathbf{x})^{\delta+1}} \prod_{i=1}^k \frac{x_i}{1 - x_i} = \frac{H_{1,\omega}(\mathbf{x})^\delta \prod_{i=1}^k x_i (1 - x_i)^\delta}{[H_{2,\omega}(\mathbf{x}) \prod_{i=1}^k (1 - x_i)]^{\delta+1}} = \frac{F_{N,\omega}(\mathbf{x})}{[F_{D,\omega}(\mathbf{x})]^{\delta+1}}. \tag{5.3}$$

From now on, we restrict to the case $\omega = \omega \mathbf{1}$. By Remark 32 we have

$$H_{1,\omega \mathbf{1}}(\mathbf{x}) = x_1 \sum_{i=0}^{k-1} e_i(\widehat{\mathbf{x}}_1), \quad H_{2,\omega \mathbf{1}}(\mathbf{x}) = \sum_{i=0}^k (1 - \omega i) e_i(\mathbf{x}).$$

In this case, the reduced denominator $F_{D,\omega \mathbf{1}}(\mathbf{x})$ of $F_{\omega \mathbf{1}}(\mathbf{x})$ is symmetric with respect to the variables x_i and coincides with the denominator of the generating function obtained in (Ekhad and Zeilberger, 2016, Proposition 1) when $\omega = 1$. The differences with (Ekhad and Zeilberger, 2016, Proposition 1) are the numerator and the exponent $\delta + 1$ in the denominator.

Our goal is to fix δ and study the asymptotic behavior of $d(n, \delta, \omega) := d(n \mathbf{1}, \delta, \omega \mathbf{1})$ for $n \rightarrow \infty$. This can be done applying the next result by Raichev and Wilson (2008). We refer to that paper for the definitions of strictly minimal, critical, isolated, and non-degenerate point needed in the statement.

Theorem 35. (Raichev and Wilson, 2008, Theorem 3.2) Let $k \geq 2$ and let $G = G_N/G_D^{\delta+1}$, whose Taylor expansion in a neighborhood of the origin is $\sum_{\alpha \in \mathbb{N}^k} g_\alpha \mathbf{x}^\alpha$. Suppose $\mathbf{c} \in \mathcal{V} = \{G_D(\mathbf{x}) = 0\}$ is smooth with $c_k \partial_k G_D(\mathbf{c}) \neq 0$, strictly minimal, critical, isolated, and non-degenerate. Then, for all $N \in \mathbb{N}$, as $n \rightarrow +\infty$,

$$g_{n\mathbf{1}} = \mathbf{c}^{-n\mathbf{1}} \left[\left((2\pi n)^{k-1} \det \tilde{g}''(0) \right)^{-1/2} \sum_{i=0}^{\delta} \sum_{j < N} \frac{(n+1)^{\delta-i}}{(\delta-i)! i!} n^{-j} L_j(\tilde{u}_i, \tilde{g}) + O\left(n^{\delta-(k-1)/2-N}\right) \right],$$

where $r^{\bar{s}} := r(r+1) \cdots (r+s-1)$. In the original formula of Raichev-Wilson, we substituted $\alpha = \mathbf{1}$ and $p = \delta + 1$.

The functions \tilde{g} , \tilde{u}_i and L_j appearing in Theorem 35 are defined in (Raichev and Wilson, 2008, Definition 3.1) and in the statement of (Raichev and Wilson, 2008, Theorem 3.2).

Proposition 36. The point $\mathbf{c} = \left(\frac{1}{\omega k-1}, \dots, \frac{1}{\omega k-1}\right)$ is smooth for $\mathcal{V} = \{F_{D,\omega\mathbf{1}}(\mathbf{x}) = 0\}$ with $c_k \partial_k F_{D,\omega\mathbf{1}}(\mathbf{c}) \neq 0$, strictly minimal, critical, isolated, and non-degenerate.

Proof. The statement is an immediate consequence of (Ottaviani et al., 2021, Remark 3.10) and (Pantone, 2017, Propositions 2.2, ..., 2.7). \square

Proposition 37. The following identity holds true:

$$L_0(\tilde{u}_0, \tilde{g}) = \tilde{u}_0(\mathbf{c}) = \frac{(\omega k - 1)^{k-\delta-1}}{(\omega k)^{k-\delta-2}(\omega k - 2)^k}.$$

Proof. Recall that $L_j(\tilde{u}_i, \tilde{g})$ is defined in (Raichev and Wilson, 2008, Theorem 3.2). For $j = 0$, we have $L_0(\tilde{u}_i, \tilde{g}) = \tilde{u}_i(\mathbf{c})$. By (Raichev and Wilson, 2008, Proposition 4.3), we have $\tilde{u}_0(\mathbf{c}) = \frac{F_{N,\omega\mathbf{1}}(\mathbf{c})}{(-c_k \partial_k F_{D,\omega\mathbf{1}}(\mathbf{c}))^{\delta+1}}$, where

$$\begin{aligned} F_{N,\omega\mathbf{1}}(\mathbf{c}) &= \left[c_1 \sum_{i=0}^{k-1} e_i(\hat{\mathbf{c}}_1) \right]^\delta \prod_{i=1}^k c_i (1 - c_i)^\delta \\ &= \left[\frac{1}{\omega k - 1} \sum_{i=0}^{k-1} \binom{k-1}{i} \left(\frac{1}{\omega k - 1}\right)^i \right]^\delta \left(\frac{1}{\omega k - 1}\right)^k \left(\frac{\omega k - 2}{\omega k - 1}\right)^{\delta k} \\ &= \left[\frac{1}{\omega k - 1} \left(\frac{\omega k}{\omega k - 1}\right)^{k-1} \right]^\delta \left(\frac{1}{\omega k - 1}\right)^k \left(\frac{\omega k - 2}{\omega k - 1}\right)^{\delta k} = \frac{(\omega k)^{(k-1)\delta} (\omega k - 2)^{k\delta}}{(\omega k - 1)^{k(2\delta+1)}}, \end{aligned}$$

and, similarly as in the proof of (Pantone, 2017, Proposition 2.7), $-c_k \partial_k F_{D,\omega\mathbf{1}}(\mathbf{c}) = \frac{(\omega k)^{k-2} (\omega k - 2)^k}{(\omega k - 1)^{2k-1}}$. Hence

$$\begin{aligned} \tilde{u}_0(\mathbf{c}) &= \frac{F_{N,\omega\mathbf{1}}(\mathbf{c})}{(-c_k \partial_k F_{D,\omega\mathbf{1}}(\mathbf{c}))^{\delta+1}} = \frac{(\omega k)^{(k-1)\delta} (\omega k - 2)^{k\delta}}{(\omega k - 1)^{k(2\delta+1)}} \left[\frac{(\omega k - 1)^{2k-1}}{(\omega k)^{k-2} (\omega k - 2)^k} \right]^{\delta+1} \\ &= \frac{(\omega k - 1)^{k-\delta-1}}{(\omega k)^{k-\delta-2} (\omega k - 2)^k}. \quad \square \end{aligned}$$

We have the necessary tools to prove Theorem 3.

Proof of Theorem 3. By Theorem 35, for $N = 1$ and $\mathbf{c} = (\frac{1}{\omega k - 1}, \dots, \frac{1}{\omega k - 1})$, we have

$$d(n, \delta, \omega) = (\omega k - 1)^{kn} \left[\frac{1}{(2\pi n)^{\frac{k-1}{2}} \det \tilde{g}''(0)^{\frac{1}{2}}} \sum_{i=0}^{\delta} \frac{(n+1)^{\delta-i}}{(\delta-i)! i!} L_0(\tilde{u}_i, \tilde{g}) + O\left(\frac{1}{n^{\frac{k+1}{2}-\delta}}\right) \right],$$

as $n \rightarrow \infty$. Define $\eta_k := (2\pi)^{-\frac{k-1}{2}} \det \tilde{g}''(0)^{-\frac{1}{2}}$. Similarly as in (Pantone, 2017, Proposition 2.7), one verifies that $\det \tilde{g}''(0) = \frac{(\omega k - 2)^{k-1}}{(\omega k)^{k-2}} \neq 0$. Then

$$\begin{aligned} d(n, \delta, \omega) &= (\omega k - 1)^{kn} \left[\frac{\eta_k L_0(\tilde{u}_0, \tilde{g}) n^\delta}{\delta! n^{\frac{k-1}{2}}} + O\left(\frac{1}{n^{\frac{k+1}{2}-\delta}}\right) \right] \\ &= (\omega k - 1)^{kn} \left[\frac{\eta_k L_0(\tilde{u}_0, \tilde{g}) n^\delta}{\delta! n^{\frac{k-1}{2}}} + \frac{1}{n^{\frac{k-1}{2}-\delta}} O\left(\frac{\eta_k L_0(\tilde{u}_0, \tilde{g})}{n}\right) \right] \\ &= (\omega k - 1)^{kn} \left[\frac{\eta_k L_0(\tilde{u}_0, \tilde{g}) n^\delta}{n^{\frac{k-1}{2}}} + \frac{\eta_k L_0(\tilde{u}_0, \tilde{g})}{n^{\frac{k-1}{2}-\delta}} O\left(\frac{1}{n}\right) \right] \\ &= \eta_k L_0(\tilde{u}_0, \tilde{g}) \frac{(\omega k - 1)^{kn}}{n^{\frac{k-1}{2}-\delta}} \left[1 + O\left(\frac{1}{n}\right) \right]. \end{aligned}$$

Conclusion follows by plugging in the identity for $L_0(\tilde{u}_0, \tilde{g})$ in Proposition 37. \square

Remark 38. When $\omega = 1$, we obtain the following $O(1/n)$ -approximations for $d(n, \delta, 1)$:

$$d(n, \delta, 1) \approx \begin{cases} \frac{2}{\sqrt{3}\pi} \left(\frac{3}{2}\right)^\delta 8^n n^{\delta-1} & \text{if } k = 3 \\ \frac{27}{2^9\pi\sqrt{\pi}} \left(\frac{4}{3}\right)^\delta 81^n n^{\delta-\frac{3}{2}} & \text{if } k = 4. \end{cases}$$

6. Kalman varieties of partially symmetric singular vector k -tuples

In this final section, we restrict to the case $V_1 = \dots = V_k = V$ for some n -dimensional complex vector space V and we assume that all isotropic quadrics Q_i coincide with the quadric $Q \subset \mathbb{P}(V)$. We also use the notations $\mathbb{P} = \mathbb{P}(V)^{\times k}$ and $Q^k = \prod_{i=1}^k Q \subset \mathbb{P}$. Our goal is to introduce another kind of Kalman variety. To motivate this construction, let us consider the matrix case. In this context, one may wonder what is the closure of the locus of matrices $A \in V \otimes V$ possessing a non-isotropic singular vector pair of the form (\mathbf{x}, \mathbf{x}) for some $\mathbf{x} \in V$. For $n = 2$, this locus coincides with the subspace $S^2V \subset V \otimes V$ of 2×2 symmetric matrices. As we shall see, for $n \geq 3$ this locus is not a linear subspace and contains the subspace S^2V as a proper closed subset.

Definition 39. The *normalized symmetric Kalman variety* is the variety

$$\kappa_{n,k}^{nor} := \overline{\{T \in \mathbb{P}(V^{\otimes k}) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}], \dots, [\mathbf{x}]) \text{ for some } [\mathbf{x}] \in \mathbb{P}(V) \setminus Q\}}. \tag{6.1}$$

In complete analogy with the generalized Kalman variety introduced in Definition 15, one can define the *generalized symmetric Kalman variety*

$$\kappa_{n,k} := \{T \in \mathbb{P}(V^{\otimes k}) \mid T \text{ has a singular vector } k\text{-tuple } ([\mathbf{x}], \dots, [\mathbf{x}]) \text{ for some } [\mathbf{x}] \in \mathbb{P}(V)\}. \tag{6.2}$$

As shown in Example 41, the generalized symmetric Kalman variety $\kappa_{n,k}$ may be reducible already for matrices.

Table 1
Codimension and degree of $\kappa_{n,2}^{nor}$ for small n .

n	$\text{codim}(\kappa_{n,2}^{nor})$	$\text{deg}(\kappa_{n,2}^{nor})$	generators of the ideal $\mathcal{I}(\kappa_{n,2}^{nor})$
2	1	1	1 linear
3	2	7	3 cubics
4	3	24	1 quadric, 1 quartic, 11 sextics
5	4	86	5 quartics, 5 quintics, 5 sextics, 31 septics
6	5	314	1 cubic, 1 quintic, 86 sextics

Theorem 40. *The variety $\kappa_{n,k}^{nor}$ is irreducible of codimension $(k - 1)(n - 1)$ in $\mathbb{P}(V^{\otimes k})$.*

Proof. Consider the spectral variety $\Sigma_{n,1,1}$ along with the two projections α and β of $\Sigma_{n,1,1}$ onto $\mathbb{P}(V^{\otimes k})$ and \mathbb{P} , as in diagram (2.6). By Theorem 13, every fiber of β at each point of $\mathbb{P} \setminus Q^k$ is a linear subspace of codimension $k(n - 1)$ in $\mathbb{P}(V^{\otimes k})$.

Consider the Segre embedding $\sigma : \mathbb{P} \rightarrow \mathbb{P}(V^{\otimes k})$, $\sigma([\mathbf{x}_1], \dots, [\mathbf{x}_k]) = [\mathbf{x}_1 \otimes \dots \otimes \mathbf{x}_k]$. Denote by $v_{n,k}$ the k -th Veronese embedding of $\mathbb{P}(V)$, as a subvariety of $\mathbb{P}(V^{\otimes k})$. The variety $\kappa_{n,k}^{nor}$ has the following description in terms of the diagram mentioned above:

$$\kappa_{n,k}^{nor} = \overline{\alpha(\beta^{-1}(\sigma^{-1}(v_{n,k}) \setminus Q^k))},$$

where $\sigma^{-1}(v_{n,k}) = \mathbb{P} \cap \Delta$ and Δ denotes the diagonal. Let $Y = \overline{\beta^{-1}(\sigma^{-1}(v_{n,k}) \setminus Q^k)}$. By Lemma 17, Y is irreducible and so is $\kappa_{n,k}^{nor}$. Moreover, we have $\dim(Y) = \dim(\mathbb{P} \cap \Delta) + \dim(\mathbb{P}(V^{\otimes k})) - (n - 1)k = n^k - 1 - (k - 1)(n - 1)$.

The general fiber of α restricted to Y is finite. Indeed, any point of $\sigma^{-1}(v_{n,k}) \setminus Q^k$ is the equivalence class of a tuple of the same non-isotropic vector. The group $\text{SO}(V)^{\times k}$ acts transitively on equivalence classes of non-isotropic vectors in \mathbb{P} . Thus a general tensor in the fiber under β of any point in the diagonal $\sigma^{-1}(v_{n,k}) \setminus Q^k$ has a finite number of singular vector k -tuples. This implies the finiteness of α restricted to Y . The variety $\kappa_{n,k}^{nor}$ is irreducible and has dimension $\dim(Y)$. Hence $\kappa_{n,k}^{nor}$ has codimension $(k - 1)(n - 1)$ in $\mathbb{P}(V^{\otimes k})$. \square

Example 41. Consider $k = 2$, $V = \mathbb{C}^n$ and $Q = \mathcal{V}(x_1^2 + \dots + x_n^2) \subset \mathbb{P}(V)$. We have $\text{codim}(\kappa_{n,2}^{nor}) = n - 1$. In particular, one finds $\kappa_{2,2}^{nor} = \mathbb{P}(S^2\mathbb{C}^2)$. The first non-trivial case is the normalized symmetric Kalman variety $\kappa_{3,2}^{nor}$ which is a subvariety of $\mathbb{P}(\mathbb{C}^3 \otimes \mathbb{C}^3)$ of codimension 2 and degree 7. By Remark 7, $\kappa_{3,2}^{nor}$ coincides with the locus of matrices admitting a usual algebraic singular vector pair. This ideal is generated by three cubics that may be recovered with the following construction, suggested to us by Jan Draisma.

Let $A = (a_{ij}) \in \mathbb{C}^n \otimes \mathbb{C}^n$ and suppose (\mathbf{y}, \mathbf{y}) is a (non-isotropic) singular vector pair of A . Up to scaling \mathbf{y} , we may write $A = \mathbf{y}\mathbf{y}^T + C$ and $A^T = \mathbf{y}\mathbf{y}^T + C^T$. The matrix $B = A - A^T$ has matrix rank at most $n - 1$. Note that, since the latter matrix is skew-symmetric, when n is odd and for general A , one has $\text{rk}(B) = n - 1$. In such a case, the cofactor matrix $\text{cof}(B)$ has rank one and satisfies $B \cdot \text{cof}(B)^T = 0$. So the image of $\text{cof}(B)$ is spanned by the vector \mathbf{y} above.

To derive the cubic equations for $n = 3$, given $\mathbf{x} = (x_1, x_2, x_3)^T \in \mathbb{C}^3$, we have $\mathbf{y} = \text{cof}(B)\mathbf{x}$. Since \mathbf{y} is then an eigenvector of A , let \mathcal{K} be the ideal of 2×2 minors of the matrix $[A\mathbf{y} \mid \mathbf{y}]$. We verified symbolically in Macaulay2 that $\mathcal{K} = \mathcal{I}(\kappa_{3,2}^{nor}) \cap \mathcal{J}^2$, where $\mathcal{I}(\kappa_{3,2}^{nor})$ is the ideal of $\kappa_{3,2}^{nor}$ generated by the desired three cubics, and where $\mathcal{J} = \langle x_3a_{12} - x_2a_{13} - x_3a_{21} + x_1a_{23} + x_2a_{31} - x_1a_{32} \rangle$ defines the hyperplane of matrices orthogonal (with respect to the Frobenius inner product) to the skew-symmetric matrix

$$\begin{pmatrix} 0 & x_3 & -x_2 \\ -x_3 & 0 & x_1 \\ x_2 & -x_1 & 0 \end{pmatrix}.$$

Moreover, $\kappa_{3,2}^{nor}$ is arithmetically Cohen-Macaulay and its reduced singular locus is $\mathbb{P}(S^2\mathbb{C}^3)$. The generalized symmetric Kalman variety $\kappa_{3,2}$ is reducible of degree 15 and codimension 2. One of its irreducible components is $\kappa_{3,2}^{nor}$. Further numerical data for the varieties $\kappa_{n,2}^{nor}$ are summarized in Table 1 for small values of n .

Example 42. In the binary tensor case ($n = 2$), we obtain that $\text{codim}(\kappa_{2,k}^{nor}) = k - 1$. The first non-trivial case is $\kappa_{2,3}^{nor}$ which is a subvariety of $\mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)$ of codimension 2 and degree 5. Its ideal is generated by one quadric and four quartics.

Question 43. Table 1 reports the degrees of the varieties $\kappa_{n,2}^{nor}$ for small values of n . What is the degree of $\kappa_{n,k}^{nor}$ in general?

It is clear that the construction above can be carried out taking into account partial symmetries of singular vector k -tuples. For simplicity, we remain in the hypercubical format $n \times k$. To this aim, let $\omega = (\omega_1, \dots, \omega_t)$ be a partition of k , namely $\omega_i \geq 1$ for all i and $\omega := \omega_1 + \dots + \omega_t = k$. Without loss of generality, we assume that $\omega_1 \geq \dots \geq \omega_t$. We denote by $S_{n,k}^\omega$ the Segre-Veronese embedding of $\mathbb{P}(V)^{\times t}$ in $\mathbb{P}(S^\omega V)$, and we consider it as a subvariety of $\mathbb{P}(V^{\otimes k})$. Define

$$\kappa_{n,k,\omega}^{nor} := \overline{\{T \in \mathbb{P}(V^{\otimes k}) \mid T \text{ has a singular vector } k\text{-tuple } (\underbrace{[\mathbf{x}_1], \dots, [\mathbf{x}_1]}_{\omega_1}, \dots, \underbrace{[\mathbf{x}_t], \dots, [\mathbf{x}_t]}_{\omega_t}) \in \mathbb{P} \setminus Q^k\}} \tag{6.3}$$

With similar arguments as in the proof of Theorem 40, one proves the following result.

Theorem 44. Let $\omega = (\omega_1, \dots, \omega_t)$ be a partition of k . The variety $\kappa_{n,k,\omega}^{nor}$ is irreducible of codimension $(k - t)(n - 1)$ in $\mathbb{P}(V^{\otimes k})$. So the codimension depends only on the number of parts t of ω .

Example 45. In the binary tensor case ($n = 2$), we see that $\text{codim}(\kappa_{2,k,\omega}^{nor}) = k - t$. The first non-trivial case not considered before is $\kappa_{2,3,(2,1)}^{nor}$ which is a hypersurface of $\mathbb{P}(\mathbb{C}^2 \otimes \mathbb{C}^2 \otimes \mathbb{C}^2)$ of degree 8. If $\omega = (2, 1^{k-2})$, we have $t = k - 1$ and therefore $\kappa_{2,k,\omega}^{nor}$ is always a hypersurface. Moreover, note that, for instance, $\text{codim}(\kappa_{2,4,(2,2)}^{nor}) = \text{codim}(\kappa_{2,4,(3,1)}^{nor}) = 2$, even though $\kappa_{2,4,(2,2)}^{nor}$ and $\kappa_{2,4,(3,1)}^{nor}$ have different degrees.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

References

Beorchia, Valentina, Galuppi, Francesco, Venturolo, Lorenzo, 2021. Eigenschemes of ternary tensors. *SIAM J. Appl. Algebra Geom.* 5 (4), 620–650.
 Draisma, Jan, Horobeţ, Emil, Ottaviani, Giorgio, Sturmfels, Bernd, Thomas, Rekha R., 2016. The Euclidean distance degree of an algebraic variety. *Found. Comput. Math.* 16 (1), 99–149.
 Ekhad, Shalosh B., Zeilberger, Doron, 2016. On the number of singular vector tuples of hyper-cubical tensors. *Pers. J. Shalosh B. Ekhad and Doron Zeilberger*.
 Friedland, Shmuel, Ottaviani, Giorgio, 2014. The number of singular vector tuples and uniqueness of best rank-one approximation of tensors. *Found. Comput. Math.* 14 (6), 1209–1242.
 Gel'fand, Israel M., Kapranov, Mikhail M., Zelevinsky, Andrey V., 1994. *Discriminants, Resultants, and Multidimensional Determinants*. Mathematics: Theory & Applications. Birkhäuser Boston, Inc., Boston, MA.
 Grayson, Daniel, Stillman, Michael, 1997. *Macaulay 2—a system for computation in algebraic geometry and commutative algebra*.
 Hartshorne, Robin, 1977. *Algebraic Geometry*. Graduate Texts in Mathematics, vol. 52. Springer-Verlag, New York-Heidelberg.
 Hogben, Leslie (Ed.), 2007. *Handbook of Linear Algebra*. Discrete Mathematics and Its Applications (Boca Raton). Chapman & Hall/CRC, Boca Raton, FL.

- Holme, Audun, 1988. The geometric and numerical properties of duality in projective algebraic geometry. *Manuscr. Math.* 61 (2), 145–162.
- Huang, Hang, 2020. Equations of Kalman varieties. *Proc. Am. Math. Soc.* <https://doi.org/10.1090/proc/15351>.
- MacMahon, Percy Alexander, 1915. *Combinatory Analysis*, vol. 1. Cambridge University Press.
- Ottaviani, Giorgio, Sturmfels, Bernd, 2013. Matrices with eigenvectors in a given subspace. *Proc. Am. Math. Soc.* 141 (4), 1219–1232.
- Ottaviani, Giorgio, Shahidi, Zahra, 2021. Tensors with eigenvectors in a given subspace. *Rend. Circ. Mat. Palermo, II. Ser.*, 1–12.
- Ottaviani, Giorgio, Sodomaco, Luca, Ventura, Emanuele, 2021. Asymptotics of degrees and ED degrees of Segre products. *Adv. Appl. Math.* 130, 102242.
- Pantone, Jay, 2017. The asymptotic number of simple singular vector tuples of a cubical tensor. *Online J. Anal. Comb.* (12), 11.
- Pólya, George, 1954. *Induction and Analogy in Mathematics. Mathematics and Plausible Reasoning*, vol. I. Princeton University Press, Princeton, N. J.
- Qi, Liqun, Luo, Ziyang, 2017. *Tensor Analysis. Spectral Theory and Special Tensors*. Society for Industrial and Applied Mathematics, Philadelphia, PA.
- Raichev, Alexander, Wilson, Mark C., 2008. Asymptotics of coefficients of multivariate generating functions: improvements for smooth points. *Electron. J. Comb.* 15 (1), 89.
- Sam, Steven V., 2012. Equations and syzygies of some Kalman varieties. *Proc. Am. Math. Soc.* 140 (12), 4153–4166.
- Sodomaco, Luca, 2018. The product of the eigenvalues of a symmetric tensor. *Linear Algebra Appl.* 554, 224–248.
- Sodomaco, Luca, 2020. *The Distance Function from the Variety of partially symmetric rank-one Tensors*. PhD thesis. Università degli Studi di Firenze. Available at <https://sites.google.com/view/luca-sodomaco/home/publications>.
- Stanley, Richard P., 2012. *Enumerative Combinatorics. Volume 1*, second edition. Cambridge Studies in Advanced Mathematics, vol. 49. Cambridge University Press, Cambridge.
- Teixera Turatti, Ettore, 2021. On tensors that are determined by their singular tuples. arXiv:2104.03686.
- Vakil, Ravi, 2017. *The rising sea. Foundations of algebraic geometry*. <https://math216.wordpress.com/>.
- Wilf, Herbert S., 1990. *generatingfunctionology*. Academic Press, Inc., Boston, MA.