

Vanishing Hessian, wild forms and their border VSP

Original

Vanishing Hessian, wild forms and their border VSP / Huang, H.; Michalek, M.; Ventura, E.. - In: MATHEMATISCHE ANNALEN. - ISSN 0025-5831. - 378:3-4(2020), pp. 1505-1532. [10.1007/s00208-020-02080-8]

Availability:

This version is available at: 11583/2958169 since: 2022-03-11T16:51:42Z

Publisher:

Springer

Published

DOI:10.1007/s00208-020-02080-8

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Vanishing Hessian, wild forms and their border VSP

Hang Huang¹ · Mateusz Michałek² · Emanuele Ventura³ 

Received: 25 June 2020 / Revised: 23 July 2020 / Published online: 14 September 2020
© The Author(s) 2020

Abstract

Wild forms are homogeneous polynomials whose smoothable rank is strictly larger than their border rank. The discrepancy between these two ranks is caused by the difference between the limit of spans of a family of zero-dimensional schemes and the span of their flat limit. For concise forms of minimal border rank, we show that the condition of vanishing Hessian is equivalent to being wild. This is proven by making a detour through structure tensors of smoothable and Gorenstein algebras. The equivalence fails in the non-minimal border rank regime. We exhibit an infinite series of minimal border rank wild forms of every degree $d \geq 3$ as well as an infinite series of wild cubics. Inspired by recent work on border apolarity of Buczyńska and Buczyński, we study the border varieties of sums of powers VSP of these forms in the corresponding multigraded Hilbert schemes.

Mathematics Subject Classification Primary 14C05 · Secondary 15A69

1 Introduction

Notions of ranks abound in the literature, perhaps because of their natural appearance in the realms of algebra and geometry, and in numerous applications thereof; see [18,22] and references therein for an introduction to the subject.

Communicated by Vasudevan Srinivas.

✉ Emanuele Ventura
emanuele.ventura@math.unibe.ch

Hang Huang
hhuang@math.tamu.edu

Mateusz Michałek
mateusz.michalek@mis.mpg.de

¹ Department of Mathematics, Texas A&M University, College Station, TX 77843-3368, USA

² Max Planck Institute for Mathematics in the Sciences, Leipzig, Germany

³ Mathematical Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland

These ranks vastly generalize matrix rank and yet they are very classical, dating back to the pioneering work of Sylvester. His work featured Waring ranks of binary forms; see [18] for a historical account on the subject. Since then, ever growing research efforts have been devoted to understanding ranks with respect to some special projective varieties X of interest. Last decades have witnessed steady progress on tensor and Waring ranks, i.e. the cases when the projective varieties are the classical Segre and Veronese varieties.

These results have been developed in parallel in their geometric and algebraic aspects. The first are naturally related to secant varieties of X [29, Chapter 1], whereas the second to Macaulay's theory of apolarity and inverse systems [18, §1.1].

Interestingly, scheme-theoretic versions of X -ranks have been introduced and studied as well. These latter ones take into account more general zero-dimensional schemes, besides the reduced zero-dimensional ones featured in the X -ranks. This more general framework naturally leads to new notions of X -rank: the *smoothable X -rank*, and the *cactus X -rank*; the latter was originally called *scheme length* [18, Definition 5.1]. We recall their definitions in Sect. 2.

One subtle phenomenon is that, for special points, perhaps unexpectedly smoothable ranks may be larger than border ranks. This discrepancy is caused by the difference between the limit of spans of a family of zero-dimensional schemes and the span of their flat limit. The difference between smoothable and border ranks does not appear for general points (forms) of fixed border rank. Therefore, it is a natural and interesting problem to investigate the structure of the instances where these two differ.

As far as we know, Buczyńska and Buczyński [4] were the first authors to bring the difference between these ranks to attention. They introduced the notion of *wild forms*, i.e. those whose smoothable rank is strictly larger than their border rank. They gave one such a form [4, §4], up to concise minimal border rank direct summands.

On another direction, in recent groundbreaking work, Buczyńska and Buczyński [3] expanded the apolarity theory of X -ranks to *border apolarity*, which is devised to provide information about border X -ranks. Along the way, they introduced the *border varieties of sums of powers VSP*, mirroring the classical varieties of sums of powers [26,28].

Inspired by [3, §5.3], we establish a new result on wild forms. To state it, let V be a complex finite-dimensional vector space; given a form $F \in S^d V^*$, let $\text{Hess}(F)$ denote the determinant of its Hessian matrix. Forms with identically vanishing Hessian have many remarkable geometric and algebraic properties; see [29, Chapter 7] for a detailed and updated exposition.

Forms with vanishing Hessian were originally studied by Hesse in two classical papers [16,17], where the author tried to prove that these homogeneous polynomials are necessarily not concise (or, in more geometric terms, that the hypersurfaces they define are cones). Thereafter, in their important work [12], Gordan and Noether showed that Hesse's claim is true in the regime of at most four variables, whereas there exist infinitely many counterexamples afterwards. The easiest counterexample is perhaps the *Perazzo cubic hypersurface* [27] (in Perazzo's words "*un esempio semplicissimo*", i.e. "a very easy example"), which appears in [4, §4] as an instance of wild cubic form: that is the point of departure of our article. Our main result is Theorem 3.4, which connects wild forms and vanishing Hessian, following the lines paved by Ottaviani's

remark [3, Remark 5.1]. (For definitions of the ranks involved in the statement, see Sect. 2.)

Theorem *Let $d \geq 3$ and $F \in S^d V^*$ be a concise form of minimal border rank. Then:*

$$\text{Hess}(F) = 0 \iff \text{cr}(F) > \underline{r}(F) \iff \text{sr}(F) > \underline{r}(F) \iff F \text{ is wild.}$$

The equivalence fails when assuming non-minimal border rank: there exist concise wild forms of non-minimal border rank whose Hessian is not vanishing, see Example 3.9.

Let $T = \bigoplus_{d \geq 0} S^d V$. Given $F \in S^d V^*$, let $\text{Ann}(F) \subset T$ be the annihilator or apolar ideal of F , see Sect. 2. Let $A = T/\text{Ann}(F)$ be the Artinian Gorenstein \mathbb{C} -algebra of (a concise) F . Let $\{\alpha_i^{(k)}\}$ be a basis of A_k . Then the determinant

$$\text{Hess}^k(F) = \det \left(\alpha_i^{(k)} \alpha_j^{(k)} F \right)$$

is the k -th Hessian of F . By definition, $\text{Hess}^1(F) = \text{Hess}(F)$.

The *Strong Lefschetz Property* (SLP) of A is characterized in terms of these higher Hessians of F : A has SLP if and only if $\text{Hess}^k(F) \neq 0$ for every $k = 1, \dots, \lfloor d/2 \rfloor$ [24, Theorem 3.1]. Then the result above reads:

Theorem *Let $d \geq 3$ and let $F \in S^d V^*$ be a concise form of minimal border rank. Then:*

$$F \text{ is wild} \implies T/\text{Ann}(F) \text{ does not have SLP.}$$

For $d = 3$ and $d = 4$, this is an equivalence.

To prove our main result we rely on the interplay between tensors and algebras. In particular, we prove the following:

Theorem *A finite-dimensional \mathbb{C} -algebra A is Gorenstein if and only if its 3-way structure tensor (i.e. the tensor associated to the multiplication map $A \times A \rightarrow A$) is symmetric if and only if its d -way structure tensor for some $d > 2$ (equivalently, for every d) is symmetric.*

We also provide a possible extension of the results of Bläser and Lysikov [2] to d -way tensors:

Theorem *Let T be a symmetric d -way tensor of minimal border rank. Suppose there exists a contraction $T(\ell^{\otimes d-2})$ which is a full-rank symmetric matrix. Then T is the structure tensor of a smoothable Gorenstein algebra.*

As an application of our main result, we exhibit two infinite series of (concise) wild forms. In Sect. 6, we give a series of wild forms G_d of every degree $d \geq 3$, and in Sect. 7 a series of wild cubics F_n . Both of them are of minimal border rank. In particular, this shows the next

Theorem *There exist concise minimal border rank wild forms of any degree $d \geq 3$.*

Employing Buczyńska-Buczyński's border apolarity theory, we offer a study of border varieties of sums of powers $\underline{\text{VSP}}$'s of these forms in the corresponding multi-graded Hilbert schemes. To our knowledge, this is the first attempt to describe such varieties for some forms. For the first series G_d , we show that they are projective spaces; see Theorem 6.6.

For the second series F_n and $n \geq 10$, in Theorem 7.9 we prove that they are reducible. This is achieved by relying on the (usual) Hilbert schemes of zero-dimensional schemes on a *chain of lines*, see Sect. 7. We point out that this result on reducibility is also motivated by the fact that establishing this property is usually a delicate and interesting issue even in the context of the classical varieties of sums of powers VSP 's.

Structure of the paper

In Sect. 2, we introduce notation and recall the definitions of ranks we need throughout the article. Section 3 is devoted to the proof of the first part of our main result, i.e., vanishing Hessian implies wild for concise forms of minimal border rank, Theorem 3.4.

In Sect. 4, we make a detour through structure tensors of smoothable and Gorenstein algebras to establish Theorem 4.9. The latter gives the remaining part of our main result.

In Sect. 5, we give the definition of a limiting scheme of a border rank decomposition and its relations with $\underline{\text{VSP}}$. Theorem 5.3 recalls that the saturation of an ideal in $\underline{\text{VSP}}$ is the ideal of a limiting scheme of a border rank decomposition. Theorem 5.4 shows the correspondence between ideals and border decompositions, in the regime of minimal border rank.

In Sect. 6, we introduce the infinite series of concise degree d forms G_d . We show that they are wild in Corollary 6.2. Moreover, we prove that their $\underline{\text{VSP}}$'s are isomorphic to projective spaces; this is achieved in Theorem 6.6.

In Sect. 7, we introduce the infinite series of concise cubics F_n . Corollary 7.7 states that they are wild. We show that when $n \geq 10$, their $\underline{\text{VSP}}$'s are reducible; see Theorem 7.9.

2 Preliminaries

Here we introduce notation and definitions we use throughout the paper. We work over the complex numbers. Let $V \cong \mathbb{C}^{n+1}$ and $V^* = \langle x_0, \dots, x_n \rangle$. Let $\mathbb{P}^n = \mathbb{P}(V)$ denote the projectivization of V . Let $S = S^\bullet V^* \cong \mathbb{C}[x_0, \dots, x_n]$ be its homogeneous coordinate ring, and $T = \mathbb{C}[y_0, \dots, y_n]$ be its dual ring, i.e. T acts by differentiation on S with $y_i \circ x_j = \delta_{i,j}$.

For a homogeneous ideal $\mathcal{J} \subset T$, let \mathcal{J}_d denote its degree d homogeneous component. For $\mathcal{J} \subset T$, let \mathcal{J}^{sat} denote its saturation. The *Hilbert function* of \mathcal{J} is the numerical function

$$\text{HF}(T/\mathcal{J}, d) = \dim S^d V - \dim \mathcal{J}_d = \dim (T/\mathcal{J})_d.$$

Given a form $F \in S^d V^*$, $\text{Ann}(F) \subset T$ denotes its annihilator or apolar ideal

$$\text{Ann}(F) = \{h \in T \mid h \circ F = 0\}.$$

The algebra $T/\text{Ann}(F)$ is a graded Artinian Gorenstein \mathbb{C} -algebra; see [18, §2.3] or [29, Theorem 7.2.15].

Let $N_d = \binom{n+d}{d} - 1$ and $X = \nu_d(\mathbb{P}^n) \subset \mathbb{P}^{N_d}$ be the d -th Veronese embedding of \mathbb{P}^n . We only consider ranks with respect to the Veronese variety, although the ensuing definitions may be more generally introduced for any projective variety.

Let R be a zero-dimensional scheme over \mathbb{C} ; then $R = \text{Spec}(A)$ for some finite-dimensional \mathbb{C} -algebra A . The *length* of R is $\text{length}(R) = \dim_{\mathbb{C}} A$.

For a zero-dimensional scheme $R \subset \mathbb{P}^n = \mathbb{P}(V)$, let $\langle R \rangle$ denote its span, i.e.

$$\langle R \rangle = \mathbb{P}((V^*/\mathcal{I}_1)^*) \subset \mathbb{P}^n,$$

where \mathcal{I} is the saturated ideal defining R in \mathbb{P}^n .

Definition 2.1 (*Border rank*) For a point $F \in \mathbb{P}^{N_d}$, the *border rank* of F is the minimal integer r such that $F \in \sigma_r(X)$, the r -th secant variety of X . The border rank of F is denoted $\underline{\mathbf{r}}(F)$.

Definition 2.2 A form $F \in S^d V^*$ is *concise* if its annihilator $\text{Ann}(F)$ does not contain linear forms.

Definition 2.3 A form $F \in S^d V^*$ is said to be of *minimal border rank* when

$$\underline{\mathbf{r}}(F) = \dim V.$$

Remark 2.4 The border rank of a concise form $F \in S^d V^*$ satisfies $\underline{\mathbf{r}}(F) \geq \dim V$. This explains the adjective *minimal* in Definition 2.3.

In general, given $F \in S^d V^*$, it might be challenging to produce a border rank decomposition $F = \lim_{t \rightarrow 0} \frac{1}{t^s} (L_1(t)^d + \dots + L_{\underline{\mathbf{r}}(F)}(t)^d)$. To determine border ranks for our infinite series of forms G_d and F_n , we employ a useful criterion:

Proposition 2.5 [4, Proposition 2.6] *Let X be as above. Suppose there exist points $z_1, \dots, z_r \in X$ such that $\dim \langle z_1, \dots, z_r \rangle < r - 1$. Then the span of the affine cones of Zariski tangent spaces at these points is contained in the r -th secant variety $\sigma_r(X)$, i.e.*

$$\langle \mathbb{P}\widehat{T}_{z_1}, \dots, \mathbb{P}\widehat{T}_{z_r} \rangle \subset \sigma_r(X).$$

We now recall the scheme-theoretic ranks attached to X .

Definition 2.6 (*Smoothable rank*) The *smoothable rank* of $F \in \mathbb{P}^{N_d}$ is the minimal integer r such that there exists a finite scheme $R \subset X$ of length r which is *smoothable* (in X) and $F \in \langle R \rangle$. Equivalently, there exists a finite smoothable scheme $R \subset X$ of length r whose saturated ideal satisfies $\mathcal{J}_R \subset \text{Ann}(F)$. (This is in analogy with the classical *Apolarity lemma* [18, Lemma 1.15].) The smoothable rank of F is denoted $\text{sr}(F)$.

Remark 2.7 Smoothable and border ranks satisfy $\underline{\mathbf{r}}(F) \leq \text{sr}(F)$; see [4, §2.1]. Equality holds for most of the points on a secant variety. The difference arises in general from the *failure* of the equality

$$\left\langle \lim_{t \rightarrow 0} R(t) \right\rangle = \lim_{t \rightarrow 0} \langle R(t) \rangle,$$

where $R(t)$ is a family of zero-dimensional schemes over the base $\text{Spec}(\mathbb{C}[t^{\pm 1}])$ and $\lim_{t \rightarrow 0} R(t)$ denotes its flat limit; see for instance [6, II.3.4].

Definition 2.8 (*Cactus rank*) The *cactus rank* of $F \in \mathbb{P}^{Nd}$ is the minimal integer r such that there exists a finite scheme $R \subset X$ of length r such that $F \in \langle R \rangle$. Equivalently, there exists a finite scheme $R \subset X$ of length r whose saturated ideal satisfies $\mathcal{J}_R \subset \text{Ann}(F)$. (This is in analogy with the classical *Apolarity lemma* [18, Lemma 1.15].) The cactus rank of F is denoted $\text{cr}(F)$.

The cactus rank was originally called *scheme length* [18, Definition 5.1].

Remark 2.9 From definitions, one has $\text{cr}(F) \leq \text{sr}(F)$. Moreover, $\text{cr}(F)$ and $\underline{\mathbf{r}}(F)$ are incomparable. Theorem 3.4 produces infinitely many examples where $\text{cr}(F) > \underline{\mathbf{r}}(F)$; however one has $\text{cr}(F) < \underline{\mathbf{r}}(F)$ as well in several instances; see e.g. [1].

Definition 2.10 A form $F \in S^d V^*$ is a *form with vanishing Hessian* if the determinant of its Hessian matrix $\text{Hess}(F) = \det \left(\left[\frac{\partial^2 F}{\partial x_i \partial x_j} \right] \right)$ vanishes identically.

See [29, Chapter 7] and references therein for a complete introduction to several remarkable algebraic and geometric properties that forms with vanishing Hessian possess.

For any form $F \in S^d V^*$, Buczyńska and Buczyński [3, §4.1] introduced the *border variety of sums of powers* $\underline{\text{VSP}}(F, \underline{\mathbf{r}}(F))$. These varieties live in *multigraded Hilbert schemes*, which were introduced by Haiman and Sturmfels [14]. We now recall the definition of an irreducible component of the multigraded Hilbert scheme we are concerned with; see [3, §3] for a detailed discussion.

Definition 2.11 An ideal $\mathcal{J} \subset T$, whose Hilbert polynomial is equal to $r \in \mathbb{N}$, is said to have a *generic Hilbert function* if its Hilbert function satisfies

$$\text{HF}(T/\mathcal{J}, d) = \min\{r, \dim S^d V\}, \text{ for } d \geq 0.$$

Let $\text{Slip}_{r, \mathbb{P}^n}$ be the irreducible component of the multigraded Hilbert scheme $\text{Hilb}_T^{h_r, \mathbb{P}^n}$ containing the radical ideals of r distinct points with a generic Hilbert function. Therefore, every ideal in $\text{Slip}_{r, \mathbb{P}^n}$ has a generic Hilbert function being a flat limit of such ideals.

Definition 2.12 Let $F \in S^d V^*$. The *border variety of sums of powers*, or *border VSP*, of F is

$$\underline{\text{VSP}}(F, r) = \{ \mathcal{J} \in \text{Slip}_{r, \mathbb{P}^n} \mid \mathcal{J} \subset \text{Ann}(F) \subset T \}.$$

One case of interest is when $r = \underline{\mathbf{r}}(F)$.

3 Wildness

Definition 3.1 A form $F \in S^d V^*$ is *wild* if $sr(F) > \underline{\mathbf{r}}(F)$.

Remark 3.2 A consequence of wildness of a form F is that all the ideals in $\underline{\mathbf{VSP}}(F, \underline{\mathbf{r}}(F))$ are *not saturated*.

Before proving the next result, we introduce another piece of notation. Let $W \subset U$ be finite-dimensional vector spaces. Then $W^\perp = \{h \in U^* \mid h(w) = 0 \text{ for all } w \in W\}$ is the *annihilator* of W .

Proposition 3.3 Let $R \subset \mathbb{P}^n$ be a projective scheme defined by the saturated ideal \mathcal{J} such that $\langle R \rangle = \mathbb{P}^n$. Then, for any $d \geq 1$, the affine variety $\mathbb{C}[(\mathcal{J}_d)^\perp]$ has dimension $n + 1$, i.e. the linear space $(\mathcal{J}_d)^\perp$ is spanned by at least $n + 1$ algebraically independent forms.

Proof Choose a linear form z such that $\mathcal{J} : z = \mathcal{J}$, i.e. \mathcal{J} is saturated with respect to z , or equivalently, there is no associated prime containing z . Such a linear form exists by assumption.

Let $S = \mathbb{C}[z, x_1, \dots, x_n]$ be the homogeneous coordinate ring of the ambient projective space. Up to change of basis, we can present the homogeneous saturated ideal of R as follows:

$$\mathcal{J} = \left\langle z^{d_1-1} \ell_1 + h_1, \dots, z^{d_k-1} \ell_k + h_k \right\rangle,$$

where the ℓ_i are (possibly zero) linear forms and the h_j are forms of degree d_j that are quadratic in the variables x_i .

Let $W = \mathcal{J}_d$. Note that $z^d \in W^\perp$ and let $V = \langle z^{d-1}x_1, \dots, z^{d-1}x_n \rangle$. By assumption $\mathcal{J}_1 = 0$. Notice that $V \cap W = 0$. Indeed, if $z^{d-1}l(x_1, \dots, x_n) \in W \subset \mathcal{J}$ with l a linear form, then $l(x_1, \dots, x_n) \in \mathcal{J}$, which is in contradiction with our assumption.

The condition $V \cap W = 0$ implies that we have a surjection

$$S_d^* \supset W^\perp \twoheadrightarrow S_d^*/V^\perp = V^* \subset S_d^*.$$

Thus we can lift the basis of $V^* \subset S_d^*$ consisting of the vectors $z^{d-1}x_1, \dots, z^{d-1}x_n$ (by abuse of notation, the duals of $z^{d-1}x_i$ are denoted in the same way) to an independent set in W^\perp . Therefore:

$$(\mathcal{J}_d)^\perp \supseteq \left\langle z^d, z^{d-1}x_1 + g_1, \dots, z^{d-1}x_n + g_n \right\rangle.$$

To show that the forms on the right-hand side are algebraically independent, we dehomogenize them and look at the affine map they induce:

$$\begin{aligned} \varphi : \mathbb{C}^n &\longrightarrow \mathbb{C}^n, \\ (x_1, \dots, x_n) &\longmapsto (x_1 + \tilde{g}_1, \dots, x_n + \tilde{g}_n). \end{aligned}$$

Notice that the differential of φ at the origin $\mathbf{0} \in \mathbb{C}^n$ is the identity. Thus there is a local isomorphism between tangent spaces and so the dimension of the image is n . Therefore the dimension of the affine variety $\mathbb{C}[(\mathcal{J}_d)^\perp]$ is $n + 1$. This is equivalent to the fact that the linear space $(\mathcal{J}_d)^\perp$ is spanned by at least $n + 1$ algebraically independent degree d forms. □

Theorem 3.4 *Let $d \geq 3$ and $F \in S^d V^*$ be a concise form of minimal border rank. Then:*

$$\text{Hess}(F) = 0 \implies \text{cr}(F) > \underline{r}(F) = n + 1 \implies F \text{ is wild.}$$

Proof Let $W = \langle \frac{\partial}{\partial x_i} F \rangle \subset \mathbb{P}(S^{d-1} V^*)$ be the linear space spanned by the first derivatives of F . Let $\mathcal{I} = \langle \text{Ann}(F)_{d-1} \rangle$ be the homogeneous ideal generated by the degree $(d - 1)$ homogeneous piece of the annihilator of F .

Note that $W^\perp = \mathcal{I}_{d-1}$. Indeed, the inclusion $\mathcal{I}_{d-1} \subset W^\perp$ is clear by definition. To see the converse, let $h \in W^\perp$. Hence $h \circ \left(\frac{\partial}{\partial x_i} F \right) = 0$ for every i ; the latter implies $\frac{\partial}{\partial x_i} (h \circ F) = 0$ for every i , where $h \circ F$ is a linear form. Thus $h \circ F = 0$ and so $h \in \mathcal{I}_{d-1}$.

Let R^W be the projective scheme in \mathbb{P}^n defined by \mathcal{I}^{sat} , which is a priori possibly empty. As $(\mathcal{I}^{sat})_{d-1} \supseteq \mathcal{I}_{d-1}$, we have:

$$W = W^{\perp\perp} = (\mathcal{I}_{d-1})^\perp \supseteq ((\mathcal{I}^{sat})_{d-1})^\perp.$$

Assume the saturated ideal \mathcal{I}^{sat} does not contain any linear form. By definition, this is equivalent to the subscheme $R^W \subset \mathbb{P}^n$ spanning the whole \mathbb{P}^n . By Proposition 3.3, $((\mathcal{I}^{sat})_{d-1})^\perp$ is spanned by at least $n + 1$ algebraically independent forms. However, W has dimension exactly $n + 1$, so $W = ((\mathcal{I}^{sat})_{d-1})^\perp$ and a basis of W consists of $n + 1$ algebraically independent forms. Therefore, the derivatives of F must be algebraically independent and so $\text{Hess}(F) \neq 0$; see [29, §7.2]. This shows that whenever F is concise and has vanishing Hessian, \mathcal{I}^{sat} must contain a linear form.

Suppose F is concise with $\text{Hess}(F) = 0$ and of minimal border rank. Then the Hilbert function of $\text{Ann}(F)$ is as follows:

$$\text{HF}(T/\text{Ann}(F)) : 1 \ (n + 1) \ \dots \ (n + 1) \ 1$$

Now, we show by contradiction the first implication in the statement. Suppose the cactus rank of F satisfies $\text{cr}(F) \leq n + 1$.

Let $\mathcal{J} \subset \text{Ann}(F)$ be any saturated ideal evincing the cactus rank of F , i.e. the zero-dimensional scheme defined by \mathcal{J} has degree $\text{cr}(F)$. Since its Hilbert function $\text{HF}(T/\mathcal{J})$ is non-decreasing until it stabilizes to the constant polynomial $\text{cr}(F) \in \mathbb{N}$ [18, Theorem 1.69], one has

$$\dim (T/\mathcal{J})_{d-1} \leq n + 1 = \dim (T/\mathcal{I})_{d-1}.$$

On the other hand, $\mathcal{J}_{d-1} \subset \mathcal{I}$ and so

$$\dim (T/\mathcal{J})_{d-1} \geq \dim (T/\mathcal{I})_{d-1} .$$

The inequalities imply $\mathcal{J}_{d-1} = \mathcal{I}_{d-1}$.

Now, $\mathcal{I}^{sat} \subset \mathcal{J}^{sat} = \mathcal{J}$. Hence \mathcal{J} contains a linear form, i.e.

$$\dim (T/\mathcal{J})_1 \leq n .$$

On the other hand, since $\mathcal{J} \subset \text{Ann}(F)$, one has:

$$n + 1 = \dim (T/\text{Ann}(F))_1 \leq \dim (T/\mathcal{J})_1 \leq n ,$$

which is a contradiction. Therefore $\text{sr}(F) \geq \text{cr}(F) > n + 1 = \underline{\mathbf{r}}(F)$. Hence F is wild. □

Remark 3.5 Keep the notation from Theorem 3.4. It is clear that for $d = 2$ and any $n \geq 1$, the condition $\text{Hess}(F) = 0$ is equivalent to F being not concise. Also, for $d = 3$ and $n \leq 3$, $\text{Hess}(F) = 0$ is equivalent to F being not concise [29, Theorem 7.1.4]. A complete classification is known up to $n \leq 6$; see [29, §7.6] for a detailed discussion.

Remark 3.6 (Limits of catalecticants and vector bundles)

Let $F \in S^d V^*$ be a form with annihilator $\mathcal{I} = \text{Ann}(F) \subset T$. For each $0 \leq i \leq d$, F induces a linear map

$$\text{Cat}_{i,d-i} : S^i V \longrightarrow S^{d-i} V^* , \quad h \mapsto h \circ F .$$

The map $\text{Cat}_{i,d-i}$ is the i -th *catalecticant*. Its kernel satisfies the equality $\text{Ker}(\text{Cat}_{i,d-i}) = \mathcal{I}_i$, and therefore $\text{HF}(T/\mathcal{I}, i)$ is the rank of the matrix representing the i -th catalecticant.

The rank of any catalecticant is a lower bound to $\underline{\mathbf{r}}(F)$, the border rank of F . Gałazka [7] showed that degeneracy conditions of vector bundles vanish on cactus varieties. In our context, this means that the rank of any catalecticant (and so any value $\text{HF}(T/\mathcal{I}, i)$) gives a lower bound to $\text{cr}(F)$, the cactus rank of F . It is inherently difficult to detect the true cactus rank when it is not witnessed by the rank of some catalecticant (which is the case, in the situation of Theorem 3.4), because one has to choose suitable (strictly contained) linear subspaces of the vector spaces \mathcal{I}_i , which is very hard in practice. Furthermore, one has to construct linear subspaces so that the resulting ideal of a zero-dimensional scheme has the minimal degree allowed. In conclusion, for explicit forms (and, even more so, for a sequence of forms), calculating cactus rank in several variables is usually a daunting task.

Moreover, the smoothable rank of a form is a priori even more difficult to compute explicitly: the general obstacle to overcome is being able to recognize ideals of smoothable schemes of some length r in a given projective space \mathbb{P}^n (whenever the Hilbert scheme $\text{Hilb}_r(\mathbb{P}^n)$ is reducible). However, these are largely unknown.

Example 3.7 The assumption of minimal border rank is independent from the vanishing Hessian condition. Let $F = v_0u_0^3 + v_1u_0^2u_1 + v_2u_0u_1^2 \in S^4\mathbb{C}^{5*}$. So $\text{Hess}(F) = 0$, as the derivatives with respect to v_i are algebraically dependent. One has

$$\text{HF}(T/\text{Ann}(F)) = 1 \ 5 \ 6 \ 5 \ 1.$$

By Remark 3.6, we have $\underline{\mathbf{r}}(F) \geq 6$ and so F is not of minimal border rank.

For a cubic example, let $n \geq 8$. Consider $F = x_0x_{n-1}^2 + x_1x_{n-1}x_n + x_2x_n^2 + G(x_3, x_4, \dots, x_n) \in S^3\mathbb{C}^{(n+1)*}$, with $G = G(x_3, x_4, \dots, x_n) \in S^3\mathbb{C}^{(n-2)*}$ being a general cubic form. By an algebraic relation among the derivatives of F with respect to x_0, x_1, x_2 , one has $\text{Hess}(F) = 0$. Moreover, we have $\underline{\mathbf{r}}(F) \geq \underline{\mathbf{r}}(G)$, for G is a degeneration of F . Whence the border rank of F is generically higher than minimal.

In the *non-minimal* border rank regime, we give the following examples.

Example 3.8 (Wild + Vanishing Hessian) Let $H_5 = v_0u_0^4 + v_1u_0^2u_1^2 + v_2u_1^4 \in S^5(\mathbb{C}^5)^*$. Since the partial derivatives with respect to the v_i are algebraically dependent, one has $\text{Hess}(H_5) = 0$. A computer algebra calculation reveals

$$\text{HF}(T/\text{Ann}(H_5)) : 1 \ 5 \ 7 \ 7 \ 5 \ 1.$$

By Lemma 6.1, the form G_5 satisfies $\underline{\mathbf{r}}(G_5) = 7$. Since H_5 is a degeneration of G_5 (i.e. some of the v_i appearing in G_5 are sent to zero), $\underline{\mathbf{r}}(H_5) \leq \underline{\mathbf{r}}(G_5) = 7$. On the other hand, since the values of the Hilbert function of $\text{Ann}(F)$ give a lower bound for $\underline{\mathbf{r}}(F)$ (see Remark 3.6), one derives the equality $\underline{\mathbf{r}}(H_5) = 7$.

As recalled in Remark 3.6, the values of the Hilbert function of $\text{Ann}(F)$ give a lower bound for the cactus rank $\text{cr}(F)$ as well. Therefore $\text{cr}(H_5) \geq 7$. We now show that $\text{cr}(H_5) > 7 = \underline{\mathbf{r}}(H_5)$, thus proving the wildness of H_5 .

For the sake of contradiction, assume $\text{cr}(H_5) = 7$ and let \mathcal{J} be the saturated ideal of a zero-dimensional scheme of length 7 such that $\mathcal{J} \subset \text{Ann}(H_5)$.

Since $\mathcal{J} \subset \text{Ann}(H_5)$, we have $\text{HF}(T/\mathcal{J}, 2) \geq \text{HF}(T/\text{Ann}(H_5), 2) = 7$ and $\text{HF}(T/\mathcal{J}, 3) \geq \text{HF}(T/\text{Ann}(H_5), 3) = 7$. Since \mathcal{J} is the ideal of a zero-dimensional scheme of length 7, by the stabilization of its Hilbert function, we thus conclude $\text{HF}(T/\mathcal{J}, 2) = \text{HF}(T/\mathcal{J}, 3) = 7$. Therefore we have the equalities $\mathcal{J}_2 = \text{Ann}(H_5)_2$ and $\mathcal{J}_3 = \text{Ann}(H_5)_3$, and hence $\mathcal{I} = \text{Ann}(H_5)_{\leq 3} \subset \mathcal{J}$.

Using a computer algebra system, one checks that the saturation of \mathcal{I} satisfies the equality $\mathcal{I}^{\text{sat}} = (x_0, x_1, x_2)$, where x_i is the dual form to v_i . Since $\mathcal{I}^{\text{sat}} \subset \mathcal{J}^{\text{sat}} = \mathcal{J}$, in particular the ideal \mathcal{J} contains a linear form. As in the very last part of the proof of Theorem 3.4, this leads to a contradiction. In conclusion, $\text{sr}(H_5) \geq \text{cr}(H_5) > 7 = \underline{\mathbf{r}}(H_5)$ and H_5 is wild.

Example 3.9 (Wild + Non-vanishing Hessian) Let $F = v_0u_0^3u_1 + v_1u_0u_1^3 + v_0^3v_1^2 \in S^5\mathbb{C}^{4*}$. This satisfies $\text{Hess}(F) \neq 0$. Let $A = T/\text{Ann}(F)$ and let $\{\alpha_i^{(2)}\}$ be a basis of A_2 . Then the second Hessian $\det(\alpha_i^{(2)}\alpha_j^{(2)}F) = \text{Hess}^2(F) = 0$. This is a classical example due to Ikeda [19], and further revisited by Maeno and Watanabe [24, Example 5.3]. This example was independently found by Dias and Gondim [5, Example 3.15],

along with other interesting families of instances with the property of being wild and yet having non-vanishing Hessian.

Now, turning to details, a computer algebra system computation reveals:

$$\text{HF}(T/\text{Ann}(F)) = 1 \ 4 \ 10 \ 10 \ 4 \ 1.$$

Note that $\text{Ann}(F)_2 = 0$. We first show that $\underline{\mathbf{r}}(F) = 10$. From the Hilbert function values and by Remark 3.6, we have $\underline{\mathbf{r}}(F) \geq 10$. To see the upper bound, let us divide the monomials of F as follows:

$$F = L_5 + v_0^3 v_1^2.$$

Since border rank is subadditive, we have $\underline{\mathbf{r}}(F) \leq \underline{\mathbf{r}}(L_5) + \underline{\mathbf{r}}(v_0^3 v_1^2)$. A straightforward computation gives $\underline{\mathbf{r}}(v_0^3 v_1^2) = 3$. Now, Lemma 6.1 below shows that $\underline{\mathbf{r}}(G_5) = 7$. Note that L_5 is a degeneration of G_5 (i.e. some of the v_i appearing in G_5 are sent to zero). Thus $\underline{\mathbf{r}}(L_5) \leq 7$. In fact, we see that

$$\text{HF}(T/\text{Ann}(L_5)) = 1 \ 4 \ 7 \ 7 \ 4 \ 1,$$

and hence $\underline{\mathbf{r}}(L_5) = 7$. In conclusion, the equality $\underline{\mathbf{r}}(F) = 10$ follows.

For the sake of contradiction, suppose $\text{cr}(F) = 10$ and let $\mathcal{J} \subset \text{Ann}(F)$ be the saturated ideal of a zero-dimensional scheme of length 10. Recall that $\text{Ann}(F)_2 = 0$ and so is \mathcal{J}_2 . Since $\mathcal{J} \subset \text{Ann}(F)$, one has $\text{HF}(T/\mathcal{J}, 3) \geq \text{HF}(T/\text{Ann}(F), 3) = 10$. By the stabilization property of the Hilbert function, it follows that $\text{HF}(T/\mathcal{J}, 3) = 10$ and so $\mathcal{I} = \text{Ann}(F)_{\leq 3} \subset \mathcal{J}$. Therefore $\mathcal{I}^{\text{sat}} \subset \mathcal{J}^{\text{sat}} = \mathcal{J}$.

It is a direct calculation to show that $\mathcal{I}^{\text{sat}} \subset T$ contains a quadric and so does \mathcal{J} . This leads to a contradiction as $\mathcal{J}_2 = 0$. In conclusion, we must have $\text{sr}(F) \geq \text{cr}(F) > 10 = \underline{\mathbf{r}}(F)$, and hence F is wild.

Example 3.10 (Non-wild + Vanishing Hessian) Let $F = v_0 u_0^3 + v_1 u_1^3 + v_2 (u_0 + u_1)^3 \in S^4(\mathbb{C}^5)^*$. Then it is straightforward to check that $\text{Hess}(F) = 0$ and $\underline{\mathbf{r}}(F) = 6$. Note that F is in the span of a scheme R consisting of three 2-jets and so smoothable. Thus $\text{sr}(F) = 6$. Therefore F is not wild.

Proposition 3.11 *Keep the notation from Theorem 3.4. Let F be a concise and minimal border rank cubic. If F is not a wild cubic and R^W is reduced, then F is a Fermat cubic (up to scaling variables) $F = x_0^3 + \dots + x_n^3$, and $\underline{\text{VSP}}(F, n + 1)$ is a single point.*

Proof Since F is concise, of minimal border rank and it is not wild, the proof of Theorem 3.4 yields that \mathcal{I}^{sat} does not contain any linear form. This is equivalent to $\langle R^W \rangle = \mathbb{P}^n$, where $W = \langle \frac{\partial}{\partial x_i} F \rangle \subset \mathbb{P}(S^2 V^*)$. Since R^W is a reduced scheme by assumption, we may find a reduced zero-dimensional subscheme $Z \subset R^W$ of length $n + 1$ such that $\langle Z \rangle = \mathbb{P}^n$. Up to change of basis, we have:

$$((\mathcal{I}_Z)_2)^\perp = \langle x_0^2, \dots, x_n^2 \rangle \subseteq ((\mathcal{I}^{\text{sat}})_2)^\perp \subseteq W.$$

Since $\dim W = n + 1$, we have $W = \langle x_0^2, \dots, x_n^2 \rangle$ and

$$\mathcal{I} = \text{Ann}(F)_2 = W^\perp = \langle x_0^2, \dots, x_n^2 \rangle^\perp.$$

Thus $\text{Ann}(F)_3 = \langle x_0^3, \dots, x_n^3 \rangle^\perp$. Then, since every cubic monomial divisible by two distinct variables is in $\text{Ann}(F)_3$, F is a Fermat cubic up to the action of a diagonal matrix. Now, the Hilbert function of \mathcal{I} is:

$$\text{HF}(T/\mathcal{I}) : 1 \ (n + 1) \ (n + 1) \ (n + 1) \ \dots$$

For $k = 0, 1, 2$, the dimension of \mathcal{I}_k is clear from definitions. To see the dimension of \mathcal{I}_k for $k \geq 3$, note that $y_0^k, \dots, y_n^k \notin \mathcal{I}_k$ and they are the only missing monomials. Let $\mathcal{J} \in \underline{\text{VSP}}(F, n + 1)$. Then its Hilbert function is

$$\text{HF}(T/\mathcal{J}) : 1 \ (n + 1) \ (n + 1) \ (n + 1) \ \dots,$$

because $\mathcal{J} \in \text{Slip}_{n+1, \mathbb{P}^n}$. Since $\mathcal{J} \subset \text{Ann}(F)$, we have the equality $\mathcal{I} = \mathcal{J}$. □

Repeating part of the proof above, one shows:

Proposition 3.12 *Keep the notation from Theorem 3.4. Let F be a concise and minimal border rank cubic. If F is not a wild cubic and $\mathcal{I} = \text{Ann}(F)_2$ is saturated of degree $n + 1$, then $\underline{\text{VSP}}(F, n + 1) = \{\mathcal{I}\}$.*

Example 3.13 Let $F = x_0^2x_1 \in S^3\mathbb{C}^{2*}$. In this case, $\mathcal{I} = \text{Ann}(F)_2 = \langle y_1^2 \rangle \subset T$ is the ideal of a 2-jet on a \mathbb{P}^1 . Proposition 3.12 gives $\underline{\text{VSP}}(F, 2) = \{\mathcal{I}\}$.

Example 3.14 Let $F_{\text{fg}} = x_1(x_0^2 + x_1x_2) \in S^3\mathbb{C}^{3*}$ (a conic with a tangent line) or $F_{\text{cusp}} = x_1^2x_2 - x_0^3 \in S^3\mathbb{C}^{3*}$ (a cuspidal cubic). In both cases, $\mathcal{I} = \text{Ann}(F)_2$ is saturated of degree 3. For F_{fg} , the scheme defined by \mathcal{I} is the 2-fat point (of length 3) in \mathbb{P}^2 ; for F_{cusp} , the scheme defined by \mathcal{I} is the union of a simple point and a 2-jet.

In both cases, Proposition 3.12 yields $\underline{\text{VSP}}(F, 3) = \{\mathcal{I}\}$.

4 Smoothable algebras, structure tensors and wild forms

In this section, for the ease of notation, we regard $F \in S^dV$ as forms (instead of using duals). A tensor in $V^{\otimes d}$ is called a d -way tensor.

Recall that a form $F \in S^dV$ is a symmetric tensor $T_F \in V^{\otimes d}$ (the identification is defined in characteristic zero). Here, we bring tensors into the picture in order to establish the converse to Theorem 3.4. With this aim at hand, we start making a detour through smoothable algebras and structure tensors. We shall demonstrate Theorem 4.9, providing a classification of wild forms of minimal border rank, thus complementing Theorem 3.4.

In the rest, a finite-dimensional \mathbb{C} -algebra A is a finite-dimensional \mathbb{C} -vector space with an associative unital multiplication.

Definition 4.1 (*Smoothable algebras*) Let A be a (commutative) \mathbb{C} -algebra of dimension zero as a ring and $n + 1$ as a \mathbb{C} -vector space. The algebra A is said to be *smoothable* if it is a degeneration of the standard algebra structure on \mathbb{C}^{n+1} . Equivalently, in scheme theory terminology, $\text{Spec}(A)$ is a smoothable zero-dimensional scheme. The dimension of A as a \mathbb{C} -vector space is the length of $\text{Spec}(A)$.

Since we deal with smoothable algebras A , henceforth we restrict our discussion to commutative algebras. For details about the subsequent material, we refer to [9, Lecture 8] or [20, §3 and §16].

A finite-dimensional algebra A is an Artinian ring, so A has finitely many maximal ideals. Let A be a local finite-dimensional \mathbb{C} -algebra (i.e., A is a local ring) with maximal ideal \mathfrak{m} . Its *socle* is the set of all ring elements $a \in A$ such that $a \in (0 : \mathfrak{m})$, i.e. $a \cdot \mathfrak{m} = 0$.

Definition 4.2 (*Gorenstein algebras*)

Let A be a local finite-dimensional \mathbb{C} -algebra. The algebra A is (local) *Gorenstein* if one of the following two equivalent conditions hold true:

- (i) its socle is a one-dimensional \mathbb{C} -vector space;
- (ii) there exists a perfect pairing $p : A \times A \rightarrow \mathbb{C}$, defined by $p(a, b) = e(ab)$ for a linear form $e : A \rightarrow \mathbb{C}$; see e.g. [20, Theorem 3.15].

A finite-dimensional algebra A is said to be *Gorenstein* if every localization $A_{\mathfrak{m}}$ at a maximal ideal $\mathfrak{m} \subset A$ is a local Gorenstein algebra.

Definition 4.3 (*The d -way structure tensor of an algebra*) Let A be a unital, commutative finite-dimensional \mathbb{C} -algebra. The multiplication map $m_A : A \times \dots \times A \rightarrow A$ given by $m_A(a_1, a_2, \dots, a_{d-1}) = a_1 a_2 \dots a_{d-1}$ is multilinear and symmetric. Therefore, one may regard it as a partially symmetric tensor $T_A \in S^{d-1}A^* \otimes A \subseteq A^* \otimes \dots \otimes A^* \otimes A$. The tensor T_A is the *d -way structure tensor* of A . In particular, when $d = 3$, this is the usual *structure tensor* of A [2].

Let V be a finite-dimensional \mathbb{C} -vector space. An (abstract) tensor $T \in V^* \otimes \dots \otimes V^* \otimes V$ is the *d -way structure tensor* of an algebra A if T is isomorphic to T_A , i.e. if there exist d linear isomorphisms $\varphi_1, \dots, \varphi_d$ such that $(\varphi_1 \otimes \varphi_2 \otimes \dots \otimes \varphi_d)(T) = T_A$, where φ_i is a linear isomorphism among the i -th factors.

We summarize the previously known results about (3-way) structure tensors.

Definition 4.4 (*1-generic tensors*) Let $T \in V \otimes V \otimes V$ be a tensor. Then T is *1-generic* if its contraction in every factor $T(V^*)$ is a linear space containing a full-rank matrix $M \in T(V^*) \subset V \otimes V$. If T is symmetric, it is enough to require that a contraction in only one factor has the desired property.

Remark 4.5 Let V_1, V_2 and V_3 be three finite-dimensional \mathbb{C} -vector spaces. Suppose $T \in V_1 \otimes V_2 \otimes V_3$ is a structure tensor of some algebra A . Then the linear spaces $T(V_1^*) \subset V_2 \otimes V_3$ and $T(V_2^*) \subset V_1 \otimes V_3$ contain full rank elements. (T is said to be *binding* [2, Lemma 3.5], or 1_{V_1} - and 1_{V_2} -generic [21, §1].)

Bläser and Lysikov characterized 3-way structure tensors of algebras of minimal border rank as the ones coming from smoothable algebras [2, Theorem 3.2, Corollary 3.3]:

Theorem 4.6 *A 3-way tensor has minimal border rank and is 1_{V_1} - and 1_{V_2} -generic if and only if it is isomorphic to a structure tensor T_A of a smoothable algebra A .*

Now we establish a similar result for d -way structure tensors, for arbitrary d , in Proposition 4.8 through the following lemma.

Lemma 4.7 *Let $F \in S^d V$ be a concise form regarded as a symmetric concise tensor $T_F \in V^{\otimes d}$. The following statements are equivalent:*

- (i) $\text{Hess}(F) \neq 0$;
- (ii) *there exists $\ell \in V^*$ such that the contraction $T_F(\ell^{\otimes d-2})$ is a full-rank symmetric matrix.*

In particular, a cubic $F \in S^3 V$ has non-vanishing Hessian if and only if T_F is a 1-generic symmetric tensor.

Proof Let $A = T/\text{Ann}(F)$ be the Artinian Gorenstein \mathbb{C} -algebra of F . The condition $\text{Hess}(F) \neq 0$ is equivalent to the fact that there exists a linear form $\ell \in A_1$ such that $\varphi_{\ell^{d-2}} : A_1 \rightarrow A_{d-1}$ is an isomorphism (the map here is multiplication by ℓ^{d-2}); see the proof of [24, Theorem 3.1] or [29, Theorem 7.2.20]. The map $\varphi_{\ell^{d-2}}$ is an isomorphism if and only if the quadric $Q = \ell^{d-2}(F) \in S^2 V$ is non-degenerate. \square

We will use the previous equivalence for concise forms of minimal border rank having non-vanishing Hessian. Before proceeding further, notice that if $F \in S^d V$ is a concise form of minimal border rank, then its corresponding tensor T_F has minimal border (tensor) rank.

Proposition 4.8 *Let V be an $(n + 1)$ -dimensional \mathbb{C} -vector space. Let $F \in S^d V$ be a concise form of minimal border rank, and such that $\text{Hess}(F) \neq 0$. Then T_F is the d -way structure tensor of a smoothable Gorenstein algebra A .*

Moreover, if T is the structure tensor of an $(n + 1)$ -dimensional smoothable algebra A , then T is (isomorphic to) a symmetric tensor if and only if A is a Gorenstein algebra.

Proof By Lemma 4.7, we can find $\ell \in V^*$ such that the contraction $T_F(\ell^{\otimes d-2})$ is a full-rank symmetric matrix. We first fix a linear basis $v_1 = \ell, v_2, \dots, v_{n+1}$ of V^* . We may regard $T_F(\ell^{\otimes d-2})$ as a bilinear map $Q : V^* \times V^* \rightarrow \mathbb{C}$. From here, we fix another linear basis $v'_1, v'_2, \dots, v'_{n+1}$ of V^* such that $Q(v_i, v'_j) = \delta_{ij}$. Now, we define $T_F^{i_1 i_2 \dots i_d}$ to be $T_F(v_{i_1} \otimes v_{i_2} \otimes \dots \otimes v_{i_{d-1}} \otimes v'_{i_d})$. By definition, $T_F^{i_1 i_2 \dots i_d}$ is symmetric with respect to the indices $(i_1 i_2 \dots i_{d-1})$ and $T_F^{11 \dots 1kl} = \delta_{kl}$.

We define the matrices $M_i := T_F^{1,1, \dots, 1, \bullet, i, \bullet}$ for $1 \leq i \leq n + 1$. We will prove the following identities that are the defining properties of the d -way structure tensor of an algebra.

- **Commutativity:**

$$M_i M_j = M_j M_i.$$

- **Closed under composition:**

$$M_i M_j = \sum_k T_F^{11 \dots 1ijk} M_k.$$

• **Structure property:**

$$M_{i_1} M_{i_2} \dots M_{i_{d-1}} = \sum_{j=1}^{n+1} T_F^{i_1 i_2 \dots i_{d-1} j} M_j.$$

Commutativity Regard $T_F \in (V^{\otimes d-2}) \otimes V \otimes V$ as a 3-way tensor. Since F is concise of minimal border rank, the border rank of T_F as a 3-way tensor is also $n + 1 = \dim V$. The commutativity of $\{M_i\}$ then follows from [21, Lemma 2.6].

Closed under composition Regard $T_F \in (V^{\otimes d-2}) \otimes V \otimes V$ as a 3-way tensor and apply [21, Proposition 2.10].

Structure property Regarding once again T_F as a 3-way tensor, T_F satisfies Strassen’s commutator equations (see [30] or [21, §2.1]), which in the given coordinates are:

$$\sum_k T_F^{i_1 i_2 \dots i_{d-2} j_1 k} T_F^{i'_1 i'_2 \dots i'_{d-2} k j_2} = \sum_k T_F^{i'_1 i'_2 \dots i'_{d-2} j_1 k} T_F^{i_1 i_2 \dots i_{d-2} k j_2}.$$

Using the symmetry of T_F in the first $(d - 1)$ indices, the equality $T_F^{11\dots 1kl} = \delta_{kl}$, and Strassen’s equations, we have the following identities:

$$\begin{aligned} \sum_k T_F^{1j_1 j_2 \dots j_{d-2} k} T_F^{11\dots 1k j_{d-1} j_d} &= \sum_k T_F^{j_1 j_2 \dots j_{d-2} 1k} T_F^{11\dots 1j_{d-1} k j_d} \\ &= \sum_k T_F^{11\dots 1j_{d-1} 1k} T_F^{j_1 j_2 \dots j_{d-2} k j_d} \\ &= \sum_k T_F^{11\dots 1j_{d-1} k} T_F^{j_1 j_2 \dots j_{d-2} k j_d} \\ &= T_F^{j_1 j_2 \dots j_{d-1} j_d}. \end{aligned}$$

Repeatedly applying the above identities, we have the following equalities:

$$\begin{aligned} (M_{i_1} M_{i_2} \dots M_{i_{d-2}})_{ab} &= \sum_{k_1, \dots, k_{d-3}} T_F^{11\dots 1a i_1 k_1} T_F^{11\dots 1k_1 i_2 k_2} T_F^{11\dots 1k_2 i_3 k_3} \dots T_F^{11\dots 1k_{d-3} i_{d-2} b} \\ &= \sum_{k_2, \dots, k_{d-3}} \left(\sum_{k_1} T_F^{11\dots 1a i_1 k_1} T_F^{11\dots 1k_1 i_2 k_2} \right) T_F^{11\dots 1k_2 i_3 k_3} \dots T_F^{11\dots 1k_{d-3} i_{d-2} b} \\ &= \sum_{k_3, \dots, k_{d-3}} \left(\sum_{k_2} T_F^{11\dots 1a i_1 i_2 k_2} T_F^{11\dots 1k_2 i_3 k_3} \right) \dots T_F^{11\dots 1k_{d-3} i_{d-2} b} \\ &= \sum_{k_{d-3}} T_F^{1a i_1 i_2 \dots i_{d-3} k_{d-3}} T_F^{11\dots 1k_{d-3} i_{d-2} b} \end{aligned}$$

$$\begin{aligned}
 &= T_F^{ai_1i_2\dots i_{d-2}b} \\
 &= T_F^{i_1i_2\dots i_{d-2}ab} \\
 &= \sum_k T_F^{i_1i_2\dots i_{d-2}1k} T_F^{11\dots 1akb} \\
 &= \sum_k T_F^{1i_1i_2\dots i_{d-2}k} T_F^{11\dots 1akb} \\
 &= \left(\sum_k T_F^{1i_1i_2\dots i_{d-2}k} M_k \right)_{ab}.
 \end{aligned}$$

Therefore, using the last equality, we find

$$\begin{aligned}
 (M_{i_1} M_{i_2} \cdots M_{i_{d-2}}) \cdot M_{i_{d-1}} &= \left(\sum_k T_F^{1i_1i_2\dots i_{d-2}k} M_k \right) \cdot M_{i_{d-1}} \\
 &= \sum_j \sum_k T_F^{1i_1i_2\dots i_{d-2}k} T_F^{11\dots 1ki_{d-1}j} M_j \\
 &= \sum_j T_F^{i_1i_2\dots i_{d-2}i_{d-1}j} M_j,
 \end{aligned}$$

where in the second line we use the identity in the **Closed under composition** property. This establishes the **Structure property**.

Let $M_n(\mathbb{C})$ be the algebra of $n \times n$ complex matrices. Consider the map

$$\varphi: \mathbb{C}[x_1, \dots, x_{n+1}] \rightarrow M_n(\mathbb{C}),$$

defined by $\varphi(x_i) = M_i$. Because of the three properties above of the matrices $\{M_i\}$, this is a ring map whose image is an $(n + 1)$ -dimensional vector space.

Define the algebra $A := \mathbb{C}[x_1, \dots, x_{n+1}]/\ker \varphi$. Then by definition and by the three properties above, T_F is the d -way structure tensor of A . As T_F is of minimal border rank, so is the 3-way structure tensor of the algebra A . Thus A is smoothable by Theorem 4.6. This finishes the proof of the first part of the proposition (the Gorenstein property will be proven as corollary of the second part).

For the second part, assume T is a d -way structure tensor of an $(n + 1)$ -dimensional smoothable algebra A . Let $u \in A$ be the identity element in A . Then $T(u^{\otimes d-2}): A \rightarrow A$ is the identity matrix. Hence contracting T with respect to the $(d - 2)$ left-most factors using only tensors of rank 1 produces a full-rank element.

Suppose T is symmetric. Then contracting T with respect to the $(d - 2)$ right-most factors using only tensors of rank 1 can produce a full-rank element. Regarding $T \in (A^*)^{\otimes d-1} \otimes A$, we may find $a_3 \otimes a_4 \otimes \cdots \otimes a_{d-1} \otimes \alpha^k \in A^{\otimes d-3} \otimes A^*$ such that $T(a_3 \otimes \cdots \otimes a_{d-1} \otimes \alpha^k): A \times A \rightarrow \mathbb{C}$ is a perfect pairing. Define $e: A \rightarrow \mathbb{C}$ to be $e(a) = \alpha^k(aa_3a_4 \cdots a_{d-1})$. Then $p = T(a_3 \otimes \cdots \otimes a_{d-1} \otimes \alpha^k)$ and the above e give us the required properties in Definition 4.2(ii). Hence A is Gorenstein.

Suppose A is a Gorenstein algebra. Then, by Definition 4.2(ii), there exists a perfect pairing $p : A \times A \rightarrow \mathbb{C}$, where $p(a, b) = e(ab)$ for some linear form $e : A \rightarrow \mathbb{C}$.

Let us fix a nonzero vector $\alpha^k \in A^*$. Since $T \in (A^*)^{\otimes d-1} \otimes A$, its coordinates are functions in the dual vector space, i.e. $a_{i_1} \otimes \dots \otimes a_{i_{d-1}} \otimes \alpha^k \in A^{\otimes d-1} \otimes A^*$ is a function on T . The $a_{i_1} \otimes \dots \otimes a_{i_{d-1}} \otimes \alpha^k$ -coordinate of T is $\alpha^k(a_{i_1} \dots a_{i_{d-1}})$ because T is the d -way structure tensor of A . Recall we have a perfect pairing $p : A \times A \rightarrow \mathbb{C}$ inducing an isomorphism (which we denote with the same name) $p : A \rightarrow A^*$. Letting $c = p^{-1}(\alpha^k)$, by definition one has $\alpha^k(a) = e(ca)$ for all $a \in A$. Now, we choose α^k so that $\alpha^k(a_{i_1} \dots a_{i_{d-1}}) = e(a_k a_{i_1} \dots a_{i_{d-1}})$ for every k , i.e. $\alpha^k = p(a_k)$. This identifies T as a symmetric tensor in $(A^*)^{\otimes d}$. \square

Theorem 4.9 *Let V be an $(n + 1)$ -dimensional \mathbb{C} -vector space. Let $F \in S^d V$ be a concise form of minimal border rank. Then:*

$$\text{cr}(F) = n + 1 \iff \text{Hess}(F) \neq 0 \iff \text{sr}(F) = n + 1.$$

Proof By Theorem 3.4, if $\text{cr}(F) = n + 1$ then $\text{Hess}(F) \neq 0$. We now show:

$$\text{Hess}(F) \neq 0 \implies \text{sr}(F) = n + 1.$$

Note that this is enough to prove the statement, as $\text{sr}(F) = n + 1$ implies $\text{cr}(F) = n + 1$.

Suppose $\text{Hess}(F) \neq 0$. By Proposition 4.8, the symmetric tensor T_F corresponding to F is the d -way structure tensor of a smoothable Gorenstein algebra A . Let $p(a, b) = e(ab)$ be the perfect pairing on A , inducing an isomorphism $p : A \rightarrow A^*$. As in the proof of Proposition 4.8, we may realize T_F as a symmetric tensor in $(A^*)^{\otimes d}$. We fix a linear basis $a_1 = 1, a_2, \dots, a_{n+1}$ of A . The (i_1, \dots, i_d) -th entry of the tensor T_F equals $e(a_{i_1} \dots a_{i_d})$.

We embed the affine scheme $\text{Spec}(A)$ in \mathbb{C}^n via the surjective ring map:

$$\varphi_A : \mathbb{C}[x_2, \dots, x_{n+1}] \rightarrow A,$$

where $x_i \mapsto a_i$. Next, we embed \mathbb{C}^n in $\mathbb{P}^n = \mathbb{P}(A^*)$ and apply the d -th Veronese embedding to $\mathbb{P}(S^d A^*)$. Let $L \subset \mathbb{P}(S^d A^*)$ be the affine subspace given by the complement of the zero locus of the function $1 \cdot 1 \dots 1 \in S^d A$. Write $L = \text{Spec}(\mathbb{C}[y_{i_1 i_2 \dots i_d}])$, where $\{i_1, i_2, \dots, i_d\}$ is a cardinality d multisubset of $\{1, \dots, n + 1\}$ distinct from $\{1, 1, \dots, 1\}$. The embedding $\nu_d(\text{Spec}(A))$ in $\nu_d(\mathbb{C}^n) \subset L \subset \mathbb{P}(S^d A^*)$ is defined by the ring map $y_{i_1 i_2 \dots i_d} \mapsto a_{i_1} a_{i_2} \dots a_{i_d} \in A$.

We claim that T_F belongs to the linear span of $\nu_d(\text{Spec}(A))$. To see this, consider any linear form $h = \sum \lambda_{i_1 i_2 \dots i_d} y_{i_1 i_2 \dots i_d}$ vanishing on $\nu_d(\text{Spec}(A))$. This means that $\sum \lambda_{i_1 i_2 \dots i_d} a_{i_1} a_{i_2} \dots a_{i_d} = 0 \in A$. Applying the linear form e we obtain:

$$0 = e(0) = e\left(\sum \lambda_{i_1 i_2 \dots i_d} a_{i_1} a_{i_2} \dots a_{i_d}\right) = \sum \lambda_{i_1 i_2 \dots i_d} e(a_{i_1} a_{i_2} \dots a_{i_d}) = h(T_F).$$

Hence the linear form h vanishes on T_F . This shows that T_F belongs to the linear span of $\nu_d(\text{Spec}(A))$, which implies $\text{sr}(F) = n + 1$, thus finishing the proof. \square

5 The limiting scheme

We start with the definition of limiting scheme:

Definition 5.1 Let $F \in S^d V^*$ be a form. Suppose we are given a border rank decomposition for F , i.e.

$$F = \lim_{t \rightarrow 0} \frac{1}{t^s} \left(L_1(t)^d + \dots + L_{\underline{r}(F)}(t)^d \right), \quad s \geq 0, \tag{1}$$

where $L_i(t)$ are linear forms. The reduced zero-dimensional scheme whose (closed) points are the $L_j(t)$ is denoted $R(t)$. For each $t \neq 0$, the radical ideal defining $R(t)$ is denoted $\mathcal{I}_{R(t)}$. The flat limit $Z = \lim_{t \rightarrow 0} R(t)$ is called the *limiting scheme* of (1). Note that Z and $R(t)$ ($t \neq 0$) have the same Hilbert polynomial; see e.g. [15, Theorem III.9.9].

We have the following corollary from the proof of [4, Proposition 2.6]:

Corollary 5.2 *Keep the assumptions from Proposition 2.5. Assume*

$$F \in \langle \mathbb{P}\widehat{T}_{z_1}, \dots, \mathbb{P}\widehat{T}_{z_r} \rangle \subset \sigma_r(X).$$

Then we can find a border rank decomposition for F whose limiting scheme is the smooth scheme supported at the r points $\{z_1, \dots, z_r\}$.

Proof Assume $F = \sum_{i=1}^r v_{z_i}$ where $[v_{z_i}] \in \mathbb{P}\widehat{T}_{z_i}$. Let $\hat{z}_i \in \mathbb{C}^{N_{d+1}}$ such that $[\hat{z}_i] = z_i$ and $\sum_i \hat{z}_i = 0$. We can find curves $\hat{z}_1(t), \dots, \hat{z}_r(t)$ in the affine cone \widehat{X} over X such that $\hat{z}_i(0) = \hat{z}_i$ and $\frac{d\hat{z}_i}{dt}(0) = v_{z_i}$. Then we have the following border rank decomposition for F :

$$F = \lim_{t \rightarrow 0} \frac{1}{t} \sum_{i=1}^r \hat{z}_i(t).$$

Since $\{z_1, \dots, z_r\}$ are r distinct points on X , the limiting scheme corresponding to the border rank decomposition above is the smooth scheme supported at the r points $\{z_1, \dots, z_r\}$. □

The next result is a consequence of Buczyńska-Buczyński’s theory:

Theorem 5.3 *The saturation of any ideal in $\underline{\text{VSP}}(F, \underline{r}(F))$ coincides with the ideal of a limiting scheme of a border rank decomposition.*

Proof Any ideal $\mathcal{J} \in \underline{\text{VSP}}(F, \underline{r}(F))$ comes from some border rank decomposition (1); see the proof of [3, Theorem 3.15]. Such a border rank decomposition (1) determines a family of zero-dimensional schemes $R(t)$, each of length $\underline{r}(F)$, such that their ideals $\mathcal{I}_{R(t)}$ have the generic Hilbert function and $\mathcal{J} = \lim_{t \rightarrow 0} \mathcal{I}_{R(t)}$.

By definition of limits, we have

$$\mathcal{J} = \lim_{t \rightarrow 0} \mathcal{I}_{R(t)} \subset \mathcal{I}_{\lim_{t \rightarrow 0} R(t)}.$$

Let $Z = \lim_{t \rightarrow 0} R(t)$ be the limiting scheme of the given border rank decomposition (1). Since Z has length $\underline{r}(F)$, its ideal has the same degree (or Hilbert polynomial) as \mathcal{J} . Since $\mathcal{J} \subset \mathcal{I}_Z$, their saturations coincide. Since \mathcal{I}_Z is saturated by definition, $\mathcal{J}^{sat} = \mathcal{I}_Z$. □

Theorem 5.4 *Suppose $F \in S^d V^*$ is concise and of minimal border rank $n + 1$. Then every border rank decomposition of F determines an ideal in $\underline{VSP}(F, n + 1)$.*

Proof Given any border rank decomposition $F = \lim_{t \rightarrow 0} H(t)$, where each $H(t) = \frac{1}{t^s} (L_1(t)^d + \dots + L_{n+1}(t)^d)$ has rank $n + 1$, one has that $\mathcal{I}_{R(t)} \subset \text{Ann}(H(t))$ ($t \neq 0$) by the classical Apolarity lemma [18, Lemma 1.15]. Since F is concise, we can find $t \neq 0$ such that $H(t)$ is concise. In this case, $R(t)$ consists of $n + 1$ linearly independent points. Therefore $\mathcal{I}_{R(t)}$ has the generic Hilbert function of $n + 1$ points in \mathbb{P}^n . Hence $\mathcal{J} := \lim_{t \rightarrow 0} \mathcal{I}_{R(t)} \in \underline{VSP}(F, n + 1)$. □

6 Wild forms of higher degree and their VSP

Let $d \geq 3$ and define the following infinite series of concise forms of degree d :

$$G_d = \sum_{i=0}^{d-1} v_i u_0^i u_1^{d-1-i}.$$

For $d = 3$, this coincides up to change of variables with the wild cubic form found in [4, §4]. This infinite series has a classical geometric significance: they are the equations of the dual hypersurfaces of the rational ruled surfaces $\mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(1) \oplus \mathcal{O}_{\mathbb{P}^1}(d - 1))$ embedded with the tautological bundle in \mathbb{P}^{d+1} ; see [10, Theorem 3.1] and [8, Chapter 3, Example 3.6]. (In the suggestive classical terminology, these rational surfaces are called *rational normal scrolls with line directrix*.)

Note that, for every $d \geq 3$, the partial derivatives of G_d with respect to the variables v_i are algebraically dependent: their relations coincide with the equations of the usual Veronese embedding of degree $d - 1$ of \mathbb{P}^1 in \mathbb{P}^{d-1} . Thus $\text{Hess}(G_d) = 0$ for $d \geq 3$. Let $\text{Ann}(G_d) \subset T = \mathbb{C}[x_0, \dots, x_{d-1}, y_0, y_1]$ be its annihilator, where x_i is dual to v_i and y_j is dual to u_j .

Lemma 6.1 *For every $d \geq 3$, the form G_d has minimal border rank $d + 2$.*

Proof Let $\ell_0^d, \dots, \ell_d^d$ be $d + 1$ pairwise distinct linear forms in u_0, u_1 . They may be viewed as $d + 1$ distinct points on the degree d rational normal curve $\nu_d(\mathbb{P}^1) \subset \mathbb{P}^d$. They are linearly independent. (This is well-known and can be explicitly checked, for instance, by calculating the corresponding Wronskian matrix at the origin and show it is full rank.) Any other form ℓ_{d+1}^d is linearly dependent to those above, because $\dim S^d \langle u_0, u_1 \rangle = d + 1$.

Up to change of bases one has $G_d = \sum_{i=0}^{d-1} v_i \ell_i^{d-1}$. Let $\mathbb{P}\widehat{T}_{\ell_i^d}$ denote the affine cone of the Zariski tangent space to $v_d(\mathbb{P}^1)$ at ℓ_i^d . Therefore

$$G_d \in \left\langle \mathbb{P}\widehat{T}_{\ell_i^d}, 0 \leq i \leq d-1 \right\rangle.$$

Since $\ell_0^d, \dots, \ell_d^d, \ell_{d+1}^d$ are linearly dependent, Proposition 2.5 yields the inequality $\underline{r}(G_d) \leq d + 2$. Since G_d is concise, $\underline{r}(G_d) \geq d + 2$. Thus equality holds. \square

Corollary 6.2 *The forms G_d are wild.*

Proof By Lemma 6.1, G_d has minimal border rank. Moreover, as noticed above, $\text{Hess}(G_d) = 0$. Theorem 3.4 shows that the degree d forms G_d are wild. \square

Remark 6.3 For every d , one has

$$d + 2 = \underline{r}(G_d) < \text{cr}(G_d) \leq \text{sr}(G_d) \leq 2d,$$

as every 2-jet is smoothable and so is their union.

This corollary complements [4, Theorem 1.3], as follows:

Theorem 6.4 *For every $d \geq 3$, there exist wild forms of degree d .*

Proposition 6.5 *Let $\mathcal{J} \subset T$ be an ideal. Then $\mathcal{J} \in \underline{\text{VSP}}(G_d, d + 2)$ if and only if $\mathcal{J} = \text{Ann}(G_d)_{\leq d-1} + \mathcal{Q}$, where \mathcal{Q} is the principal ideal generated by a form $q \in S^{d+2}(y_0, y_1)$.*

Proof Let $\mathcal{J} \in \underline{\text{VSP}}(G_d, d + 2)$ and let $\mathcal{I} = \text{Ann}(G_d)_{\leq d-1}$.

Then $\mathcal{J} \in \text{Slip}_{d+2, \mathbb{P}^{d+1}}$ and, by definition, the Hilbert function of \mathcal{J} is the generic Hilbert function of $d + 2$ points in \mathbb{P}^{d+1} . Since $\mathcal{J} \subset \text{Ann}(G_d)$, it follows that $\mathcal{J} \supset \mathcal{I}$, as they must coincide up to degree $d - 1$.

Now, consider the Hilbert function of \mathcal{I} . The following relations hold in T/\mathcal{I} :

$$\begin{aligned} x_i x_j &\equiv 0 \pmod{\mathcal{I}} \quad \text{for } 0 \leq i, j \leq d-1, \\ c_i x_i y_1 &\equiv x_{i+1} y_0 \pmod{\mathcal{I}} \quad \text{for } -1 \leq i \leq d-1, \end{aligned}$$

where c_i is a non-zero constant for every i and $x_{-1} = x_d := 0$. Then it is a direct computation to show that $\text{HF}(T/\mathcal{I}, d) = \text{HF}(T/\mathcal{I}, d + 1) = d + 2$ and $\text{HF}(T/\mathcal{I}, d + 2) = d + 3$. Moreover, one has

$$(T/\mathcal{I})_{d+2} = \left\langle y_0^{d+2}, y_0^{d+1} y_1, \dots, y_1^{d+2} \right\rangle = S^{d+2}(y_0, y_1).$$

Thus if $\mathcal{J} \in \underline{\text{VSP}}(G_d, d + 2)$ then \mathcal{J} contains \mathcal{I} and an ideal \mathcal{Q} generated by some element $q \in (T/\mathcal{I})_{d+2}$.

For the converse, let $\mathcal{J}_{\mathcal{Q}} = \text{Ann}(G_d)_{\leq d-1} + \mathcal{Q}$, for some $\mathcal{Q} = \langle q \rangle$ such that $q \in S^{d+2}(y_0, y_1)$. Note that $\text{HF}(T/\mathcal{J}_{\mathcal{Q}}, d + 2) = d + 2$. We can apply Gotzmann’s

Persistence Theorem [13, Theorem 3.8] to conclude that $\text{HF}(T/\mathcal{J}_Q, j) = d + 2$ for all $j \geq d + 2$.

To show that all such \mathcal{J}_Q are in fact in $\underline{\text{VSP}}(G_d, d + 2)$, consider $Q = \langle q \rangle$ where q is a form with $d + 2$ distinct roots. This gives us $d + 2$ distinct points $z_1, \dots, z_{d+2} \in \mathbb{P}(\langle y_0, y_1 \rangle^*)$. By Lemma 6.1 and Corollary 5.2, we may find a border rank decomposition defining a family of zero-dimensional schemes $R(t)$, whose limiting scheme is the smooth scheme Z supported at z_1, \dots, z_{d+2} .

Let \mathcal{I}_Z be the radical ideal defining the limiting scheme Z . By Theorem 5.4, there is a corresponding ideal $\mathcal{J}' \in \underline{\text{VSP}}(G_d, d + 2)$ such that $\mathcal{J}'^{\text{sat}} = \mathcal{I}_Z$. Note that $\mathcal{I}_Z = \mathcal{J}_Q^{\text{sat}}$.

Since $\mathcal{J}' \in \underline{\text{VSP}}(G_d, d + 2)$, by the first part of this proof, $\mathcal{J}' = \mathcal{J}_{Q'}$, for some $Q' = \langle q' \rangle$. Therefore one has $\mathcal{J}_Q^{\text{sat}} = \mathcal{I}_Z$. Since $\mathcal{J}_Q^{\text{sat}} = \mathcal{J}_Q^{\text{sat}}$, we must have $Q = Q'$. In conclusion, we derive $\mathcal{J}_Q = \mathcal{J}' \in \underline{\text{VSP}}(G_d, d + 2)$.

Now, consider the morphism

$$\begin{aligned} \psi_d : \underline{\text{VSP}}(G_d, d + 2) &\longrightarrow \mathbb{P}((T/\mathcal{I})_{d+2}), \\ \mathcal{J}_Q &\longmapsto [q]. \end{aligned}$$

This morphism is proper and hence closed. We have shown that the generic point of $\mathbb{P}((T/\mathcal{I})_{d+2})$ lies in the image, therefore ψ_d is surjective. □

Let $S^{d+2}\mathbb{P}^1$ denote the $(d + 2)$ -fold symmetric product of the projective line. Proposition 6.5 yields the following:

Theorem 6.6 *The projective variety $\underline{\text{VSP}}(G_d, d + 2)$ is isomorphic to the projective space $\mathbb{P}^{d+2} \cong \mathbb{P}(S^{d+2}\mathbb{C}^2) \cong S^{d+2}\mathbb{P}^1$.*

Proof The morphism ψ_d in the proof of Proposition 6.5 is surjective. It is also injective because the point $[q]$ uniquely determines the ideal \mathcal{I}_Q . Since \mathbb{P}^{d+2} is smooth and so normal, the map ψ_d is an isomorphism by a variant of Zariski’s Main Theorem [23, Corollary 4.6]. The isomorphism follows from the description of the vector space $(T/\mathcal{I})_{d+2}$ given in the proof of Proposition 6.5. □

7 An infinite series of wild cubics and their VSP

For every $k \geq 1$ and $n = 3k + 1$, we introduce the following infinite series of concise cubic forms:

$$\begin{aligned} F_n &= x_0x_1^2 + x_1x_2x_4 + x_3x_4^2 + x_4x_5x_7 + x_6x_7^2 \\ &\quad + x_8x_7x_{10} + x_9x_{10}^2 + \dots + x_{n-4}x_{n-3}^2 \\ &\quad + x_{n-3}x_{n-2}x_n + x_{n-1}x_n^2. \end{aligned}$$

Remark 7.1 This infinite series is inspired by the examples appeared in [11], which in turn are a generalization of the *Perazzo cubic hypersurface* $\{F_4 = 0\} \subset \mathbb{P}^4$. The latter is exactly the wild cubic found in [4, §4].

Let $\text{Ann}(F_n) \subset T = \mathbb{C}[y_0, \dots, y_n]$ be the annihilator of F_n with y_i being dual to x_i .

Remark 7.2 Note that F_4 coincides with G_3 from Sect. 6 up to change of basis. So $\text{VSP}(F_4, 5) \cong \mathbb{P}^5$.

The following combinatorial arrangement of lines is important for us to study the infinite series F_n ; Proposition 7.4 provides the motivation for looking at it.

Definition 7.3 (*Chains of lines*) A chain of lines is a collection of distinct lines $C^1, \dots, C^m = \mathbb{P}^1$ such that (up to reindexing) for $1 \leq i, j \leq m$:

$$C^i \cap C^{i+1} \neq \emptyset, \text{ and}$$

$$C^i \cap C^j = \emptyset \text{ otherwise.}$$

Proposition 7.4 *The projective scheme whose ideal is $(\text{Ann}(F_n)_2)^{\text{sat}}$ is a chain of lines.*

Proof Let $\mathcal{I} = \text{Ann}(F_n)_2$ and let \mathcal{I}^{sat} denote its saturation. Let $n = 3k + 1$ for $k \geq 1$. We divide the proof according to the residue (mod 3) of each index $0 \leq j \leq n$ in y_j .

- $j \equiv 0 \pmod{3}$. The monomial $y_j y_i \in \mathcal{I}$ for all $i \neq j + 1$. Moreover, $y_j y_{j+1}^3 \in \mathcal{I}$, therefore $y_j \in \mathcal{I}^{\text{sat}}$.
- $j \equiv 2 \pmod{3}$. In this case, one has $y_j y_i \in \mathcal{I}$ for all $i \neq j - 1, j + 2$. Note that $y_j y_{j-1}^2, y_j y_{j+2}^2 \in \mathcal{I}$. Therefore $y_j \in \mathcal{I}^{\text{sat}}$.
- $j \equiv 1 \pmod{3}$. In this case, $y_j y_{j+3h} \in \mathcal{I} \subset \mathcal{I}^{\text{sat}}$ for $h \geq 2$. For $i \equiv 1 \pmod{3}$, $y_i \notin \mathcal{I}^{\text{sat}}$. We show that $y_i^k \notin \mathcal{I}$ for any $k \geq 2$. Assuming on the contrary that $y_i^k \in \mathcal{I}$, one has $y_i^k = \sum_{j=0}^s m_j h_j$, where the h_j are the generators of \mathcal{I} and $m_j \in T$. However, on the right-hand side, every monomial that is divisible by y_i is divisible by some other distinct y_j as well. We show that $y_i y_{i+3} \notin \mathcal{I}^{\text{sat}}$. On the contrary, suppose $y_i y_{i+3} \in \mathcal{I}^{\text{sat}}$. Thus, there exists $k > 1$ such that $(y_i y_{i+3})^k \in \mathcal{I}$. Using the same argument as above, we see that every monomial in \mathcal{I} , that is divisible by $y_i y_{i+3}$, must be divisible by some other distinct y_j as well.

In conclusion, the saturated ideal \mathcal{I}^{sat} is generated by:

$$\mathcal{I}^{\text{sat}} = \langle y_{3h}, y_{3h+2}, y_{3h+1} y_{3(h+s)+1}, h \geq 0, s \geq 2 \rangle$$

The projective scheme defined by \mathcal{I}^{sat} is a chain of k lines C_k , as in Definition 7.3. Its irreducible components are lines $L_{i,j}$, such that $|i - j| = 3$ and $i \equiv 1 \pmod{3}$, whose ideal is defined by

$$\mathcal{J}_{i,j} = \langle y_k \mid k \neq i, j \rangle.$$

This concludes the proof. □

Notation In the following, let C_k denote the chain of lines appearing in the proof of Proposition 7.4, corresponding to the cubic F_n when $n = 3k + 1$. Moreover, C_k comes equipped with an ordering on its components C_k^h induced by the lexicographic order on the variables $y_1 > y_4 > \dots > y_{3k+1}$. Therefore, the h -th component of C_k refers to the line where the homogeneous coordinates are y_{3h-2} and y_{3h+1} .

Proposition 7.5 *Let $n \geq 4$. The form F_n has minimal border rank $n + 1$.*

Proof By conciseness, $\mathbf{r}(F_n) \geq n + 1$. We show the opposite inequality as follows. Let $n = 3k + 1$ and consider the powers of linear forms

$$\begin{aligned} &\ell_{1,1}^3, \ell_{1,2}^3, \ell_{1,3}^3, \ell_{1,4}^3, \\ &\ell_{2,2}^3, \ell_{2,3}^3, \ell_{2,4}^3, \\ &\dots \dots \dots \\ &\ell_{k-1,2}^3, \ell_{k-1,3}^3, \ell_{k-1,4}^3, \\ &\ell_{k,1}^3, \ell_{k,2}^3, \ell_{k,3}^3, \ell_{k,4}^3, \end{aligned}$$

where $\ell_{i,j}$ is a linear form defined on the i -th component of the chain of lines C_k , i.e. $\ell_{i,j}$ depends only on corresponding two variables. Note that there are $n + 1$ of such forms. Moreover, we require $\ell_{i,j}^3 \neq x_{3i-2}^3, x_{3i+1}^3$ for $1 \leq i \leq k - 1$, where the latter cubes correspond to the intersections between the i -th component of C_k and the other lines in C_k . This may be regarded as a configuration of points on the lines C_k^i of the chain C_k ; an instance of this is depicted in Fig. 1.

Write $\mathbb{P}\widehat{T}_{\ell^3}$ for the affine cone of the Zariski tangent space to $v_3(\mathbb{P}^n)$ at the point ℓ^3 . As in the proof of Proposition 6.1, for $d = 3$, we have

$$F_4 = x_0x_1^2 + x_1x_2x_4 + x_3x_4^2 \in \langle \mathbb{P}\widehat{T}_{\ell_{i,j}^3} \text{ for } 2 \leq j \leq 4 \rangle.$$

More generally, for each $1 \leq i \leq k$, one has:


$$H_i = x_{3i-3}x_{3i-2}^2 + x_{3i-2}x_{3i-1}x_{3i+1} + x_{3i}x_{3i+1}^2 \in \langle \mathbb{P}\widehat{T}_{\ell_{i,j}^3} \text{ for } 2 \leq j \leq 4 \rangle.$$

Thus $F_n \in \langle \mathbb{P}\widehat{T}_{\ell_{i,j}^3} \text{ for } 1 \leq i \leq k, 2 \leq j \leq 4 \rangle$.

The forms $\ell_{i,j}^3$ are linearly dependent. Indeed, consider the forms on the first component C_k^1 of the chain: $\ell_{1,1}^3, \ell_{1,2}^3, \ell_{1,3}^3$ and $\ell_{1,4}^3$ are linearly independent and span every cubic form defined on C_k^1 . Thus the cube x_4^3 may be written as

$$x_4^3 = \sum_{j=1}^4 \lambda_j \ell_{1,j}^3.$$

Fig. 1 The chain of lines C_k and a configuration of points featured in the proof of Proposition 7.5, for $k = 4$ and $n = 13$



On the line C_k^2 , we have the linear relation:

$$\mu_1 x_4^3 + \left(\sum_{j=2}^4 \mu_j \ell_{2,j}^3 \right) + \mu_5 x_7^3 = 0.$$

So we may rewrite the cube x_7^3 as a linear combination of the $\ell_{i,j}^3$ with $1 \leq i \leq 2$. Proceed similarly up to C_k^{k-1} , where we express x_{3k-2}^3 as a linear combination of $\ell_{i,j}^3$ with $1 \leq i \leq k - 1$. Finally, on the line C_k^k use the linear relation among the five cubic forms $x_{3k-2}^3, \ell_{k,1}^3, \ell_{k,2}^3, \ell_{k,3}^3,$ and $\ell_{k,4}^3$. This process gives a linear relation among the $\ell_{i,j}^3$.

Now, to conclude, use Proposition 2.5 which yields $\underline{r}(F_n) \leq n + 1$. □

Remark 7.6 For every $k \geq 1$ and $n = 3k + 1$, one has

$$n + 1 = \underline{r}(F_n) < \text{cr}(F_n) \leq \text{sr}(F_n) \leq 6k = 2(n - 1),$$

as every 2-jet is smoothable and so is their union.

Corollary 7.7 *The cubics F_n are wild.*

Proof Their Hessian is vanishing, as the partial derivatives are algebraically dependent. By Proposition 7.5, F_n has minimal border rank. Theorem 3.4 shows that the cubics F_n are wild. □

Lemma 7.8 *Let $n = 3k + 1$ and let C_k be the chain of lines from Proposition 7.4. Let $\mathcal{J} \in \underline{\text{VSP}}(F_n, n + 1)$. Then:*

- (i) *there exist forms q_1, \dots, q_s in the variables y_{3h+1} 's such that $\mathcal{J} = \text{Ann}(F)_2 + \langle q_1, \dots, q_s \rangle$;*
- (ii) *the ideal \mathcal{J}^{sat} defines a projective scheme that is a zero-dimensional scheme of length $n + 1$ supported on C_k ;*
- (iii) *each form q_j has degree at most $n + 1$;*
- (iv) *there exists a proper map*

$$\psi_k : \underline{\text{VSP}}(F_n, n + 1) \rightarrow \text{Hilb}_{n+1}(C_k),$$

where the latter is the Hilbert scheme of zero-dimensional schemes of length $n + 1$ supported on the reducible curve $C_k \subset \mathbb{P}^n$.

- Proof** (i) From the description of $\mathcal{I} = \text{Ann}(F)_2$ in the proof of Proposition 7.4, it is straightforward to see that each degree $d \geq 4$ graded piece of the quotient ring T/\mathcal{I} has a basis of monomials $\{m_1, \dots, m_h\}$, where each m_j is a monomial in two variables y_{3h-2} and y_{3h+1} ; each such a pair of variables corresponds to a unique line in the chain C_k . Since a necessary condition for membership of an ideal \mathcal{J} in $\text{VSP}(F_n, n + 1)$ is possessing a generic Hilbert function, we add forms q_j (in the variables y_{3h+1} 's) to \mathcal{J} until the ideal reaches a generic Hilbert function.
- (ii) Let $\mathcal{I} = \text{Ann}(F)_2$. Note that the Hilbert polynomial of \mathcal{J}^{sat} is $n + 1$. Moreover, $\mathcal{J}^{sat} \supset \mathcal{I}^{sat}$. By Proposition 7.4, \mathcal{I}^{sat} is the ideal defining the projective scheme C_k . Therefore \mathcal{J}^{sat} defines a zero-dimensional scheme of length $n + 1$ supported on C_k .
- (iii) The Castelnuovo–Mumford regularity of a zero-dimensional scheme of length $n + 1$ is at most $n + 1$ [18, Theorem 1.69], and the degrees of the generators q_j are bounded above by the regularity.
- (iv) By (ii), the projective scheme defined by \mathcal{J}^{sat} is a zero-dimensional scheme of length $n + 1$ supported on C_k . Define the map:

$$\begin{aligned} \psi_k : \text{VSP}(F_n, n + 1) &\rightarrow \text{Hilb}_{n+1}(C_k), \\ \mathcal{J} &\mapsto \mathcal{J}^{sat}. \end{aligned}$$

Then this is a well-defined morphism. It is proper because it is projective. □

Theorem 7.9 *Let $n = 3k + 1 \geq 10$. The variety $\text{VSP}(F_n, n + 1)$ is reducible.*

Proof Since ψ_k is proper by Lemma 7.8(iv), if $\text{VSP}(F_n, n + 1)$ were irreducible, then the image $\psi_k(\text{VSP}(F_n, n + 1))$ would be closed and irreducible. We show next that $\psi_k(\text{VSP}(F_n, n + 1))$ has at least two irreducible components.

As in proof of Proposition 7.5, F_n is in the span of the affine cones of Zariski tangent spaces of $v_3(\mathbb{P}^n)$ at the following points:

$$\begin{aligned} &\ell_{1,2}^3, \ell_{1,3}^3, \ell_{1,4}^3, \\ &\ell_{2,2}^3, \ell_{2,3}^3, \ell_{2,4}^3, \\ &\dots \dots \dots \\ &\ell_{a,1}^3, \ell_{a,2}^3, \ell_{a,3}^3, \ell_{a,4}^3, \\ &\dots \dots \dots \\ &\ell_{b,1}^3, \ell_{b,2}^3, \ell_{b,3}^3, \ell_{b,4}^3, \\ &\dots \dots \dots \\ &\ell_{k,2}^3, \ell_{k,3}^3, \ell_{k,4}^3. \end{aligned}$$

Here we generalize the original configuration described in Proposition 7.5, where the forms $\ell_{1,1}^3$ and $\ell_{k,1}^3$ are replaced by $\ell_{a,1}^3$ and $\ell_{b,1}^3$, with $1 \leq a < b \leq n - 3$ and $a \equiv b \equiv 1 \pmod{3}$. Note that, if $n \geq 10$, there are at least two such pairs (a, b) . To see that the forms $\ell_{i,j}^3$ above are linearly dependent, perform the same procedure presented in the proof of Proposition 7.5, starting from the a -th component and ending at the b -th component of C_k . This produces a linear relation among the $\ell_{i,j}^3$ with $a \leq i \leq b$.

Thus, by Corollary 5.2 and Theorem 5.4, we may find a border rank decomposition given by a family of schemes $R(t)$ over the base $\text{Spec}(\mathbb{C}[t^{\pm 1}])$, whose limiting scheme is supported on the points $\ell_{i,j}^3$ above, with a corresponding ideal $\mathcal{J}_{(a,b)} \in \underline{\text{VSP}}(F_n, n+1)$. Thus $\psi_k(\mathcal{J}_{(a,b)}) \in \text{Hilb}_{n+1}(C_k)$.

As before, let C_k^h denote the h -th component of C_k . Let $\text{Hilb}_{n+1}(C_k)^{(a,b)}$ be the irreducible component of $\text{Hilb}_{n+1}(C_k)$ defined by:

$$\begin{aligned} &\text{Hilb}_{n+1}(C_k)^{(a,b)} \\ &= \overline{\{Z \subset C_k \mid Z \text{ smooth, } Z \cap C_k^a = Z \cap C_k^b = 4, Z \cap C_k^j = 3, j \neq a, b\}}. \end{aligned}$$

Consider the components $\text{Hilb}_{n+1}(C_k)^{(a,b)}$. Notice that $\dim \text{Hilb}_{n+1}(C_k)^{(a,b)} = 3k + 2 = n + 1$. To see this, let $Z \in \text{Hilb}_{n+1}(C_k)^{(a,b)}$ be a general point and so Z is a smooth zero-dimensional scheme. The normal bundle N_{Z/C_k} has $n + 1$ global sections (an affine coordinate at each smooth point of Z). Therefore

$$\begin{aligned} n + 1 &= h^0(N_{Z/C_k}) = \dim T_Z \text{Hilb}_{n+1}(C_k) \\ &= \dim T_Z \text{Hilb}_{n+1}(C_k)^{(a,b)} = \dim \text{Hilb}_{n+1}(C_k)^{(a,b)}. \end{aligned}$$

Next, we show that the components $\text{Hilb}_{n+1}(C_k)^{(1,2)}$ and $\text{Hilb}_{n+1}(C_k)^{(1,3)}$ are distinct. The general element $Z \in \text{Hilb}_{n+1}(C_k)^{(1,2)}$ is such that $Z \cap C_k^2$ is a smooth scheme of length four supported outside $C_k^1 \cap C_k^2$ and $C_k^2 \cap C_k^3$.

However, any flat limit R of zero-dimensional schemes $R(t) \in \text{Hilb}_{n+1}(C_k)^{(1,3)}$, with $\text{length}(R \cap C_k^2) = 4$, is such that

$$\text{either } \text{Supp}(R) \cap C_k^1 \cap C_k^2 \neq \emptyset \quad \text{or} \quad \text{Supp}(R) \cap C_k^2 \cap C_k^3 \neq \emptyset.$$

Thus Z cannot be in the component $\text{Hilb}_{n+1}(C_k)^{(1,3)}$.

Since $\text{Hilb}_{n+1}(C_k)^{(1,2)}$ and $\text{Hilb}_{n+1}(C_k)^{(1,3)}$ have the same maximal dimension, they are two distinct irreducible components of the Hilbert scheme $\text{Hilb}_{n+1}(C_k)$. As the image $\psi_k(\underline{\text{VSP}}(F_n, n + 1))$ is closed, it contains both of them; therefore the image is reducible. In conclusion, $\underline{\text{VSP}}(F_n, n + 1)$ must be reducible. \square

Remark 7.10 The variety $\underline{\text{VSP}}(F_7, 8)$ is at least 8-dimensional. Let $\mathcal{I} = \text{Ann}(F)_2$. One can check there is a generically injective map

$$\begin{aligned} \rho : \underline{\text{VSP}}(F_7, 8) &\longrightarrow \mathbb{P}((T/\mathcal{I})_4) = \mathbb{P}^8, \\ \mathcal{J} &\longmapsto \mathcal{J}_4, \end{aligned}$$

i.e. the generic fiber of ρ is a single point. We do not know whether $\underline{\text{VSP}}(F_7, 8)$ is irreducible or not. The locus where the morphism is injective is then birational to \mathbb{P}^8 . If it is irreducible and 8-dimensional, it cannot be isomorphic to \mathbb{P}^8 : the isomorphic fibers $\rho^{-1}(x_1^4)$ and $\rho^{-1}(x_7^4)$ both contain a linear space \mathbb{P}^4 . Thus these two linear spaces do not intersect.

Question 7.11 Is $\underline{\text{VSP}}(F_7, 8)$ irreducible? Are the irreducible components of the border varieties $\underline{\text{VSP}}(F_n, n + 1)$ rational? More generally, it would be interesting to analyze rationality and unirationality of (the irreducible components of) border varieties of sums of powers $\underline{\text{VSP}}$'s alike in the context of VSP 's; see for instance [25] for several results in this direction.

Acknowledgements We thank Edoardo Ballico, Weronika Buczyńska, Jarosław Buczyński, J. M. Landsberg, Giorgio Ottaviani, and Francesco Russo for very useful discussions. We thank Giorgio Ottaviani for his remark [3, Remark 5.1]: this stimulated the curiosity that led to the present article. We thank an anonymous referee for useful comments and for posing interesting and stimulating questions. The third author is supported by the Vici Grant 639.033.514 of Jan Draisma from the Netherlands Organisation for Scientific Research.

Funding Open access funding provided by University of Bern

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Bernardi, A., Ranestad, K.: On the cactus rank of cubic forms. *J. Symb. Comput.* **50**, 291–297 (2013)
2. Bläser, M., Lysikov, V.: On degeneration of tensors and algebras. In: 41st International Symposium on Mathematical Foundations of Computer Science, LIPIcs. Leibniz Int. Proc. Inform., vol. 58, Schloss Dagstuhl. Leibniz-Zent. Inform., Wadern, pp. Art. No. 19, 11 (2016)
3. Buczyńska, W., Buczyński, J.: *Apolarity, border rank and multigraded Hilbert scheme*. [arXiv:1910.01944](https://arxiv.org/abs/1910.01944) (2019) (**preprint**)
4. Buczyńska, W., Buczyński, J.: On differences between the border rank and the smoothable rank of a polynomial. *Glasg. Math. J.* **57**(2), 401–413 (2015)
5. Dias, T., Gondim, R.: Waring problems and the Lefschetz properties. [arXiv:2006.10865](https://arxiv.org/abs/2006.10865) (2020) (**preprint**)
6. Eisenbud, D., Harris, J.: *The Geometry of Schemes*. Graduate Texts in Mathematics, vol. 197. Springer, Berlin (2000)
7. Gałazka, M.: Vector bundles give equations of cactus varieties. *Linear Algebra Appl.* **521**, 254–262 (2017)
8. Gel'fand, I.M., Kapranov, M.M., Zelevinsky, A.V.: *Discriminants. Resultants and Multidimensional Determinants*. Birkhäuser, Boston (1994)
9. Geramita, A.V.: Inverse systems of fat points: Waring's problem, secant varieties of Veronese varieties and parameter spaces for Gorenstein ideals. In: *The Curves Seminar at Queen's*, vol. X (Kingston, ON, 1995). Queen's University, Kingston, pp. 2–114 (1996)
10. Gondim, R., Russo, F., Staglianò, G.: Hypersurfaces with vanishing hessian via Dual Cayley Trick. *J. Pure Appl. Algebra* **224**, 1215–1240 (2020)
11. Gondim, R., Russo, F.: Cubic hypersurfaces with vanishing Hessian. *J. Pure Appl. Algebra* **219**, 779–806 (2015)
12. Gordan, P., Noether, M.: Über die algebraischen Formen, deren Hesse'sche Determinante identisch verschwindet. *Math. Ann.* **10**, 547–568 (1876)
13. Green, M.: Generic initial ideals, in *Six Lectures on Commutative Algebra*. Mod. Birkhäuser Class., pp. 119–186. Birkhäuser Verlag, Basel (2010)
14. Haiman, M., Sturmfels, B.: Multigraded Hilbert schemes. *J. Algebr. Geom.* **13**(4), 725–769 (2004)

15. Hartshorne, R.: Algebraic Geometry. Springer, Berlin (1977)
16. Hesse, O.: Zur Theorie der ganzen homogenen Functionen. *J. Reine Angew. Math.* **56**, 263–269 (1859)
17. Hesse, O.: Über die Bedingung, unter welche eine homogene ganze Function von n unabhängigen Variablen durch Lineäre Substitutionen von n andern unabhängigen Variablen auf eine homogene Function sich zurück-führen lässt, die eine Variable weniger enthält. *J. Reine Angew. Math.* **42**, 117–124 (1851)
18. Iarrobino, A., Kanev, V.: Power Sums, Gorenstein Algebras, and Determinantal Loci. Lecture notes in Mathematics, vol. 1721. Springer, Berlin (1999)
19. Ikeda, H.: Results on Dilworth and Rees numbers of Artinian local rings. *Jpn. J. Math.* **22**, 147–158 (1996)
20. Lam, T.-Y.: Lectures on Modules and Rings. Graduate Texts in Mathematics, vol. 189. Springer, New York (1999)
21. Landsberg, J.M., Michalek, M.: Abelian tensors. *J. Math. Pures Appl.* (9) **108**, 333–371 (2017)
22. Landsberg, J.M.: Tensors: Geometry and Applications. Graduate Studies in Mathematics, vol. 128. AMS, Providence (2011)
23. Liu, Q.: Algebraic Geometry and Arithmetic Curves. Oxford Graduate Texts in Mathematics, vol. 6. Oxford Science Publications, Oxford University Press, Oxford (2002)
24. Maeno, T., Watanabe, J.: Lefschetz elements of artinian Gorenstein algebras and Hessians of homogeneous polynomials. III. *J. Math.* **53**, 593–603 (2009)
25. Massarenti, A., Mella, M.: Birational aspects of the geometry of varieties of sums of powers. *Adv. Math.* **243**, 187–202 (2013)
26. Mukai, S.: Fano 3- folds. *Lond. Math. Soc. Lect. Note Ser.* **179**, 255–263 (1992)
27. Perazzo, U.: Sulle varietà cubiche la cui hessiana svanisce identicamente. *G. Mat. Battaglini* **38**, 337–354 (1900)
28. Ranestad, K., Schreyer, F.-O.: Varieties of sums of powers. *J. Reine Angew. Math.* **525**, 147–181 (2000)
29. Russo, F.: On the Geometry of Some Special Projective Varieties. Lecture Notes of the Unione Matematica Italiana, vol. 18. Springer, Cham; Unione Matematica Italiana, Bologna (2016)
30. Strassen, V.: Rank and optimal computation of generic tensors. *Linear Algebra Appl.* **52**(53), 645–685 (1983)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.