POLITECNICO DI TORINO Repository ISTITUZIONALE

An optimal multiplier theorem for Grushin operators in the plane, II

Original

An optimal multiplier theorem for Grushin operators in the plane, II / Dall'Ara, Gian Maria; Martini, Alessio. - In: JOURNAL OF FOURIER ANALYSIS AND APPLICATIONS. - ISSN 1069-5869. - ELETTRONICO. - 28:2(2022). [10.1007/s00041-022-09931-9]

Availability: This version is available at: 11583/2959160 since: 2022-03-23T07:52:06Z

Publisher: Springer

Published DOI:10.1007/s00041-022-09931-9

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright Springer postprint/Author's Accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: http://dx.doi.org/10.1007/s00041-022-09931-9

(Article begins on next page)

AN OPTIMAL MULTIPLIER THEOREM FOR GRUSHIN OPERATORS IN THE PLANE, II

GIAN MARIA DALL'ARA AND ALESSIO MARTINI

ABSTRACT. In a previous work we proved a spectral multiplier theorem of Mihlin–Hörmander type for two-dimensional Grushin operators $-\partial_x^2 - V(x)\partial_y^2$, where V is a doubling single-well potential, yielding the surprising result that the optimal smoothness requirement on the multiplier is independent of V. Here we refine this result, by replacing the L^{∞} -Sobolev condition on the multiplier with a sharper L^2 -Sobolev condition. As a consequence, we obtain the sharp range of L^1 -boundedness for the associated Bochner–Riesz means. The key new ingredient of the proof is a precise pointwise estimate in the transition region for eigenfunctions of one-dimensional Schrödinger operators with doubling single-well potentials.

1. INTRODUCTION

1.1. **Statement of the results.** In this paper we continue the analysis, begun in [DM21], of two-dimensional Grushin operators

$$\mathcal{L} := -\partial_x^2 - V(x)\partial_y^2, \tag{1.1}$$

where $V : \mathbb{R} \to [0, \infty)$ is a "single-well potential" satisfying a scale-invariant regularity condition of order $1 + \theta$. More precisely, we assume that V is continuous, not identically zero, C^1 off the origin, and that, for some $\theta \in (0, 1)$, the estimates

$$V(-x) \simeq V(x) \simeq x V'(x), \tag{1.2a}$$

$$|V'(xe^h) - V'(x)| \lesssim |V'(x)| |h|^{\theta}$$
 (1.2b)

hold for all $x \in \mathbb{R} \setminus \{0\}$ and $h \in [-1, 1]$. Here we use the standard notation $A \leq B$ to denote the estimate $A \leq CB$ for some positive constant C, and $A \simeq B$ to denote the conjunction of $A \leq B$ and $B \leq A$; below we will also write $A \leq_s B$ or $A \simeq_s B$ to indicate that the implicit constants may depend on a parameter s. We refer to the introduction of [DM21] for a discussion of the scope of the assumptions (1.2); here we limit ourselves to pointing out that they are satisfied by power laws $V(x) = |x|^d$ of any degree d > 0 and appropriate perturbations thereof.

In [DM21] we proved a spectral multiplier theorem of Mihlin–Hörmander type for \mathcal{L} , whose smoothness requirement is independent of V and formulated in terms of an L^{∞} -Sobolev norm of order s > 2/2, that is, half the topological dimension of the underlying manifold. The independence from V of the smoothness requirement is particularly striking when compared, e.g., to the classical results based on heat kernel bounds [Heb95, DOS02, RS08], which would give instead a condition of order s > (2 + d/2)/2 in the case $V(x) \simeq |x|^d$. We refer to the introduction of [DM21] for an extensive discussion of the relevance of such result, in the context of a programme (see also [MMN21]) aimed at understanding the optimal smoothness requirement in multiplier theorems for sub-elliptic operators.

 $^{2020\} Mathematics\ Subject\ Classification.\ 34L20,\ 35J70,\ 35H20,\ 42B15.$

Key words and phrases. Grushin operator, spectral multiplier, Schrödinger operator.

The authors are members of the Gruppo Nazionale per l'Analisi Matematica, la Probabilità e le loro Applicazioni (GNAMPA) of the Istituto Nazionale di Alta Matematica (INdAM).

While the result of [DM21] is optimal, in the sense that the smoothness threshold 2/2 cannot be lowered, it is still possible to refine it, by replacing the L^{∞} -Sobolev norm with an L^2 -Sobolev norm. This is the main result of the present paper. We write $L_s^q(\mathbb{R})$ to denote the L^q -Sobolev space of (fractional) order s on \mathbb{R} .

Theorem 1.1. Let \mathcal{L} be the Grushin operator (1.1) on \mathbb{R}^2 , where the coefficient V satisfies the estimates (1.2). Let s > 2/2.

(i) For all $m : \mathbb{R} \to \mathbb{C}$ such that supp $m \subseteq [-1, 1]$,

$$\sup_{t>0} \|\mathbf{m}(t\mathcal{L})\|_{L^1 \to L^1} \lesssim_s \|\mathbf{m}\|_{L^2_s}.$$

(ii) Let $\eta \in C_c^{\infty}((0,\infty))$ be nonzero. For all $m : \mathbb{R} \to \mathbb{C}$ and $p \in (1,\infty)$,

 $\|\mathbf{m}(\mathcal{L})\|_{L^{1}\to L^{1,\infty}} \lesssim_{s} \sup_{t>0} \|\mathbf{m}(t\cdot)\eta\|_{L^{2}_{s}}, \qquad \|\mathbf{m}(\mathcal{L})\|_{L^{p}\to L^{p}} \lesssim_{s,p} \sup_{t>0} \|\mathbf{m}(t\cdot)\eta\|_{L^{2}_{s}}.$

To appreciate the nature of the improvement, one may notice that Theorem 1.1 gives the sharp L^1 -boundedness range for Bochner–Riesz means associated with the Grushin operator \mathcal{L} , a result that cannot be deduced from the multiplier theorem of [DM21].

Corollary 1.2. Under the same assumptions as in Theorem 1.1, the Bochner–Riesz means $(1 - r\mathcal{L})^{\lambda}_{+}$ of order λ associated with \mathcal{L} are bounded on $L^{1}(\mathbb{R}^{2})$ uniformly in $r \geq 0$ whenever $\lambda > 1/2$.

To deduce Corollary 1.2 it is sufficient to apply part (i) of Theorem 1.1 to the function $\mathbf{m} = (1 - \cdot)_{+}^{\lambda} \chi$, where $\chi \in C^{\infty}(\mathbb{R})$ is identically 1 on $[0, \infty)$ and zero on $(-\infty, -1]$, and observe that this m belongs to $L_s^2(\mathbb{R})$ whenever $\lambda > s - 1/2$.

The sharpness of Theorem 1.1 and Corollary 1.2 follows by a standard "transplantation" technique (cf. [Mit74, KST82]; see also [Mar17, Theorem 5.2]). Indeed \mathcal{L} is elliptic (its principal symbol is a positive definite quadratic form) where $x \neq 0$, and therefore the ranges of indices s and λ for which the boundedness results in Theorem 1.1 and Corollary 1.2 hold cannot be larger than the analogous ranges when \mathcal{L} is replaced by the Euclidean Laplacian $-\partial_x^2 - \partial_y^2$ on \mathbb{R}^2 . Theorem 1.1 is already known under more restrictive assumptions on V. Namely,

Theorem 1.1 is already known under more restrictive assumptions on V. Namely, the case $V(x) = x^2$ is in [MS12, MM14] and the case V(x) = |x| is in [CS13]. Moreover, in a previous joint paper [DM20], we established the same result when V is convex, C^3 off the origin, and, for some $d \in (1, 2]$, the estimates

$$|x^{2}V''(x)| + |x^{3}V'''(x)| \lesssim xV'(x) \simeq V(x) = V(-x) \simeq |x|^{d}$$

hold for all $x \in \mathbb{R} \setminus \{0\}$. This appears to have been the first optimal multiplier theorem for a nonelliptic (sub-elliptic) operator enjoying some form of stability under perturbations of the coefficients of the operator. However, the restriction on the power *d* cannot be removed using the methods of [DM20], and the desire to overcome this limitation has been the main motivation for the development of a new proof strategy in [DM21] and in the present paper. Notice that the aforementioned works [RS08, MS12, MM14, CS13, DM20] treat also higher-dimensional cases, and, as a matter of fact, some higher-dimensional cases could be treated by adapting the methods used here too. However, in the same spirit as in [DM21], here we consider only two-dimensional Grushin operators.

1.2. Strategy of the proof. In order to present the main ideas of the paper, it is convenient to recall the notation for the classes of single-well potentials defined in [DM21, Definitions 7.5 and 8.3], which express the assumptions (1.2) in a quantitative form.

 $\mathbf{2}$

Definition 1.3. Let $\kappa \geq 1$ and $\theta \in (0, 1)$. We denote by $\mathcal{P}_1(\kappa)$ the class of nonidentically zero continuous functions $V : \mathbb{R} \to [0, \infty)$ which are C^1 off the origin and such that

$$\kappa^{-1}V(x) \le xV'(x) \le \kappa V(x), \qquad V(-x) \le \kappa V(x)$$

for all $x \neq 0$. We denote by $\mathcal{P}_{1+\theta}(\kappa)$ the class of the $V \in \mathcal{P}_1(\kappa)$ that satisfy the additional inequality

$$|V'(e^h x) - V'(x)| \le \kappa |h|^{\theta}$$

for all $x \neq 0$ and $h \in [-1, 1]$.

As in other works on the subject, Theorem 1.1 will be deduced from an appropriate "weighted Plancherel estimate". In the present case, in light of [DM21, Theorem 4.1], it will be enough to prove that for all $V \in \mathcal{P}_{1+\theta}(\kappa)$, $\gamma \in [0, 1/2)$, r > 0, and all continuous functions $m : \mathbb{R} \to \mathbb{C}$ with supp $m \subseteq [1/4, 1]$,

$$\underset{z' \in \mathbb{R}^2}{\text{ess sup } r^{2-2\gamma} \max\{V(r), V(x')\}^{1/2-\gamma} \int_{\mathbb{R}^2} |y - y'|^{2\gamma} \left| \mathcal{K}_{\mathrm{m}(r^2 \mathcal{L})}(z, z') \right|^2 dz } \\ \lesssim_{\theta, \kappa, \gamma} \|\mathbf{m}\|_{L^2}^2.$$
 (1.3)

Here z := (x, y) and z' := (x', y'), while $\mathcal{K}_{\mathrm{m}(r^2\mathcal{L})}$ denotes the integral kernel of the operator $\mathrm{m}(r^2\mathcal{L})$. Indeed, the estimate (1.3) proves assumption (A) of [DM21, Theorem 4.1] for q = 2, while assumption (B) is already proved in [DM21, Theorem 9.1]. We point out that, in the special case $V(x) = x^2$, the above estimate is proved in [MM14], while the techniques of [MS12, CS13, DM20] lead to a different Plancherel estimate, with a weight depending only on x, x' in place of $|y - y'|^{2\gamma}$ and L^2 in place of L^2_{γ} in the right-hand side.

Our proof of the weighted Plancherel estimate (1.3) largely follows the lines of the analogous estimate proved in [DM21, Theorem 9.1], with the addition of a key new ingredient: universal pointwise estimates for eigenfunctions of one-dimensional Schrödinger operators with potentials in the class $\mathcal{P}_{1+\theta}(\kappa)$. As in [DM21, Section 7], we consider the Schrödinger operator $\mathcal{H}[V] := -\partial_x^2 + V$ on \mathbb{R} with potential $V \in \mathcal{P}_{1+\theta}(\kappa)$, and we denote by $E_n(V)$ and $\psi_n(\cdot; V)$ $(n \geq 1)$ the corresponding eigenvalues and normalised eigenfunctions. The eigenfunction estimates that we need here have the form

$$|\psi_n(x;V)| \lesssim |\{V \le E_n(V)\}|^{-1/2} \min\{n^{\delta/2}, E_n(V)^{\beta/2}|V(x) - E_n(V)|^{-\beta/2}\} \quad (1.4)$$

for some $\delta, \beta \in (0, 1)$, and they have the crucial feature that the implicit constant depends only on κ and θ and not on the specific potential V. The "universality" of an estimate such as (1.4) lies in the fact that the right-hand side is simply expressed in terms of natural quantities such as $V, E_n(V), n$ and universal exponents δ, β , and does not depend, e.g., on the degree of polynomial growth of V.

In the regions where $V \ll E_n(V)$ and $V \gg E_n(V)$, the estimate (1.4) is already contained in estimates proved in [DM21], which actually hold for all $V \in \mathcal{P}_1(\kappa)$. What is crucial for our present purposes is that (1.4) also covers the "transition region" $\{V \simeq E_n(V)\}$, where the eigenfunction $\psi_n(\cdot; V)$ exhibits a change in behaviour from oscillatory to decaying. Various techniques are available to deal with the more general problem of approximating eigenfunctions in the transition region (e.g., Olver's method [Olv74, Chapter 11] and the WKB method, both yielding approximations in terms of the Airy function), but it does not seem possible to use any of them as a black box to prove (1.4) in the required generality. The method used here is in fact substantially different and of a more direct nature, establishing the upper bound (1.4) via a monotonicity argument inspired by what is dubbed the "Sonin's function" method in [Kr08], which in turn refers it back to the work of Szegő on orthogonal polynomials [Sz75, §7.31 and §7.6]. The importance of the pointwise estimate (1.4) is that from it one can deduce a variant of the "spectral projector bound" proved in [DM21, Theorem 8.5], which plays a fundamental role in the proof of the weighted Plancherel estimate (1.3). Specifically, the desired spectral projector bound (Theorem 2.2 below) is obtained by summing instances of the eigenfunction estimate (1.4) corresponding to different values of n and suitably scaled versions τV of the potential V, where the scaling parameter τ depends on n. In order to bound the resulting sum, another important ingredient is an approximated Bohr–Sommerfeld identity with logarithmic error term (Proposition 2.6 below) valid for Schrödinger operators with potentials in the class $\mathcal{P}_1(\kappa)$, which provides precise information on the "gaps" between the quantities $E_n(\tau V)$ involved in the estimate.

1.3. Structure of the paper. In Section 2 we prove the spectral projector bound in a conditional form, namely, by assuming that suitable pointwise eigenfunction estimates of the form (1.4) hold.

Section 3 is devoted to the proof of the required pointwise eigenfunction estimates. As discussed in that section, suitable pointwise estimates can be proved for a larger class than $\mathcal{P}_{1+\theta}(\kappa)$. Indeed, several variants of the above eigenfunction estimates (1.4) are discussed, which may be of independent interest, with different values of δ and β corresponding to different assumptions on the potential V.

Finally, in Section 4, we prove the weighted Plancherel estimate (1.3) with L^2 -Sobolev norm, which, in light of [DM21, Theorem 4.1], implies our main result.

1.4. Notation. $\mathbf{1}_A$ denotes the characteristic function of the set A. We set $\mathbb{R}^+ := (0, \infty)$ and $\mathbb{R}_0^+ := [0, \infty)$. \mathbb{N} denotes the set of natural numbers (including zero), while $\mathbb{N}_+ := \mathbb{N} \setminus \{0\}$ is the set of the positive integers. For an invertible function V, we write V^{\leftarrow} to denote its compositional inverse. # I denotes the number of elements of a finite set I. For a measurable subset $A \subseteq \mathbb{R}$ we denote by |A| its Lebesgue measure. We write \mathcal{K}_T to denote the integral kernel of the operator T.

2. A VARIANT OF THE SPECTRAL PROJECTOR BOUND

2.1. Summary of the results. As before, let $E_n(V)$ and $\psi_n(\cdot; V)$ $(n \ge 1)$ be the eigenvalues and normalised eigenfunctions of the Schrödinger operator $\mathcal{H}[V] := -\partial_x^2 + V$ on \mathbb{R} . We begin by recording an immediate consequence of the "virial theorem" in [DM21, Theorem 7.3]. Under more restrictive assumptions on V, analogous estimates can be found in [DM20, eq. (5.5)].

Proposition 2.1. Let $V \in \mathcal{P}_1(\kappa)$ and $n \in \mathbb{N}_+$. Then the function

$$\mathbb{R}^+ \ni \tau \mapsto E_n(\tau V) \in \mathbb{R}^+$$

is a strictly increasing, real analytic bijection, and

$$E_n(\tau V) \lesssim_{\kappa} \tau \partial_{\tau} E_n(\tau V) \le E_n(\tau V)$$

for all $\tau \in \mathbb{R}^+$. Moreover, if $\Xi_n(\cdot; V) : \mathbb{R}^+ \to \mathbb{R}^+$ denotes its inverse, then

$$\Xi_n(\lambda; V) \simeq_{\kappa} \lambda \partial_{\lambda} \Xi_n(\lambda; V).$$

for all $\lambda \in \mathbb{R}^+$.

The aim of this section is the proof of the following bound, which should be compared to the "spectral projector bound" of [DM21, Theorem 8.5]. In the statement below, by a *conic subset* of $\mathcal{P}_1(\kappa)$ we mean a subset of $\mathcal{P}_1(\kappa)$ closed under multiplication by positive scalars.

Theorem 2.2. Let $\kappa, a > 1$ and $\theta, \delta \in (0, 1)$. Let $\tilde{\mathcal{P}}$ be a conic subset of $\mathcal{P}_1(\kappa)$ such that the eigenfunction estimate

 $|\psi_n(x;V)| \le a |\{V \le E_n(V)\}|^{-1/2} \min\{n^{\delta/2}, E_n(V)^{\theta/2}|V(x) - E_n(V)|^{-\theta/2}\}$ holds for all $V \in \tilde{\mathcal{P}}$, $n \in \mathbb{N}_+$, and $x \in \mathbb{R}$. Then, for all $V \in \tilde{\mathcal{P}}$ and $\lambda, A \in \mathbb{R}^+$,

$$\sum_{\substack{n \in \mathbb{N}_+\\\lambda/\Xi_n(\lambda;V) \in [A,2A]}} \psi_n(x;\Xi_n(\lambda;V)V)^2 \lesssim_{\kappa,a,\theta,\delta} \lambda^{1/2} (\mathbf{1}_{V \le 8A} + e^{-c\lambda^{1/2}|x|} \mathbf{1}_{V > 8A}),$$

where $c = c(\kappa)$.

The main difference between the previous result and [DM21, Theorem 8.5] is that the above sum involves eigenfunctions corresponding to different potentials (that is, potentials τV where τ depends on the summation index n), so cannot be immediately related to properties of the spectral decomposition of a single Schrödinger operator. A similar bound can be found in [DM20, Proposition 5.8], under more restrictive assumptions on V.

The rest of the section is devoted to the proof of Theorem 2.2.

2.2. A summation lemma. The following elementary summation lemma will be a key tool in the proof of the spectral projector bound.

Lemma 2.3. Let $c \in \mathbb{R}^+$, $\kappa \in [1, \infty)$, $\theta, \beta \in [0, 1)$. Let $I \subseteq \mathbb{N}_+$ and, for all $n \in I$, let $t_n \in [\kappa^{-1}, \infty)$ be such that

$$|t_n - cn| \le \kappa n^\beta. \tag{2.1}$$

Then

$$\sup_{\substack{a>0\\0
(2.2)$$

The proof of Lemma 2.3 should be compared to that of [MM14, Lemma 10]. In the case $\beta = 0$, the condition (2.1) implies that the t_n are essentially equispaced, and the estimate (2.2) could be obtained, e.g., by using [DM20, Lemma 5.7] to estimate a sum with the corresponding integral. The point of this lemma is to show that a similar estimate can be obtained even when $\beta > 0$, that is, under a weaker assumption on the gaps between the t_n , by taking advantage of the stronger uniform bound $a^{-\beta}$ in the left-hand side of (2.2).

Proof of Lemma 2.3. Note that $t_n, n \gtrsim_{\kappa} 1$ for all $n \in I$. Hence, from the assumption (2.1) and the fact that $\beta < 1$, we deduce that $t_n \simeq_{\kappa,c,\beta} n$ for all $n \in I$.

For a given a > 0, from the condition $t_n \leq \kappa a$ and $t_n \simeq_{\kappa,c,\beta} n$, we deduce that

$$n \lesssim_{\kappa,c,\beta} a$$
 (2.3)

as well. Therefore, if $I_a := \{n \in I : t_n \leq \kappa a\}$, then (2.1) implies that

$$|t_n - cn| \le E$$

for all $n \in I_a$, where

$$E = E(\kappa, c, \beta, a) \lesssim_{\kappa, c, \beta} a^{\beta}.$$
(2.4)

We now split I_a into the three subsets

$$\begin{split} I_{-} &:= \{ n \in I_{a} \, : \, cn + E < b - c \}, \\ I_{+} &:= \{ n \in I_{a} \, : \, cn - E > b + c \}, \\ I_{0} &:= I_{a} \setminus (I_{+} \cup I_{-}). \end{split}$$

Then, for all $n \in I_-$,

$$t_n \le cn + E < b - c,$$

and therefore

$$|t_n - b|^{-\theta} \le \inf_{t \in [cn + E, cn + E + c]} |t - b|^{-\theta} \le \frac{1}{c} \int_{cn + E}^{cn + E + c} |t - b|^{-\theta} dt$$

(here we use that $t \mapsto |t-b|^{-\theta}$ is increasing for t < b) and

$$\sum_{n \in I_{-}} |t_n - b|^{-\theta} \le \frac{1}{c} \int_0^0 |t - b|^{-\theta} dt \lesssim_{c,\theta} b^{1-\theta} \lesssim_{\kappa,\theta} a^{1-\theta},$$
(2.5)

since $\theta < 1$. In a similar way, one proves that

$$\sum_{n \in I_+} |t_n - b|^{-\theta} \le \frac{1}{c} \int_b^{\kappa a} |t - b|^{-\theta} dt \lesssim_{\kappa, c, \theta} a^{1-\theta}.$$
 (2.6)

Finally, if $n \in I_0$, then

$$|b - cn| \le E + c \lesssim_{\kappa, c, \beta} a^{\beta}$$

(here we used (2.4) and the fact that, by (2.3), $a^{\beta} \gtrsim_{\kappa,c,\beta} n^{\beta} \geq 1 \gtrsim_c c$), which implies that

$$\# I_0 \lesssim_{\kappa,c,\beta} a^{\beta}. \tag{2.7}$$

The estimate (2.2) follows by combining (2.5), (2.6) and (2.7).

2.3. A consequence of Lagrange's Mean Value Theorem. Let $\kappa \geq 1$. Recall from [DM21, Definition 6.1] the class $\mathcal{HP}_1(\kappa)$ of the C^1 functions $W : \mathbb{R}^+ \to \mathbb{R}^+$ such that

$$\kappa^{-1}W(x) \le xW'(x) \le \kappa W(x) \tag{2.8}$$

for all $x \in \mathbb{R}^+$. In other words, an element of $\mathcal{HP}_1(\kappa)$ is "half of a potential" in the class $\mathcal{P}_1(\kappa)$. Indeed, if $V \in \mathcal{P}_1(\kappa)$, then $V_{\oplus}, V_{\ominus} \in \mathcal{HP}_1(\kappa)$, where $V_{\oplus}, V_{\ominus} : \mathbb{R}^+ \to \mathbb{R}^+$ are defined by

$$V_{\oplus}(x) := V(x), \qquad V_{\ominus}(x) := V(-x)$$
 (2.9)

for all $x \in \mathbb{R}^+$.

We record here some useful properties of functions in the class $\mathcal{HP}_1(\kappa)$, including an elementary consequence of Lagrange's Mean Value Theorem, which will be used multiple times later.

Lemma 2.4. Let $W \in \mathcal{HP}_1(\kappa)$.

(i) For all $x \in \mathbb{R}^+$ and $\lambda \ge 1$,

$$\lambda^{1/\kappa} W(x) \le W(\lambda x) \le \lambda^{\kappa} W(x)$$

(ii) W is strictly increasing and invertible, and $W^{\leftarrow} \in \mathcal{HP}_1(\kappa)$ too. (iii) For all $x, y \in \mathbb{R}^+$, if $x \ge y$ then

$$W(x) - W(y) \simeq_{\kappa} \frac{W(x)}{x}(x-y).$$

Proof. Parts (i) and (ii) are proved in [DM21, Propositions 6.4 and 6.5].

As for part (iii), if $x \ge 2y$, then $W(x) \ge 2^{1/\kappa}W(y)$ by part (i), whence

$$x - y \simeq x$$
, $W(x) - W(y) \simeq_{\kappa} W(x)$

and the desired estimate follows. If instead $x \leq 2y$, then, by Lagrange's Mean Value Theorem,

$$W(x) - W(y) = W'(\xi)(x - y)$$

for some $\xi \in (y, x)$, and moreover

$$W'(\xi) \simeq_{\kappa} \frac{W(\xi)}{\xi} \simeq_{\kappa} \frac{W(x)}{x}$$

by (2.8) and part (i), as $x \simeq y \simeq \xi$ in this case, whence the desired estimate again follows.

2.4. Bohr–Sommerfeld approximation with logarithmic error. Let us recall from [DM21, Theorem 7.6 and Proposition 7.11] some useful estimates involving eigenvalues and sublevel sets of the potential of one-dimensional Schrödinger operators.

Proposition 2.5. Let $V \in \mathcal{P}_1(\kappa)$. Then

$$E_n(V)^{1/2} |\{V \le E_n(V)\}| \simeq_{\kappa} n.$$
 (2.10)

for all $n \in \mathbb{N}_+$. Moreover, for all $E, \lambda \in \mathbb{R}^+$,

$$|\{V \le \lambda E\}| \simeq_{\kappa,\lambda} |\{V \le E\}| \simeq_{\kappa} |\{V_{\oplus} \le E\}| \simeq_{\kappa} |\{V_{\ominus} \le E\}|.$$

In what follows we will need a sharper version of the estimate (2.10).

Proposition 2.6. Let $V \in \mathcal{P}_1(\kappa)$. Then, for all $n \in \mathbb{N}_+$,

$$\left| \int_{\mathbb{R}} (E_n(V) - V)_+^{1/2} - \pi n \right| \lesssim_{\kappa} \log(1+n)$$

The proof of the above estimate follows the lines of [Tit62, $\S7.4$]. In the case V is convex, the logarithmic divergence in the right-hand side can be replaced by a constant, as shown in [Tit62, \$7.5] and [DM20, Theorem 4.2]; however the weaker logarithmic bound does not require convexity and will be enough for our purposes.

Proof of Proposition 2.6. Let $x_n^{\pm} \in \mathbb{R}^+$ be such that $V(\pm x_n^{\pm}) = E_n := E_n(V)$; in other words, the points $\pm x_n^{\pm}$ are the transition points corresponding to the energy level E_n . Let $y_n^{\pm} \in (0, x_n^{\pm})$ be points to be fixed later, and define $Q_n(x) := (E_n - V(x))^{1/2}$ for $x \in (-x_n^-, x_n^+)$.

By classical Sturm-Liouville theory, $\psi_n := \psi_n(\cdot; V)$ has n-1 zeros, which are all in the interval $(-x_n^-, x_n^+)$. Note now that $V - E_n \ge V(y_n^\pm) - E_n$ on $\pm [y_n^\pm, \infty)$, with strict inequality away from $\pm y_n^\pm$. Hence, by Sturm's comparison theorem (see, e.g., [BS91, Chapter 2, Theorem 3.2]), if u is any nontrivial solution of $-u'' + (V(y_n^\pm) - E_n)u = 0$ on an interval contained in $\pm [y_n^\pm, \infty)$, then we can find a zero of u between any two zeros of ψ_n . This implies in particular that on $\pm [y_n^\pm, x_n^\pm)$ there are at most

$$1 + (x_n^{\pm} - y_n^{\pm})Q_n(\pm y_n^{\pm})/\pi$$

zeros of ψ_n . Note also that

$$\int_{\pm [y_n^{\pm}, x_n^{\pm}]} (E_n - V)^{1/2} \le (x_n^{\pm} - y_n^{\pm})Q_n(\pm y_n^{\pm}).$$

Hence, if Z_n^{\pm} denotes the number of zeros of ψ_n in $\pm (0, y_n^{\pm})$, then

$$\left| \int_{\mathbb{R}} (E_n - V)_{\pm}^{1/2} - \pi n \right| \\ \leq 3\pi + 2 \sum_{\pm} (x_n^{\pm} - y_n^{\pm}) Q_n(\pm y_n^{\pm}) + \sum_{\pm} \left| \int_{\pm (0, y_n^{\pm})} (E_n - V)^{1/2} - \pi Z_n^{\pm} \right|.$$

On the other hand, by [Tit62, §7.3, Lemma] (see also [DM20, Appendix]),

$$\left| \int_{\pm(0,y_n^{\pm})} (E_n - V)^{1/2} - \pi Z_n^{\pm} \right| \le \pi + \frac{1}{2} \int_{\pm(0,y_n^{\pm})} \frac{|Q'_n|}{Q_n}.$$

Since Q'_n is increasing on $(-y_n^{\pm}, 0)$ and decreasing on $(0, y_n^{\pm})$,

$$\int_{\pm(0,y_n^{\pm})} \frac{|Q'_n|}{Q_n} = \log \frac{Q_n(0)}{Q_n(\pm y_n^{\pm})} = \frac{1}{2} \log \frac{E_n}{E_n - V(\pm y_n^{\pm})},$$

and therefore

$$\left| \int_{\mathbb{R}} (E_n - V)_+^{1/2} - \pi n \right| \le 5\pi + \sum_{\pm} \left[2(x_n^{\pm} - y_n^{\pm})Q_n(\pm y_n^{\pm}) + \frac{1}{4}\log\frac{E_n}{E_n - V(\pm y_n^{\pm})} \right].$$

We now choose $y_n^{\pm} := x_n^{\pm} - c(x_n^{\pm}/E_n)^{1/3}$ for a suitable c > 0. Note that

$$\left(\frac{x_n^{\pm}}{E_n}\right)^{1/3} = \frac{x_n^{\pm}}{\left(x_n^{\pm} E_n^{1/2}\right)^{2/3}} \simeq_{\kappa} \frac{x_n^{\pm}}{n^{2/3}},$$

by Proposition 2.5; so, by choosing $c = c(\kappa)$ sufficiently small, we can ensure that

$$y_n^{\pm} \simeq_{\kappa} x_n^{\pm}, \qquad x_n^{\pm} - y_n^{\pm} \simeq_{\kappa} \left(\frac{x_n^{\pm}}{E_n}\right)^{1/3}.$$

Hence, by Lemma 2.4, we deduce that

$$Q_n(\pm y_n^{\pm})^2 = E_n - V(\pm y_n^{\pm})$$

$$\simeq_{\kappa} \frac{V(\pm x_n^{\pm})}{x_n^{\pm}} (x_n^{\pm} - y_n^{\pm}) \simeq_{\kappa} \frac{E_n}{x_n^{\pm}} \left(\frac{x_n^{\pm}}{E_n}\right)^{1/3} = \left(\frac{E_n}{x_n^{\pm}}\right)^{2/3}$$

and therefore

$$(x_n^{\pm} - y_n^{\pm})Q_n(\pm y_n^{\pm}) \simeq_{\kappa} 1, \qquad \frac{E_n}{E_n - V(\pm y_n^{\pm})} \simeq_{\kappa} n^{2/3},$$

which proves the desired estimate.

2.5. A useful change of variables. The lemma below will be used to extract and exploit the "rough gap information" from Proposition 2.6.

Lemma 2.7. For $V \in \mathcal{P}_1(\kappa)$, define $K_V : \mathbb{R}^+ \to \mathbb{R}^+$ by

$$K_V(t) := t^{-1/2} \int_{\mathbb{R}} (t - V)_+^{1/2}$$

for all t > 0. Then

$$K_V(t) \simeq_{\kappa} t K'_V(t) \simeq_{\kappa} |\{V \le t\}|.$$

Proof. Note first that, if V_{\oplus}, V_{\ominus} are defined as in (2.9), then

$$|\{V \leq t\}| = |\{V_{\oplus} \leq t\}| + |\{V_{\ominus} \leq t\}| = V_{\oplus}^{\leftarrow}(t) + V_{\ominus}^{\leftarrow}(t) + V_{\Box}^{\leftarrow}(t) + V_{\Box}$$

and moreover

$$K_V = K_{V_{\oplus}} + K_{V_{\ominus}},$$

where, for $W : \mathbb{R}^+ \to \mathbb{R}^+$, we define

$$K_W(t) := t^{-1/2} \int_0^\infty (t - W)_+^{1/2}$$

It is then enough to prove that, for all $W \in \mathcal{HP}_1(\kappa)$,

$$K_W(t) \simeq_{\kappa} t K'_W(t) \simeq_{\kappa} W^{\leftarrow}(t).$$

Now, for all t > 0,

$$K_W(t) = t^{-1/2} \int_0^{W^{\leftarrow}(t)} (t - W)^{1/2} \simeq_{\kappa} t^{-1/2} \int_0^{W^{\leftarrow}(t)} \frac{xW'(x)}{W(x)} (t - W(x))^{1/2} dx,$$

and the change of variables $\tau = W(x)/t$ yields

$$K_W(t) \simeq_{\kappa} \int_0^1 W^{\leftarrow}(\tau t) \, (1-\tau)^{1/2} \frac{d\tau}{\tau} \simeq_{\kappa} W^{\leftarrow}(t);$$

the last equivalence is consequence of the fact (see Lemma 2.4) that $\tau^{\kappa}W^{\leftarrow}(t) \leq W^{\leftarrow}(\tau t) \leq \tau^{1/\kappa}W^{\leftarrow}(t)$ for $\tau \in (0, 1)$.

Similarly, one readily sees that

$$2tK'_W(t) = t^{-1/2} \int_0^{W^{\leftarrow}(t)} \frac{W}{(t-W)^{1/2}} \simeq_{\kappa} t^{-1/2} \int_0^{W^{\leftarrow}(t)} \frac{xW'(x)}{(t-W(x))^{1/2}} dx$$

and again the change of variables $\tau = W(x)/t$ yields

$$tK'_W(t) \simeq_{\kappa} \int_0^1 W^{\leftarrow}(\tau t) \, (1-\tau)^{-1/2} \, d\tau \simeq_{\kappa} W^{\leftarrow}(t),$$

as desired.

2.6. **Proof of the variant of the spectral projector bound.** Here we prove Theorem 2.2, that is, the estimate

$$\sum_{\substack{n \in \mathbb{N}_+\\\lambda/\Xi_n(\lambda;V) \in [A,2A]}} \psi_n(x;\Xi_n(\lambda;V)V)^2 \lesssim_{\kappa,a,\theta,\delta} \lambda^{1/2} (\mathbf{1}_{V \le 8A} + e^{-c\lambda^{1/2}|x|} \mathbf{1}_{V > 8A})$$

for all $V \in \tilde{\mathcal{P}}, \lambda, A \in \mathbb{R}^+, x \in \mathbb{R}$.

Recall from [DM21, Theorem 7.7] that, for all $V \in \mathcal{P}_1(\kappa)$, there exists $c = c(\kappa)$ such that

$$|\psi_n(x;V)| \lesssim_{\kappa} |\{V \le E_n(V)\}|^{-1/2} \exp(-c|x|\sqrt{V(x)})$$
 (2.11)

whenever $n \in \mathbb{N}_+$ and $x \in \{V \ge 4E_n\}$. Recall moreover that, by assumption, $\tilde{\mathcal{P}}$ is a conic subset of $\mathcal{P}_1(\kappa)$ such that, for some $\theta, \delta \in (0, 1)$ and a > 1,

$$|\psi_n(x;V)| \le a |\{V \le E_n(V)\}|^{-1/2} \min\{n^{\delta/2}, E_n(V)^{\theta/2}|V(x) - E_n(V)|^{-\theta/2}\}$$
(2.12)

for all $V \in \mathcal{P}$, $x \in \mathbb{R}$, $n \in \mathbb{N}_+$. For simplicity, in the rest of the proof, we will write \lesssim and \simeq instead of $\lesssim_{\kappa,a,\theta,\delta}$ and $\simeq_{\kappa,a,\theta,\delta}$.

Fix $V \in \tilde{\mathcal{P}}$ and let $\Xi_n := \Xi_n(\cdot; V)$. First note that, if V(x) > 8A and $\lambda/\Xi_n(\lambda) \in [A, 2A]$, then $\Xi_n(\lambda)V(x) > 4\lambda$, and therefore by (2.11) we deduce that

$$\psi_n(x;\Xi_n(\lambda)V)^2 \lesssim |\{V \le \lambda/\Xi_n(\lambda)\}|^{-1} \exp(-2c|x|\sqrt{\Xi_n(\lambda)V(x)})$$
$$\le |\{V \le A\}|^{-1} \exp(-4c\lambda^{1/2}|x|).$$

On the other hand, by Proposition 2.5,

$$\lambda^{1/2}|\{V \le A\}| \simeq \lambda^{1/2}|\{V \le \lambda/\Xi_n(\lambda)\}| \simeq n \tag{2.13}$$

so the number of summands is $\lesssim \lambda^{1/2} | \{ V \leq A \} |$, and we deduce that

$$\sum_{\substack{n \in \mathbb{N}_+\\\lambda/\Xi_n(\lambda;V) \in [A,2A]}} \psi_n(x;\Xi_n(\lambda)V)^2 \lesssim \lambda^{1/2} \exp(-4c\lambda^{1/2}|x|)$$

whenever V(x) > 8A.

It remains to prove the uniform bound on $\{V \leq 8A\}$. For this, we use (2.12) to obtain that

$$\psi_n(x;\Xi_n(\lambda)V)^2 \lesssim |\{V \le \lambda/\Xi_n(\lambda)\}|^{-1} \min\{n^{\delta}, \lambda^{\theta}|\Xi_n(\lambda)V(x) - \lambda|^{-\theta}\} \simeq |\{V \le A\}|^{-1} \min\{n^{\delta}, A^{\theta}|V(x) - \lambda/\Xi_n(\lambda)|^{-\theta}\}.$$
(2.14)

Define K_V as in Lemma 2.7, and recall that

$$K_V(t) \simeq t K'_V(t) \simeq |\{V \le t\}|.$$
 (2.15)

Since $\lambda/\Xi_n(\lambda) \simeq A \gtrsim V(x)$, we deduce, by Lemma 2.4, that

$$\begin{aligned} |V(x) - \lambda/\Xi_n(\lambda)| &\simeq \frac{A}{K_V(A)} |K_V(V(x)) - K_V(\lambda/\Xi_n(\lambda))| \\ &= \frac{A}{\lambda^{1/2} K_V(A)} |\lambda^{1/2} K_V(V(x)) - \lambda^{1/2} K_V(\lambda/\Xi_n(\lambda))|. \end{aligned}$$

Set now $a := \lambda^{1/2} K_V(A)$, $b := \lambda^{1/2} K_V(V(x))$, $t_n := \lambda^{1/2} K_V(\lambda/\Xi_n(\lambda))$, and observe that $t_n \simeq a \simeq n \gtrsim b$ by (2.13) and (2.15), so the bound (2.14) can be rewritten as

$$\psi_n(x;\Xi_n(\lambda)V)^2 \lesssim \lambda^{1/2} \min\{a^{\delta-1}, a^{\theta-1}|b-t_n|^{-\theta}\}.$$

Furthermore,

$$t_n = \lambda^{1/2} K_V(\lambda/\Xi_n(\lambda)) = \int_{\mathbb{R}} (\lambda - \Xi_n(\lambda)V)_+^{1/2},$$

and therefore

$$|t_n - \pi n| \lesssim \log(1+n) \lesssim n^{1-\delta}$$

by Proposition 2.6 (applied to the potential $\Xi_n(\lambda)V$) and the fact that $\delta < 1$. As a consequence, we can apply Lemma 2.3 and obtain that

$$\sum_{\substack{n\in\mathbb{N}_+\\\lambda/\Xi_n(\lambda;V)\in[A,2A]}}\psi_n(x;\Xi_n(\lambda)V)^2\lesssim\lambda^{1/2}\sum_{\substack{n\in\mathbb{N}_+\\t_n\simeq a}}\min\{a^{\delta-1},a^{\theta-1}|b-t_n|^{-\theta}\}\lesssim\lambda^{1/2},$$

as desired.

3. Pointwise eigenfunction estimates in the transition region

3.1. Summary of the results. Let $\kappa \geq 1$. Let us introduce the following subclasses of $\mathcal{P}_1(\kappa)$. Recall that a *modulus of continuity* is a function $\omega : [0, \infty] \to [0, \infty]$ such that $\lim_{t\to 0} \omega(t) = 0$.

Definition 3.1. If ω is a modulus of continuity, let $\mathcal{P}_{1,\mathrm{uc}}(\kappa,\omega)$ be the class of potentials $V \in \mathcal{P}_1(\kappa)$ such that

$$\left|\log(V'(\pm e^t)/V'(\pm e^{t'}))\right| \le \omega(|t - t'|) \quad \text{for all } t, t' \in \mathbb{R}.$$

In other words, ω is a modulus of continuity for the functions $t \mapsto \log |V'(\pm e^t)|$.

Remark 3.2. It is easy to see that, for all $\theta \in (0,1)$, $\mathcal{P}_{1+\theta}(\kappa) \subseteq \mathcal{P}_{1,\mathrm{uc}}(\kappa,\omega_{\kappa,\theta})$, where $\omega_{\kappa,\theta}$ is a suitable modulus of continuity such that $\omega_{\kappa,\theta}(t) \simeq_{\kappa,\theta} t^{\theta}$ for t small.

Definition 3.3. Let $\mathcal{P}_{1,cv}(\kappa)$ be the class of the convex potentials in $\mathcal{P}_1(\kappa)$.

Definition 3.4. For $k \geq 2$, let $\mathcal{P}_k(\kappa)$ be the class of the potentials $V \in \mathcal{P}_1(\kappa)$ which are C^k on $\mathbb{R} \setminus \{0\}$ and satisfy the estimates

$$|x^{\ell}V^{(\ell)}(x)| \le \kappa V(x)$$
 for all $x \ne 0$ and $\ell = 2, \dots, k$.

The aim of this section is to prove the following pointwise estimates for the eigenfunctions of $\mathcal{H}[V] := -\partial_x^2 + V$.

Theorem 3.5. For all $x \in \mathbb{R}$ and $n \in \mathbb{N}_+$, the estimates

$$\begin{aligned} |\psi_n(x;V)| \lesssim_{\bar{\kappa},\alpha} \frac{1}{|\{V \le E_n(V)\}|^{1/2}} \min\{n^{2\alpha/3}, |1-V(x)/E_n(V)|^{-\alpha}\}, \\ |\psi'_n(x;V)| \lesssim_{\bar{\kappa},\alpha} \frac{E_n(V)^{1/2}}{|\{V \le E_n(V)\}|^{1/2}} \max\{n^{(2\alpha-1)/3}, (1-V(x)/E_n(V))_+^{1/2-\alpha}\}. \end{aligned}$$

hold in the following cases:

- (i) with $\bar{\kappa} = \kappa$ and $\alpha = 1/2$, whenever $V \in \mathcal{P}_1(\kappa)$;
- (ii) with $\bar{\kappa} = (\kappa, \omega)$ and $\alpha \in (1/4, 1/2)$, whenever $V \in \mathcal{P}_{1,uc}(\kappa, \omega)$;
- (iii) with $\bar{\kappa} = \kappa$ and $\alpha = 1/4$, whenever $V \in \mathcal{P}_{1,cv}(\kappa)$;
- (iv) with $\bar{\kappa} = \kappa$ and $\alpha = 1/4$, whenever $V \in \mathcal{P}_3(\kappa)$.

10

We point out that, well inside the classical region (say, where $V(x) \leq E_n(V)/2$), the above bounds reduce to the uniform bound stated, e.g., in [DM21, Proposition 6.2], which applies to any $V \in \mathcal{P}_1(\kappa)$; similarly, far from the classical region (say, where $V(x) \geq 4E_n(V)$), a much better (exponentially decaying) bound is known to hold, again for arbitrary $V \in \mathcal{P}_1(\kappa)$ (see, e.g., [DM21, Theorem 7.7]). As anticipated in the introduction, the relevance of the above bounds is therefore their validity in the transition region, where $V(x) \simeq E_n(V)$.

We also point out that the bound for ψ_n in Theorem 3.5(iv) matches the one obtained in [DM20, Proposition 3.4] under the additional assumption $V(x) \simeq |x|^d$ for some d > 1. The method used in [DM20] is based on a theorem by Olver [Olv74], which essentially allows one to approximate ψ_n with a suitably rescaled Airy function, so the bounds for ψ_n can be reduced to known bounds for the Airy function. The method presented here, instead, does not go through such an approximation, but yields the desired bounds directly. Moreover it allows us to treat potentials $V(x) \simeq |x|^d$ for arbitrary d > 0, or even potentials that are not comparable to a single power law (provided they belong to one of the classes of potentials defined above); see also the discussion in the introduction of [DM21].

Here and in the following sections, we shall write E_n and ψ_n in place of $E_n(V)$ and $\psi_n(\cdot; V)$ when the potential V is clear from the context.

3.2. Pointwise estimate in the classical region: C^1 potentials. Here we assume that $V \in \mathcal{P}_1(\kappa)$ and prove the pointwise estimate

$$\psi_n^2 + \frac{(\psi_n')^2}{E_n - V} \lesssim_\kappa \frac{E_n (E_n - V)^{-1}}{|\{V \le E_n\}|}$$
(3.1)

in the classical region $\{V < E_n\}$.

The key to the proof is the monotonicity information provided by the following elementary identity, valid on $\mathbb{R} \setminus \{0\}$:

$$\left((E_n - V)\psi_n^2 + (\psi_n')^2 \right)' = -V'\psi_n^2 \tag{3.2}$$

(cf. [DM21, Proposition 5.7]). As the right-hand side is positive on $(-\infty, 0)$ and negative on $(0, \infty)$, we conclude that

$$(E_n - V)\psi_n^2 + (\psi_n')^2 \le E_n \psi_n(0)^2 + \psi_n'(0)^2$$
(3.3)

on the whole \mathbb{R} .

To bound the right-hand side of the latter, we use another monotonicity argument, based on the following counterpart to (3.2):

$$\left(\psi_n^2 + \frac{(\psi_n')^2}{E_n - V}\right)' = \frac{V'}{(E_n - V)^2} (\psi_n')^2 \tag{3.4}$$

on the region $\{V \neq E_n\} \setminus \{0\}$ (cf. [Tit62, §8.3]). As the right-hand side is negative on $(-\infty, 0)$ and positive on $(0, \infty)$, we can control (cf. [DM21, Section 6.4]) the value of $\psi_n^2 + \frac{(\psi_n)^2}{E_n - V}$ at 0 with its average on $\{V \leq E_n/2\}$, thus obtaining that

$$\psi_n(0)^2 + \frac{\psi_n'(0)^2}{E_n} \le \frac{1}{|\{V \le E_n/2\}|} \left(\|\psi_n\|_2^2 + \frac{2}{E_n} \|\psi_n'\|_2^2 \right) \lesssim_{\kappa} \frac{1}{|\{V \le E_n\}|}$$
(3.5)

by Proposition 2.5.

The desired estimate (3.1) then follows by combining (3.3) and (3.5).

We record here an elementary consequence of (3.1), that is, a uniform estimate which is valid well within the classical region: namely, for any $\theta \in (0, 1)$,

$$\psi_n^2 + \frac{(\psi_n')^2}{E_n - V} \lesssim_{\kappa,\theta} \frac{1}{|\{V \le E_n\}|} \quad \text{on } \{V \le (1 - \theta)E_n\}.$$
 (3.6)

3.3. Pointwise estimate in the classical region: C^3 potentials. In this section we assume that $V \in \mathcal{P}_3(\kappa)$, and prove the following "improvement" of (3.1):

$$\psi_n^2 + \left(1 + \frac{n^{-2/3}E_n}{E_n - V}\right)^{-3} \frac{(\psi_n')^2}{E_n - V} \lesssim_\kappa \frac{E_n^{1/2}(E_n - V)^{-1/2}}{|\{V \le E_n\}|}$$
(3.7)

in the classical region $\{V < E_n\}$.

As in Section 3.2 above, the proof is based on a monotonicity argument. Specifically, the method used here is inspired by what is referred to as the "Sonin's function" method in [Kr08]. The main idea is to consider the function

$$f_n := (E_n - V)^{1/4} \psi_n,$$

which is well defined and C^2 in the punctured classical region $\{V < E_n\} \setminus \{0\}$. From the differential equation

$$-\psi_n'' + V\psi_n = E_n\psi_n \tag{3.8}$$

satisfied by ψ_n , one readily obtains that

$$f_n'' - 2A_n f_n' + B_n f_n = 0 ag{3.9}$$

in the punctured classical region, where

$$A_n := -\frac{1}{4} \frac{V'}{E_n - V}, \qquad B_n := E_n - V + \frac{5}{16} \frac{(V')^2}{(E_n - V)^2} + \frac{1}{4} \frac{V''}{E_n - V}$$

We now consider the "Sonin's function" for f_n , namely,

$$S_n := f_n^2 + \frac{(f_n')^2}{B_n}$$

which is defined and C^1 in the subset $\{V < E_n, B_n \neq 0\}$ of the punctured classical region. A quick computation shows that

$$S'_{n} = \frac{4A_{n}B_{n} - B'_{n}}{B_{n}^{2}}(f'_{n})^{2}, \qquad (3.10)$$

that is, the derivative of S_n has the same sign as

$$4A_nB_n - B'_n = -\frac{15}{16}\frac{(V')^3}{(E_n - V)^3} - \frac{9}{8}\frac{V'V''}{(E_n - V)^2} - \frac{1}{4}\frac{V'''}{E_n - V}$$

To study the sign of B_n and $4A_nB_n - B'_n$, we rewrite them as

$$B_n = E_n - V + \frac{5}{16} \frac{(V')^2}{(E_n - V)^2} \left[1 + \frac{4}{5} \frac{V''}{(V')^2} (E_n - V) \right],$$

$$4A_n B_n - B'_n = -\frac{15}{16} \frac{(V')^3}{(E_n - V)^3} \left[1 + \frac{6}{5} \frac{V''}{(V')^2} (E_n - V) + \frac{4}{15} \frac{V'''}{(V')^3} (E_n - V)^2 \right].$$

This is convenient because, for $V \in \mathcal{P}_3(\kappa)$,

$$|V''/(V')^2| \lesssim_{\kappa} 1/V, \qquad |V'''/(V')^3| \lesssim_{\kappa} 1/V^2.$$

Hence, we can choose $\epsilon=\epsilon(\kappa)>0$ sufficiently small to guarantee that

$$B_n \simeq E_n - V + \frac{(V')^2}{(E_n - V)^2}, \qquad 4A_n B_n - B'_n \simeq -\frac{(V')^3}{(E_n - V)^3}$$
 (3.11)

in the region $\Omega_n := \{(1 - \epsilon)E_n \leq V < E_n\}$. In particular, $B_n > 0$ there, and S'_n has the same sign as -V'.

Define now $y_n^{\pm} > 0$ as the points such that $V(\pm y_n^{\pm}) = (1 - \epsilon)E_n$. Then, using the fact that S_n is decreasing on $\Omega_n \cap (0, \infty)$ and increasing on $\Omega_n \cap (-\infty, 0)$, we conclude that

$$f_n^2 + \frac{(f_n')^2}{B_n} \le \max_{\pm} \left(f_n(\pm y_n^{\pm})^2 + \frac{f_n'(\pm y_{\pm})^2}{B_n(\pm y_n^{\pm})} \right) \quad \text{on } \Omega_n.$$
(3.12)

Note now that

$$f_n = (E_n - V)^{1/4} \psi_n, \qquad f'_n = (E_n - V)^{1/4} \left[\psi'_n - \frac{1}{4} \frac{V'}{E_n - V} \psi_n \right]$$

In particular, by (3.11),

$$f_n^2 + \frac{(f_n')^2}{B_n} \simeq (E_n - V)^{1/2} \left[\psi_n^2 + \frac{(\psi_n')^2}{B_n} \right]$$
(3.13)

in the region Ω_n , and

$$f_n(\pm y_n^{\pm})^2 + \frac{f'_n(y_n^{\pm})^2}{B_n(y_n^{\pm})} \lesssim (\epsilon E_n)^{1/2} \left[\psi_n(\pm y_n^{\pm})^2 + \frac{\psi'_n(\pm y_n^{\pm})^2}{\epsilon E_n} \right] \lesssim_{\kappa} \frac{E_n^{1/2}}{|\{V \le E_n\}|};$$

the last inequality is consequence of the fact that $\pm y_n^{\pm} \in \{V \leq (1-\epsilon)E_n\}$, that is, $\pm y_n^{\pm}$ are well inside the classical region, so the uniform estimate (3.6) applies. From (3.12) and (3.13) we then deduce that

$$\psi_n^2 + \frac{(\psi_n')^2}{B_n} \lesssim_{\kappa} \frac{E_n^{1/2} (E_n - V)^{-1/2}}{|\{V \le E_n\}|} \quad \text{on } \Omega_n.$$
 (3.14)

As $|V'| \simeq_{\kappa} |\{V \leq E_n\}|^{-1} E_n \simeq_{\kappa} n^{-1} E_n^{3/2}$ on Ω_n by Proposition 2.5, from (3.11) we deduce that

$$B_n \simeq E_n - V + \frac{(V')^2}{(E_n - V)^2} \simeq_{\kappa} (E_n - V) \left(1 + \frac{n^{-2/3}E_n}{E_n - V}\right)^3,$$

so from (3.14) the desired estimate (3.7) follows on Ω_n . Actually the same estimate (3.7) holds on the whole classical region $\{V < E_n\}$, because on $\{V \le (1 - \epsilon)E_n\}$ it simply reduces to the uniform estimate (3.6).

3.4. Pointwise estimate in the classical region: convex potentials and potentials with uniformly continuous derivative. A variation of the method presented in the previous section allows us to show that, if $V \in \mathcal{P}_{1,uc}(\kappa, \omega)$, then the following pointwise estimate holds for all $\alpha \in (1/4, 1/2)$:

$$\psi_n^2 + \left(1 + \frac{n^{-2/3} E_n}{E_n - V}\right)^{-3} \frac{(\psi_n')^2}{E_n - V} \lesssim_{\bar{\kappa}, \alpha} \frac{E_n^{2\alpha} (E_n - V)^{-2\alpha}}{|\{V \le E_n\}|}$$
(3.15)

in the classical region $\{V < E_n\}$, where $\bar{\kappa} = (\kappa, \omega)$; in addition, we will prove the analogous estimate for $\alpha = 1/4$ and $\bar{\kappa} = \kappa$ in the case $V \in \mathcal{P}_{1,cv}(\kappa)$.

As in Section 3.3, we apply the Sonin's function method. We only discuss the estimates for $x \ge 0$, as the case $x \le 0$ can be be treated analogously. Let $x_n \in \mathbb{R}^+$ denote the positive transition point, i.e., $V(x_n) = E_n$.

Let $\alpha \in \mathbb{R}^+$, and consider the function

$$f_n(x) := (x_n - x)^{\alpha} \psi_n(x),$$

which is well defined and C^2 in the classical region. From the differential equation (3.8) satisfied by ψ_n , one readily obtains that f_n satisfies (3.9) in the classical region, where

$$A_n := -\alpha (x_n - x)^{-1}, \qquad B_n := E_n - V + \alpha (\alpha + 1)(x_n - x)^{-2}.$$
(3.16)

Since $\alpha > 0$, clearly $B_n > 0$ in the classical region. We now consider the Sonin's function for f_n , namely,

$$S_n := f_n^2 + \frac{(f_n')^2}{B_n},$$

which is defined and C^1 in the punctured classical region. By arguing as in (3.10), we deduce that the derivative of S_n has the same sign as

$$4A_nB_n - B'_n = V' - 4\alpha \frac{E_n - V}{x_n - x} - 2\alpha(\alpha + 1)(2\alpha + 1)(x_n - x)^{-3}.$$

To study the sign of $4A_nB_n - B'_n$, we observe that, by Lagrange's Mean Value Theorem,

$$V'(x) - 4\alpha \frac{E_n - V(x)}{x_n - x} = V'(x) - 4\alpha V'(\xi) = -V'(x)(4\alpha V'(\xi)/V'(x) - 1) \quad (3.17)$$

for some $\xi \in (x, x_n)$.

Now, if $V \in \mathcal{P}_{1,\mathrm{uc}}(\kappa,\omega)$, then

$$\left|\log(V'(\xi)/V'(x))\right| \le \omega(\log(\xi/x))$$

and therefore

 $4\alpha V'(\xi)/V'(x) = \exp(\log(4\alpha) + \log(V'(\xi)/V'(x))) \ge \exp(\log(4\alpha) - \omega(\log(\xi/x))).$ If we take $\alpha \in (1/4, 1/2)$, then $\log(4\alpha) > 0$. Since $\lim_{t\to 0} \omega(t) = 0$, we can find $\delta = \delta(\alpha, \omega) > 0$ such that

 $\omega(t) \leq \log(4\alpha)$ whenever $0 < t \leq \delta$.

Consequently, whenever $0 < x_n/x \le e^{\delta}$, we have $4\alpha V'(\xi) - V'(x) \ge 0$, and therefore $4A_nB_n - B'_n < 0$. As a consequence, S_n is decreasing in the interval $[e^{-\delta}x_n, x_n)$.

If instead $V \in \mathcal{P}_{1,\mathrm{cv}}(\kappa)$, then an even simpler argument applies. Indeed, one can go back to (3.17), and observe that $V'(\xi)/V'(x) \geq 1$ in this case, because V' is increasing; consequently we obtain that $4A_nB_n - B'_n < 0$ on the whole $\mathbb{R}^+ \cap \{V < E_n\}$ whenever $\alpha \geq 1/4$. In particular, we can take $\alpha = 1/4$ and $\delta = 1$ in this case, and again conclude that S_n is decreasing in the interval $[e^{-\delta}x_n, x_n)$.

The fact that S_n is decreasing on $[e^{-\delta}x_n, x_n)$ implies that

$$f_n^2 + \frac{(f_n')^2}{B_n} \le f_n (e^{-\delta} x_n)^2 + \frac{f_n' (e^{-\delta} x_n)^2}{B_n (e^{-\delta} x_n)} \quad \text{on } [e^{-\delta} x_n, x_n).$$
(3.18)

Note now that

$$f_n = (x_n - x)^{\alpha} \psi_n, \qquad f'_n = (x_n - x)^{\alpha} \left[\psi'_n - \alpha (x_n - x)^{-1} \psi_n \right].$$

In particular, by (3.16),

$$f_n^2 + \frac{(f_n')^2}{B_n} \simeq_\alpha (x_n - x)^{2\alpha} \left[\psi_n^2 + \frac{(\psi_n')^2}{B_n} \right]$$
(3.19)

on $[e^{-\delta}x_n, x_n)$, and

$$f_{n}(e^{-\delta}x_{n})^{2} + \frac{f_{n}'(e^{-\delta}x_{n})^{2}}{B_{n}(e^{-\delta}x_{n})} \lesssim_{\bar{\kappa},\alpha} x_{n}^{2\alpha} \left[\psi_{n}(e^{-\delta}x_{n})^{2} + \frac{\psi_{n}'(e^{-\delta}x_{n})^{2}}{E_{n}} \right]$$
$$\lesssim_{\bar{\kappa},\alpha} \frac{x_{n}^{2\alpha}}{|\{V \le E_{n}\}|};$$

these inequalities are consequence of the fact that $e^{-\delta}x_n$ is well within the classical region (see Lemma 2.4(i)), so $E_n - V(e^{-\delta}x_n) \simeq_{\bar{\kappa},\alpha} E_n$ and the uniform estimate (3.6) applies. From (3.18) and (3.19) we then deduce that

$$\psi_n^2 + \frac{(\psi_n')^2}{B_n} \lesssim_{\bar{\kappa},\alpha} \frac{x_n^{2\alpha} (x_n - x)^{-2\alpha}}{|\{V \le E_n\}|} \quad \text{on } [e^{-\delta} x_n, x_n).$$
(3.20)

We now observe that, by Proposition 2.5,

$$x_n \simeq_{\kappa} |\{V \le E_n\}|$$

and, by Lemma 2.4,

$$E_n - V(x) \simeq_{\kappa} \frac{E_n}{|\{V \le E_n\}|} (x_n - x)$$

on $[e^{-\delta}x_n, x_n]$; hence, from (3.16) and Proposition 2.5 we deduce that

$$B_n \simeq_{\kappa} (E_n - V) \left(1 + \frac{n^{-2/3} E_n}{E_n - V} \right)^3$$

on $[e^{-\delta}x_n, x_n)$. As a consequence, the estimate (3.20) gives (3.15) on $[e^{-\delta}x_n, x_n)$. On the other hand, the interval $[0, e^{-\delta}x_n)$ is well within the classical region, so on that interval the estimate (3.15) follows from the uniform estimate (3.6). In conclusion, (3.15) is proved on $[0, x_n) = \{V < E_n\} \cap [0, \infty)$, as desired.

3.5. Zeros and local extrema of eigenfunctions and their derivatives. In this section we prove the estimates for ψ'_n of Theorem 3.5 on the whole \mathbb{R} , as well as the corresponding estimates for ψ_n within the classical region. These estimates will be derived from those proved in the previous sections, combined with information on the location of the extremum points of ψ_n and ψ'_n .

Assume at first that $V \in \mathcal{P}_1(\kappa)$. We recall a few basic facts about zeros and local extrema of ψ_n and ψ'_n , which are easy consequences of the fact that ψ_n is a square-integrable solution of (3.8).

- (a) ψ_n and ψ'_n do not vanish simultaneously at any point.
- (b) The zeros of ψ_n and ψ'_n are contained in the classical region $\{V < E_n\}$; outside of the classical region, $x\psi_n(x)\psi'_n(x) < 0$, which implies that both ψ_n^2 and $(\psi'_n)^2$ are strictly increasing on $\{V \ge E_n\} \cap (-\infty, 0)$ and strictly decreasing on $\{V \ge E_n\} \cap (0, \infty)$.
- (c) ψ_n has n-1 zeros in the classical region, which are all simple, so ψ_n changes sign at each zero.
- (d) The zeros of ψ''_n are the zeros of ψ_n and the two transition points (i.e., the points where $V = E_n$); these are the inflexion points of ψ_n .
- (e) Between two consecutive zeros of ψ''_n , the function ψ_n is strictly concave or strictly convex according to whether ψ_n is positive or negative.
- (f) Between two consecutive zeros of ψ''_n , there is exactly one zero of ψ'_n ; these are all the zeros of ψ'_n , which has n zeros, and they are all simple.
- (g) Similarly, between two consecutive zeros of ψ'_n , there is exactly one zero of ψ_n .
- (h) The zeros of ψ'_n are the local maximum/minimum points of ψ_n , that is, the local maximum points of ψ_n^2 .
- (i) Similarly, the zeros of ψ''_n are the local maximum/minimum points of ψ'_n , that is, the local maximum points of $(\psi'_n)^2$.

Since $|\psi'_n|$ attains its maximum in the classical region $\{V < E_n\}$, from (3.1) we derive the following uniform estimate for ψ'_n :

$$\|\psi'_n\|_{\infty} = \sup_{\{V < E_n\}} |\psi'_n| \lesssim_{\kappa} \frac{E_n^{1/2}}{|\{V \le E_n\}|^{1/2}}.$$
(3.21)

This proves the estimate for ψ'_n in Theorem 3.5(i), and moreover implies a rough uniform estimate for ψ_n :

$$\|\psi_n\|_{\infty} = \sup_{\{V < E_n\}} |\psi_n| \le |\psi_n(0)| + \int_{\{V < E_n\}} |\psi'_n| \lesssim_{\kappa} \frac{n}{|\{V \le E_n\}|^{1/2}}, \quad (3.22)$$

where (3.5) and Proposition 2.5 were used. Notice that the estimate (3.22) is worse than the uniform estimate for ψ_n in Theorem 3.5(i); to prove the latter, a more careful analysis of the local extrema of ψ_n and ψ'_n is needed. Further important information about local extrema is deduced from the monotonicity identities (3.2) and (3.4). These identities give us precise information on the sign of the derivatives of the two functions $(E_n - V)\psi_n^2 + (\psi'_n)^2$ and $\psi_n^2 + \frac{(\psi'_n)^2}{E_n - V}$; by evaluation at the zeros of $\psi''_n = (V - E_n)\psi_n$ and ψ'_n , these yield the following information.

- (j) The local maxima of ψ_n^2 (that is, the values of ψ_n^2 at the zeros of ψ_n') on $[0,\infty)$ form a strictly increasing sequence, while on $(-\infty, 0]$ they form a strictly decreasing sequence.
- (k) The local maxima of $(\psi'_n)^2$ (that is, the values of $(\psi'_n)^2$ at the zeros of ψ''_n) on $[0,\infty)$ form a strictly decreasing sequence, while on $(-\infty,0]$ they form a strictly increasing sequence.

In particular, the global maximum of ψ_n^2 is attained at an *outermost* zero of ψ_n' (that is, a zero closest to one of the two transition points). Similarly, for n > 1, the global maximum of $(\psi_n')^2$ is attained at an *innermost* zero of ψ_n (that is, the origin if $\psi_n(0) = 0$, or the positive and negative zeros of ψ_n that are closest to the origin if $\psi_n(0) \neq 0$). For this reason, it is useful to investigate the location of the zeros of ψ_n and ψ_n' within the classical region.

To this purpose, as in [Sz75, §6.31], we can fruitfully use Sturm's comparison theorem. Namely, for any $\tilde{E} \in (0, E_n)$, we have that $V - E_n \leq \tilde{E} - E_n$ on $\{V \leq \tilde{E}\}$. Hence, we can find a zero of ψ_n between any two zeros of a nontrivial solution of $-u'' + (\tilde{E} - E_n)u = 0$ on $\{V \leq \tilde{E}\}$; in other words, we have proved the following.

(1) For all $\tilde{E} < E_n$, there is a zero of ψ_n in any interval of length $\pi/\sqrt{E_n - \tilde{E}}$ fully contained in $\{V \leq \tilde{E}\}$.

In order to be able to apply this result, we need to ensure that such an interval exists, that is, we need to choose \tilde{E} so that

$$|\{V \le \tilde{E}\}| \sqrt{E_n - \tilde{E}} \ge \pi.$$

We now observe that, by Proposition 2.5,

$$\left(\frac{E_n}{n^{2/3}}\right)^{1/2} |\{V \le E_n/2\}| \simeq_{\kappa} n^{2/3}.$$

This means that there exists $n_0 = n_0(\kappa) \in \mathbb{N}_+$ sufficiently large that

$$\frac{E_n}{n^{2/3}} \le \frac{E_n}{2}, \quad \left(\frac{E_n}{n^{2/3}}\right)^{1/2} |\{V \le E_n/2\}| \ge \pi \quad \text{for all } n \ge n_0.$$

Consequently, if we take $\tilde{E} := E_n - \frac{E_n}{n^{2/3}}$, then

$$|\{V \le \tilde{E}\}|\sqrt{E_n - \tilde{E}} \ge \left(\frac{E_n}{n^{2/3}}\right)^{1/2} |\{V \le E_n/2\}| \ge \pi \quad \text{for all } n \ge n_0,$$

and the previous result can be applied.

We now observe that, if $x_n^{\pm}, y_n^{\pm} \in (0, \infty)$ are such that $V(\pm x_n^{\pm}) = E_n$ and $V(\pm y_n^{\pm}) = \tilde{E}$, then, by Lemma 2.4 and Proposition 2.5,

$$x_n^{\pm} - y_n^{\pm} \simeq_{\kappa} \frac{x_n^{\pm}}{E_n} (E_n - \tilde{E}) \simeq_{\kappa} \frac{|\{V \le E_n\}|}{n^{2/3}} \simeq_{\kappa} \frac{n^{1/3}}{E_n^{1/2}} \simeq \frac{\pi}{\sqrt{E_n - \tilde{E}}}$$

In conclusion, for all $n \ge n_0 = n_0(\kappa)$:

- (m) There are $\gtrsim_{\kappa} n^{2/3}$ zeros of ψ_n within the region $\{V \leq E_n E_n/n^{2/3}\}$, and the outermost of them have distance $\simeq_{\kappa} n^{-2/3} |\{V \leq E_n\}|$ from the transition point with the same sign.
- (n) Any two consecutive zeros of ψ_n within the classical region have distance $\lesssim_{\kappa} n^{-2/3} |\{V \le E_n\}|$

By using the above information, we can improve, for all $V \in \mathcal{P}_1(\kappa)$, the uniform estimate (3.22). Indeed, we know that the maximum of $|\psi_n|$ is attained at one of the outermost zeros of ψ'_n ; let us call this point w_n , and let ζ_n be the outermost zero of ψ_n with the same sign. From the above discussion, we deduce that, for all $n \ge n_0$, $|\zeta_n - w_n| \lesssim_{\kappa} n^{-2/3} |\{V \le E_n\}| \simeq_{\kappa} n^{1/3} / E_n^{1/2}$, hence, by (3.21),

$$\|\psi_n\|_{\infty} \le \left|\int_{\zeta_n}^{w_n} \psi'_n\right| \le |\zeta_n - w_n| \|\psi'_n\|_{\infty} \lesssim_{\kappa} \frac{n^{1/3}}{|\{V \le E_n\}|^{1/2}}.$$
 (3.23)

The same estimate for $n < n_0$ is already contained in (3.22), since $n \simeq_{\kappa} n^{1/3}$ for $n < n_0$. Combining this estimate with (3.1) proves the validity of the estimate for ψ_n of Theorem 3.5(i) within the classical region.

The above information on the location of the zeros and extrema of ψ_n and ψ'_n can also be used to prove the estimates for ψ'_n of parts (ii) to (iv) of Theorem 3.5 on the whole real line, as well as the corresponding estimates for ψ_n in the classical region. Indeed, under the assumptions on V and α in any of parts (ii) to (iv) of Theorem 3.5, we know from (3.7) and (3.15) that the improved pointwise estimate

$$\psi_n^2 + \left(1 + \frac{n^{-2/3}E_n}{E_n - V}\right)^{-3} \frac{(\psi_n')^2}{E_n - V} \lesssim_{\bar{\kappa},\alpha} \frac{E_n^{2\alpha}(E_n - V)^{-2\alpha}}{|\{V \le E_n\}|}$$
(3.24)

holds in the classical region $\{V < E_n\}$. In particular,

$$(\psi'_n)^2 \lesssim_{\bar{\kappa},\alpha} \frac{E_n^{2\alpha} (E_n - V)^{1-2\alpha}}{|\{V \le E_n\}|} \quad \text{on } \{V \le E_n - E_n/n^{2/3}\}.$$
 (3.25)

If we now apply this estimate at the two outermost zeros $\pm z_n^{\pm}$ of ψ_n within the region $\{V \leq E_n - E_n/n^{2/3}\}$, we obtain that, for all $n \geq n_0$,

$$\psi_n'(\pm z_n^{\pm})^2 \lesssim_{\bar{\kappa},\alpha} \frac{E_n^{2\alpha} (E_n - V(\pm z_n^{\pm}))^{1-2\alpha}}{|\{V \le E_n\}|} \simeq_{\kappa} \frac{E_n}{|\{V \le E_n\}|} n^{-2(1-2\alpha)/3},$$

where we used that $E_n - V(\pm z_n^{\pm}) \simeq_{\kappa} n^{-2/3} E_n$, due to the fact that the distance between $\pm z_n^{\pm}$ and the transition point of the same sign is $\simeq_{\kappa} n^{-2/3} |\{V \leq E_n\}|$ (see (m) and (n) above).

As previously discussed, the $\pm z_n^{\pm}$ are local maximum points of $(\psi'_n)^2$, and because of the monotonicity properties of the sequence of local maxima of $(\psi'_n)^2$ (see (k) above), we conclude that, for all $n \ge n_0$,

$$(\psi'_n)^2 \le \max_{\pm} \psi'_n (\pm z_n^{\pm})^2 \lesssim_{\bar{\kappa},\alpha} \frac{E_n}{|\{V \le E_n\}|} n^{-2(1-2\alpha)/3} \quad \text{on } \mathbb{R} \setminus (-z_n^-, z_n^+); \quad (3.26)$$

here we can go beyond the classical region, because $(\psi'_n)^2$ is increasing on $\{V \ge E_n\} \cap (-\infty, 0)$ and decreasing on $\{V \ge E_n\} \cap (0, \infty)$ (see (b) above). By combining (3.25) and (3.26) we deduce the estimates for ψ'_n of Theorem 3.5(ii)-(iv) on the whole \mathbb{R} .

As for the bound on ψ_n , we can argue as in (3.23), but use the improved bound on ψ'_n from (3.26). Namely, let w_n be an outermost zero of ψ'_n where $|\psi_n|$ attains its maximum, and ζ_n be the outermost zero of ψ_n with the same sign. Then $\zeta_n, w_n \in \mathbb{R} \setminus (-z_n^-, z_n^+)$ and $|z_n - w_n| \lesssim_{\kappa} n^{-2/3} |\{V \leq E_n\}| \simeq_{\kappa} n^{1/3} / E_n^{1/2}$, so

$$\|\psi_n\|_{\infty} = |\psi_n(w_n)| \le \left| \int_{\zeta_n}^{w_n} \psi_n' \right| \lesssim_{\kappa} \frac{n^{2\alpha/3}}{|\{V \le E_n\}|^{1/2}}$$
(3.27)

for all $n \ge n_0$. The same estimate for $n < n_0$ is already contained in (3.22), since $n \simeq_{\kappa,\alpha} n^{2\alpha/3}$ for $n < n_0$. Combining this estimate with (3.24) proves the validity of the estimates for ψ_n of Theorem 3.5(ii)-(iv) within the classical region.

3.6. Estimate outside the classical region. In order to complete the proof of Theorem 3.5, it remains to prove the estimate

$$|\psi_n(x)| \lesssim_{\bar{\kappa},\alpha} \frac{1}{|\{V \le E_n\}|^{1/2}} (V(x)/E_n - 1)^{-c}$$

outside the classical region. We only discuss the case x > 0, since the case x < 0 is treated analogously. So we need to prove the above estimates for $x > x_n^+$, where x_n^+ is the positive transition point.

Here we use the estimate from [Tit62, §8.2],

$$|\psi_n(x)| \le |\psi_n(x_n^+)| \exp\left(-\int_{x_n^+}^x (V - E_n)^{1/2}\right) \le \|\psi_n\|_{\infty} \exp\left(-\int_{x_n^+}^x (V - E_n)^{1/2}\right),$$

valid for all $x \ge x_n^+$, together with the estimates for $\|\psi_n\|_{\infty}$ obtained previously. Let $\tilde{x}_n^+ > 0$ be such that $V(\tilde{x}_n^+) = 4E_n$. Then, for $x \ge \tilde{x}_n^+$,

$$\int_{x_n^+}^x (V - E_n)^{1/2} \simeq_{\kappa} x \sqrt{V(x)} \ge x_n^+ E_n^{1/2} \sqrt{V(x)/E_n} \simeq_{\kappa} n \sqrt{V(x)/E_n}$$

(see [DM21, eq. (6.11)] and Proposition 2.5) and both factors in the last product are greater than or equal to 1. Hence, for some $c = c(\kappa)$, if we use the estimate for $\|\psi_n\|_{\infty}$ from (3.22), then we deduce that

$$\begin{aligned} |\psi_n(x)| \lesssim_{\kappa} \frac{n}{|\{V \le E_n\}|^{1/2}} \exp(-cn) \exp(-c\sqrt{V(x)/E_n}) \\ \lesssim_{\kappa,N} \frac{1}{|\{V \le E_n\}|^{1/2}} (V(x)/E_n)^{-N} \simeq_N \frac{1}{|\{V \le E_n\}|^{1/2}} (V(x)/E_n - 1)^{-N}, \end{aligned}$$

for any N > 0, since $V(x)/E_n \ge 4$ for $x \ge \tilde{x}_n^+$. For $x \in (x_n^+, \tilde{x}_n^+)$, instead,

$$\int_{x_n^+}^x (V - E_n)^{1/2} \simeq_{\kappa} \frac{|\{V \le E_n\}|}{E_n} \int_{x_n^+}^x (V - E_n)^{1/2} V'$$
$$\simeq_{\kappa} \frac{|\{V \le E_n\}|}{E_n} (V(x) - E_n)^{3/2}$$
$$\simeq_{\kappa} n \left(V(x)/E_n - 1\right)^{3/2},$$

by Proposition 2.5. So, if we use the estimate for $\|\psi_n\|_{\infty}$ from (3.23) and (3.27), then

$$\begin{aligned} |\psi_n(x)| \lesssim_{\bar{\kappa},\alpha} \frac{n^{2\alpha/3}}{|\{V \le E_n\}|^{1/2}} \exp\left(-cn\left(V(x)/E_n - 1\right)^{3/2}\right) \\ \lesssim_\alpha \frac{1}{|\{V \le E_n\}|^{1/2}} (V(x)/E_n - 1)^{-\alpha}, \end{aligned}$$

as desired.

4. Proof of the sharpened weighted Plancherel estimate

We are finally in a position to prove the desired sharpened version of the weighted Plancherel estimate of [DM21, Theorem 9.1]. We restate it as a separate theorem.

Theorem 4.1. Assume that $V \in \mathcal{P}_{1+\theta}(\kappa)$ for some $\theta \in (0,1)$. Let $m : \mathbb{R} \to \mathbb{C}$ be a bounded Borel function such that supp $m \subseteq [1/4, 1]$. Then, for all $\gamma \in [0, 1/2)$ and all r > 0,

$$\underset{z' \in \mathbb{R}^2}{\text{ess sup }} r^{2-2\gamma} \max\{V(r), V(x')\}^{1/2-\gamma} \int_{\mathbb{R}^2} |y - y'|^{2\gamma} \left| \mathcal{K}_{\mathrm{m}(r^2 \mathcal{L})}(z, z') \right|^2 \, dz \\ \lesssim_{\theta, \kappa, \gamma} \|\mathbf{m}\|_{L^2_{\gamma}}^2,$$

where z := (x, y) and z' := (x', y').

Proof. We follow the set-up and notation of [DM21, Section 9], but assume additionally that supp $\mathbf{m} \subseteq [1/4, 1]$. For $A \in \mathbb{R}^+$, define $G_A(\lambda, \tau) := \mathbf{m}(\lambda)\chi(A\tau)$, where $\chi \in C_c^{\infty}([1/4, 1])$ is such that $\sum_{j \in \mathbb{Z}} \chi(2^j \cdot) = 1$ on \mathbb{R}^+ , and set $K_{G_A} := \mathcal{K}_{G_A(\mathcal{L}, -\partial_y^2)}$. Then

$$\int_{\mathbb{R}^2} |K_{G_A}(z',z)|^2 dz' \lesssim A^{-1/2} \int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_1(\tau).\vec{\psi}(x;\tau V)\|^2 \frac{d\tau}{\tau}$$

$$\int_{\mathbb{R}^2} (y'-y)^2 |K_{G_A}(z',z)|^2 dz' \lesssim A^{1/2} \sum_{j=1}^4 \int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_j(\tau).\vec{\psi}(x;\tau V)\|^2 \frac{d\tau}{\tau}$$
(4.1)

(see [DM21, eqs. (9.11) and (9.15)]), where

$$\begin{split} \mathbf{M}_1(\tau) &:= \operatorname{diag}(\mathbf{m}(E(\tau V))), \\ \mathbf{M}_2(\tau) &:= \operatorname{diag}\mathbf{m}'(\vec{E}(\tau V)) \odot \operatorname{diag} \vec{F}(\tau V), \\ \mathbf{M}_3(\tau) &:= \mathbf{N} \odot \mathbf{A}(\tau V) \odot \operatorname{inc} \mathbf{m}(\vec{E}(\tau V)), \\ \mathbf{M}_4(\tau) &:= \mathbf{F} \odot \mathbf{A}(\tau V) \odot \operatorname{inc} \mathbf{m}(\vec{E}(\tau V)). \end{split}$$

Here $\vec{E}(\tau V) := (E_n(\tau V))_n$, $\vec{F}(\tau V) := (\tau \partial_\tau E_n(\tau V))_n$, $\vec{\psi}(\cdot; \tau V) := (\psi_n(\cdot; \tau V))_n$; the matrices $\mathbf{A}(\tau V)$, $\mathbf{P}(\tau V)$, \mathbf{N} , \mathbf{F} , are given by $\mathbf{A}_{nm}(\tau V) := \langle \tau \partial_\tau \psi_n(\cdot; \tau V), \psi_m(\cdot; \tau V) \rangle$, $\mathbf{P}_{nm}(\tau V) := \langle \tau V \psi_n(\cdot; \tau V), \psi_m(\cdot; \tau V) \rangle$, $\mathbf{N}_{nm} := \mathbf{1}_{n/2 \le m \le 2n}$, $\mathbf{F}_{nm} := \mathbf{1}_{n>2m} + \mathbf{1}_{m>2n}$; moreover \odot is the Schur product between matrices, $\|\cdot\|$ is the ℓ^2 -norm, and

diag
$$\vec{f} := (f_n \delta_{nm})_{n,m}$$
, inc $\vec{f} := (f_n - f_m)_{n,m}$.

In the proof of [DM21, Theorem 9.1], the integrals in $\frac{d\tau}{\tau}$ in (4.1) are bounded by the corresponding suprema, which eventually results in estimates involving L^{∞} -Sobolev norms of m. In order to obtain sharper estimates with L^2 -Sobolev norms, here instead we crucially take advantage of the integration in τ .

Let us first consider the term involving the diagonal matrix $\mathbf{M}_1(\tau)$:

$$\int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_{1}(\tau).\vec{\psi}(x;\tau V)\|^{2} \frac{d\tau}{\tau}$$

$$= \int_{A^{-1}}^{4A^{-1}} \sum_{n} |\mathbf{m}(E_{n}(\tau V))|^{2} \psi_{n}(x;\tau V)^{2} \frac{d\tau}{\tau}$$

$$= \int_{0}^{\infty} |\mathbf{m}(\lambda)|^{2} \sum_{n:1/\Xi_{n}(\lambda)\in[A/4,A]} \psi_{n}(x;\Xi_{n}(\lambda)V)^{2} \frac{\lambda \Xi_{n}'(\lambda)}{\Xi_{n}(\lambda)} \frac{d\lambda}{\lambda}$$

$$\lesssim_{\kappa} \int_{0}^{\infty} |\mathbf{m}(\lambda)|^{2} \sum_{n:\lambda/\Xi_{n}(\lambda)\in[A/16,A]} \psi_{n}(x;\Xi_{n}(\lambda)V)^{2} \frac{d\lambda}{\lambda}$$

by Proposition 2.1. In light of Theorem 3.5(ii) and Remark 3.2, we can apply Theorem 2.2 with $\tilde{\mathcal{P}} = \mathcal{P}_{1+\theta}(\kappa)$ to bound the above sum, and deduce that

$$\int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_{1}(\tau).\vec{\psi}(x;\tau V)\|^{2} \frac{d\tau}{\tau} \lesssim_{\theta,\kappa} \|\mathbf{m}\|_{2}^{2} (\mathbf{1}_{V \le 4A} + e^{-c|x|} \mathbf{1}_{V > 4A}).$$
(4.2)

As observed in [DM21, Section 9.3], $\mathbf{M}_2(\tau)$ is also a diagonal matrix, with diagonal entry $\mathbf{m}'(E_n(\tau V))\tau \partial_{\tau} E_n(\tau V)$, and

$$|\mathbf{m}'(E_n(\tau V))\tau \partial_\tau E_n(\tau V)| \le |\widetilde{\mathbf{m}}(E_n(\tau))|,$$

where $\widetilde{m}(\lambda) := \lambda m'(\lambda)$ (see Proposition 2.1). So the same argument as above, with \widetilde{m} in place of m, yields

$$\int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_{2}(\tau).\vec{\psi}(x;\tau V)\|^{2} \frac{d\tau}{\tau} \lesssim_{\theta,\kappa} \|\mathbf{m}'\|_{2}^{2} (\mathbf{1}_{V \le 4A} + e^{-c|x|} \mathbf{1}_{V > 4A}).$$
(4.3)

We now deal with the "near-diagonal" term $\mathbf{M}_3(\tau)$. As discussed in [DM21, Section 9.5], the absolute value of the (n, m) entry of $\mathbf{M}_3(\tau)$ is

 $\mathbf{1}_{n/2 \le m \le 2n} |\mathbf{A}_{nm}(\tau V)| |\mathbf{m}(E_n(\tau V)) - \mathbf{m}(E_m(\tau V))|.$

Note that, since supp $m \subseteq [1/4, 1]$ and $E_n(\tau V) \simeq_{\kappa} E_m(\tau V)$ for $n/2 \leq m \leq 2n$, there is $S = S(\kappa) \geq 1$ such that the above entry vanishes unless $E_m(\tau V) \in [S^{-1}, S]$. Now,

$$|\mathbf{m}(E_n(\tau V)) - \mathbf{m}(E_m(\tau V))| = \left| \int_{E_m(\tau V)}^{E_n(\tau V)} \mathbf{m}'(\lambda) \, d\lambda \right|$$
$$\leq |E_n(\tau V) - E_m(\tau V)| \, \mathcal{M}(\mathbf{m}')(E_m(\tau V)),$$

where \mathcal{M} denotes the uncentred Hardy–Littlewood maximal function on \mathbb{R} . Consequently, if we set $\widehat{\mathbf{m}} := \mathbf{1}_{[S^{-1},S]}\mathcal{M}\mathbf{m}'$, then

$$\begin{split} \mathbf{1}_{n/2 \leq m \leq 2n} \left| \mathbf{A}_{nm}(\tau V) \right| \left| \mathbf{m}(E_n(\tau V)) - \mathbf{m}(E_m(\tau V)) \right| \\ \leq \mathbf{1}_{n/2 \leq m \leq 2n} \left| \mathbf{P}_{nm}(\tau V) \right| \widehat{\mathbf{m}}(E_m(\tau V)) \\ \lesssim_{\kappa,\theta} \frac{1}{1 + |m - n|^{1+\epsilon}} \widehat{\mathbf{m}}(E_m(\tau V)), \end{split}$$

where $\epsilon = \epsilon(\kappa, \theta)$, and we applied [DM21, Proposition 8.1 and Theorem 8.4]. Since the matrix $((1 + |m - n|^{1+\epsilon})^{-1})_{n,m \ge 1}$ is ℓ^2 -bounded, we conclude that

$$\|\mathbf{M}_{3}(\tau).\vec{\psi}(x;\tau V)\| \lesssim_{\kappa,\theta} \|\operatorname{diag}(\widehat{\mathbf{m}}(\vec{E}(\tau V))).\vec{\psi}(x;\tau V)\|.$$

So, the same argument that proves (4.2), applied with \widehat{m} in place of m, yields, for some $T_1 = T_1(\kappa)$,

$$\int_{4A^{-1}}^{A^{-1}} \|\mathbf{M}_{3}(\tau).\vec{\psi}(x;\tau V)\|^{2} \frac{d\tau}{\tau} \lesssim_{\theta,\kappa} \|\widehat{\mathbf{m}}\|_{2}^{2} (\mathbf{1}_{V \leq T_{1}A} + e^{-c|x|} \mathbf{1}_{V > T_{1}A}) \\
\lesssim \|\mathbf{m}'\|_{2}^{2} (\mathbf{1}_{V \leq T_{1}A} + e^{-c|x|} \mathbf{1}_{V > T_{1}A}),$$
(4.4)

where the last bound follows from the L^2 -boundedness of \mathcal{M} .

Finally, from [DM21, eq. (9.19)], we already know that

$$\int_{A^{-1}}^{4A^{-1}} \|\mathbf{M}_{4}(\tau).\vec{\psi}(x;\tau V)\| \frac{d\tau}{\tau} \lesssim_{\kappa} \|\mathbf{m}\|_{\infty}^{2} (\mathbf{1}_{V \le T_{2}A} + e^{-c|x|} \mathbf{1}_{V > T_{2}A})
\lesssim (\|\mathbf{m}\|_{2} + \|\mathbf{m}'\|_{2})^{2} (\mathbf{1}_{V \le T_{2}A} + e^{-c|x|} \mathbf{1}_{V > T_{2}A})$$
(4.5)

for some $T_2 = T_2(\kappa) \ge 4$, where the last estimate follows from Sobolev's embedding. In conclusion, from (4.1), (4.2), (4.3), (4.4) and (4.5), we deduce that

$$\int_{\mathbb{R}^2} |K_{G_A}(z',z)|^2 dz' \lesssim_{\kappa} A^{-1/2} \|\mathbf{m}\|_2^2 (\mathbf{1}_{V \le T_3 A} + e^{-c|x|} \mathbf{1}_{V > T_3 A}),$$
$$\int_{\mathbb{R}^2} (y'-y)^2 |K_{G_A}(z',z)|^2 dz' \lesssim_{\kappa,\theta} A^{1/2} (\|\mathbf{m}\|_2 + \|\mathbf{m}'\|_2)^2 (\mathbf{1}_{V \le T_3 A} + e^{-c|x|} \mathbf{1}_{V > T_3 A}),$$

where $T_3 := \max\{T_1, T_2\}$. These two estimates are analogous to the ones stated at the beginning of [DM21, Section 9.6], with L^2 -norms of m and m' instead of L^{∞} -norms. As in [DM21, Section 9.6], by interpolating these two estimates and then

summing them for $A = 2^j$, $j \in \mathbb{Z}$, one eventually deduces that, for all $\gamma \in [0, 1/2)$,

$$\left(\int_{\mathbb{R}^2} |y - y'|^{2\gamma} |\mathcal{K}_{\mathrm{m}(\mathcal{L})}(z', z)|^2 \, dz'\right)^{1/2} \lesssim_{\theta, \kappa, \gamma} \|\mathbf{m}\|_{L_2^{\gamma}} \max\{V(1), V(x)\}^{\gamma/2 - 1/4},$$

which is the case r = 1 of the estimate in Theorem 4.1. The estimate for arbitrary r > 0 follows by rescaling, that is, by replacing V(x) with $V_r(x) := r^2 V(rx)$, as explained at the end of [DM21, Section 9.6].

References

- [BS91] F. A. Berezin and M. A. Shubin, The Schrödinger equation, Mathematics and its Applications (Soviet Series), vol. 66, Kluwer Academic Publishers Group, Dordrecht, 1991.
- [CS13] P. Chen and A. Sikora, Sharp spectral multipliers for a new class of Grushin type operators, J. Fourier Anal. Appl. 19 (2013), no. 6, 1274–1293.
- [DM20] G. M. Dall'Ara and A. Martini, A robust approach to sharp multiplier theorems for Grushin operators, Trans. Amer. Math. Soc. 373 (2020), 7533–7574.
- [DM21] G. M. Dall'Ara and A. Martini, An optimal multiplier theorem for Grushin operators in the plane, I, preprint (2021). arXiv:2107.12015
- [DOS02] X. T. Duong, E. M. Ouhabaz, and A. Sikora, *Plancherel-type estimates and sharp spectral multipliers*, J. Funct. Anal. **196** (2002), no. 2, 443–485.
- [Heb95] W. Hebisch, Functional calculus for slowly decaying kernels, preprint (1995), available at http://www.math.uni.wroc.pl/~hebisch/.
- [KST82] C. E. Kenig, R. J. Stanton, and P. A. Tomas, Divergence of eigenfunction expansions, J. Funct. Anal. 46 (1982), no. 1, 28–44.
- [Kr08] I. Krasikov, On the Erdélyi-Magnus-Nevai conjecture for Jacobi polynomials, Constr. Approx. 28 (2008), 113–125.
- [Mar17] A. Martini, Joint functional calculi and a sharp multiplier theorem for the Kohn Laplacian on spheres, Math. Z. 286 (2017), 1539–1574.
- [MM14] A. Martini and D. Müller, A sharp multiplier theorem for Grushin operators in arbitrary dimensions, Rev. Mat. Iberoam. 30 (2014), no. 4, 1265–1280.
- [MMN21] A. Martini, D. Müller, and S. Nicolussi Golo, Spectral multipliers and wave equation for sub-Laplacians: lower regularity bounds of Euclidean type, J. Eur. Math. Soc. (JEMS), to appear. doi:10.4171/JEMS/1191
- [MS12] A. Martini and A. Sikora, Weighted Plancherel estimates and sharp spectral multipliers for the Grushin operators, Math. Res. Lett. 19 (2012), no. 5, 1075–1088.
- [Mit74] B. S. Mitjagin, Divergenz von Spektralentwicklungen in L_p-Räumen, Linear operators and approximation, II (Proc. Conf., Oberwolfach Math. Res. Inst., Oberwolfach, 1974), Internat. Ser. Numer. Math., Vol. 25, Birkhäuser, Basel, 1974, 521–530.
- [Olv74] F. W. J. Olver, Asymptotics and special functions, Academic Press, New York-London, 1974.
- [RS08] D. W. Robinson and A. Sikora, Analysis of degenerate elliptic operators of Grušin type, Math. Z. 260 (2008), no. 3, 475–508.
- [Sz75] G. Szegő, Orthogonal polynomials, fourth ed., American Mathematical Society, Providence, R.I., 1975.
- [Tit62] E. C. Titchmarsh, Eigenfunction expansions associated with second-order differential equations. Part I, Second Edition, Clarendon Press, Oxford, 1962.

(G. M. Dall'Ara) Istituto Nazionale di Alta Matematica "Francesco Severi", Research Unit Scuola Normale Superiore, Piazza dei Cavalieri 7, 56126 Pisa, Italy

ORCID: 0000-0003-1168-3815

 $Email \ address: \verb"dallara@altamatematica.it"$

(A. Martini) School of Mathematics, University of Birmingham, Edgbaston, Birmingham, B15 2TT, United Kingdom

ORCID: 0000-0002-1568-704X

Current address: Dipartimento di Scienze Matematiche "G. L. Lagrange", Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Torino, Italy

Email address: alessio.martini@polito.it