# POLITECNICO DI TORINO Repository ISTITUZIONALE

Production of muons from heavy-flavour hadron decays at high transverse momentum in Pb–Pb collisions at sNN=5.02 and 2.76 TeV

**Original** 

Production of muons from heavy-flavour hadron decays at high transverse momentum in Pb–Pb collisions at sNN=5.02 and 2.76 TeV / Acharya, S; Adamova, D.; Adolfson, J; Aggarwal, M.; Rinella, G.; Agnello, M.; Agrawal, N; Bufalino, S.; Concas, M.; Grosa, F.; Ravasenga, I; Balbino, A.; Catalano, F.; Fecchio, P.. - In: PHYSICS LETTERS. SECTION B. - ISSN 0370-2693. - STAMPA. - 820:(2021), p. 136558. [10.1016/j.physletb.2021.136558]

Availability: This version is available at: 11583/2962447 since: 2022-05-02T21:50:57Z

Publisher: Elsevier B.V.

Published DOI:10.1016/j.physletb.2021.136558

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Contents lists available at [ScienceDirect](http://www.ScienceDirect.com/)

## Physics Letters B



[www.elsevier.com/locate/physletb](http://www.elsevier.com/locate/physletb)

Production of muons from heavy-flavour hadron decays at high transverse momentum in Pb–Pb collisions at  $\sqrt{s_{NN}} = 5.02$  and 2.76 TeV

## .ALICE Collaboration *-*

#### A R T I C L E I N F O A B S T R A C T

*Article history:* Received 8 December 2020 Received in revised form 30 June 2021 Accepted 28 July 2021 Available online 2 August 2021 Editor: M. Doser

Measurements of the production of muons from heavy-flavour hadron decays in Pb–Pb collisions at √<sup>*S*NN</sub> = 5.02 and 2.76 TeV using the ALICE detector at the LHC are reported. The nuclear modification</sup> factor  $R_{AA}$  at  $\sqrt{s_{NN}}$  = 5.02 TeV is measured at forward rapidity (2.5 < *y* < 4) as a function of transverse momentum  $p<sub>T</sub>$  in central, semi-central, and peripheral collisions over a wide  $p<sub>T</sub>$  interval,  $3 < p_T < 20$  GeV/*c*, in which muons from beauty-hadron decays are expected to take over from charm as the dominant source at high  $p_T$  ( $p_T > 7$  GeV/*c*). The  $R_{AA}$  shows an increase of the suppression of the yields of muons from heavy-flavour hadron decays with increasing centrality. A suppression by a factor of about three is observed in the 10% most central collisions. The  $R_{AA}$  at  $\sqrt{s_{NN}} = 5.02$  TeV is similar to that at  $\sqrt{s_{NN}}$  = 2.76 TeV. The precise  $R_{AA}$  measurements have the potential to distinguish between model predictions implementing different mechanisms of parton energy loss in the high-density medium formed in heavy-ion collisions. They place important constraints for the understanding of the heavy-quark interaction with the hot and dense QCD medium.

© 2021 European Organization for Nuclear Research, ALICE. Published by Elsevier B.V. This is an open access article under the CC BY license [\(http://creativecommons.org/licenses/by/4.0/\)](http://creativecommons.org/licenses/by/4.0/). Funded by  $SCOAP<sup>3</sup>$ .

#### **1. Introduction**

The study of ultra-relativistic heavy-ion collisions aims to investigate a state of strongly-interacting matter at high energy density and temperature. Under these extreme conditions, quantum chromodynamics (QCD) calculations on the lattice predict the formation of a quark–gluon plasma (QGP), where quarks and gluons are deconfined, and chiral symmetry is partially restored [1–4].

Heavy quarks (charm and beauty) are key probes of the QGP properties in the laboratory. They are predominantly created in hard-scattering processes at the early stage of the collision on a timescale shorter than the formation time of the QGP of ∼ 0*.*1–1 fm/*c* [5,6]. Therefore, they experience the full evolution of the hot and dense QCD medium. During their propagation through the medium, they lose energy via radiative and collisional processes [7–12]. Quarks are expected to lose less energy than gluons due to the colour-charge dependence of the strong interaction. Furthermore, several mass-dependent effects can also influence the energy loss. Due to the dead-cone effect [8,9,13], the heavyquark radiative energy loss is reduced compared to that of light quarks and the energy loss of beauty quarks is expected to be

smaller than that of charm quarks. The collisional heavy-quark energy loss is also expected to be reduced since the spatial diffusion coefficient, which controls the momentum exchange with the medium, is predicted to scale with the inverse of the quark mass [14]. In addition to the heavy-quark energy loss, modifications of the hadronisation process via fragmentation and/or recombination [15,16] and initial-state effects such as the modification of the parton distribution functions (PDF) inside the nucleus [17–19] can also change the particle yields and phase-space distributions. The medium effects can be quantified using the nuclear modification factor  $R_{AA}$ , which is the ratio between the  $p_T$ and *y*-differential particle yields in nucleus-nucleus (AA) collisions  $(d^2N_{AA}/dp_Tdy)$  and the corresponding production cross section in pp collisions (d<sup>2</sup>*σ*<sub>pp</sub>/d*p*<sub>T</sub>d*y*) scaled by the average nuclear overlap function  $\langle T_{AA} \rangle$ :

$$
R_{AA}(p_T, y) = \frac{1}{\langle T_{AA} \rangle} \times \frac{d^2 N_{AA}/dp_T dy}{d^2 \sigma_{pp}/dp_T dy}.
$$
 (1)

The  $\langle T_{AA} \rangle$  is defined as the ratio between the average number of nucleon–nucleon collisions  $\langle N_{\text{coll}} \rangle$  and the inelastic nucleon– nucleon cross section [20].

Evidence of a strong suppression of open heavy-flavour yields was observed in central Au–Au and Cu–Cu collisions at  $\sqrt{s_{NN}}$  = 200 GeV by the PHENIX and STAR collaborations at RHIC

*<sup>-</sup> E-mail address:* [alice-publications@cern.ch](mailto:alice-publications@cern.ch).

<https://doi.org/10.1016/j.physletb.2021.136558>

<sup>0370-2693/© 2021</sup> European Organization for Nuclear Research, ALICE. Published by Elsevier B.V. This is an open access article under the CC BY license ([http://creativecommons.org/licenses/by/4.0/\)](http://creativecommons.org/licenses/by/4.0/). Funded by SCOAP3.

and in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV by the ALICE, ATLAS, and CMS collaborations at the LHC (see [5] and references therein, and [21–24]). Recently, the ALICE and CMS collaborations reported a significant suppression of the prompt D-meson yields measured at midrapidity in the 10% most central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV with respect to the scaled pp reference, reaching a factor of about 5–6 in the interval  $8 < p_T < 12$  GeV/*c* [25,26]. A strong suppression of the yields of high- $p<sub>T</sub>$  electrons from heavy-flavour hadron decays was also observed by the ALICE collaboration at midrapidity in the 0–10% centrality class, where the measured *R*<sub>AA</sub> is about 0.3 at  $p<sub>T</sub> \sim 7$  GeV/*c* [27]. The suppression is similar to that observed for prompt D mesons and leptons from heavy-flavour hadron decays at  $\sqrt{s_{NN}}$  = 2.76 TeV [21,24,28]. The nuclear modification factor of  $B^{\pm}$  mesons, reconstructed via the exclusive decay channel  $B^{\pm} \to J/\psi K^{\pm} \to \mu^+ \mu^- K^{\pm}$  with the CMS detector for  $|y| < 2.4$  and  $7 < p_T < 50$  GeV/*c*, indicates a suppression of about a factor two in Pb–Pb collisions (0–100% centrality class) at  $\sqrt{s_{NN}}$  = 5.02 TeV [29] compatible with that of J/*ψ* from b-hadron decays (non-prompt J/*ψ*) [30]. A similar suppression as for  $B^{\pm}$  mesons and non-prompt  $J/\psi$  is also observed for non-prompt  $D^0$  mesons in the kinematic region  $|y| < 2.4$  and  $2 < p_T < 100$  GeV/*c* [31]. The suppression of B mesons is weaker than that of prompt  $D^0$  mesons at about  $p_T = 10$  GeV/*c*, in line with the expected quark-mass ordering of energy loss.

This letter presents the first measurement of open heavyflavour production via muons from semi-leptonic decays of charm and beauty hadrons in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV with the ALICE detector at the LHC. These measurements are carried out in the forward rapidity region  $(2.5 < y < 4)$ , presently only covered by the ALICE experiment at the LHC in Pb–Pb collisions. They extend the measurement of open heavy-flavour production from mid to forward rapidities, providing a tomography of the QGP medium in broader phase space region. The analysis of muontriggered events and large branching ratios (∼ 10%) allow us to perform high precision measurements of the  $p_T$ -differential  $R_{AA}$  of these muons over a broad  $p<sub>T</sub>$  interval, extended for the first time to  $p_T = 20$  GeV/*c* in central (0 – 10%), semi-central (20 – 40%), and peripheral  $(60 - 80%)$  collisions. This gives access to the investigation of medium effects in a new kinematic regime where the contribution of muons originating from beauty hadrons is dominant at high  $p_T$  ( $p_T > 7$  GeV/*c*). New measurements in central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV, with a significantly extended  $p_T$  coverage and a higher precision compared to the previous ALICE publication [32], are reported and compared to the results at  $\sqrt{s_{NN}}$  = 5.02 TeV. The computation of the  $R_{AA}$  makes use of the measured pp references published in [32,33]. Detailed comparisons with model calculations with different implementations of in-medium energy loss are discussed as well.

#### **2. Experimental apparatus and data samples**

The ALICE apparatus and its performance are described in [34, 35]. The analysis is based on the detection of muons in the forward muon spectrometer covering the pseudorapidity interval −<sup>4</sup> *< η <* −2*.*5. Note that the muon spectrometer covers <sup>a</sup> negative *η* range in the ALICE reference frame and consequently a negative *y* range. The results are chosen to be presented with a positive *y* notation, due to the symmetry of the collision system. The muon spectrometer consists of a front absorber of 10 nuclear interaction lengths  $(\lambda_I)$  filtering hadrons, followed by five tracking stations, each composed of two planes of Cathode Pad Chambers, with the third station inside a dipole magnet with a field integral of 3  $T \times m$ . The tracking system is complemented with two trigger stations, each equipped with two planes of Resistive Plate Chambers downstream an iron wall of 7  $\lambda$ <sub>I</sub>. Finally, a conical absorber shields the muon spectrometer against secondary particles produced by the

interaction of primary particles at large *η* in the beam pipe. The Silicon Pixel Detector (SPD), made of two cylindrical layers covering the pseudorapidity intervals  $|\eta| < 2$  and  $|\eta| < 1.4$ , is employed for the reconstruction of the primary vertex. Two V0-scintillator arrays covering  $2.8 < \eta < 5.1$  and  $-3.7 < \eta < -1.7$  provide a minimum bias (MB) trigger defined as the coincidence of signals from the two hodoscopes. The V0 detectors are also used to classify events according to their centrality, determined from a fit of the total signal amplitude based on a two-component particle production model connected to the collision geometry using the Glauber formalism [36]. The centrality intervals are defined as percentiles of the Pb–Pb hadronic cross section. The V0 and the Zero Degree Calorimeters (ZDC), placed at  $\pm$ 112.5 m from the interaction point along the beam direction, are used for the event selection.

The results presented in this letter are based on the data sample recorded with the ALICE detector during the 2015 Pb–Pb run at a centre-of-mass energy  $\sqrt{s_{NN}}$  = 5.02 TeV. For the comparison with measurements at lower energy,  $\sqrt{s_{NN}}$  = 2.76 TeV, the 2011 data sample is used in order to extend the  $p_T$  coverage with respect to the published results from the 2010 data sample [32]. The analysis of the two data samples is based on muon-triggered events requiring a MB trigger and at least one track segment in the muon trigger system with a  $p_T$  larger than a programmable threshold [34]. Data were collected with two  $p_T$ -trigger thresholds of about 1 (0.5) and 4.2 (4.2) GeV/*c* at  $\sqrt{s_{NN}}$  = 5.02 TeV  $(\sqrt{s_{NN}}$  = 2.76 TeV). The  $p_{\text{T}}$  threshold of the trigger algorithm is set such that the corresponding efficiency for muon tracks is 50%. In the following, the low- and high- $p<sub>T</sub>$  trigger-threshold samples are referred to as MSL and MSH, respectively. The beam-induced background is reduced offline using the V0 and ZDC timing information, and electromagnetic interactions are removed by requiring a minimum energy deposited in the ZDC [37,38]. Only events with a primary vertex within  $\pm 10$  cm along the beam line are analysed. Finally, the measurements are done in the three representative centrality classes 0–10%, 20–40% and 60–80% to investigate the evolution of the *R*<sub>AA</sub> with the collision centrality. After the event selection, the data samples correspond to integrated luminosities of about 21*.*9 (224*.*8) μb−<sup>1</sup> and 4*.*0 (71*.*0) μb−<sup>1</sup> for MSL- (MSH- ) triggered events at  $\sqrt{s_{NN}}$  = 5.02 and 2.76 TeV, respectively. The integrated luminosity is derived from the number of muontriggered events. These muon-triggered events are normalised by a factor, inversely proportional to the probability of having a muon trigger in a MB event in a given centrality class, calculated from the relative count rate between the muon and MB triggers.

### **3. Analysis procedure**

#### *3.1. Measurement of muons from heavy-flavour hadron decays*

Standard selection criteria are applied to the muon candidates [33]. Tracks in the muon spectrometer are reconstructed within the pseudorapidity range  $-4 < \eta < -2.5$  and they are required to have a polar angle measured at the exit of the absorber in the interval  $170^\circ < \theta_{\text{abs}} < 178^\circ$ . Furthermore, tracks are identified as muons if they match a track segment in the trigger system. Finally, the remaining beam-induced background is reduced by requiring the distance of the track to the primary vertex measured in the transverse plane (DCA, distance of closest approach) weighted with its momentum (*p*),  $p \times DCA$ , to be smaller than  $6 \times \sigma_{pDCA}$ , where  $\sigma_{pDCA}$  is the width of the distribution.

The nuclear modification factor R<sub>AA</sub> of muons from heavyflavour hadron decays is measured down to  $p_T = 3$  GeV/ $c$  and up to  $p_T = 20$  GeV/*c* in all centrality classes at  $\sqrt{s_{NN}} = 5.02$  TeV and in the 0–10% centrality class at  $\sqrt{s_{NN}}$  = 2.76 TeV. The  $R_{AA}$  is computed for  $p_T > 3$  GeV/*c* in order to limit the systematic uncertainty on the subtraction of the background of muons from light-hadron

decays, which increases with decreasing  $p<sub>T</sub>$ . These measurements are performed by using MSL-triggered events up to  $p_T = 7$  GeV/*c* and MSH-triggered events for  $p_T > 7$  GeV/*c*. In the selected  $p_T$ interval, after the selection criteria are implemented, the main background contributions to the muon yields consist of muons from primary charged-pion and kaon decays for  $p_T < 6$  GeV/*c*, and muons from W-boson, Z-boson, and  $\gamma^{\star}$  (Drell-Yan process) decays for  $p_T > 13$  GeV/*c*. Two additional small contributions of muons from secondary (charged) light-hadron decays in the interval  $3 < p_T < 5$  GeV/*c*, resulting from the interaction of light hadrons with the material of the front absorber and of muons from  $J/\psi$  decays over the entire  $p_T$  range, are also considered. Therefore, the  $p_T$ -differential  $R_{AA}$  of muons from heavy-flavour hadron decays in a given centrality class is expressed as

$$
R_{AA}(p_T, y) = \frac{\left(\frac{d^2 N^{\mu^{\pm}}}{dp_T dy} - \sum_{\text{non-HF}\to\mu^{\pm}} \frac{d^2 N^{\text{non-HF}\to\mu^{\pm}}}{dp_T dy}\right)_{\text{Pb}-\text{Pb}}}{\langle T_{AA} \rangle \times \left(\frac{d^2 \sigma^{c,b\to\mu^{\pm}}}{dp_T dy}\right)_{\text{pp}}},
$$
(2)

where  $d^2N^{\mu^{\pm}}/dp_Tdy$  is the differential yield of inclusive muons and  $\sum_{\text{non-HF}\to\mu^{\pm}} d^2N^{\text{non-HF}\to\mu^{\pm}}/dp_Tdy$  refers to the differential yields of muons from various non heavy-flavour sources in Pb– Pb collisions, as indicated above Eq. (2). In the denominator, d2*σ*<sup>c</sup>*,*b→*μ*<sup>±</sup> */*d*p*Td*y* is the pp differential production cross section of muons from heavy-flavour hadron decays at the same centreof-mass energy and in the same kinematic region (see [32,33]) as in Pb–Pb collisions.

## 3.2. *Pb–Pb* collisions at  $\sqrt{s_{NN}} = 5.02$  *TeV*

#### *3.2.1. Efficiency corrections*

The inclusive muon yields in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5*.*02 TeV are corrected for detector acceptance and detection efficiencies ( $A \times \varepsilon$ ) using the procedure described in previous publications [32,33]. In peripheral collisions,  $A \times \varepsilon$  amounts to about 90% with almost no  $p<sub>T</sub>$  dependence in the region of interest for MSL-triggered events, while for MSH-triggered events the *<sup>A</sup>* × *ε* increases with  $p_T$  from 75% at  $p_T = 7$  GeV/*c* towards a plateau at a value close to 90% for  $p<sub>T</sub> > 14$  GeV/*c*. The dependence of the trigger and tracking efficiency on the detector occupancy is determined by embedding simulated muons from heavy-flavour hadron decays in measured MB Pb–Pb events. A decrease in the efficiency of 6% from peripheral (60–80%) to central (0–10%) collisions, independent of  $p<sub>T</sub>$  is observed.

#### *3.2.2. Estimation of the muon background sources*

The estimation of the contribution of muons from primary *π*<sup>±</sup> and  $K^{\pm}$  decays is based on a data-tuned Monte Carlo cocktail. The procedure uses the midrapidity ( $|\eta|$  < 0.8)  $\pi^{\pm}$  and K<sup> $\pm$ </sup> spectra measured by the ALICE collaboration up to  $p_T = 20$  GeV/*c* [39] in pp and Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV. They are further extrapolated to higher  $p<sub>T</sub>$ , up to  $p<sub>T</sub> = 40$  GeV/*c*, by means of a power-law fit to extend the  $p_T$  coverage to the  $p_T$  interval relevant for the estimation of the decay muons up to  $p_T = 20$  GeV/*c*. Then, the extrapolation to forward rapidities is performed assuming the same suppression of primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> yields from midrapidity up to  $y = 4$  according to

$$
\left[\frac{d^2 N^{\pi^{\pm}(K^{\pm})}}{dp_T dy}\right]_{AA} = \langle N_{\text{coll}} \rangle \times \left[ R_{AA}^{\pi^{\pm}(K^{\pm})} \right]^{\text{mid}-y}
$$

$$
\times [F_{\text{extrap}}^{\pi^{\pm}(K^{\pm})}(p_{T}, y)]_{\text{pp}} \times \left[\frac{d^{2}N^{\pi^{\pm}(K^{\pm})}}{dp_{T}dy}\right]_{\text{pp}}^{\text{mid}-y}.
$$
\n(3)

Equation (3) can be also expressed as

$$
\left[\frac{d^2 N^{\pi^{\pm}(K^{\pm})}}{dp_T dy}\right]_{AA} = [F_{\text{extrap}}^{\pi^{\pm}(K^{\pm})}(p_T, y)]_{\text{pp}} \times \left[\frac{d^2 N^{\pi^{\pm}(K^{\pm})}}{dp_T dy}\right]_{AA}^{\text{mid}-y}, (4)
$$

where  $[F_{\text{extrap}}^{\pi^{\pm}(K^{\pm})}(p_{T}, y)]_{\text{pp}}$  is the  $p_{T}$ - and y-dependent extrapolation factor in pp collisions at  $\sqrt{s}$  = 5.02 TeV, discussed in [33], which is based on Monte Carlo simulations. The systematic uncertainty due to the unknown suppression at forward rapidity will be discussed below. The PYTHIA 6*.*4 [40] and PHOJET [41] event generators are employed for the rapidity extrapolation, while PYTHIA 8*.*2 simulations [42] with various colour reconnection (CR) options are performed to take into account the rapidity dependence of the  $p_T$  extrapolation and its uncertainty. The  $p_T$  and *y* distributions of muons from primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays in Pb–Pb collisions are generated according to a fast detector simulation of the decay kinematics and of the effect of the front absorber [33] using as input the extrapolated  $\pi^{\pm}$  and K<sup> $\pm$ </sup> spectra. For each centrality class, the yields are further subtracted from the inclusive muon distribution. The total contribution of muons from primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays decreases with increasing  $p_T$  from about 21% (13%) at  $p_T = 3$  GeV/*c* down to about 7% (4%) at  $p_T = 20$  GeV/*c* in the 60–80% (0–10%) centrality class, with a weak  $p<sub>T</sub>$  dependence for  $p_T > 10$  GeV/*c*.

The estimation of the background muons from secondary  $\pi^{\pm}$ and  $K^{\pm}$  decays produced in the front absorber is based on Monte Carlo simulations using the HIJING event generator [43] and the GEANT3 transport package [44]. These simulation results indicate that in the  $p_T$  interval of interest, the relative contribution of secondary muons with respect to muons from primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays is about 9%, independently of both  $p_T$  and the collision centrality. Given the estimated contamination of muons from primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays, the contribution of these secondary muons relative to the total muon yield decreases with increasing  $p<sub>T</sub>$  from about 2% (1%) at  $p_T = 3$  GeV/*c* in the 60-80% (0-10%) centrality class to less than 1% at  $p_T = 5$  GeV/*c* for all centrality classes.

The estimation of the contribution of muons from W-boson decays and dimuons from Z-boson and  $\gamma^*$  decays, which is relevant in the high- $p<sub>T</sub>$  region, is based on the POWHEG NLO event generator [45] combined with PYTHIA 6*.*4*.*25 [40] for the parton shower, which reproduces within uncertainties the W- and Z-boson production in various LHC experiments [46–50]. These simulations include the CT10 PDF set [51] and the EPS09 NLO parameterisation [17] of the nuclear modification of the PDFs. In order to account for isospin effects, muons from W-boson decays and dimuons from Z-boson decays and  $\gamma^*$  decays are simulated separately in pp, np, pn, and nn collisions. A weighted sum of the production cross sections in the four systems is performed to obtain the production cross section per nucleon–nucleon collision for the Pb–Pb system. The latter is further scaled with  $\langle T_{AA} \rangle$  in a given centrality class in order to estimate the corresponding relative contribution of W and  $Z/\gamma^*$  with respect to inclusive muons. The relative contribution of muons from W and  $Z/\gamma^*$  with respect to inclusive muons is negligible for  $p<sub>T</sub> < 13$  GeV/*c* and it increases with  $p_T$  and the collision centrality from about 3% (6%) at  $p_T = 14$  GeV/*c* up to 18% (36%) at  $p_T = 20$  GeV/*c* in the 60-80% (0–10%) centrality class.

The contribution of muons from J*/ψ* decays is estimated by extrapolating the  $J/\psi$   $p_T$  and *y* spectra measured by ALICE at forward rapidity  $(2.5 < y < 4)$  in the interval of  $p_T < 12$  GeV/*c* [52]. The  $J/\psi$   $p_T$  and rapidity spectra are extrapolated by means of a power-law and Gaussian function up to  $p<sub>T</sub> = 50$  GeV/*c* and

 $|y| = 6.5$ , respectively. Then, the decay muon distributions are estimated with a fast detector simulation using the extrapolated J*/ψ* distributions as inputs, similar to pp collisions [33]. In the 10% most central collisions, the relative contribution to the inclusive muon distribution varies between 0*.*5 and 4%, with the maximum fraction at intermediate  $p_T$  (4 <  $p_T$  < 6 GeV/*c*).

#### *3.2.3. Systematic uncertainties*

The systematic uncertainties of the  $R_{AA}$  of muons from heavyflavour hadron decays at  $\sqrt{s_{NN}}$  = 5.02 TeV are evaluated considering the following sources: uncertainties of the inclusive muon yields and background contributions in Pb–Pb collisions, the pp reference, and the normalisation in both pp and Pb–Pb collisions.

The procedure to determine the systematic uncertainty on the inclusive muon yields is similar to that described in [33] and includes the following contributions: i) the muon tracking efficiency (1*.*5%), ii) the muon trigger efficiency resulting from the intrinsic efficiency of the muon trigger chambers and the response of the trigger algorithm (1*.*4% (3%) for the MSL (MSH) data sample), and iii) the choice of the  $\chi^2$  selection used in defining the matching of tracks reconstructed in the tracking system with those in the trigger system (0*.*5%). These systematic uncertainties are approximately independent of centrality and  $p<sub>T</sub>$  in the region of interest. The systematic uncertainty arising from the dependence of *<sup>A</sup>* × *ε* on the detector occupancy, obtained from a fit with a constant of the  $p_T$ -differential ratio of the efficiency in a given centrality class to that in peripheral collisions, increases up to 0*.*5% when going from peripheral to central collisions. Finally, the systematic uncertainty due to the tracking chamber resolution and alignment is based on a Monte Carlo simulation modelling the tracker response with a parameterisation of the tracking chamber resolution and misalignment effects, as described in [33,50]. This systematic uncertainty is negligible for  $p<sub>T</sub> < 7$  GeV/*c* and increases up to 12% in the interval  $18 < p_T < 20$  GeV/*c*.

The estimation of the yields of muons from primary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays is subject to systematic uncertainties arising, as described in [33], from i) the uncertainties of the measured midrapidity spectra of  $\pi^{\pm}$  (K<sup> $\pm$ </sup>) and their  $p_T$  extrapolation, which increase from about 3% (6%) to 6% (13%), ii) the rapidity extrapolation which results in a systematic uncertainty of about 8*.*5% (6%) for muons from  $\pi^{\pm}$  (K<sup> $\pm$ </sup>) decays obtained by comparing the results with PYTHIA 6 and PHOJET generators, iii) the rapidity dependence of the  $p_T$  extrapolation with a systematic uncertainty, obtained from the PYTHIA 8 generator with different CR options, increasing up to about 4% (2%) at  $p_T = 20$  GeV/*c* for  $\pi^{\pm}$  (K<sup> $\pm$ </sup>), and iv) the simulation of hadronic interactions in the absorber which leads to a systematic uncertainty of 4% independently of the muon origin, as reported in [33]. Adding in quadrature the uncertainties coming from each source, the total systematic uncertainty ranges from about 9% (10%) to 13% (15%) as a function of the  $p<sub>T</sub>$  of muons from primary  $\pi^{\pm}$  (K<sup> $\pm$ </sup>) decays. Finally, there is a contribution related to the assumption on the rapidity dependence of the suppression of  $\pi^{\pm}$  and K<sup> ${\pm}$ </sup>. Based on ATLAS measurements in Pb–Pb collisions at  $√s_{NN}$  = 2.76 TeV, which indicate no significant *η* dependence of the charged-particle *R*<sub>AA</sub> up to  $|\eta|$  < 2 [53], the suppression of  $\pi^{\pm}$ and  $K^{\pm}$  is considered to be independent of rapidity up to  $y = 4$ , and the *R*<sub>AA</sub> of  $\pi^{\pm}$  and K<sup> $\pm$ </sup> is varied conservatively within  $\pm$ 50%. This uncertainty is propagated to the decay muons and the difference between the upper and lower limits is further divided by  $\sqrt{12}$ , corresponding to the RMS of a uniform distribution. Furthermore, the effect of the transport code is conservatively evaluated by varying the estimated yield of muons from secondary  $\pi^{\pm}$  and  $K^{\pm}$  decays by  $\pm 100\%$  and dividing also the difference between lower and upper limits by  $\sqrt{12}$ .

The systematic uncertainty of the extracted muon yields from W and Z/ $\gamma^*$  decays is obtained considering the CT10 PDF uncertainty [51] and a different nuclear modification of the PDF (EKS98 [54–56] was used as well). It amounts to 5*.*9% (13*.*2%) for muons from W (Z/*γ*<sup>\*</sup>) decays.

The systematic uncertainty of the estimated yields of muons from J*/ψ* decays reflects the uncertainty of the measured J*/ψ* spectra at forward rapidity and their extrapolation to a wider kinematic region. It varies from about 9% at  $p_T = 3$  GeV/*c* to 34% at  $p_T = 20$  GeV/*c* in central collisions.

Two sources contribute to the systematic uncertainty on the normalisation, the systematic uncertainty of  $\langle T_{AA} \rangle$  values [20] and the systematic uncertainty of the normalisation factor needed to calculate the number of equivalent MB events in the muon samples. The latter is evaluated comparing the values from the nominal procedure (see section 2) with those calculated by applying the muon-trigger condition in the analysis of MB events [33].

The sources of systematic uncertainty affecting the measurement of the pp reference production cross section were evaluated in [33]. The total systematic uncertainty ranges from 2*.*1% to 15*.*1%, depending on  $p_T$ . A global pp normalisation uncertainty of 2.1%, discussed in [33], is considered as well. When computing the nuclear modification factor, the systematic uncertainty on track resolution and misalignment is considered to be partially correlated between the pp and Pb–Pb measurements because the pp data were collected just before the Pb–Pb run at  $\sqrt{s_{NN}}$  = 5.02 TeV and the detector conditions remained unchanged. The other sources of systematic uncertainties are treated as uncorrelated. The systematic uncertainty on the  $p_T$ -differential production cross section in pp collisions without including the correlated part of the uncertainty varies from 2*.*1% to 4*.*2%. The uncorrelated part of the uncertainty on track resolution and misalignment is due to the different shapes of the  $p_T$  distribution between pp and Pb–Pb collisions. It is estimated by comparing the results with and without correcting the residual misalignment between data and Monte Carlo when calculating the  $R_{AA}$ , as detailed in  $[33]$ .

The various systematic uncertainties are propagated to the measurement of the yields or nuclear modification factors of muons from heavy-flavour hadron decays and added in quadrature, except for the systematic uncertainties on normalisation which are shown separately.

Table 1 presents a summary of the relative systematic uncertainties assigned to the  $p_T$ -differential yields of muons from heavy-flavour hadron decays in Pb–Pb collisions. The systematic uncertainty on the pp reference, needed for the computation of the  $R_{AA}$ , is also reported.

## 3.3. *Pb–Pb collisions*  $at \sqrt{s_{NN}} = 2.76 \text{ TeV}$

For a direct comparison with lower energy measurements in the same  $p_T$  interval, the Pb–Pb data sample at  $\sqrt{s_{NN}} = 2.76$  TeV, collected in 2011, was analysed in order to significantly extend the  $p_T$  interval of the published  $R_{AA}$  measurements of muons from heavy-flavour hadron decays, which was limited to  $4 < p<sub>T</sub>$ 10 GeV/*c* [32]. Such an improvement is possible due to the larger integrated luminosity (4  $\mu$ b<sup>-1</sup> and 71  $\mu$ b<sup>-1</sup> for MSL- and MSHtriggered collisions compared to 2*.*7 *μ*b<sup>−</sup>1) and the use of a high- $p_T$  muon trigger.

The strategy to extract the yields of muons from heavy-flavour hadron decays in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV is similar to that just discussed for  $\sqrt{s_{NN}}$  = 5.02 TeV. Compared to the latter case, the  $A \times \varepsilon$  exhibits the same trend as a function of  $p_{\text{t}}$ , although the values are smaller due to the status of the tracking chambers (larger number of inactive channels). The factor  $A \times \varepsilon$ saturates at a value close to 80% in the high- $p<sub>T</sub>$  region for peripheral collisions (60–80% centrality class). A decrease of the efficiency of 4% from peripheral collisions to the 10% most central collisions, due to the detector occupancy, is seen. The fractions of the vari-

#### **Table 1**

Summary of the relative systematic uncertainties of the  $p_T$ -differential yields of muons from heavy-flavour hadron decays at forward rapidity (2.5 <  $y$  < 4) in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (second and third columns) and 2.76 TeV (fourth column). The systematic uncertainties of the pp reference are also summarised. For the  $p_T$ -dependent uncertainties, the minimum and maximum values are reported and correspond to the lowest and highest  $p<sub>T</sub>$  interval with the exception of the background of muons from light-hadron decays and the  $R_{AA}^{\pi^{\pm}(K^{\pm})}(y)$  assumption, where this is the opposite. See the text for details.

Source	$\sqrt{s_{NN}}$ = 5.02 TeV 0-10% centrality class	60–80% centrality class	$\sqrt{s_{NN}}$ = 2.76 TeV 0-10% centrality class
Tracking efficiency	1.5%	1.5%	2.5%
Trigger efficiency	1.4% (MSL), 3% (MSH)	1.4% (MSL), 3% (MSH)	1.4% (MSL), 2.3% (MSH)
Matching efficiency	0.5%	0.5%	0.5%
$A \times \varepsilon$	0.5%	$\Omega$	1%
Resolution and alignment	$0-12\%$ (0-4.1% on $R_{AA}$ )	$0-12\%$ (0-4.1% on $R_{AA}$ )	$1\% \times p_{\text{T}}$ ( $p_{\text{T}}$ in GeV/c)
Background subtraction $\mu \leftarrow \pi$	< 1.6%	< 2.5%	< 1.8%
Background subtraction $\mu \leftarrow K$	< 1.6%	< 2.5%	< 4%
$R_{AA}^{\pi^{\pm}(K^{\pm})}(y)$ assumption	$1.3 - 4.8%$	$1.5 - 7.8%$	$1.8 - 5.2%$
Background subtraction $\mu \leftarrow \sec \pi / K$	$0 - 0.8%$	$0 - 1.4%$	$0 - 0.9%$
Background subtraction $\mu \leftarrow W/Z/\gamma^*$	$0 - 1.6%$	$0 - 0.7%$	$0 - 3.1%$
Background subtraction $\mu \leftarrow J/\psi$	${<}0.4%$	${<}0.4%$	${<}0.3%$
Normalisation factor	$0.3\%$ (MSL), $0.7\%$ (MSH)	$0.3\%$ (MSL), $0.7\%$ (MSH)	$0.4\%$ (MSL), $1.6\%$ (MSH)
$\langle T_{AA} \rangle$	0.7%	2.5%	0.9%
pp reference for $R_{AA}$	$2.1 - 4.2%$	$2.1 - 4.2%$	15–18% (3 < $p_T$ < 10 GeV/c data)
			30–34% (10 $< p_T < 20$ GeV/c extrapolation)
pp reference (global) for $R_{AA}$	2.1%	2.1%	1.9%

ous background sources with respect to the inclusive muon yields at  $\sqrt{s_{NN}}$  = 2.76 TeV are compatible with the ones measured at  $\sqrt{s_{NN}}$  = 5.02 TeV. The fraction of muons from primary  $\pi^{\pm}$  and  $K^{\pm}$  decays with respect to inclusive muons varies between about 3% and 14% in the 0–10% centrality class, the largest values being obtained at  $p_T = 3$  GeV/*c*. On the other hand, the fraction of muons from secondary  $\pi^{\pm}$  and K<sup> $\pm$ </sup> decays reaches about 1% at  $p_T = 3$  GeV/*c*. The fraction of muons from electroweak-boson decays is significant at high  $p<sub>T</sub>$ , where it reaches about 30% in the interval  $16.5 < p_T < 20$  GeV/*c* for central collisions. Finally, the component of muons from  $J/\psi$  decays is small over the whole  $p_T$ interval with a maximum of 4% at intermediate  $p_T$  (~ 6 GeV/*c*) in central collisions. The same sources of systematic uncertainties as for the  $\sqrt{s_{NN}}$  = 5.02 TeV analysis are considered and same methods to estimate them are employed, except the systematic uncertainty of the tracking chamber resolution and alignment which varies linearly with  $p_T$  as  $1\% \times p_T$  ( $p_T$  in GeV/*c*) [32]. The  $p_T$ differential cross section of muons from heavy-flavour hadron decays in pp collisions at  $\sqrt{s}$  = 2.76 TeV measured in the intervals  $2.5 < y < 4$  and  $3 < p_T < 10$  GeV/*c* is used for the  $R_{AA}$  computation [32]. The measured production cross section is extrapolated up to  $p_T = 20$  GeV/*c* using fixed-order plus next-to-leading logarithms (FONLL) calculations [57,58]. The systematic uncertainty of the  $p_T$ -differential production cross section in pp collisions at  $√s$  = 2.76 TeV varies within 15–18% in 3 <  $p_T$  < 10 GeV/*c*. At higher  $p<sub>T</sub>$ , the systematic uncertainty, which also includes the systematic uncertainty on the FONLL calculations, reaches 30–34%.

A summary of all systematic uncertainties taken into account in the measurement of the  $p<sub>T</sub>$ -differential yields of muons from heavy-flavour hadron decays at  $\sqrt{s_{NN}}$  = 2.76 TeV is reported in Table 1, including the uncertainties of the pp reference.

#### **4. Results and model comparisons**

The  $p_T$ -differential yields of muons from heavy-flavour hadron decays normalised to the equivalent number of MB events at forward rapidity  $(2.5 < y < 4)$  in central, semi-central and peripheral Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV are shown in Fig. 1 (upper panel). The same observable measured in central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV is displayed in the lower panel of Fig. 1. The measurements are performed over a wide  $p<sub>T</sub>$  range from 3 to 20 GeV/*c* for all centrality classes.



**Fig. 1.** The  $p_T$ -differential yields of muons from heavy-flavour hadron decays at forward rapidity  $(2.5 < y < 4)$  in central  $(0-10\%)$ , semi-central  $(20-40\%)$ , and peripheral (60–80%) Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (upper panel), and in central (0–10%) Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV (lower panel). Statistical uncertainties (vertical bars) and systematic uncertainties (open boxes) are shown. The additional systematic uncertainty on normalisation in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5*.*02 *(*2*.*76*)* TeV for MSL- and MSH-triggered events, respectively, is not included in the uncertainty boxes (see Table 1).

The *p*T-differential *R*<sub>AA</sub> of muons from heavy-flavour hadron decays at forward rapidity  $(2.5 < y < 4)$  in Pb-Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV is presented in Fig. 2 for the same centrality classes as in Fig. 1. An increasing reduction of the yield of muons from heavy-flavour hadron decays with increasing centrality with respect to the pp reference scaled by the average nuclear overlap function is clearly seen. The suppression is largest at in-



Fig. 2. The *p*<sub>T</sub>-differential nuclear modification factor  $R_{AA}$  of muons from heavyflavour hadron decays at forward rapidity  $(2.5 < y < 4)$  in central  $(0-10\%$ , top), semi-central  $(20-40\%$ , middle), and peripheral  $(60-80\%$ , bottom) Pb-Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (symbols). Statistical (vertical bars) and systematic uncertainties (open boxes) are shown. The filled boxes centered at  $R_{AA} = 1$  represent the normalisation uncertainty of pp and Pb–Pb measurements. Horizontal bars reflect the bin widths and the values are shown at the centre of the bin. Left: the measured *R*<sub>AA</sub> is compared with the TAMU and SCET models [59,60] displayed with their uncertainty bands. Right: the measured R<sub>AA</sub> is compared with MC@sHQ+EPOS2 model calculations with pure collisional energy loss (dashed lines) and a combination of collisional and radiative energy loss (full lines) [61,62].

termediate  $p_T$ , in the interval from about 6 to 10 GeV/ $c$ , and reaches a factor of about three in the 10% most central collisions. Such behaviour is more pronounced in central and semi-central collisions, while moving towards peripheral collisions, the suppression presents no significant  $p<sub>T</sub>$  dependence. In minimum bias p–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, where the formation of an extended QGP is not expected, the nuclear modification factor  $R_{\text{pPb}}$  of muons from heavy-flavour hadron decays is consistent with unity at  $p_T > 6$  GeV/*c* [63]. The latter measurement confirms that the strong suppression observed in Pb–Pb collisions results from final-state interactions of charm and beauty quarks with the QGP. The evolution of R<sub>AA</sub> as a function of centrality is compatible with the dependence of the heavy-quark energy loss on the medium density and the average path length in the medium, both of which are larger in central than in peripheral collisions.

The measured  $R_{AA}$  is compared with various model predictions such as TAMU [59] and SCET [60] (Fig. 2, left), and MC@sHQ+EPOS2 [61,62] (Fig. 2, right). In the TAMU model, the interactions are described by elastic collisions only. The perturbative QCD (pQCD)-based SCET model implements medium-induced gluon radiation via modified splitting functions with finite quark masses. These SCET calculations depend on the coupling constant *g* which describes the coupling strength between hard partons and the QGP medium. Its value is  $g = 1.9-2$ . In the MC@sHQ+EPOS2 model, two different options are considered, energy loss from



**Fig. 3.** Comparison of the  $p_T$ -differential nuclear modification factor of muons from heavy-flavour hadron decays at forward rapidity (2*.*5 *< y <* 4) in central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (green symbols) and  $\sqrt{s_{NN}}$  = 2.76 TeV (red symbols). Statistical (vertical bars) and systematic uncertainties (open boxes) are shown. The filled boxes centered at  $R_{AA} = 1$  are the normalisation uncertainties. Horizontal bars represent the bin widths.

medium-induced gluon radiation and collisional (elastic) processes or only collisional energy loss. In the scenario with pure collisional energy loss, the scattering rates are scaled by a global factor *K* larger than unity ( $K = 1.5$ ) in order to reproduce the  $R_{AA}$  and elliptic flow of open heavy-flavour hadrons measured at midrapidity at the LHC [61]. With a combination of collisional and radiative energy loss, the scaling factor is  $K = 0.8$ . All these models also consider a nuclear modification of the PDF (EPS09) [17]. Note that in the MC@sHQ+EPOS2 model shadowing is not considered for beauty-quark production. In addition to independent fragmentation, a contribution of hadronisation via quark recombination is included in all models with the exception of SCET. The SCET model is based on pQCD calculations of high- $p_T$  parton energy loss and provides a fair description of the data in central collisions, but it deviates from the data in non-central collisions. The TAMU calculations, which do not include radiative energy loss processes, underestimate the suppression at  $p_T > 6$  GeV/*c* in central and semi-central collisions, in particular. Both versions of the MC@sHQ+EPOS2 model, without and with radiative energy loss, describe the measurement within uncertainties for all centrality classes over the entire  $p<sub>T</sub>$  interval.

The results obtained at forward rapidity for muons from heavyflavour hadron decays at  $\sqrt{s_{NN}} = 5.02$  TeV complement those obtained at midrapidity for the electrons from heavy-flavour hadron decays [27] by the ALICE collaboration as well as the prompt Dmeson  $\left[25,26\right]$  and beauty measurements via B<sup> $\pm$ </sup> mesons [31], nonprompt  $D^0$  [31] and J/ $\psi$  [30] by the ALICE and CMS collaborations. The measured *R*<sub>AA</sub> of muons from heavy-flavour hadron decays for  $p_T > 8$  GeV/*c* is compatible with that obtained for beauty ( $D^0$ and J/ $\psi$  from beauty hadrons, B<sup> $\pm$ </sup>) for  $p_{\rm T}$ <sup>hadron</sup>  $> 10$  GeV/ $c$  [30,31] within uncertainties, although in a different kinematic region (different  $p_T$  and *y* intervals).

A comparison of the *R*<sub>AA</sub> of muons from heavy-flavour hadron decays in the 10% most central Pb–Pb collisions at  $\sqrt{s_{NN}} = 2.76$ and 5*.*02 TeV is presented in Fig. 3. The comparison illustrates the improvement of the precision of the measurement at  $\sqrt{s_{NN}}$  = 5.02 TeV with respect to that at  $\sqrt{s_{NN}}$  = 2.76 TeV. The total systematic uncertainty on the  $R_{AA}$  at  $\sqrt{s_{NN}} = 5.02$  TeV is reduced by a factor of about 3 to 6, depending on  $p<sub>T</sub>$ , compared to the same measurement at  $\sqrt{s_{NN}}$  = 2.76 TeV using the 2011 data sample. The reasons for such an improvement are twofold. The detector conditions were more stable during the  $\sqrt{s_{NN}}$  = 5.02 TeV than the  $\sqrt{s_{NN}}$  = 2.76 TeV data taking campaign and therefore better described in the simulations. Moreover, as the pp data at  $\sqrt{s}$  = 5.02 TeV were collected just a few days before the Pb–Pb run at  $\sqrt{s_{NN}}$  = 5.02 TeV, the detector conditions

were comparable and the systematic uncertainty on alignment and resolution between the two systems partially cancel when computing *R*<sub>AA</sub>, as discussed in section 3. The present measurement at  $\sqrt{s_{NN}}$  = 2.76 TeV is in agreement with the published results obtained at the same centre-of-mass energy in a smaller  $p<sub>T</sub>$  interval (4 <  $p<sub>T</sub>$  < 10 GeV/*c*) with larger uncertainties [32]. The precision is increased by a factor 1*.*1–1*.*6, mainly due to a better understanding of the detector response and a new datadriven strategy for the estimation of the contribution of muons from primary light-hadron decays. The comparison between the results obtained at the two centre-of-mass energies indicates that the suppression of heavy quarks at  $\sqrt{s_{NN}}$  = 5.02 TeV is similar to that at  $\sqrt{s_{NN}}$  = 2.76 TeV, as already observed in the midrapidity region for electrons from heavy-flavour hadron decays [22,27] and prompt D mesons  $[25]$ . This similarity between the  $R_{AA}$  measurements at the two energies may result from the interplay of the following two effects as discussed in [64]: a flattening of the  $p_T$ spectra of charm and beauty quarks with increasing collision energy, and a medium temperature estimated to be higher by about 7% at  $\sqrt{s_{NN}}$  = 5.02 TeV than at 2.76 TeV. The former would decrease the heavy-quark suppression (increase the *R*<sub>AA</sub>) by about 5% if the medium temperature remains unchanged, while the latter would increase the suppression (decrease the *R*<sub>AA</sub>) by about 10% (5%) for charm (beauty) quarks.

The measured  $R_{AA}$  at  $\sqrt{s_{NN}}$  = 2.76 TeV is compatible with that measured for muons from heavy-flavour hadron decays in  $|\eta|$  < 1 with the ATLAS detector [21] and for electrons from heavyflavour hadron decays in the interval  $|y| < 0.6 - 0.8$  by the AL-ICE collaboration [24]. The same behaviour is also observed at  $\sqrt{s_{NN}}$  = 5.02 TeV when comparing the  $R_{AA}$  of muons from heavy-flavour hadron decays with that measured at midrapidity for electrons from heavy-flavour hadron decays [27]. This confirms that heavy quarks suffer a strong in-medium energy loss over a wide rapidity interval. The similarity of the suppression in the two rapidity regions does not imply that heavy quarks lose similar energy. The observed trend may also result from the interplay of several effects such as the shape of initial heavy-quark  $p_T$  spectra and the path-length dependence of the heavy-quark energy loss, as discussed in [65]. Indeed, the properties of the QGP medium differ between mid and forward rapidity. The measured chargedparticle multiplicity densities are smaller at forward rapidity than at midrapidity [66]. The created medium is also smaller and consequently the travelled path length is shorter at forward rapidity.

The  $p_T$  distributions of muons from heavy-flavour hadron decays are sensitive to energy loss of both charm and beauty quarks. Due to the decay kinematics and the charm- and beauty-quark  $p_T$ differential production cross sections, one expects that for  $p_{\rm T}\lesssim$ 5 GeV/*c* the distributions are predominantly sensitive to the charm in-medium energy loss. FONLL calculations [57,58] predict that in pp collisions at  $\sqrt{s}$  = 5.02 TeV more than 70% of muons from heavy-flavour hadron decays originate from beauty quarks in the high- $p_T$  region ( $p_T > 10$  GeV/ $c$ ) and this contribution reaches 75% in the interval  $18 < p_T < 20$  GeV/*c*. Therefore, the strong suppression of muons from heavy-flavour hadron decays in the high- $p<sub>T</sub>$ region is expected to be dominated by the in-medium energy loss of beauty quarks. In order to further interpret the results, Fig. 4 shows a comparison with MC@sHQ+EPOS2 predictions for muons from charm- and beauty-hadron decays, separately, and for muons from the combination of the two, in central (0–10%) Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (top) and 2.76 TeV (bottom). The predictions considering the combination of elastic and radiative energy loss and pure elastic energy loss are shown in the left and right panels, respectively. Both versions of the MC@sHQ+EPOS2 model provide a fair description of the measured *R*<sub>AA</sub> of muons from heavy-flavour hadron decays in central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV within uncertainties. A similar agreement be-



**Fig. 4.** Comparison of the  $p_T$ -differential nuclear modification factors  $R_{AA}$  of muons from heavy-flavour hadron decays at forward rapidity (2*.*5 *< y <* 4) in central Pb– Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV (top) and  $\sqrt{s_{NN}}$  = 2.76 TeV (bottom) with MC@sHQ+EPOS2 calculations [61,62] with different scenarios considering either a combination of collisional and radiative energy loss (left) or a pure collisional energy loss (right). The predictions are shown for muons from heavy-flavour hadron decays, muons from only charm-hadron decays and muons from only beauty-hadron decays.

tween data and MC@sHQ+EPOS2 is achieved at  $\sqrt{s_{NN}}$  = 2.76 TeV although the model tends to slightly overestimate the measured  $R_{AA}$  at low/intermediate  $p_T$ . The measured  $R_{AA}$  at large  $p_T$  is closer to the model calculations for muons from beauty-hadron decays than for muons from charm-hadron decays when considering both elastic and radiative energy loss. For the scenario involving only collisional energy loss, the predicted difference between the suppression of muons from charm and beauty-hadron decays is less pronounced. The predicted ratio of the  $p_T$ -differential  $R_{AA}$  of muons from beauty-hadron decays to that of muons from charmhadron decays for  $p_T > 10$  GeV/*c* is in the range 1.2–1.4 for the scenario involving only collisional energy loss and in the range 2*.*5–2*.*8 when considering both elastic and radiative energy loss, depending on  $p_T$  and centre-of-mass energy. It is worth mentioning that the MC@sHQ+EPOS2 model is characterised by a large running coupling constant  $\alpha_s$  and a reduced Debye mass in the elastic heavy-quark scattering generating the radiation  $[67]$ . As a consequence, the radiative energy loss neglects finite path-length effects due to the gluon formation outside the QGP and is overestimated at high  $p_T$ . Such an effect is expected to be more pronounced for charm quarks than for beauty quarks due to the dead-cone effect [8].

#### **5. Conclusions**

In summary, the  $p_T$ -differential normalised yield and the nuclear modification factor R<sub>AA</sub> of muons from semi-leptonic decays of charm and beauty hadrons was measured at forward rapidity  $(2.5 < y < 4)$  for the first time over the wide  $p<sub>T</sub>$  interval  $3 < p_T < 20$  GeV/*c* in central, semi-central, and peripheral Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, and in central Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV with reduced systematic uncertainties compared to previous measurements.

The measured  $R_{AA}$  shows a clear evidence of a strong suppression, up to a factor of three in the 10% most central collisions with respect to the binary-scaled pp reference, for both collision energies. This suppression pattern is compatible with a large heavy-quark in-medium energy loss. The strong suppression which persists in the high- $p_T$  region, up to  $p_T = 20$  GeV/*c*, indicates that beauty quarks lose a significant fraction of their energy in the medium. The suppression becomes weaker from central to peripheral collisions. The evolution of R<sub>AA</sub> with the collision centrality reflects the dependence of energy loss on the path length in the QGP and the QGP energy density.

The *R*<sub>AA</sub> measurements have the potential to discriminate between different model calculations. The *RAA* is in fair agreement with transport model calculations that consider both collisional and radiative energy loss. The MC@sHQ+EPOS2 transport model including a hydrodynamic description of the medium, coupled with different implementations of the in-medium parton energy loss, describes the measured  $R_{AA}$  well over the whole  $p_T$  interval in central, semi-central, and peripheral collisions within uncertainties. This comparison brings new constraints on the relative inmedium energy loss of charm and beauty quarks.

The suppression is compatible with that measured at central rapidity for electrons from heavy-flavour hadron decays. These new precise  $R_{AA}$  measurements carried out over a wide  $p_T$  interval at forward rapidity in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV with smaller uncertainties with respect to same measurements at  $\sqrt{s_{NN}}$  = 2.76 TeV, currently only accessible by ALICE in central collisions, bring significant constraints on the modelling of the longitudinal dependence of the open heavy-flavour *R*<sub>AA</sub>. Therefore, the obtained results provide further insight on the in-medium parton energy loss mechanisms and, ultimately, will help determining the transport properties of the hot and dense deconfined QCD medium in the full phase space.

### **Declaration of competing interest**

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

#### **Acknowledgements**

The ALICE Collaboration would like to thank all its engineers and technicians for their invaluable contributions to the construction of the experiment and the CERN accelerator teams for the outstanding performance of the LHC complex. The ALICE Collaboration gratefully acknowledges the resources and support provided by all Grid centres and the Worldwide LHC Computing Grid (WLCG) collaboration. The ALICE Collaboration acknowledges the following funding agencies for their support in building and running the ALICE detector: A. I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation (ANSL), State Committee of Science and World Federation of Scientists (WFS), Armenia; Austrian Academy of Sciences, Austrian Science Fund (FWF): [M 2467-N36] and Nationalstiftung für Forschung, Technologie und Entwicklung, Austria; Ministry of Communications and High Technologies, National Nuclear Research Center, Azerbaijan; Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Financiadora de Estudos e Projetos (Finep), Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) and Universidade Federal do Rio Grande do Sul (UFRGS), Brazil; Ministry of Education of China (MOEC), Ministry of Science & Technology of China (MSTC) and National Natural Science Foundation of China (NSFC), China; Ministry of Science and Education and Croatian Science Foundation, Croatia; Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Cubaenergía, Cuba; Ministry of Education, Youth and Sports of the Czech Republic, Czech Republic; The Danish Council for Independent Research | Natural Sciences, the Villum Fonden and Danish National Research Foundation (DNRF),

Denmark; Helsinki Institute of Physics (HIP), Finland; Commissariat à l'Énergie Atomique (CEA) and Institut National de Physique Nucléaire et de Physique des Particules (IN2P3) and Centre National de la Recherche Scientifique (CNRS), France; Bundesministerium für Bildung und Forschung (BMBF) and GSI Helmholtzzentrum für Schwerionenforschung GmbH, Germany; General Secretariat for Research and Technology, Ministry of Education, Research and Religions, Greece; National Research, Development and Innovation Office, Hungary; Department of Atomic Energy, Government of India (DAE), Department of Science and Technology, Government of India (DST), University Grants Commission, Government of India (UGC) and Council of Scientific and Industrial Research (CSIR), India; Indonesian Institute of Sciences, Indonesia; Istituto Nazionale di Fisica Nucleare (INFN), Italy; Institute for Innovative Science and Technology, Nagasaki Institute of Applied Science (IIST), Japanese Ministry of Education, Culture, Sports, Science and Technology (MEXT) and Japan Society for the Promotion of Science (JSPS) KAKENHI, Japan; Consejo Nacional de Ciencia (CONACYT) y Tecnología, through Fondo de Cooperación Internacional en Ciencia y Tecnología (FONCICYT) and Dirección General de Asuntos del Personal Academico (DGAPA), Mexico; Nederlandse Organisatie voor Wetenschappelijk Onderzoek (NWO), Netherlands; The Research Council of Norway, Norway; Commission on Science and Technology for Sustainable Development in the South (COMSATS), Pakistan; Pontificia Universidad Católica del Perú, Peru; Ministry of Science and Higher Education, National Science Centre and WUT ID-UB, Poland; Korea Institute of Science and Technology Information and National Research Foundation of Korea (NRF), Republic of Korea; Ministry of Education and Scientific Research, Institute of Atomic Physics and Ministry of Research and Innovation and Institute of Atomic Physics, Romania; Joint Institute for Nuclear Research (JINR), Ministry of Education and Science of the Russian Federation, National Research Centre Kurchatov Institute, Russian Science Foundation and Russian Foundation for Basic Research, Russia; Ministry of Education, Science, Research and Sport of the Slovak Republic, Slovakia; National Research Foundation of South Africa, South Africa; Swedish Research Council (VR) and Knut & Alice Wallenberg Foundation (KAW), Sweden; European Organization for Nuclear Research, Switzerland; Suranaree University of Technology (SUT), National Science and Technology Development Agency (NSDTA) and Office of the Higher Education Commission under NRU project of Thailand, Thailand; Turkish Atomic Energy Agency (TAEK), Turkey; National Academy of Sciences of Ukraine, Ukraine; Science and Technology Facilities Council (STFC), United Kingdom; National Science Foundation of the United States of America (NSF) and United States Department of Energy, Office of Nuclear Physics (DOE NP), United States of America.

#### **References**

- [1] [Wuppertal-Budapest](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib90492710D3BBD7CF8CC8F1E74A1CB92As1) Collaboration, S. Borsanyi, Z. Fodor, C. Hoelbling, S.D. Katz, S. Krieg, C. Ratti, K.K. Szabo, Is there still any  $T_c$  [mystery](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib90492710D3BBD7CF8CC8F1E74A1CB92As1) in lattice QCD? Results with physical masses in the [continuum](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib90492710D3BBD7CF8CC8F1E74A1CB92As1) limit III, J. High Energy Phys. 09 (2010) 073, [arXiv:1005.3508 \[hep-lat\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib90492710D3BBD7CF8CC8F1E74A1CB92As1)
- [2] A. Bazavov, et al., Chiral and [deconfinement](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib40134703CC744C7C2E25B4EEB4BCA124s1) aspects of the QCD transition, Phys. Rev. D 85 (2012) 054503, [arXiv:1111.1710 \[hep-lat\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib40134703CC744C7C2E25B4EEB4BCA124s1)
- [3] S. Borsanyi, Z. Fodor, C. [Hoelbling,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B6A4EFA9CB7B3D9D1CBEE51E9FCB0BFs1) S.D. Katz, S. Krieg, K.K. Szabo, Full result for the QCD [equation](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B6A4EFA9CB7B3D9D1CBEE51E9FCB0BFs1) of state with 2+1 flavors, Phys. Lett. B 730 (2014) 99–104, [arXiv:1309.5258 \[hep-lat\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B6A4EFA9CB7B3D9D1CBEE51E9FCB0BFs1)
- [4] HotQCD [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAE239A4CCC891FC3597D7F292846A5B6s1) A. Bazavov, et al., Chiral crossover in QCD at zero and non-zero chemical potentials, Phys. Lett. B 795 (2019) 15–21, [arXiv:1812.08235](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAE239A4CCC891FC3597D7F292846A5B6s1) [\[hep-lat\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAE239A4CCC891FC3597D7F292846A5B6s1)
- [5] A. Andronic, et al., [Heavy-flavour](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB3673105415D9E34BD731A0DFD2DD50Fs1) and quarkonium production in the LHC era: from [proton-proton](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB3673105415D9E34BD731A0DFD2DD50Fs1) to heavy-ion collisions, Eur. Phys. J. C 76 (3) (2016) 107, [arXiv:1506.03981 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB3673105415D9E34BD731A0DFD2DD50Fs1)
- [6] F.-M. Liu, S.-X. Liu, [Quark-gluon](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAB23F4EB3D7F26313AF4150EE7798081s1) plasma formation time and direct photons from heavy ion collisions, Phys. Rev. C 89 (3) (2014) 034906, [arXiv:1212.6587](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAB23F4EB3D7F26313AF4150EE7798081s1) [\[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAB23F4EB3D7F26313AF4150EE7798081s1)
- [7] R. Baier, Y.L. [Dokshitzer,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib51A1F760C1B98F999DC0526BEFEAE4C8s1) A.H. Mueller, S. Peigne, D. Schiff, Radiative energy loss and  $p_T$  broadening of [high-energy](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib51A1F760C1B98F999DC0526BEFEAE4C8s1) partons in nuclei, Nucl. Phys. B 484 (1997) 265–282, [arXiv:hep-ph/9608322 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib51A1F760C1B98F999DC0526BEFEAE4C8s1)
- [8] Y.L. Dokshitzer, D.E. Kharzeev, Heavy quark [colorimetry](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib15E0329F58F5CADE770C30A31FCB08EBs1) of QCD matter, Phys. Lett. B 519 (2001) 199–206, [arXiv:hep-ph/0106202 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib15E0329F58F5CADE770C30A31FCB08EBs1)
- [9] M. [Djordjevic,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib5434D7BE33A31BDFECF5617CD95A4E2Cs1) M. Gyulassy, Heavy quark radiative energy loss in QCD matter, Nucl. Phys. A 733 (2004) 265–298, [arXiv:nucl-th/0310076 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib5434D7BE33A31BDFECF5617CD95A4E2Cs1)
- [10] B.-W. Zhang, E. Wang, X.-N. Wang, Heavy quark energy loss in a [nuclear](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAD5B15790600E2FA5808850F8AB56D3As1) medium, Phys. Rev. Lett. 93 (2004) 072301, [arXiv:nucl-th/0309040.](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAD5B15790600E2FA5808850F8AB56D3As1)
- [11] S. Wicks, W. Horowitz, M. [Djordjevic,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEF0B87FC0CC08664934589D4D505727Bs1) M. Gyulassy, Heavy quark jet quenching with collisional plus radiative energy loss and path length [fluctuations,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEF0B87FC0CC08664934589D4D505727Bs1) Nucl. Phys. A 783 (2007) 493–496, [arXiv:nucl-th/0701063 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEF0B87FC0CC08664934589D4D505727Bs1)
- [12] P.B. Gossiaux, J. Aichelin, T. Gousset, V. Guiho, [Competition](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibDAD04D456B25B7FF530174D5F2DEC50As1) of heavy quark radiative and collisional energy loss in [deconfined](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibDAD04D456B25B7FF530174D5F2DEC50As1) matter, J. Phys. G 37 (2010) 094019, [arXiv:1001.4166 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibDAD04D456B25B7FF530174D5F2DEC50As1)
- [13] N. Armesto, C.A. Salgado, U.A. [Wiedemann,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib5F4AA9BD7460E8D472E01EC09DC1966Bs1) Medium induced gluon radiation off massive quarks fills the dead cone, Phys. Rev. D 69 (2004) [114003,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib5F4AA9BD7460E8D472E01EC09DC1966Bs1) arXiv: [hep-ph/0312106 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib5F4AA9BD7460E8D472E01EC09DC1966Bs1)
- [14] H. van Hees, V. Greco, R. Rapp, [Heavy-quark](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib498FC2AEC967E1C273B21041AD57EF2Cs1) probes of the quark-gluon plasma and [interpretation](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib498FC2AEC967E1C273B21041AD57EF2Cs1) of recent data taken at the BNL relativistic heavy ion collider, Phys. Rev. C 73 (2006) 034913, [arXiv:nucl-th/0508055 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib498FC2AEC967E1C273B21041AD57EF2Cs1)
- [15] V. Greco, C.M. Ko, R. Rapp, Quark [coalescence](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib619E5F8AC22CE923715229B2CD2BA24As1) for charmed mesons in ul[trarelativistic](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib619E5F8AC22CE923715229B2CD2BA24As1) heavy ion collisions, Phys. Lett. B 595 (2004) 202–208, arXiv: [nucl-th/0312100 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib619E5F8AC22CE923715229B2CD2BA24As1)
- [16] A. Andronic, P. [Braun-Munzinger,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib223EDEA11545DA282566D91FBA15D992s1) K. Redlich, J. Stachel, Statistical hadronization of charm in heavy ion [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib223EDEA11545DA282566D91FBA15D992s1) at SPS, RHIC and LHC, Phys. Lett. B 571 (2003) 36–44, [arXiv:nucl-th/0303036 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib223EDEA11545DA282566D91FBA15D992s1)
- [17] K.J. Eskola, H. [Paukkunen,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3AC6EA9D723C8FDCC0EFC074474ECAA7s1) C.A. Salgado, EPS09: a new generation of NLO and LO nuclear parton [distribution](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3AC6EA9D723C8FDCC0EFC074474ECAA7s1) functions, J. High Energy Phys. 04 (2009) 065, [arXiv:0902.4154 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3AC6EA9D723C8FDCC0EFC074474ECAA7s1)
- [18] B.Z. [Kopeliovich,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib77FD79096481E38004751E3308420B80s1) J. Nemchik, A. Schafer, A.V. Tarasov, Cronin effect in hadron production off nuclei, Phys. Rev. Lett. 88 (2002) 232303, [arXiv:hep-ph/0201010](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib77FD79096481E38004751E3308420B80s1) [\[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib77FD79096481E38004751E3308420B80s1)
- [19] I. Vitev, [Non-Abelian](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibFABB201BBBF6F557120DD8477503FFC3s1) energy loss in cold nuclear matter, Phys. Rev. C 75 (2007) 064906, [arXiv:hep-ph/0703002 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibFABB201BBBF6F557120DD8477503FFC3s1)
- [20] ALICE Collaboration, Centrality determination in heavy ion collisions, ALICE-PUBLIC-2018-011, <http://cds.cern.ch/record/2636623>, 2018.
- [21] ATLAS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib7A866520C9DD96A9E238CD97F2E832C7s1) M. Aaboud, et al., Measurement of the suppression and azimuthal anisotropy of muons from [heavy-flavor](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib7A866520C9DD96A9E238CD97F2E832C7s1) decays in Pb+Pb collisions at √*S*<sub>NN</sub> = 2.76 TeV with the ATLAS [detector,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib7A866520C9DD96A9E238CD97F2E832C7s1) Phys. Rev. C 98 (4) (2018) 044905, [arXiv:1805.05220 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib7A866520C9DD96A9E238CD97F2E832C7s1)
- [22] ALICE Collaboration, S. Acharya, et al., [Measurements](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2DC065EC9BEEB2848638FA347412AAB4s1) of low- $p_T$  electrons from semileptonic [heavy-flavour](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2DC065EC9BEEB2848638FA347412AAB4s1) hadron decays at mid-rapidity in pp and Pb–Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2DC065EC9BEEB2848638FA347412AAB4s1) at  $\sqrt{s_{NN}}$  = 2.76 TeV, J. High Energy Phys. 10 (2018) 061, arXiv: [1805.04379 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2DC065EC9BEEB2848638FA347412AAB4s1)
- [23] STAR [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB0AB9630027566D290C290D24F12DD25s1) J. Adam, et al., Centrality and transverse momentum dependence of D<sup>0</sup>-meson production at [mid-rapidity](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB0AB9630027566D290C290D24F12DD25s1) in Au+Au collisions at  $\sqrt{s_{NN}}$  = 200 GeV, Phys. Rev. C 99 (3) (2019) 034908, [arXiv:1812.10224 \[nucl](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB0AB9630027566D290C290D24F12DD25s1)[ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB0AB9630027566D290C290D24F12DD25s1)
- [24] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib72B986B365F2A8F94C50F555486B6FEAs1) J. Adam, et al., Measurement of the production of high- $p_T$ electrons from [heavy-flavour](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib72B986B365F2A8F94C50F555486B6FEAs1) hadron decays in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2*.*76 TeV, Phys. Lett. B 771 (2017) 467–481, [arXiv:1609.07104 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib72B986B365F2A8F94C50F555486B6FEAs1)
- [25] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibC39B39E2120EAE119BDAC3176A8D38C4s1) S. Acharya, et al., Measurement of  $D^0$ ,  $D^+$ ,  $D^{*+}$  and  $D_s^+$ [production](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibC39B39E2120EAE119BDAC3176A8D38C4s1) in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, J. High Energy Phys. 10 (2018) 174, [arXiv:1804.09083 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibC39B39E2120EAE119BDAC3176A8D38C4s1)
- [26] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3A125FA4DE14766664380A8472C3EF59s1) A.M. Sirunyan, et al., Nuclear modification factor of  $D^0$ mesons in Pb–Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3A125FA4DE14766664380A8472C3EF59s1) at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Lett. B 782 (2018) 474–496, [arXiv:1708.04962 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib3A125FA4DE14766664380A8472C3EF59s1)
- [27] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib651EA521D2AE9B3E825E78C591903568s1) S. Acharya, et al., Measurement of electrons from semileptonic [heavy-flavour](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib651EA521D2AE9B3E825E78C591903568s1) hadron decays at midrapidity in pp and Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Lett. B 804 (2020) 135377, [arXiv:1910.09110 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib651EA521D2AE9B3E825E78C591903568s1)
- [28] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCB5610223675E3C966933A9C20A492C9s1) J. Adam, et al., Transverse momentum dependence of Dmeson [production](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCB5610223675E3C966933A9C20A492C9s1) in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV, J. High Energy Phys. 03 (2016) 081, [arXiv:1509.06888 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCB5610223675E3C966933A9C20A492C9s1)
- [29] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib88BCE0F78BB2770F93A2AF501670060Cs1) A.M. Sirunyan, et al., Measurement of the  $B^{\pm}$  meson nuclear [modification](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib88BCE0F78BB2770F93A2AF501670060Cs1) factor in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Rev. Lett. 119 (15) (2017) 152301, [arXiv:1705.04727 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib88BCE0F78BB2770F93A2AF501670060Cs1)
- [30] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib12E6E322AC9DCFCE99F288F7C985F4FEs1) A.M. Sirunyan, et al., Measurement of prompt and nonprompt charmonium suppression in [Pb–Pb collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib12E6E322AC9DCFCE99F288F7C985F4FEs1) at 5.02 TeV, Eur. Phys. J. C 78 (6) (2018) 509, [arXiv:1712.08959 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib12E6E322AC9DCFCE99F288F7C985F4FEs1)
- [31] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2C8E02C79CC148BDB92291E7912B0EBAs1) A.M. Sirunyan, et al., Studies of beauty suppression via nonprompt D<sup>0</sup> mesons in Pb–Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2C8E02C79CC148BDB92291E7912B0EBAs1) at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Rev. Lett. 123 (2) (2019) 022001, [arXiv:1810.11102 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2C8E02C79CC148BDB92291E7912B0EBAs1)
- [32] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib976CD2076F6EF18C18E45C611A7E7E68s1) B. Abelev, et al., Production of muons from heavy flavour decays at forward rapidity in pp and Pb–Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib976CD2076F6EF18C18E45C611A7E7E68s1) at  $\sqrt{s_{NN}}$  = 2.76 TeV, Phys. Rev. Lett. 109 (2012) 112301, [arXiv:1205.6443 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib976CD2076F6EF18C18E45C611A7E7E68s1)
- [33] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAC7531F039724AD81165D3694A9386FAs1) S. Acharya, et al., Production of muons from heavy-flavour hadron decays in pp [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAC7531F039724AD81165D3694A9386FAs1) at  $\sqrt{s}$  = 5.02 TeV, J. High Energy Phys. 09 (2019) 008, [arXiv:1905.07207 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAC7531F039724AD81165D3694A9386FAs1)
- [34] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib74324F3A25C36872BACFC3C10296A2D0s1) K. Aamodt, et al., The ALICE experiment at the CERN LHC, J. [Instrum.](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib74324F3A25C36872BACFC3C10296A2D0s1) 3 (2008) S08002.
- [35] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1C40B21E544639126654DEEB3C9BBDA7s1) B.B. Abelev, et al., Performance of the ALICE experiment at the CERN LHC, Int. J. Mod. Phys. A 29 (2014) 1430044, [arXiv:1402.4476 \[nucl](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1C40B21E544639126654DEEB3C9BBDA7s1)[ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1C40B21E544639126654DEEB3C9BBDA7s1)
- [36] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1A4604424520927248410E567267BB43s1) J. Adam, et al., Centrality dependence of the chargedparticle multiplicity density at [midrapidity](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1A4604424520927248410E567267BB43s1) in Pb–Pb collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Rev. Lett. 116 (22) (2016) 222302, [arXiv:1512.06104 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib1A4604424520927248410E567267BB43s1)
- [37] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAD596D16A7453F1400FD6FB1A2941CEFs1) B. Abelev, et al., Measurement of the cross section for elec[tromagnetic](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAD596D16A7453F1400FD6FB1A2941CEFs1) dissociation with neutron emission in Pb-Pb collisions at  $\sqrt{s_{NN}}$  = 2*.*76 TeV, Phys. Rev. Lett. 109 (2012) 252302, [arXiv:1203.2436 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibAD596D16A7453F1400FD6FB1A2941CEFs1)
- [38] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib6978EE5EDFD728183194F55E93435D2Bs1) P. Cortese, Performance of the ALICE Zero Degree [Calorimeters](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib6978EE5EDFD728183194F55E93435D2Bs1) and upgrade strategy, J. Phys. Conf. Ser. 1162 (1) (2019) 012006.
- [39] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib14603E84BF56F9E23284D1187DDDA4ADs1) S. Acharya, et al., Production of charged pions, kaons and [\(anti-\)protons](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib14603E84BF56F9E23284D1187DDDA4ADs1) in Pb-Pb and inelastic pp collisions at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Rev. C 101 (4) (2020) 044907, [arXiv:1910.07678 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib14603E84BF56F9E23284D1187DDDA4ADs1)
- [40] T. [Sjostrand,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8E0BC180543D5F603283642FE92FCE87s1) S. Mrenna, P.Z. Skands, PYTHIA 6.4 physics and manual, J. High Energy Phys. 05 (2006) 026, [arXiv:hep-ph/0603175 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8E0BC180543D5F603283642FE92FCE87s1)
- [41] R. Engel, J. Ranft, S. Roesler, Hard diffraction in [hadron-hadron](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib626311B09A1B8A20B3F4A9177188C309s1) interactions and in [photoproduction,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib626311B09A1B8A20B3F4A9177188C309s1) Phys. Rev. D 52 (1995) 1459–1468.
- [42] T. Sjöstrand, S. Ask, J.R. [Christiansen,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib52E0141B59A5D3E5BF7C55D1B143BBEEs1) R. Corke, N. Desai, P. Ilten, S. Mrenna, S. Prestel, C.O. Rasmussen, P.Z. Skands, An [introduction](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib52E0141B59A5D3E5BF7C55D1B143BBEEs1) to PYTHIA 8.2, Comput. Phys. Commun. 191 (2015) 159–177, [arXiv:1410.3012 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib52E0141B59A5D3E5BF7C55D1B143BBEEs1)
- [43] X.-N. Wang, M. [Gyulassy,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib903B777E79687CAE61171DDE8C682993s1) HIJING: a Monte Carlo model for multiple jet production in pp, pA and AA collisions, Phys. Rev. D 44 (1991) [3501–3516.](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib903B777E79687CAE61171DDE8C682993s1)
- [44] R. Brun, F. Bruyant, F. Carminati, S. Giani, M. Maire, A. McPherson, G. Patrick, L. Urban, GEANT Detector Description and Simulation Tool, CERN-W5013, 1994.
- [45] S. Alioli, P. Nason, C. Oleari, E. Re, NLO [vector-boson](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCE64883F412791987BB39BF543A72333s1) production matched with shower in POWHEG, J. High Energy Phys. 07 (2008) 060, [arXiv:0805.4802 \[hep](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCE64883F412791987BB39BF543A72333s1)[ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibCE64883F412791987BB39BF543A72333s1)
- [46] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0ED1ADB8780201A886502EFD628359D3s1) S. Chatrchyan, et al., Study of Z boson production in Pb– Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0ED1ADB8780201A886502EFD628359D3s1) at  $\sqrt{s_{NN}}$  = 2.76 TeV, Phys. Rev. Lett. 106 (2011) 212301, arXiv: [1102.5435 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0ED1ADB8780201A886502EFD628359D3s1)
- [47] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0CDDA586A39EFD16D2A6E088EBE7E663s1) S. Chatrchyan, et al., Study of Z production in PbPb and pp collisions at  $\sqrt{s_{NN}}$  = 2.76 TeV in the dimuon and [dielectron](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0CDDA586A39EFD16D2A6E088EBE7E663s1) decay channels, J. High Energy Phys. 03 (2015) 022, [arXiv:1410.4825 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib0CDDA586A39EFD16D2A6E088EBE7E663s1)
- [48] ATLAS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibD8BFE38074CDB6CF1A9F245230D71FB0s1) G. Aad, et al., Measurement of the production and lepton charge [asymmetry](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibD8BFE38074CDB6CF1A9F245230D71FB0s1) of *W* bosons in Pb+Pb collisions at  $\sqrt{s_{NN}} = 2.76$  TeV with the ATLAS detector, Eur. Phys. J. C 75 (1) (2015) 23, [arXiv:1408.4674 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibD8BFE38074CDB6CF1A9F245230D71FB0s1)
- [49] CMS [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBBC2FAA391E145EDE8BD15B0F3298C90s1) V. Khachatryan, et al., Study of Z boson production in p-Pb collisions at √*SNN* = 5.02 TeV, Phys. Lett. B 759 (2016) 36–57, [arXiv:1512.](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBBC2FAA391E145EDE8BD15B0F3298C90s1) [06461 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBBC2FAA391E145EDE8BD15B0F3298C90s1)
- [50] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F9E8366B0C24A1BD529E303B5C35025s1) J. Adam, et al., W and Z boson production in p–Pb collisions at √<sup>*s*<sub>NN</sub></sup> = 5.02 TeV, J. High Energy Phys. 02 (2017) 077, [arXiv:1611.03002](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F9E8366B0C24A1BD529E303B5C35025s1) [\[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F9E8366B0C24A1BD529E303B5C35025s1)
- [51] H.-L. Lai, M. Guzzi, J. Huston, Z. Li, P.M. [Nadolsky,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBA228A27BB6BDF869899D72504DFD165s1) J. Pumplin, C.P. Yuan, New parton [distributions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBA228A27BB6BDF869899D72504DFD165s1) for collider physics, Phys. Rev. D 82 (2010) 074024, arXiv: [1007.2241 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibBA228A27BB6BDF869899D72504DFD165s1)
- [52] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F2040B03A7D89B601F6407B6711974As1) J. Adam, et al., J/*ψ* suppression at forward rapidity in Pb– Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F2040B03A7D89B601F6407B6711974As1) at  $\sqrt{s_{NN}}$  = 5.02 TeV, Phys. Lett. B 766 (2017) 212–224, arXiv: [1606.08197 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib2F2040B03A7D89B601F6407B6711974As1)
- [53] ATLAS Collaboration, G. Aad, et al., Measurement of [charged-particle](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib193F442A201F6CEA7DB49457729A884Es1) spectra in Pb+Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib193F442A201F6CEA7DB49457729A884Es1) at  $\sqrt{s_{NN}}$  = 2.76 TeV with the ATLAS detector at the LHC, J. High Energy Phys. 09 (2015) 050, [arXiv:1504.04337 \[hep-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib193F442A201F6CEA7DB49457729A884Es1)
- [54] K.J. Eskola, H. [Honkanen,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib35AF3D356037E779F7734256D04ECA03s1) V.J. Kolhinen, C.A. Salgado, Global DGLAP fit analyses of the nPDF: EKS98 and HKM, [arXiv:hep-ph/0302170 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib35AF3D356037E779F7734256D04ECA03s1)
- [55] K.J. Eskola, V.J. Kolhinen, C.A. Salgado, The scale [dependent](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF6EDC1E6F049AA98CAC8E1955C24624Fs1) nuclear effects in parton [distributions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF6EDC1E6F049AA98CAC8E1955C24624Fs1) for practical applications, Eur. Phys. J. C 9 (1999) 61–68, [arXiv:hep-ph/9807297.](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF6EDC1E6F049AA98CAC8E1955C24624Fs1)
- [56] Heavy Quarks and Quarkonia Working Group, Jet Physics Working Group, PDFs, Shadowing and p+A Working Group Photons Working Group Collaboration, M.L. Mangano, H. Satz, U.A. Wiedemann, Hard Probes in Heavy Ion Collisions at the LHC: [report of the 4 working groups], CERN Yellow Reports: Monographs, CERN, Geneva, 2004, <https://cds.cern.ch/record/815037>. Report of the 4 working groups completed by Oct 2003, following 3 plenary meetings held from Oct 2001 - Oct 2002 of the CERN Theory Workshop on Hard Probes in Heavy Ion Collisions at the LHC.
- [57] M. Cacciari, M. Greco, P. Nason, The  $p_T$  spectrum in heavy flavor [hadroproduc](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib04C88DCD3769F4E18DE421CFB368A4BBs1)tion, J. High Energy Phys. 05 (1998) 007, [arXiv:hep-ph/9803400 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib04C88DCD3769F4E18DE421CFB368A4BBs1)
- [58] M. Cacciari, S. Frixione, N. Houdeau, M.L. [Mangano,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB23AFAD89920B203204DDBFB65D93035s1) P. Nason, G. Ridolfi, Theoretical [predictions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB23AFAD89920B203204DDBFB65D93035s1) for charm and bottom production at the LHC, J. High Energy Phys. 10 (2012) 137, [arXiv:1205.6344 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibB23AFAD89920B203204DDBFB65D93035s1)
- [59] M. He, R.J. Fries, R. Rapp, Heavy flavor at the large hadron [collider](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib081B1238C10A088E01CF05487B37E5C3s1) in a strong coupling approach, Phys. Lett. B 735 (2014) 445–450, [arXiv:1401.3817 \[nucl](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib081B1238C10A088E01CF05487B37E5C3s1)th<sub>1</sub>
- [60] Z.-B. Kang, F. Ringer, I. Vitev, Effective field theory [approach](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib4FCCFBD3158BFD4FB25252F9019C7251s1) to open heavy flavor [production](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib4FCCFBD3158BFD4FB25252F9019C7251s1) in heavy-ion collisions, J. High Energy Phys. 03 (2017) 146, [arXiv:1610.02043 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib4FCCFBD3158BFD4FB25252F9019C7251s1)
- [61] M. [Nahrgang,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib78B0546F32936A10FD95CB0E8D4CDA80s1) J. Aichelin, P.B. Gossiaux, K. Werner, Influence of hadronic bound states above *T<sub>c</sub>* on [heavy-quark](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib78B0546F32936A10FD95CB0E8D4CDA80s1) observables in Pb+Pb collisions at the CERN Large Hadron Collider, Phys. Rev. C 89 (1) (2014) 014905, [arXiv:1305.6544](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib78B0546F32936A10FD95CB0E8D4CDA80s1) [\[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib78B0546F32936A10FD95CB0E8D4CDA80s1)
- [62] M. Nahrgang, J. Aichelin, P.B. Gossiaux, K. Werner, Azimuthal [correlations](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib58B946E9BCF88D66CC77666B2FB618B8s1) of heavy quarks in Pb + Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib58B946E9BCF88D66CC77666B2FB618B8s1) at  $\sqrt{s}$  = 2.76 TeV at the CERN Large Hadron Collider, Phys. Rev. C 90 (2) (2014) 024907, [arXiv:1305.3823 \[hep-ph\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib58B946E9BCF88D66CC77666B2FB618B8s1)
- [63] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B3937A226FDC81D5E38C7E433256E5Es1) S. Acharya, et al., Production of muons from heavy-flavour hadron decays in p-Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B3937A226FDC81D5E38C7E433256E5Es1) at  $\sqrt{s}_{NN}$  = 5.02 TeV, Phys. Lett. B 770 (2017) 459–472, [arXiv:1702.01479 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib8B3937A226FDC81D5E38C7E433256E5Es1)
- [64] M. Djordjevic, M. Djordjevic, Predictions of [heavy-flavor](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib6FD37C71FBEC285935700E28668A9032s1) suppression at 5.1 TeV Pb+Pb [collisions](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib6FD37C71FBEC285935700E28668A9032s1) at the CERN Large Hadron Collider, Phys. Rev. C 92 (2) (2015) 024918, [arXiv:1505.04316 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib6FD37C71FBEC285935700E28668A9032s1)

### **ALICE Collaboration**

S. Acharya <sup>142</sup>, D. Adamová <sup>96</sup>, A. Adler <sup>74</sup>, J. Adolfsson <sup>81</sup>, G. Aglieri Rinella <sup>34</sup>, M. Agnello <sup>30</sup>, N. Agrawal <sup>54, 9</sup>, Z. Ahammed <sup>142</sup>, S. Ahmad <sup>16</sup>, S.U. Ahn <sup>76</sup>, Z. Akbar <sup>51</sup>, A. Akindinov <sup>93</sup>, M. Al-Turany <sup>108</sup>, D.S.D. Albuquerque <sup>123</sup>, D. Aleksandrov <sup>89</sup>, B. Alessandro <sup>59</sup>, H.M. Alfanda <sup>6</sup>, R. Alfaro Molina <sup>71</sup>, B. Ali <sup>16</sup>, Y. Ali <sup>14</sup>, A. Alici <sup>26, 9, 54</sup>, N. Alizadehvandchali <sup>126</sup>, A. Alkin <sup>34</sup>, J. Alme <sup>21</sup>, T. Alt <sup>68</sup>, L. Altenkamper <sup>21</sup>, I. Altsybeev  $^{114}$ , M.N. Anaam  $^6$ , C. Andrei  $^{48}$ , D. Andreou  $^{91}$ , A. Andronic  $^{145}$ , M. Angeletti  $^{34}$ , V. Anguelov  $^{105}$ , T. Antičić  $^{109}$ , F. Antinori  $^{57}$ , P. Antonioli  $^{54}$ , N. Apadula  $^{80}$ , L. Aphecetche  $^{116}$ , H. Appelshäuser 68, S. Arcelli 26, R. Arnaldi 59, M. Arratia 80, I.C. Arsene 20, M. Arslandok <sup>147</sup>*,*105, A. Augustinus  $^{34}$ , R. Averbeck  $^{108}$ , S. Aziz  $^{78}$ , M.D. Azmi  $^{16}$ , A. Badalà  $^{56}$ , Y.W. Baek  $^{41}$ , X. Bai  $^{108}$ , R. Bailhache  $^{68}$ , R. Bala  $^{102}$ , A. Balbino  $^{30}$ , A. Baldisseri  $^{138}$ , M. Ball $^{43}$ , D. Banerjee  $^3$ , R. Barbera  $^{27}$ , L. Barioglio <sup>25</sup>, M. Barlou <sup>85</sup>, G.G. Barnaföldi <sup>146</sup>, L.S. Barnby <sup>95</sup>, V. Barret <sup>135</sup>, C. Bartels <sup>128</sup>, K. Barth <sup>34</sup>, E. Bartsch 68, F. Baruffaldi 28, N. Bastid 135, S. Basu <sup>81</sup>*,*144, G. Batigne 116, B. Batyunya 75, D. Bauri 49, J.L. Bazo Alba <sup>113</sup>, I.G. Bearden <sup>90</sup>, C. Beattie <sup>147</sup>, I. Belikov <sup>137</sup>, A.D.C. Bell Hechavarria <sup>145</sup>, F. Bellini<sup>34</sup>, R. Bellwied 126, S. Belokurova 114, V. Belyaev 94, G. Bencedi <sup>69</sup>*,*146, S. Beole 25, A. Bercuci 48, Y. Berdnikov <sup>99</sup>, A. Berdnikova <sup>105</sup>, D. Berenyi <sup>146</sup>, D. Berzano <sup>59</sup>, M.G. Besoiu <sup>67</sup>, L. Betev <sup>34</sup>, P.P. Bhaduri <sup>142</sup>, A. Bhasin <sup>102</sup>, I.R. Bhat <sup>102</sup>, M.A. Bhat <sup>3</sup>, B. Bhattacharjee <sup>42</sup>, P. Bhattacharya <sup>23</sup>, A. Bianchi <sup>25</sup>, L. Bianchi <sup>25</sup>, N. Bianchi <sup>52</sup>, J. Bielčík <sup>37</sup>, J. Bielčíková <sup>96</sup>, A. Bilandzic <sup>106</sup>, G. Biro <sup>146</sup>, S. Biswas <sup>3</sup>, J.T. Blair <sup>120</sup>, D. Blau <sup>89</sup>, M.B. Blidaru <sup>108</sup>, C. Blume <sup>68</sup>, G. Boca <sup>140</sup>, F. Bock <sup>97</sup>, A. Bogdanov <sup>94</sup>, S. Boi  $^{23}$ , J. Bok $^{61}$ , L. Boldizsár  $^{146}$ , A. Bolozdynya  $^{94}$ , M. Bombara  $^{38}$ , G. Bonomi  $^{141}$ , H. Borel  $^{138}$ , A. Borissov <sup>82</sup>*,*94, H. Bossi 147, E. Botta 25, L. Bratrud 68, P. Braun-Munzinger 108, M. Bregant 122, M. Broz 37, G.E. Bruno <sup>107,33</sup>, M.D. Buckland <sup>128</sup>, D. Budnikov <sup>110</sup>, H. Buesching <sup>68</sup>, S. Bufalino <sup>30</sup>, O. Bugnon <sup>116</sup>, P. Buhler 115, P. Buncic 34, Z. Buthelezi <sup>72</sup>*,*132, J.B. Butt 14, S.A. Bysiak 119, D. Caffarri 91, A. Caliva 108, E. Calvo Villar <sup>113</sup>, J.M.M. Camacho <sup>121</sup>, R.S. Camacho <sup>45</sup>, P. Camerini <sup>24</sup>, A.A. Capon <sup>115</sup>, F. Carnesecchi <sup>26</sup>, R. Caron <sup>138</sup>, J. Castillo Castellanos <sup>138</sup>, A.J. Castro <sup>131</sup>, E.A.R. Casula <sup>55</sup>, F. Catalano <sup>30</sup>, C. Ceballos Sanchez <sup>75</sup>, P. Chakraborty <sup>49</sup>, S. Chandra  $^{142}$ , W. Chang  $^6$ , S. Chapeland  $^{34}$ , M. Chartier  $^{128}$ , S. Chattopadhyay  $^{142}$ , S. Chattopadhyay  $^{111}$ , A. Chauvin  $^{23}$ , C. Cheshkov  $^{136}$ , B. Cheynis  $^{136}$ , V. Chibante Barroso<sup>34</sup>, D.D. Chinellato<sup>123</sup>, S. Cho<sup>61</sup>, P. Chochula<sup>34</sup>, P. Christakoglou<sup>91</sup>, C.H. Christensen 90, P. Christiansen 81, T. Chujo 134, C. Cicalo 55, L. Cifarelli <sup>26</sup>*,*9, F. Cindolo 54, M.R. Ciupek 108, G. Clai <sup>54</sup>*,*II, J. Cleymans 125, F. Colamaria 53, J.S. Colburn 112, D. Colella 53, A. Collu 80, M. Colocci <sup>34</sup>*,*26, M. Concas <sup>59</sup>*,*III, G. Conesa Balbastre 79, Z. Conesa del Valle 78, G. Contin <sup>24</sup>*,*60, J.G. Contreras  $3^7$ , T.M. Cormier  $9^7$ , P. Cortese  $3^1$ , M.R. Cosentino  $124$ , F. Costa  $3^4$ , S. Costanza  $140$ , P. Crochet 135, E. Cuautle 69, P. Cui 6, L. Cunqueiro 97, T. Dahms 106, A. Dainese 57, F.P.A. Damas <sup>116</sup>*,*138, M.C. Danisch  $^{105}$ , A. Danu  $^{67}$ , D. Das  $^{111}$ , I. Das  $^{111}$ , P. Das  $^{87}$ , P. Das  $^3$ , S. Das  $^3$ , S. Dash  $^{49}$ , S. De  $^{87}$ , A. De Caro 29, G. de Cataldo 53, L. De Cilladi 25, J. de Cuveland 39, A. De Falco 23, D. De Gruttola <sup>29</sup>*,*9, N. De Marco  $^{59}$ , C. De Martin  $^{24}$ , S. De Pasquale  $^{29}$ , S. Deb  $^{50}$ , H.F. Degenhardt  $^{122}$ , K.R. Deja  $^{143}$ , S. Delsanto <sup>25</sup>, W. Deng <sup>6</sup>, P. Dhankher <sup>19,49</sup>, D. Di Bari <sup>33</sup>, A. Di Mauro <sup>34</sup>, R.A. Diaz <sup>7</sup>, T. Dietel <sup>125</sup>, P. Dillenseger  $^{68}$ , Y. Ding  $^6$ , R. Divià  $^{34}$ , D.U. Dixit  $^{19}$ , Ø. Djuvsland  $^{21}$ , U. Dmitrieva  $^{63}$ , J. Do  $^{61}$ , A. Dobrin  $^{67}$ , B. Dönigus 68, O. Dordic 20, A.K. Dubey 142, A. Dubla <sup>108</sup>*,*91, S. Dudi 101, M. Dukhishyam 87, P. Dupieux 135, T.M. Eder  $^{145}$ , R.J. Ehlers  $^{97}$ , V.N. Eikeland  $^{21}$ , D. Elia  $^{53}$ , B. Erazmus  $^{116}$ , F. Erhardt  $^{100}$ , A. Erokhin  $^{114}$ , M.R. Ersdal  $^{21}$ , B. Espagnon  $^{78}$ , G. Eulisse  $^{34}$ , D. Evans  $^{112}$ , S. Evdokimov  $^{92}$ , L. Fabbietti  $^{106}$ , M. Faggin  $^{28}$ , J. Faivre <sup>79</sup>, F. Fan <sup>6</sup>, A. Fantoni <sup>52</sup>, M. Fasel <sup>97</sup>, P. Fecchio <sup>30</sup>, A. Feliciello <sup>59</sup>, G. Feofilov <sup>114</sup>, A. Fernández Téllez <sup>45</sup>, A. Ferrero <sup>138</sup>, A. Ferretti <sup>25</sup>, A. Festanti <sup>34</sup>, V.J.G. Feuillard <sup>105</sup>, J. Figiel <sup>119</sup>, S. Filchagin <sup>110</sup>, D. Finogeev <sup>63</sup>, F.M. Fionda <sup>21</sup>, G. Fiorenza <sup>53</sup>, F. Flor <sup>126</sup>, A.N. Flores <sup>120</sup>, S. Foertsch <sup>72</sup>, P. Foka  $^{108}$ , S. Fokin  $^{89}$ , E. Fragiacomo  $^{60}$ , U. Fuchs  $^{34}$ , C. Furget  $^{79}$ , A. Furs  $^{63}$ , M. Fusco Girard  $^{29}$ ,

- [65] C.A.G. Prado, W.-J. Xing, S. Cao, G.-Y. Qin, X.-N. Wang, [Longitudinal](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEFA18ABF7D37CB366B80100CCBDC5085s1) dependence of open heavy flavor  $R_{AA}$  in [relativistic](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEFA18ABF7D37CB366B80100CCBDC5085s1) heavy-ion collisions, Phys. Rev. C 101 (6) (2020) 064907, [arXiv:1911.06527 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibEFA18ABF7D37CB366B80100CCBDC5085s1)
- [66] ALICE [Collaboration,](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF8CFC573E2CE9A66798BE0DB6387B684s1) J. Adam, et al., Centrality dependence of the pseudorapidity density [distribution](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF8CFC573E2CE9A66798BE0DB6387B684s1) for charged particles in Pb-Pb collisions at  $\sqrt{s_{NN}} = 5.02$ TeV, Phys. Lett. B 772 (2017) 567–577, [arXiv:1612.08966 \[nucl-ex\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bibF8CFC573E2CE9A66798BE0DB6387B684s1)
- [67] A. Beraudo, et al., Extraction of [heavy-flavor](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib064E9A5CB78F6CCF25F4624205A0223As1) transport coefficients in QCD matter, Nucl. Phys. A 979 (2018) 21–86, [arXiv:1803.03824 \[nucl-th\].](http://refhub.elsevier.com/S0370-2693(21)00498-6/bib064E9A5CB78F6CCF25F4624205A0223As1)

J.J. Gaardhøje <sup>90</sup>, M. Gagliardi <sup>25</sup>, A.M. Gago <sup>113</sup>, A. Gal <sup>137</sup>, C.D. Galvan <sup>121</sup>, P. Ganoti <sup>85</sup>, C. Garabatos <sup>108</sup>, J.R.A. Garcia $^{45}$ , E. Garcia-Solis $^{10}$ , K. Garg $^{116}$ , C. Gargiulo $^{34}$ , A. Garibli $^{88}$ , K. Garner $^{145}$ , P. Gasik $^{106}$ , E.F. Gauger  $^{120}$ , M.B. Gay Ducati  $^{70}$ , M. Germain  $^{116}$ , J. Ghosh  $^{111}$ , P. Ghosh  $^{142}$ , S.K. Ghosh  $^3$ , M. Giacalone 26, P. Gianotti 52, P. Giubellino <sup>108</sup>*,*59, P. Giubilato 28, A.M.C. Glaenzer 138, P. Glässel 105, V. Gonzalez  $^{144}$ , L.H. González-Trueba  $^{71}$ , S. Gorbunov  $^{39}$ , L. Görlich  $^{119}$ , S. Gotovac  $^{35}$ , V. Grabski  $^{71}$ , L.K. Graczykowski  $^{143}$ , K.L. Graham  $^{112}$ , L. Greiner  $^{80}$ , A. Grelli  $^{62}$ , C. Grigoras  $^{34}$ , V. Grigoriev  $^{94},$ A. Grigoryan <sup>1, I</sup>, S. Grigoryan <sup>75</sup>, O.S. Groettvik <sup>21</sup>, F. Grosa <sup>59</sup>, J.F. Grosse-Oetringhaus <sup>34</sup>, R. Grosso <sup>108</sup>, R. Guernane<sup>79</sup>, M. Guilbaud <sup>116</sup>, M. Guittiere <sup>116</sup>, K. Gulbrandsen <sup>90</sup>, T. Gunji <sup>133</sup>, A. Gupta <sup>102</sup>, R. Gupta <sup>102</sup>, I.B. Guzman <sup>45</sup>, R. Haake <sup>147</sup>, M.K. Habib <sup>108</sup>, C. Hadjidakis <sup>78</sup>, H. Hamagaki <sup>83</sup>, G. Hamar <sup>146</sup>, M. Hamid <sup>6</sup>, R. Hannigan <sup>120</sup>, M.R. Haque <sup>143,87</sup>, A. Harlenderova <sup>108</sup>, J.W. Harris <sup>147</sup>, A. Harton <sup>10</sup>, J.A. Hasenbichler 34, H. Hassan 97, D. Hatzifotiadou <sup>54</sup>*,*9, P. Hauer 43, L.B. Havener 147, S. Hayashi 133, S.T. Heckel  $^{106}$ , E. Hellbär  $^{68}$ , H. Helstrup  $^{36}$ , T. Herman  $^{37}$ , E.G. Hernandez  $^{45}$ , G. Herrera Corral  $^8$ , F. Herrmann  $^{145}$ , K.F. Hetland  $^{36}$ , H. Hillemanns  $^{34}$ , C. Hills  $^{128}$ , B. Hippolyte  $^{137}$ , B. Hohlweger  $^{106}$ , J. Honermann  $^{145}$ , G.H. Hong  $^{148}$ , D. Horak  $^{37}$ , A. Hornung  $^{68}$ , S. Hornung  $^{108}$ , R. Hosokawa  $^{15}$ , P. Hristov  $^{34}$ , C. Huang  $^{78}$ , C. Hughes  $^{131}$ , P. Huhn  $^{68}$ , T.J. Humanic  $^{98}$ , H. Hushnud  $^{111}$ , L.A. Husova  $^{145}$ , N. Hussain  $^{42}$ , D. Hutter 39, J.P. Iddon <sup>34</sup>*,*128, R. Ilkaev 110, H. Ilyas 14, M. Inaba 134, G.M. Innocenti 34, M. Ippolitov 89, A. Isakov <sup>37</sup>*,*96, M.S. Islam 111, M. Ivanov 108, V. Ivanov 99, V. Izucheev 92, B. Jacak 80, N. Jacazio <sup>34</sup>*,*54, P.M. Jacobs  $^{80}$ , S. Jadlovska  $^{118}$ , J. Jadlovsky  $^{118}$ , S. Jaelani  $^{62}$ , C. Jahnke  $^{122}$ , M.J. Jakubowska  $^{143}$ , P.M. Jakubowska  $^{143}$ , M.A. Janik 143, T. Janson 74, M. Jercic 100, O. Jevons 112, M. Jin 126, F. Jonas <sup>97</sup>*,*145, P.G. Jones 112, J. Jung 68, M. Jung  $^{68}$ , A. Jusko  $^{112}$ , P. Kalinak  $^{64}$ , A. Kalweit  $^{34}$ , V. Kaplin  $^{94}$ , S. Kar  $^6$ , A. Karasu Uysal  $^{77}$ , D. Karatovic <sup>100</sup>, O. Karavichev <sup>63</sup>, T. Karavicheva <sup>63</sup>, P. Karczmarczyk <sup>143</sup>, E. Karpechev <sup>63</sup>, A. Kazantsev <sup>89</sup>, U. Kebschull <sup>74</sup>, R. Keidel <sup>47</sup>, M. Keil <sup>34</sup>, B. Ketzer <sup>43</sup>, Z. Khabanova <sup>91</sup>, A.M. Khan <sup>6</sup>, S. Khan <sup>16</sup>, A. Khanzadeev  $^{99}$ , Y. Kharlov  $^{92}$ , A. Khatun  $^{16}$ , A. Khuntia  $^{119}$ , B. Kileng  $^{36}$ , B. Kim  $^{61}$ , B. Kim  $^{134}$ , D. Kim  $^{148}$ , D.J. Kim  $^{127}$ , E.J. Kim  $^{73}$ , H. Kim  $^{17}$ , J. Kim  $^{148}$ , J.S. Kim  $^{41}$ , J. Kim  $^{105}$ , J. Kim  $^{148}$ , J. Kim  $^{73}$ , M. Kim  $^{105}$ , S. Kim  $^{18}$ , T. Kim  $^{148}$ , T. Kim  $^{148}$ , S. Kirsch  $^{68}$ , I. Kisel  $^{39}$ , S. Kiselev  $^{93}$ , A. Kisiel  $^{143}$ , J.L. Klay  $^5$ , C. Klein  $^{68}$ , J. Klein <sup>34</sup>*,*59, S. Klein 80, C. Klein-Bösing 145, M. Kleiner 68, T. Klemenz 106, A. Kluge 34, A.G. Knospe 126, C. Kobdaj <sup>117</sup>, M.K. Köhler <sup>105</sup>, T. Kollegger <sup>108</sup>, A. Kondratyev <sup>75</sup>, N. Kondratyeva <sup>94</sup>, E. Kondratyuk <sup>92</sup>, J. Konig <sup>68</sup>, S.A. Konigstorfer <sup>106</sup>, P.J. Konopka <sup>34</sup>, G. Kornakov <sup>143</sup>, L. Koska <sup>118</sup>, O. Kovalenko <sup>86</sup>, V. Kovalenko <sup>114</sup>, M. Kowalski <sup>119</sup>, I. Králik <sup>64</sup>, A. Kravčáková <sup>38</sup>, L. Kreis <sup>108</sup>, M. Krivda <sup>112,64</sup>, F. Krizek <sup>96</sup>, K. Krizkova Gajdosova <sup>37</sup>, M. Kroesen <sup>105</sup>, M. Krüger <sup>68</sup>, E. Kryshen <sup>99</sup>, M. Krzewicki <sup>39</sup>, V. Kučera <sup>34</sup>, C. Kuhn <sup>137</sup>, P.G. Kuijer <sup>91</sup>, L. Kumar <sup>101</sup>, S. Kundu <sup>87</sup>, P. Kurashvili <sup>86</sup>, A. Kurepin <sup>63</sup>, A.B. Kurepin <sup>63</sup>, A. Kuryakin <sup>110</sup>, S. Kushpil <sup>96</sup>, J. Kvapil <sup>112</sup>, M.J. Kweon <sup>61</sup>, J.Y. Kwon <sup>61</sup>, Y. Kwon <sup>148</sup>, S.L. La Pointe <sup>39</sup>, P. La Rocca  $^{27}$ , Y.S. Lai  $^{80}$ , A. Lakrathok  $^{117}$ , M. Lamanna  $^{34}$ , R. Langoy  $^{130}$ , K. Lapidus  $^{34}$ , A. Lardeux  $^{20}$ , P. Larionov <sup>52</sup>, E. Laudi <sup>34</sup>, L. Lautner <sup>34</sup>, R. Lavicka <sup>37</sup>, T. Lazareva <sup>114</sup>, R. Lea <sup>24</sup>, J. Lee <sup>134</sup>, S. Lee <sup>148</sup>, J. Lehrbach  $^{39}$ , R.C. Lemmon  $^{95}$ , I. León Monzón  $^{121}$ , E.D. Lesser  $^{19}$ , M. Lettrich  $^{34}$ , P. Lévai  $^{146}$ , X. Li  $^{11}$ , X.L. Li<sup>6</sup>, J. Lien <sup>130</sup>, R. Lietava <sup>112</sup>, B. Lim <sup>17</sup>, S.H. Lim <sup>17</sup>, V. Lindenstruth <sup>39</sup>, A. Lindner <sup>48</sup>, C. Lippmann <sup>108</sup>, A. Liu  $^{19}$ , J. Liu  $^{128}$ , I.M. Lofnes  $^{21}$ , V. Loginov  $^{94}$ , C. Loizides  $^{97}$ , P. Loncar  $^{35}$ , J.A. Lopez  $^{105}$ , X. Lopez  $^{135}$ , E. López Torres <sup>7</sup>, J.R. Luhder  $^{145}$ , M. Lunardon  $^{28}$ , G. Luparello  $^{60}$ , Y.G. Ma  $^{40}$ , A. Maevskaya  $^{63}$ , M. Mager <sup>34</sup>, S.M. Mahmood <sup>20</sup>, T. Mahmoud <sup>43</sup>, A. Maire <sup>137</sup>, R.D. Majka <sup>147, I</sup>, M. Malaev <sup>99</sup>, Q.W. Malik <sup>20</sup>, L. Malinina <sup>75,IV</sup>, D. Mal'Kevich <sup>93</sup>, N. Mallick <sup>50</sup>, P. Malzacher <sup>108</sup>, G. Mandaglio <sup>32,56</sup>, V. Manko <sup>89</sup>, F. Manso  $^{135}$ , V. Manzari  $^{53}$ , Y. Mao  $^6$ , M. Marchisone  $^{136}$ , J. Mareš  $^{66}$ , G.V. Margagliotti  $^{24}$ , A. Margotti  $^{54}$ , A. Marín  $^{108}$ , C. Markert  $^{120}$ , M. Marquard  $^{68}$ , N.A. Martin  $^{105}$ , P. Martinengo  $^{34}$ , J.L. Martinez  $^{126}$ , M.I. Martínez <sup>45</sup>, G. Martínez García <sup>116</sup>, S. Masciocchi <sup>108</sup>, M. Masera <sup>25</sup>, A. Masoni <sup>55</sup>, L. Massacrier <sup>78</sup>, A. Mastroserio <sup>139,53</sup>, A.M. Mathis <sup>106</sup>, O. Matonoha <sup>81</sup>, P.F.T. Matuoka <sup>122</sup>, A. Matyja <sup>119</sup>, C. Mayer <sup>119</sup>, F. Mazzaschi 25, M. Mazzilli 53, M.A. Mazzoni 58, A.F. Mechler 68, F. Meddi 22, Y. Melikyan 63, A. Menchaca-Rocha <sup>71</sup>, C. Mengke <sup>6</sup>, E. Meninno <sup>115,29</sup>, A.S. Menon <sup>126</sup>, M. Meres <sup>13</sup>, S. Mhlanga <sup>125</sup>, Y. Miake 134, L. Micheletti 25, L.C. Migliorin 136, D.L. Mihaylov 106, K. Mikhaylov <sup>75</sup>*,*93, A.N. Mishra <sup>146</sup>*,*69, D. Miśkowiec <sup>108</sup>, A. Modak <sup>3</sup>, N. Mohammadi <sup>34</sup>, A.P. Mohanty <sup>62</sup>, B. Mohanty <sup>87</sup>, M. Mohisin Khan <sup>16, V</sup>, Z. Moravcova <sup>90</sup>, C. Mordasini <sup>106</sup>, D.A. Moreira De Godoy <sup>145</sup>, L.A.P. Moreno <sup>45</sup>, I. Morozov <sup>63</sup>, A. Morsch<sup>34</sup>, T. Mrnjavac<sup>34</sup>, V. Muccifora<sup>52</sup>, E. Mudnic<sup>35</sup>, D. Mühlheim <sup>145</sup>, S. Muhuri <sup>142</sup>, J.D. Mulligan 80, A. Mulliri <sup>23</sup>*,*55, M.G. Munhoz 122, R.H. Munzer 68, H. Murakami 133, S. Murray 125, L. Musa 34, J. Musinsky 64, C.J. Myers 126, J.W. Myrcha 143, B. Naik 49, R. Nair 86, B.K. Nandi 49, R. Nania <sup>54</sup>*,*9, E. Nappi  $^{53}$ , M.U. Naru  $^{14}$ , A.F. Nassirpour  $^{81}$ , C. Nattrass  $^{131}$ , R. Nayak  $^{49}$ , S. Nazarenko  $^{110}$ , A. Neagu  $^{20}$ ,

L. Nellen  $^{69}$ , S.V. Nesbo  $^{36}$ , G. Neskovic  $^{39}$ , D. Nesterov  $^{114}$ , B.S. Nielsen  $^{90}$ , S. Nikolaev  $^{89}$ , S. Nikulin  $^{89}$ , V. Nikulin 99, F. Noferini <sup>54</sup>*,*9, S. Noh 12, P. Nomokonov 75, J. Norman <sup>128</sup>*,*79, N. Novitzky 134, P. Nowakowski <sup>143</sup>, A. Nyanin <sup>89</sup>, J. Nystrand <sup>21</sup>, M. Ogino <sup>83</sup>, A. Ohlson <sup>81</sup>, J. Oleniacz <sup>143</sup>, A.C. Oliveira Da Silva <sup>131</sup>, M.H. Oliver <sup>147</sup>, B.S. Onnerstad <sup>127</sup>, C. Oppedisano <sup>59</sup>, A. Ortiz Velasquez <sup>69</sup>, T. Osako <sup>46</sup>, A. Oskarsson <sup>81</sup>, J. Otwinowski <sup>119</sup>, K. Oyama <sup>83</sup>, Y. Pachmayer <sup>105</sup>, V. Pacik <sup>90</sup>, S. Padhan <sup>49</sup>, D. Pagano <sup>141</sup>, G. Paić <sup>69</sup>, J. Pan <sup>144</sup>, S. Panebianco <sup>138</sup>, P. Pareek <sup>142</sup>, J. Park <sup>61</sup>, J.E. Parkkila <sup>127</sup>, S. Parmar  $^{101}$ , S.P. Pathak  $^{126}$ , B. Paul  $^{23}$ , J. Pazzini  $^{141}$ , H. Pei  $^6$ , T. Peitzmann  $^{62}$ , X. Peng  $^6$ , L.G. Pereira  $^{70}$ , H. Pereira Da Costa $^{138}$ , D. Peresunko $^{89}$ , G.M. Perez $^7$ , S. Perrin $^{138}$ , Y. Pestov $^4$ , V. Petráček $^{37}$ , M. Petrovici 48, R.P. Pezzi 70, S. Piano 60, M. Pikna 13, P. Pillot 116, O. Pinazza <sup>54</sup>*,*34, L. Pinsky 126, C. Pinto 27, S. Pisano <sup>9,52</sup>, M. Płoskoń <sup>80</sup>, M. Planinic <sup>100</sup>, F. Pliquett <sup>68</sup>, M.G. Poghosyan <sup>97</sup>, B. Polichtchouk <sup>92</sup>, N. Poljak <sup>100</sup>, A. Pop<sup>48</sup>, S. Porteboeuf-Houssais <sup>135</sup>, J. Porter <sup>80</sup>, V. Pozdniakov <sup>75</sup>, S.K. Prasad <sup>3</sup>, R. Preghenella <sup>54</sup>, F. Prino <sup>59</sup>, C.A. Pruneau <sup>144</sup>, I. Pshenichnov <sup>63</sup>, M. Puccio <sup>34</sup>, S. Qiu <sup>91</sup>, L. Quaglia <sup>25</sup>, R.E. Quishpe <sup>126</sup>, S. Ragoni <sup>112</sup>, J. Rak <sup>127</sup>, A. Rakotozafindrabe <sup>138</sup>, L. Ramello <sup>31</sup>, F. Rami <sup>137</sup>, S.A.R. Ramirez  $^{45}$ , R. Raniwala  $^{103}$ , S. Raniwala  $^{103}$ , S.S. Räsänen  $^{44}$ , R. Rath  $^{50}$ , I. Ravasenga  $^{91}$ , K.F. Read <sup>97,131</sup>, A.R. Redelbach <sup>39</sup>, K. Redlich <sup>86, VI</sup>, A. Rehman <sup>21</sup>, P. Reichelt <sup>68</sup>, F. Reidt <sup>34</sup>, R. Renfordt <sup>68</sup>, Z. Rescakova 38, K. Reygers 105, A. Riabov 99, V. Riabov 99, T. Richert <sup>81</sup>*,*90, M. Richter 20, P. Riedler 34, W. Riegler  $^{34}$ , F. Riggi  $^{27}$ , C. Ristea  $^{67}$ , S.P. Rode  $^{50}$ , M. Rodríguez Cahuantzi  $^{45}$ , K. Røed  $^{20}$ , R. Rogalev  $^{92}$ , E. Rogochaya 75, D. Rohr 34, D. Röhrich 21, P.F. Rojas 45, P.S. Rokita 143, F. Ronchetti 52, A. Rosano <sup>32</sup>*,*56, E.D. Rosas  $^{69}$ , A. Rossi  $^{57}$ , A. Rotondi  $^{140}$ , A. Roy  $^{50}$ , P. Roy  $^{111}$ , O.V. Rueda  $^{81}$ , R. Rui  $^{24}$ , B. Rumyantsev  $^{75}$ , A. Rustamov  $^{88}$ , E. Ryabinkin  $^{89}$ , Y. Ryabov  $^{99}$ , A. Rybicki  $^{119}$ , H. Rytkonen  $^{127}$ , O.A.M. Saarimaki  $^{44}$ , R. Sadek  $^{116}$ , S. Sadovsky  $^{92}$ , J. Saetre  $^{21}$ , K. Šafařík $^{37}$ , S.K. Saha $^{142}$ , S. Saha $^{87}$ , B. Sahoo  $^{49}$ , P. Sahoo  $^{49}$ , R. Sahoo 50, S. Sahoo 65, D. Sahu 50, P.K. Sahu 65, J. Saini 142, S. Sakai 134, S. Sambyal 102, V. Samsonov <sup>99</sup>*,*94, D. Sarkar 144, N. Sarkar 142, P. Sarma 42, V.M. Sarti 106, M.H.P. Sas <sup>147</sup>*,*62, J. Schambach <sup>97</sup>*,*120, H.S. Scheid 68, C. Schiaua  $^{48}$ , R. Schicker  $^{105}$ , A. Schmah  $^{105}$ , C. Schmidt  $^{108}$ , H.R. Schmidt  $^{104}$ , M.O. Schmidt  $^{105}$ , M. Schmidt 104, N.V. Schmidt <sup>97</sup>*,*68, A.R. Schmier 131, J. Schukraft 89, Y. Schutz 137, K. Schwarz 108, K. Schweda <sup>108</sup>, G. Scioli <sup>26</sup>, E. Scomparin <sup>59</sup>, J.E. Seger <sup>15</sup>, Y. Sekiguchi <sup>133</sup>, D. Sekihata <sup>133</sup>, I. Selyuzhenkov <sup>108,94</sup>, S. Senyukov <sup>137</sup>, J.J. Seo <sup>61</sup>, D. Serebryakov <sup>63</sup>, L. Šerkšnytė <sup>106</sup>, A. Sevcenco <sup>67</sup>, A. Shabanov  $^{63}$ , A. Shabetai  $^{116}$ , R. Shahoyan  $^{34}$ , W. Shaikh  $^{111}$ , A. Shangaraev  $^{92}$ , A. Sharma  $^{101}$ , H. Sharma <sup>119</sup>, M. Sharma <sup>102</sup>, N. Sharma <sup>101</sup>, S. Sharma <sup>102</sup>, O. Sheibani <sup>126</sup>, A.I. Sheikh <sup>142</sup>, K. Shigaki <sup>46</sup>, M. Shimomura  $^{84}$ , S. Shirinkin  $^{93}$ , Q. Shou  $^{40}$ , Y. Sibiriak  $^{89}$ , S. Siddhanta  $^{55}$ , T. Siemiarczuk  $^{86}$ , D. Silvermyr <sup>81</sup>, G. Simatovic <sup>91</sup>, G. Simonetti <sup>34</sup>, B. Singh <sup>106</sup>, R. Singh <sup>87</sup>, R. Singh <sup>102</sup>, R. Singh <sup>50</sup>, V.K. Singh  $^{142}$ , V. Singhal  $^{142}$ , T. Sinha  $^{111}$ , B. Sitar  $^{13}$ , M. Sitta  $^{31}$ , T.B. Skaali  $^{20}$ , M. Slupecki  $^{44}$ , N. Smirnov  $^{147}$ , R.J.M. Snellings  $^{62}$ , C. Soncco  $^{113}$ , J. Song  $^{126}$ , A. Songmoolnak  $^{117}$ , F. Soramel  $^{28}$ , S. Sorensen <sup>131</sup>, I. Sputowska <sup>119</sup>, J. Stachel <sup>105</sup>, I. Stan <sup>67</sup>, P.J. Steffanic <sup>131</sup>, S.F. Stiefelmaier <sup>105</sup>, D. Stocco  $^{116}$ , M.M. Storetvedt  $^{36}$ , L.D. Stritto  $^{29}$ , C.P. Stylianidis  $^{91}$ , A.A.P. Suaide  $^{122}$ , T. Sugitate  $^{46}$ , C. Suire <sup>78</sup>, M. Suljic <sup>34</sup>, R. Sultanov <sup>93</sup>, M. Šumbera <sup>96</sup>, V. Sumberia <sup>102</sup>, S. Sumowidagdo <sup>51</sup>, S. Swain <sup>65</sup>, A. Szabo <sup>13</sup>, I. Szarka <sup>13</sup>, U. Tabassam <sup>14</sup>, S.F. Taghavi <sup>106</sup>, G. Taillepied <sup>135</sup>, J. Takahashi <sup>123</sup>, G.J. Tambave <sup>21</sup>, S. Tang <sup>135,6</sup>, Z. Tang <sup>129</sup>, M. Tarhini <sup>116</sup>, M.G. Tarzila <sup>48</sup>, A. Tauro <sup>34</sup>, G. Tejeda Muñoz <sup>45</sup>, A. Telesca <sup>34</sup>, L. Terlizzi  $^{25}$ , C. Terrevoli  $^{126}$ , S. Thakur  $^{142}$ , D. Thomas  $^{120}$ , F. Thoresen  $^{90}$ , R. Tieulent  $^{136}$ , A. Tikhonov  $^{63}$ , A.R. Timmins <sup>126</sup>, M. Tkacik <sup>118</sup>, A. Toia <sup>68</sup>, N. Topilskaya <sup>63</sup>, M. Toppi <sup>52</sup>, F. Torales-Acosta <sup>19</sup>, S.R. Torres <sup>37</sup>*,*8, A. Trifiró <sup>32</sup>*,*56, S. Tripathy 69, T. Tripathy 49, S. Trogolo 28, G. Trombetta 33, L. Tropp 38, V. Trubnikov <sup>2</sup>, W.H. Trzaska  $^{127}$ , T.P. Trzcinski  $^{143}$ , B.A. Trzeciak  $^{62}$ , A. Tumkin  $^{110}$ , R. Turrisi  $^{57}$ , T.S. Tveter <sup>20</sup>, K. Ullaland <sup>21</sup>, E.N. Umaka <sup>126</sup>, A. Uras <sup>136</sup>, G.L. Usai <sup>23</sup>, M. Vala <sup>38</sup>, N. Valle <sup>140</sup>, S. Vallero <sup>59</sup>, N. van der Kolk $^{62}$ , L.V.R. van Doremalen $^{62}$ , M. van Leeuwen $^{62}$ , P. Vande Vyvre $^{34}$ , D. Varga $^{146}$ , Z. Varga <sup>146</sup>, M. Varga-Kofarago <sup>146</sup>, A. Vargas <sup>45</sup>, M. Vasileiou <sup>85</sup>, A. Vasiliev <sup>89</sup>, O. Vázquez Doce <sup>106</sup>, V. Vechernin  $^{114}$ , E. Vercellin  $^{25}$ , S. Vergara Limón  $^{45}$ , L. Vermunt  $^{62}$ , R. Vértesi  $^{146}$ , M. Verweij  $^{62}$ , L. Vickovic <sup>35</sup>, Z. Vilakazi <sup>132</sup>, O. Villalobos Baillie <sup>112</sup>, G. Vino <sup>53</sup>, A. Vinogradov <sup>89</sup>, T. Virgili <sup>29</sup> V. Vislavicius  $^{90}$ , A. Vodopyanov  $^{75}$ , B. Volkel  $^{34}$ , M.A. Völkl  $^{104}$ , K. Voloshin  $^{93}$ , S.A. Voloshin  $^{144},$ G. Volpe $^{33}$ , B. von Haller $^{34}$ , I. Vorobyev  $^{106}$ , D. Voscek $^{118}$ , J. Vrláková $^{38}$ , B. Wagner $^{21}$ , M. Weber $^{115}$ , S.G. Weber  $^{145}$ , A. Wegrzynek  $^{34}$ , S.C. Wenzel  $^{34}$ , J.P. Wessels  $^{145}$ , J. Wiechula  $^{68}$ , J. Wikne  $^{20}$ , G. Wilk  $^{86}$ , J. Wilkinson <sup>108</sup>*,*9, G.A. Willems 145, E. Willsher 112, B. Windelband 105, M. Winn 138, W.E. Witt 131, J.R. Wright 120, Y. Wu 129, R. Xu 6, S. Yalcin 77, Y. Yamaguchi 46, K. Yamakawa 46, S. Yang 21, S. Yano <sup>46</sup>*,*138, Z. Yin  $^6$ , H. Yokoyama  $^{62}$ , I.-K. Yoo  $^{17}$ , J.H. Yoon  $^{61}$ , S. Yuan  $^{21}$ , A. Yuncu  $^{105}$ , V. Yurchenko  $^2$ , V. Zaccolo  $^{24}$ ,

A. Zaman <sup>14</sup>, C. Zampolli <sup>34</sup>, H.J.C. Zanoli <sup>62</sup>, N. Zardoshti <sup>34</sup>, A. Zarochentsev <sup>114</sup>, P. Závada <sup>66</sup>,

N. Zaviyalov  $^{110}$ , H. Zbroszczyk  $^{143}$ , M. Zhalov  $^{99}$ , S. Zhang  $^{40}$ , X. Zhang  $^6$ , Y. Zhang  $^{129}$ , Z. Zhang  $^6$ ,

V. Zherebchevskii 114, Y. Zhi 11, D. Zhou 6, Y. Zhou 90, J. Zhu <sup>6</sup>*,*108, Y. Zhu 6, A. Zichichi <sup>9</sup>*,*26, G. Zinovjev 2,

- *A.I. Alikhanyan National Science Laboratory (Yerevan Physics Institute) Foundation, Yerevan, Armenia*
- *Bogolyubov Institute for Theoretical Physics, National Academy of Sciences of Ukraine, Kiev, Ukraine*
- <sup>3</sup> Bose Institute, Department of Physics and Centre for Astroparticle Physics and Space Science (CAPSS), Kolkata, India
- *Budker Institute for Nuclear Physics, Novosibirsk, Russia*
- *California Polytechnic State University, San Luis Obispo, CA, United States*
- *Central China Normal University, Wuhan, China*
- *Centro de Aplicaciones Tecnológicas y Desarrollo Nuclear (CEADEN), Havana, Cuba*
- *Centro de Investigación y de Estudios Avanzados (CINVESTAV), Mexico City and Mérida, Mexico*
- *Centro Fermi – Museo Storico della Fisica e Centro Studi e Ricerche "Enrico Fermi', Rome, Italy*
- *Chicago State University, Chicago, IL, United States*
- *China Institute of Atomic Energy, Beijing, China*
- *Chungbuk National University, Cheongju, Republic of Korea*
- *Comenius University Bratislava, Faculty of Mathematics, Physics and Informatics, Bratislava, Slovakia*
- *COMSATS University Islamabad, Islamabad, Pakistan*
- *Creighton University, Omaha, NE, United States*
- *Department of Physics, Aligarh Muslim University, Aligarh, India*
- *Department of Physics, Pusan National University, Pusan, Republic of Korea*
- *Department of Physics, Sejong University, Seoul, Republic of Korea*
- *Department of Physics, University of California, Berkeley, CA, United States*
- *Department of Physics, University of Oslo, Oslo, Norway*
- *Department of Physics and Technology, University of Bergen, Bergen, Norway*
- *Dipartimento di Fisica dell'Università 'La Sapienza' and Sezione INFN, Rome, Italy*
- *Dipartimento di Fisica dell'Università and Sezione INFN, Cagliari, Italy*
- *Dipartimento di Fisica dell'Università and Sezione INFN, Trieste, Italy*
- *Dipartimento di Fisica dell'Università and Sezione INFN, Turin, Italy*
- *Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Bologna, Italy*
- *Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Catania, Italy*
- *Dipartimento di Fisica e Astronomia dell'Università and Sezione INFN, Padova, Italy*
- *Dipartimento di Fisica 'E.R. Caianiello' dell'Università and Gruppo Collegato INFN, Salerno, Italy*
- *Dipartimento DISAT del Politecnico and Sezione INFN, Turin, Italy*
- <sup>31</sup> Dipartimento di Scienze e Innovazione Tecnologica dell'Università del Piemonte Orientale and INFN Sezione di Torino, Alessandria, Italy
- *Dipartimento di Scienze MIFT, Università di Messina, Messina, Italy*
- *Dipartimento Interateneo di Fisica 'M. Merlin' and Sezione INFN, Bari, Italy*
- *European Organization for Nuclear Research (CERN), Geneva, Switzerland*
- *Faculty of Electrical Engineering, Mechanical Engineering and Naval Architecture, University of Split, Split, Croatia*
- *Faculty of Engineering and Science, Western Norway University of Applied Sciences, Bergen, Norway*
- *Faculty of Nuclear Sciences and Physical Engineering, Czech Technical University in Prague, Prague, Czech Republic*
- *Faculty of Science, P.J. Šafárik University, Košice, Slovakia*
- *Frankfurt Institute for Advanced Studies, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany*
- *Fudan University, Shanghai, China*
- *Gangneung-Wonju National University, Gangneung, Republic of Korea*
- *Gauhati University, Department of Physics, Guwahati, India*
- *Helmholtz-Institut für Strahlen- und Kernphysik, Rheinische Friedrich-Wilhelms-Universität Bonn, Bonn, Germany*
- *Helsinki Institute of Physics (HIP), Helsinki, Finland*
- *High Energy Physics Group, Universidad Autónoma de Puebla, Puebla, Mexico*
- *Hiroshima University, Hiroshima, Japan*
- *Hochschule Worms, Zentrum für Technologietransfer und Telekommunikation (ZTT), Worms, Germany*
- *Horia Hulubei National Institute of Physics and Nuclear Engineering, Bucharest, Romania*
- *Indian Institute of Technology Bombay (IIT), Mumbai, India*
- *Indian Institute of Technology Indore, Indore, India*
- *Indonesian Institute of Sciences, Jakarta, Indonesia*
- *INFN, Laboratori Nazionali di Frascati, Frascati, Italy*
- *INFN, Sezione di Bari, Bari, Italy*
- *INFN, Sezione di Bologna, Bologna, Italy*
- *INFN, Sezione di Cagliari, Cagliari, Italy*
- *INFN, Sezione di Catania, Catania, Italy*
- *INFN, Sezione di Padova, Padova, Italy*
- *INFN, Sezione di Roma, Rome, Italy*
- *INFN, Sezione di Torino, Turin, Italy*
- *INFN, Sezione di Trieste, Trieste, Italy*
- *Inha University, Incheon, Republic of Korea*
- *Institute for Gravitational and Subatomic Physics (GRASP), Utrecht University/Nikhef, Utrecht, Netherlands*
- *Institute for Nuclear Research, Academy of Sciences, Moscow, Russia*
- *Institute of Experimental Physics, Slovak Academy of Sciences, Košice, Slovakia*
- *Institute of Physics, Homi Bhabha National Institute, Bhubaneswar, India*
- *Institute of Physics of the Czech Academy of Sciences, Prague, Czech Republic*
- *Institute of Space Science (ISS), Bucharest, Romania*
- *Institut für Kernphysik, Johann Wolfgang Goethe-Universität Frankfurt, Frankfurt, Germany*
- *Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México, Mexico City, Mexico*
- *Instituto de Física, Universidade Federal do Rio Grande do Sul (UFRGS), Porto Alegre, Brazil*
- *Instituto de Física, Universidad Nacional Autónoma de México, Mexico City, Mexico*
- *iThemba LABS, National Research Foundation, Somerset West, South Africa*
- *Jeonbuk National University, Jeonju, Republic of Korea*

N. Zurlo <sup>141</sup>

*Johann-Wolfgang-Goethe Universität Frankfurt Institut für Informatik, Fachbereich Informatik und Mathematik, Frankfurt, Germany*

- *Joint Institute for Nuclear Research (JINR), Dubna, Russia*
- *Korea Institute of Science and Technology Information, Daejeon, Republic of Korea*
- *KTO Karatay University, Konya, Turkey*
- *Laboratoire de Physique des 2 Infinis, Irène Joliot-Curie, Orsay, France*
- *Laboratoire de Physique Subatomique et de Cosmologie, Université Grenoble-Alpes, CNRS-IN2P3, Grenoble, France*
- *Lawrence Berkeley National Laboratory, Berkeley, CA, United States*
- *Lund University Department of Physics, Division of Particle Physics, Lund, Sweden*
- *Moscow Institute for Physics and Technology, Moscow, Russia*
- *Nagasaki Institute of Applied Science, Nagasaki, Japan*
- *Nara Women's University (NWU), Nara, Japan*
- *National and Kapodistrian University of Athens, School of Science, Department of Physics, Athens, Greece*
- *National Centre for Nuclear Research, Warsaw, Poland*
- *National Institute of Science Education and Research, Homi Bhabha National Institute, Jatni, India*
- *National Nuclear Research Center, Baku, Azerbaijan*
- *National Research Centre Kurchatov Institute, Moscow, Russia*
- *Niels Bohr Institute, University of Copenhagen, Copenhagen, Denmark*
- *Nikhef, National institute for subatomic physics, Amsterdam, Netherlands*
- *NRC Kurchatov Institute IHEP, Protvino, Russia*
- *NRC «Kurchatov» Institute – ITEP, Moscow, Russia*
- *NRNU Moscow Engineering Physics Institute, Moscow, Russia*
- *Nuclear Physics Group, STFC Daresbury Laboratory, Daresbury, United Kingdom*
- *Nuclear Physics Institute of the Czech Academy of Sciences, Rež ˇ u Prahy, Czech Republic*
- *Oak Ridge National Laboratory, Oak Ridge, TN, United States*
- *Ohio State University, Columbus, OH, United States*
- *Petersburg Nuclear Physics Institute, Gatchina, Russia*
- *Physics department, Faculty of science, University of Zagreb, Zagreb, Croatia*
- *Physics Department, Panjab University, Chandigarh, India*
- *Physics Department, University of Jammu, Jammu, India*
- *Physics Department, University of Rajasthan, Jaipur, India*
- *Physikalisches Institut, Eberhard-Karls-Universität Tübingen, Tübingen, Germany*
- *Physikalisches Institut, Ruprecht-Karls-Universität Heidelberg, Heidelberg, Germany*
- *Physik Department, Technische Universität München, Munich, Germany*
- *Politecnico di Bari and Sezione INFN, Bari, Italy*
- <sup>108</sup> Research Division and ExtreMe Matter Institute EMMI, GSI Helmholtzzentrum für Schwerionenforschung GmbH, Darmstadt, Germany
- *Rudjer Boškovi´c Institute, Zagreb, Croatia*
- *Russian Federal Nuclear Center (VNIIEF), Sarov, Russia*
- *Saha Institute of Nuclear Physics, Homi Bhabha National Institute, Kolkata, India*
- *School of Physics and Astronomy, University of Birmingham, Birmingham, United Kingdom*
- *Sección Física, Departamento de Ciencias, Pontificia Universidad Católica del Perú, Lima, Peru*
- *St. Petersburg State University, St. Petersburg, Russia*
- *Stefan Meyer Institut für Subatomare Physik (SMI), Vienna, Austria*
- *SUBATECH, IMT Atlantique, Université de Nantes, CNRS-IN2P3, Nantes, France*
- *Suranaree University of Technology, Nakhon Ratchasima, Thailand*
- *Technical University of Košice, Košice, Slovakia*
- *The Henryk Niewodniczanski Institute of Nuclear Physics, Polish Academy of Sciences, Cracow, Poland*
- *The University of Texas at Austin, Austin, TX, United States*
- *Universidad Autónoma de Sinaloa, Culiacán, Mexico*
- *Universidade de São Paulo (USP), São Paulo, Brazil*
- *Universidade Estadual de Campinas (UNICAMP), Campinas, Brazil*
- *Universidade Federal do ABC, Santo Andre, Brazil*
- *University of Cape Town, Cape Town, South Africa*
- *University of Houston, Houston, TX, United States*
- *University of Jyväskylä, Jyväskylä, Finland*
- *University of Liverpool, Liverpool, United Kingdom*
- *University of Science and Technology of China, Hefei, China*
- *University of South-Eastern Norway, Tonsberg, Norway*
- *University of Tennessee, Knoxville, TN, United States*
- *University of the Witwatersrand, Johannesburg, South Africa*
- *University of Tokyo, Tokyo, Japan*
- *University of Tsukuba, Tsukuba, Japan*
- *Université Clermont Auvergne, CNRS/IN2P3, LPC, Clermont-Ferrand, France*
- *Université de Lyon, Université Lyon 1, CNRS/IN2P3, IPN-Lyon, Villeurbanne, Lyon, France*
- *Université de Strasbourg, CNRS, IPHC UMR 7178, F-67000 Strasbourg, France*
- 138 Université Paris-Saclay Centre d'Etudes de Saclay (CEA), IRFU, Départment de Physique Nucléaire (DPhN), Saclay, France
- *Università degli Studi di Foggia, Foggia, Italy*
- *Università degli Studi di Pavia and Sezione INFN, Pavia, Italy*
- *Università di Brescia and Sezione INFN, Brescia, Italy*
- *Variable Energy Cyclotron Centre, Homi Bhabha National Institute, Kolkata, India*
- *Warsaw University of Technology, Warsaw, Poland*
- *Wayne State University, Detroit, MI, United States*
- *Westfälische Wilhelms-Universität Münster, Institut für Kernphysik, Münster, Germany*
- *Wigner Research Centre for Physics, Budapest, Hungary*
- *Yale University, New Haven, CT, United States*
- *Yonsei University, Seoul, Republic of Korea*
- <sup>I</sup> Deceased.
- II Also at: Italian National Agency for New Technologies, Energy and Sustainable Economic Development (ENEA), Bologna, Italy.
- III Also at: Dipartimento DET del Politecnico di Torino, Turin, Italy.
- IV Also at: M.V. Lomonosov Moscow State University, D.V. Skobeltsyn Institute of Nuclear, Physics, Moscow, Russia.
- $^{\vee}$  Also at: Department of Applied Physics, Aligarh Muslim University, Aligarh, India.
- <sup>VI</sup> Also at: Institute of Theoretical Physics, University of Wroclaw, Poland.