

Short-Term sequelae of Multisystem Inflammatory Syndrome in Children Assessed by CMR

*Original*

Short-Term sequelae of Multisystem Inflammatory Syndrome in Children Assessed by CMR / Bermejo, Ivan Altamar; Bautista-Rodriguez, Carles; Fraisse, Alain; Voges, Inga; Gatehouse, Peter; Kang, Heechan; Piccinelli, Enrico; Rowlinson, Giselle; Lane, Mary; Semple, Thomas; Moscatelli, Sara; Dwornik, Maria; Lota, Amrit; Di Salvo, Giovanni; Wage, Ricardo; Prasad, Sanjay K; Mohiaddin, Raad; Pennell, Dudley J; Thavendiranathan, Paladinesh; Krupickova, Sylvia. - In: JACC. CARDIOVASCULAR IMAGING. - ISSN 1936-878X. - 14:8(2021), pp. 1666-1667.

[10.1016/j.jcmg.2021.01.035]

*Availability:*

This version is available at: 11583/2970578 since: 2022-08-10T15:21:16Z

*Publisher:*

ELSEVIER SCIENCE INC

*Published*

DOI:10.1016/j.jcmg.2021.01.035

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Elsevier postprint/Author's Accepted Manuscript

© 2021. This manuscript version is made available under the CC-BY-NC-ND 4.0 license

<http://creativecommons.org/licenses/by-nc-nd/4.0/>. The final authenticated version is available online at:

<http://dx.doi.org/10.1016/j.jcmg.2021.01.035>

(Article begins on next page)



### Short-Term sequelae of Multisystem Inflammatory Syndrome in Children Assessed by CMR



Recent studies have reported a wide clinical spectrum of multisystem inflammatory syndrome in children (MIS-C) (1-3). This is the first study that describes short-term sequelae of the acute phase of MIS-C assessed by cardiac magnetic resonance (CMR) using a comprehensive imaging protocol.

A total of 44 children were referred with MIS-C (fulfilling the criteria of the U.S. Centers for Disease Control and Prevention) between April and June 2020 to our center (Royal Brompton Hospital, London, United Kingdom). A total of 20 patients were referred for CMR scanning because of myocardial involvement in the acute phase: 1) 10 (50%) patients had myocardial dysfunction (left ventricular ejection fraction  $\leq 55\%$ ); 2) 5 (25%) patients had dilated coronary arteries (Z-score  $>2.5$ ); and 3) 5 (25%) patients had increased cardiac inflammatory markers (B-type natriuretic peptide and troponin). The study was approved by the local Research Ethics Committee. The median age of the patients was 8 years (range 17 months to 14 years).

Patients under general anesthesia were scanned with a 3.0-T scanner, and the other patients were scanned with 1.5-T scanner (Vida and Aera/Avanto-FIT, Siemens, Erlangen, Germany).

Native  $T_1$  mapping was performed using a 3(3)5 modified Look-Locker imaging single-shot balanced steady-state free precession (bSSFP) sequence at the basal and midventricular levels.  $T_2$  mapping was also conducted on the basis of a single-shot sequence (1.5-T bSSFP, 3.0-T spoiled gradient recalled echo) measured at the basal, midventricular, and apical levels.

As required by the Society for Cardiovascular Magnetic Resonance consensus statements, normal-range data are available for all users of mapping at the Brompton Hospital. These data had been obtained and shared from healthy young and middle-aged individuals who were scanned in the 1.5-T (VE11C, Siemens Healthineers, Erlangen, Germany) and 3.0-T (VA11A, Siemens Healthineers) scanners, and the maximum value from all of them (approximately 20

healthy control subjects for each  $T_1$  and  $T_2$  mapping per scanner) was taken as a cutoff value to avoid overestimating the results.

Mean global  $T_1$  and  $T_2$  values for each slice and mean segmental  $T_1$  and  $T_2$  values were measured. The native  $T_1$  normal range was 975 to 1,065 ms at 1.5-T and 1,140 to 1,395 ms at a 3.0-T scanner.  $T_2$  values were considered abnormal if they were  $>60$  ms at 1.5-T and  $>49$  ms at a 3.0-T scanner. The same range was used for mean global and mean segmental values.

The results of a polymerase chain reaction nasal swab for coronavirus disease-2019 (COVID-19) were negative in the acute phase in all patients. Results of COVID-19 serological testing at the time of admission were positive in 14 (70%) patients for immunoglobulin G severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2).

The time interval between the onset of symptoms and the date of CMR was between 12 and 72 days ( $27 \pm 14$  days). Six (30%) patients required general anesthesia. Repeated echocardiogram at the time of CMR showed normal systolic function and normalized size of the coronary arteries in all but 1 patient.

All CMR conventional parameters were in the normal range (Table 1). Only 1 patient had mildly reduced left ventricular ejection fraction. Short tau inversion recovery images were negative in all patients. Two patients had small areas of late gadolinium enhancement.

Mean  $T_1$  values were normal in all but 1 patient. This patient had a slightly increased mean  $T_1$  value at the midventricular level of 1,068 ms. Altogether, 5 patients had abnormal segmental values in 1 to 5 segments. The patient with the increased  $T_1$  values in 5 segments presented in shock with moderate systolic dysfunction. Abnormally increased  $T_1$  values were associated with high C-reactive protein at the time of admission ( $p < 0.001$ ).

Mean global  $T_2$  values of basal, midventricular, and apical slices were normal in all patients. However, 1 patient showed higher  $T_2$  values in the apical lateral segment. Both basal septal values were abnormal in another patient. Abnormal  $T_2$  values were associated with elevated D-dimer levels at the time of admission ( $p = 0.042$ ) and abnormal mitral annular planar systolic excursion (MAPSE) ( $p < 0.001$ ). The number of segments with abnormal  $T_2$  values was associated with C-reactive protein at the time of admission ( $p < 0.001$ ) and with MAPSE ( $p = 0.011$ ). (The Mann-Whitney  $U$ -test and the Kruskal-Wallis test were used as appropriate.) Three patients had dilated neck

**TABLE 1 CMR Results of the Conventional and Parametric Data**

CMR Parameter	N = 20	CMR Parameter	N = 20
LV EDV index, mL/m <sup>2</sup>	67 ± 12	RV EDV index, mL/m <sup>2</sup>	67 ± 11
LV ESV index, mL/m <sup>2</sup>	24 ± 6	RV ESV index, mL/m <sup>2</sup>	26 ± 6
LV SV index, mL/m <sup>2</sup>	44 ± 10	RV SV index, mL/m <sup>2</sup>	42 ± 7
LV EF, %	64 ± 5	RV EF, %	62 ± 5
LV mass index, g/m <sup>2</sup>	45 ± 8	—	—
LA index, cm <sup>2</sup> /m <sup>2</sup>	10 ± 2	RA index, cm <sup>2</sup> /m <sup>2</sup>	12 ± 2
MAPSE lateral, mm	13 ± 3	TAPSE, mm	20 ± 4
CMR Parameter (1.5-T Scanner)	N = 14	CMR Parameter (3.0-T Scanner)	N = 6
Native T <sub>1</sub> at basal level, ms	1,016 ± 29	Native T <sub>1</sub> at basal level, ms	1,298 ± 34
Native T <sub>1</sub> at midventricular level, ms	1,008 ± 33	Native T <sub>1</sub> at midventricular level, ms	1,315 ± 35
Native T <sub>2</sub> at basal level, ms	45 ± 2	Native T <sub>2</sub> at basal level, ms	41 ± 3
Native T <sub>2</sub> at midventricular level, ms	46 ± 2	Native T <sub>2</sub> at midventricular level, ms	38 ± 2
Native T <sub>2</sub> at apical level, ms	49 ± 4	Native T <sub>2</sub> at apical level, ms	39 ± 3

Values are mean ± SD.

CMR = cardiac magnetic resonance; EDV = end-diastolic volume; EF = ejection fraction; ESV = end-systolic volume; LA = left atrial; LGE = late gadolinium enhancement; LV = left ventricular; MAPSE = mitral annular planar systolic excursion; RA = right atrial; RV = right ventricular; SV = systolic volume; TAPSE = tricuspid annular planar systolic excursion

and/or innominate veins. A splenic infarct was demonstrated in 1 of the most severely affected patients.

This study demonstrates that despite the initial life-threatening presentation of some children with MIS-C, there is rapid improvement and nearly complete normalization of all cardiac abnormalities within a few weeks after the onset of the disease. Subtle residual cardiac and extracardiac abnormalities were found in a subset of children, who will require follow-up studies.

Limitations include the large spread of patients' ages, caused by the nature of the disease and the wide range of time intervals from the onset of symptoms to CMR, that makes the interpretation of the results challenging. Statistical analysis of T<sub>1</sub> mapping is further limited by different field strengths used for the scanning. Extracellular volume, another parameter used for tissue characterization of acute myocarditis, was not used in this study because of a lack of institutional normal values.

Ivan Altamar Bermejo, MD  
Carles Bautista-Rodriguez, MD, PhD  
Alain Fraisse, MD, PhD  
Inga Voges, MD  
Peter Gatehouse, PhD  
Heechan Kang, MD  
Enrico Piccinelli, MD  
Giselle Rowlinson, PhD  
Mary Lane, MD  
Thomas Semple, MD  
Sara Moscatelli, MD

Maria Dwornik, MD  
Amrit Lota, PhD  
Giovanni Di Salvo, MD, PhD  
Ricardo Wage, DCR R, FSCMR  
Sanjay K. Prasad, PhD  
Raad Mohiaddin, PhD  
Dudley J. Pennell, PhD  
Sylvia Krupickova, MD, PhD\*

\*Department of Pediatric Cardiology

Royal Brompton Hospital  
Sydney Street  
London SW3 6NP  
United Kingdom

E-mail: [s.krupickova@rbht.nhs.uk](mailto:s.krupickova@rbht.nhs.uk)

<https://doi.org/10.1016/j.jcmg.2021.01.035>

© 2021 the American College of Cardiology Foundation. Published by Elsevier. All rights reserved.

The authors have reported that they have no relationships relevant to the contents of this paper to disclose. Drs. Bermejo and Bautista-Rodriguez are joint first authors.

Paladinesh Thavendiranathan, MD, served as Guest Editor for this paper.

The authors attest they are in compliance with human studies committees and animal welfare regulations of the authors' institutions and Food and Drug Administration guidelines, including patient consent where appropriate. For more information, visit the [Author Center](#).

## REFERENCES

1. Dufort EM, Koumans EH, Chow EJ, et al. Multisystem inflammatory syndrome in children in New York State. *N Engl J Med*. 2020;383:347-358.
2. Feldstein LR, Rose EB, Horwitz SM, et al. Multisystem inflammatory syndrome in U.S. children and adolescents. *N Engl J Med*. 2020;383:334-346.
3. Bautista-Rodriguez C, Sanchez-de-Toledo J, Clark BC, et al. Multisystem inflammatory syndrome in children: an international survey. *Pediatrics*. 2021;147:e2020024554.