

Modification and characterization of clinoptilolite for the co-immobilization of formate dehydrogenase and glycerol dehydrogenase enzymes

Original

Modification and characterization of clinoptilolite for the co-immobilization of formate dehydrogenase and glycerol dehydrogenase enzymes / Cocuzza, Clarissa; Pietricola, Giuseppe; Cauda, VALENTINA ALICE; Dosa, Melodj; Fino, Debora; Carminna, Ottone; Romero, Oscar; Tommasi, Tonia; Piumetti, Marco. - ELETTRONICO. - (2021), p. 27. (Intervento presentato al convegno XV Italian Congress of Zeolites Science and Technology (AIZ2021) tenutosi a Napoli nel 9th - 11th of December 2021).

Availability:

This version is available at: 11583/2971304 since: 2022-09-14T15:49:48Z

Publisher:

Associazione italiana zeoliti

Published

DOI:

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

P3

MODIFICATION AND CHARACTERIZATION OF CLINOPTILOLITE FOR THE CO-IMMOBILIZATION OF FORMATE DEHYDROGENASE AND GLYCEROL DEHYDROGENASE ENZYMES

Clarissa Cocuzza^a, Giuseppe Pietricola^a, Valentina Cauda^a, Melodi Dosa^a, Debora Fino^a,
Carminna Ottone^b, Oscar Romero^c, Tonia Tommasi^a, Marco Piumetti^a

^aDepartment of Applied Science and Technology, Corso Duca degli Abruzzi 24, Politecnico di Torino, I-10129 Turin, Italy *) marco.piumetti@polito.it

^bEscuela de Ingeniería Bioquímica, Pontificia Universidad Católica de Valparaíso, Av. Brasil 2085, Valparaíso, Chile

^cBioprocess Engineering and Applied Biocatalysis Group, Department of Chemical, Biological and Environmental Engineering, Universitat Autònoma de Barcelona, 08193

In the last decades there is a rising concern for the increasing concentration of carbon dioxide, considered the major responsible of Global Warming. A solution to this critical issue is the catalytic conversion of CO₂ into high value-added products. Among the different strategies that could be applied, the enzymatic process of CO₂ reduction to methanol, employing a sequence of three enzyme-catalyzed reactions¹, seems to be very promising. The simultaneous employment of formate dehydrogenase and glycerol dehydrogenase allows to reduce CO₂ to formic acid, the first of the sequential reactions, and at the same time regenerate the nicotinamide cofactor, that is very expensive. To reuse enzymes, with a consequential reduction of cost, and increased their stability, they can be immobilized on a proper support. In this context, porous materials, such as zeolites, present appropriate features to be suitable for enzymes immobilization. In particular, they are well suited for the covalent immobilization technique due to the fact that they can be functionalized with different functional groups². Natural zeolites, like Clinoptilolite have the advantage to be low-cost materials largely diffused in different part of the world. Clinoptilolite was subjected to dealumination-desilication treatments to modify the zeolite's morphology, increasing its specific surface area. According to the literature, the dealumination procedure was done with sequential acid attacks using HCl solutions. Instead, for the subsequent desilication process NaOH solution is required³. The effects of desilication-dealumination treatments were investigated through complementary techniques such as N₂ physisorption at -196 °C, XRD and SEM. The analysis revealed that the Clinoptilolite specific surface area increased by 400% following the dealumination-desilication procedure; at the same time the XRD analysis shows that the processed Clinoptilolite has the same main peaks of the unmodified one. Finally, the retained activity and the stability of the immobilized enzymes were evaluated, the results show that these aspects were enhanced by the modification through acid-alkaline attacks of the Clinoptilolite.

References:

¹Obert, R.; Dave B. C.; J. Am. Chem. Soc. **1999**, 121, 12192-12193

²Pietricola, G.; Ottone, C.; Fino, D.; Tommasi, T.; J. CO₂ Util. **2020**, 42, 101343

³Verboekend, D.; Keller, T.C.; Milina, M.; Hauert, R.; Pérez-Ramírez, J.; Chem. Mater. **2013**, 25, 1947-1959



AIZ 2021