

Boundedness of pseudodifferential operators with symbols in Wiener amalgam spaces on modulation spaces

*Original*

Boundedness of pseudodifferential operators with symbols in Wiener amalgam spaces on modulation spaces / Trapasso, salvatore ivan; D'Elia, Lorenza. - In: JOURNAL OF PSEUDO-DIFFERENTIAL OPERATORS AND APPLICATIONS. - ISSN 1662-9981. - 9:4(2018), pp. 881-890. [10.1007/s11868-017-0220-1]

*Availability:*

This version is available at: 11583/2971949 since: 2022-10-01T09:24:49Z

*Publisher:*

Birkhauser Verlag AG

*Published*

DOI:10.1007/s11868-017-0220-1

*Terms of use:*

openAccess

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

(Article begins on next page)

# BOUNDEDNESS OF PSEUDODIFFERENTIAL OPERATORS WITH SYMBOLS IN WIENER AMALGAM SPACES ON MODULATION SPACES

LORENZA D'ELIA AND S. IVAN TRAPASSO

ABSTRACT. This paper provides sufficient conditions for the boundedness of Weyl operators on modulation spaces. The Weyl symbols belong to Wiener amalgam spaces, or generalized modulation spaces, as recently renamed by their inventor Hans Feichtinger. This is the first result which relates symbols in Wiener amalgam spaces to operators acting on classical modulation spaces.

*This is an Accepted Manuscript of an article published by Springer in Journal of Pseudo-Differential Operators and Applications on 20 July 2017, available at: <https://link.springer.com/article/10.1007/s11868-017-0220-1>, DOI: 10.1007/s11868-017-0220-1.*

## 1. INTRODUCTION

In this paper we investigate the boundedness properties of pseudodifferential operators in the Weyl form. These operators arise as quantization rule proposed by Weyl in [41]. Namely, the rule assigns an operator  $Op_W(a)$  to a function  $a$  (the so-called Weyl symbol) on the phase space  $\mathbb{R}^{2d}$ :

$$a \rightarrow Op_W(a).$$

The operator  $Op_W(a)$  is called a Weyl operator or Weyl transform (cf., e.g., [42]). From a Time-frequency Analysis perspective Weyl operators can be introduced by means of the related time-frequency representation, the so-called (cross-)Wigner distribution  $W(f, g)$ , which for signals  $f, g$  in the Schwartz class  $\mathcal{S}(\mathbb{R}^d)$  is defined by

$$(1) \quad W(f, g)(x, \omega) = \int_{\mathbb{R}^d} e^{-2\pi iy\omega} f\left(x + \frac{y}{2}\right) \overline{g\left(x - \frac{y}{2}\right)} dy.$$

---

2000 *Mathematics Subject Classification.* 42B35, 35B65, 35J10, 35B40.

*Key words and phrases.* Wigner distribution, Wiener amalgam spaces, modulation spaces.

The Weyl operator  $Op_W(a)$  with symbol  $a$  in the space of tempered distribution  $\mathcal{S}'(\mathbb{R}^{2d})$  can be then defined by the formula

$$(2) \quad \langle Op_W(a)f, g \rangle = \langle a, W(g, f) \rangle, \quad f, g \in \mathcal{S}(\mathbb{R}^d).$$

The study of continuity properties for Weyl operators on different kinds of function spaces has been pursued by many authors. Depending on the properties of the symbol  $a$ , one can infer the corresponding continuity properties of the related operator  $Op_W(a)$ .

For the continuity properties of  $Op_W(a)$  on  $L^p(\mathbb{R}^d)$  spaces we refer the reader to [13, 42].

Here we focus on Banach spaces which measure the time-frequency decay of a function/distribution in the phase space. They are called modulation and Wiener amalgam spaces. Indeed, we shall study the continuity properties of the operator  $Op_W(a)$  on the modulation spaces  $M^{r_1, r_2}(\mathbb{R}^d)$   $1 \leq r_1, r_2 \leq \infty$  (cf. the following section for their definition), introduced by Hans Feichtinger in [28]. The corresponding Weyl symbol  $a$  belongs to the Wiener amalgam spaces  $W(\mathcal{F}L^p, L^q)$ ,  $1 \leq p, q \leq \infty$  (cf. Section 2). The latter spaces are often known in the literature as Wiener amalgam spaces with local component  $\mathcal{F}L^p$  and global component  $L^q$ , for  $1 \leq p, q \leq \infty$ , but nowadays their inventor Hans Feichtinger [29] is suggesting to call them simply modulation spaces, since they arise as the Fourier transform of the classical modulation spaces  $M^{p, q}$  introduced in [28] and can similarly be defined by means of the short-time Fourier transform (see Section 2 for details).

Continuity properties of Weyl operators with symbols in classical modulation spaces  $M^{p, q}$  have been investigated by many authors, starting from the earliest paper [33]. The most important contributions in this framework are contained in [1, 2, 6, 4, 5, 12, 13, 14, 24, 25, 31, 35, 38, 39, 40].

Let us also recall the many studies on the continuity properties of Fourier integral operators (FIOs) on modulation spaces [7, 10, 11, 15, 16, 17, 18, 19, 20, 21, 22, 23] which find applications principally in the study of Schrödinger equations. Pseudo-differential operators are a special case of FIOs, having phase function  $\Phi(x, \xi) = 2\pi i x \xi$ .

This study is limited to pseudodifferential operators, however a future object of our research would be to investigate the continuity properties for FIOs.

The main result of this paper can be formulated in the un-weighted case as follows (cf. the subsequent Theorem 3.1).

**Theorem 1.1.** *Assume that  $1 \leq p, q, r_1, r_2 \leq \infty$  satisfy*

$$q \leq p'$$

and

$$\max\{r_1, r_2, r'_1, r'_2\} \leq p.$$

Then every Weyl operator  $\text{Op}_W(a)$  having symbol  $a \in W(\mathcal{FL}^p, L^q)$ , from  $\mathcal{S}(\mathbb{R}^d)$  to  $\mathcal{S}'(\mathbb{R}^d)$ , extends uniquely to a bounded operator on  $M^{r_1, r_2}(\mathbb{R}^d)$ , with the estimate

$$\|\text{Op}_W(a)f\|_{M^{r_1, r_2}} \lesssim \|a\|_{W(\mathcal{FL}^p, L^q)} \|f\|_{M^{r_1, r_2}}.$$

To our knowledge, this is the first result in the literature which links symbols in Wiener amalgam spaces to operators acting on modulation spaces.

Boundedness results for Weyl operators with symbols in modulation spaces still hold for the other forms of pseudodifferential operators, the so-called  $\tau$ -operators. These operators can be either defined as a quantization rule or by means of the related time-frequency representation (cf. [3]). Here we simply recall the latter. For  $\tau \in [0, 1]$ , the (cross-)  $\tau$ -Wigner distributions is given by

$$(3) \quad W_\tau(f, g)(x, \omega) = \int_{\mathbb{R}^d} e^{-2\pi i y \zeta} f(x + \tau y) \overline{g(x - (1 - \tau)y)} dy \quad f, g \in \mathcal{S}(\mathbb{R}^d),$$

whereas the  $\tau$ -pseudodifferential operators is

$$(4) \quad \langle \text{Op}_\tau(a)f, g \rangle = \langle a, W_\tau(g, f) \rangle \quad f, g \in \mathcal{S}(\mathbb{R}^d).$$

For  $\tau = 1/2$  we recapture the Weyl operator, if  $\tau = 0$  the operator is called the Kohn-Nirenberg operator  $\text{Op}_{\text{KN}}$ . A Kohn-Nirenberg operator  $\text{Op}_{\text{KN}}$  and a Weyl operator  $\text{Op}_W$  are related by the formula

$$\text{Op}_{\text{KN}}(a) = \text{Op}_W(\mathcal{U}^{-1}a)$$

where

$$(5) \quad \mathcal{U}^{-1} = \mathcal{F}^{-1} \mathcal{N}_C \mathcal{F},$$

$\mathcal{F}$  is the Fourier transform,  $\mathcal{N}_C f(z) = e^{-\pi i z \cdot C z} f(z)$ ,  $z \in \mathbb{R}^{2d}$ , and

$$C = \begin{pmatrix} 0 & 1/2I \\ 1/2I & 0 \end{pmatrix}.$$

An easy computation (cf. [32, Corollary 14.5.5]) shows that

$$|V_\Phi(\mathcal{U}^{-1}a)(z, \zeta)| = |V_{\mathcal{U}\Phi}a(z - C\zeta, \zeta)|$$

from which we conclude that  $M^{p, q}$  is invariant under the action of  $\mathcal{U}^{-1}$  and therefore, results for Kohn-Nirenberg pseudodifferential operators with symbols in  $M^{p, q}$  still hold for Weyl operators and viceversa.

More generally, for  $\tau$ -pseudodifferential operator it was proved in [34] and in [39, Remark 1.5] that for every choice  $\tau_1, \tau_2 \in [0, 1]$ ,  $a_1, a_2 \in \mathcal{S}'(\mathbb{R}^{2d})$ ,

$$(6) \quad \text{Op}_{\tau_1}(a_1) = \text{Op}_{\tau_2}(a_2) \Leftrightarrow \widehat{a}_2(\xi_1, \xi_2) = e^{-2\pi i(\tau_2 - \tau_1)\xi_1 \xi_2} \widehat{a}_1(\xi_1, \xi_2).$$

For  $t > 0$  consider  $H_t(x, \xi) = e^{2\pi i t x \xi}$  and observe that

$$(7) \quad \mathcal{F}H_t(\zeta_1, \zeta_2) = \frac{1}{t^d} e^{-2\pi i \frac{1}{t} \zeta_1 \zeta_2}.$$

So, for  $\tau_1 \neq \tau_2$ , by (7),

$$(8) \quad a_2(x, \xi) = \frac{1}{|\tau_1 - \tau_2|^d} e^{2\pi i(\tau_2 - \tau_1)\Phi} * a_1(x, \xi),$$

where  $\Phi(x, \xi) = x\xi$ . The mapping  $a \mapsto T_\Phi a = e^{2\pi i\Phi} * a$  is a homeomorphism on  $M^{p,q}(\mathbb{R}^{2d})$ ,  $1 \leq p, q \leq \infty$ , [39, Proposition 1.2 (5)].

Coming back to Wiener amalgam spaces  $W(\mathcal{F}L^p, L^q)$ , we first observe that they are not invariant under the action of the operator  $\mathcal{U} = \mathcal{F}^{-1}\mathcal{N}_{-c}\mathcal{F}$ . This is proved in [13, Proposition 6.4]. So that boundedness results for Weyl operators do not extend automatically to Kohn-Nirenberg ones and vice-versa. This result easily extends to the case of any  $\tau$ -pseudodifferential operator. Indeed, for any  $\tau > 0$ , the same arguments as in the proof of Proposition 6.4 of [13] apply to the metaplectic operator  $\mathcal{U}_\tau := \mathcal{F}^{-1}\mathcal{N}_{-\tau c}\mathcal{F}$ . This is the reason why our main result can be stated only for Weyl operators.

We shall pursue the study of boundedness properties of  $\tau$ -pseudodifferential operators in a subsequent paper.

**Notation.** We define  $t^2 = t \cdot t$ , for  $t \in \mathbb{R}^d$ , and  $xy = x \cdot y$  is the scalar product on  $\mathbb{R}^d$ . The Schwartz class is denoted by  $\mathcal{S}(\mathbb{R}^d)$ , the space of tempered distributions by  $\mathcal{S}'(\mathbb{R}^d)$ . We use the brackets  $\langle f, g \rangle$  to denote the extension to  $\mathcal{S}(\mathbb{R}^d) \times \mathcal{S}'(\mathbb{R}^d)$  of the inner product  $\langle f, g \rangle = \int f(t)\overline{g(t)}dt$  on  $L^2(\mathbb{R}^d)$ . The Fourier transform of a function  $f$  on  $\mathbb{R}^d$  is normalized as

$$\mathcal{F}f(\xi) = \int_{\mathbb{R}^d} e^{-2\pi i x \xi} f(x) dx.$$

## 2. PRELIMINARIES

**2.1. Modulation and Wiener amalgam spaces.** Modulation and Wiener amalgam space norms are a measure of the joint time-frequency distribution of  $f \in \mathcal{S}'$ . For their basic properties we refer to [27, 28, 29] and the textbooks [26, 32].

Let  $f \in \mathcal{S}'(\mathbb{R}^d)$ . We define the short-time Fourier transform of  $f$  as

$$(9) \quad V_g f(z) = \mathcal{F}[fT_x g](\xi) = \int_{\mathbb{R}^d} f(y) \overline{g(y-x)} e^{-2\pi i y \xi} dy$$

for  $z = (x, \xi) \in \mathbb{R}^d \times \mathbb{R}^d$ .

For description of decay properties, we use weight functions on the time-frequency plane. In the sequel  $v$  will always be a continuous, positive, even, submultiplicative weight function (i.e. a submultiplicative weight), i.e.,  $v(0) = 1$ ,  $v(z) = v(-z)$ , and  $v(z_1 + z_2) \leq v(z_1)v(z_2)$ , for all  $z, z_1, z_2 \in \mathbb{R}^{2d}$ . A positive, even weight function  $m$  on  $\mathbb{R}^{2d}$  is called *v-moderate* if  $m(z_1 + z_2) \leq Cv(z_1)m(z_2)$  for all  $z_1, z_2 \in \mathbb{R}^{2d}$ . Let us denote by  $\mathcal{M}_v(\mathbb{R}^{2d})$  the space of *v-moderate* weights.

Given  $g \in \mathcal{S}(\mathbb{R}^d) \setminus \{0\}$ , a  $v$ -moderate weight function  $m$  on  $\mathbb{R}^{2d}$ ,  $1 \leq p, q \leq \infty$ , the *modulation space*  $M_m^{p,q}(\mathbb{R}^d)$  consists of all tempered distributions  $f \in \mathcal{S}'(\mathbb{R}^d)$  such that  $V_g f \in L_m^{p,q}(\mathbb{R}^{2d})$  (weighted mixed-norm spaces). The norm on  $M_m^{p,q}$  is

$$\|f\|_{M_m^{p,q}} = \|V_g f\|_{L_m^{p,q}} = \left( \int_{\mathbb{R}^d} \left( \int_{\mathbb{R}^d} |V_g f(x, \xi)|^p m(x, \xi)^p dx \right)^{q/p} d\xi \right)^{1/q}$$

(obvious modifications for  $p = \infty$  or  $q = \infty$ ). If  $p = q$ , we write  $M_m^p$  instead of  $M_m^{p,p}$ , and if  $m(z) \equiv 1$  on  $\mathbb{R}^{2d}$ , then we write  $M^{p,q}$  and  $M^p$  for  $M_m^{p,q}$  and  $M_m^{p,p}$ .

The space  $M_m^{p,q}(\mathbb{R}^d)$  is a Banach space whose definition is independent of the choice of the window  $g$ , in the sense that different non-zero window functions yield equivalent norms. The modulation space  $M^{\infty,1}$  is also called Sjöstrand's class [37].

For any  $p, q \in [1, \infty]$  and any  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$ , the inner product  $\langle \cdot, \cdot \rangle$  on  $\mathcal{S}(\mathbb{R}^d) \times \mathcal{S}(\mathbb{R}^d)$  extends to a continuous sesquilinear map  $M_m^{p,q}(\mathbb{R}^d) \times M_{1/m}^{p',q'}(\mathbb{R}^d) \rightarrow \mathbb{C}$ .

Here and elsewhere the conjugate exponent  $p'$  of  $p \in [1, \infty]$  is defined by  $1/p + 1/p' = 1$ . For any *even* weight functions  $u, w$  on  $\mathbb{R}^d$ , the Wiener amalgam spaces  $W(\mathcal{F}L_u^p, L_w^q)(\mathbb{R}^d)$  are given by the distributions  $f \in \mathcal{S}'(\mathbb{R}^d)$  such that

$$\|f\|_{W(\mathcal{F}L_u^p, L_w^q)(\mathbb{R}^d)} := \left( \int_{\mathbb{R}^d} \left( \int_{\mathbb{R}^d} |V_g f(x, \xi)|^p u^p(\xi) d\xi \right)^{q/p} w^q(x) dx \right)^{1/q} < \infty$$

(obvious modifications for  $p = \infty$  or  $q = \infty$ ). Using Parseval identity in (9), we can write the so-called fundamental identity of time-frequency analysis  $V_g f(x, \xi) = e^{-2\pi i x \xi} V_{\hat{g}} \hat{f}(\xi, -x)$ , so that

$$|V_g f(x, \xi)| = |V_{\hat{g}} \hat{f}(\xi, -x)| = |\mathcal{F}(\hat{f} T_{\xi} \bar{\hat{g}})(-x)|$$

and (recall  $u(x) = u(-x)$ )

$$\|f\|_{M_{u \otimes w}^{p,q}} = \left( \int_{\mathbb{R}^d} \|\hat{f} T_{\xi} \bar{\hat{g}}\|_{\mathcal{F}L_u^p}^q w^q(\xi) d\xi \right)^{1/q} = \|\hat{f}\|_{W(\mathcal{F}L_u^p, L_w^q)}.$$

Hence Wiener amalgam spaces are simply the image under Fourier transform of modulation spaces:

$$(10) \quad \mathcal{F}(M_{u \otimes w}^{p,q}) = W(\mathcal{F}L_u^p, L_w^q).$$

For completeness, let us recall the inclusion properties of modulation spaces. Suppose  $m_1, m_2 \in \mathcal{M}_v(\mathbb{R}^{2d})$ . Then

$$(11) \quad \mathcal{S}(\mathbb{R}^d) \subseteq M_{m_1}^{p_1, q_1}(\mathbb{R}^d) \subseteq M_{m_2}^{p_2, q_2}(\mathbb{R}^d) \subseteq \mathcal{S}'(\mathbb{R}^d), \\ p_1 \leq p_2, \quad q_1 \leq q_2, \quad m_2 \lesssim m_1.$$

We denote by  $J$  the symplectic matrix

$$(12) \quad J = \begin{pmatrix} 0_{d \times d} & I_{d \times d} \\ -I_{d \times d} & 0_{d \times d} \end{pmatrix}.$$

### 3. SYMBOLS IN WIENER AMALGAM SPACES

We need first to investigate the properties of the Wigner distribution in terms of Wiener amalgam spaces. From now on we set  $v_J(z) = v(Jz)$ , where  $J$  is the symplectic matrix in (12). We obtain the following results.

**Lemma 3.1.** *Consider  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$ ,  $1 \leq p_1, p_2 \leq \infty$ ,  $f \in M_m^{p_1, p_2}$ ,  $g \in M_{1/m}^{p'_1, p'_2}$ , then the Wigner distribution  $W(g, f) \in W(\mathcal{FL}_{1/v_J}^1, L^\infty)$ , with*

$$(13) \quad \|W(g, f)\|_{W(\mathcal{FL}_{1/v_J}^1, L^\infty)} \lesssim \|f\|_{M_m^{p_1, p_2}} \|g\|_{M_{1/m}^{p'_1, p'_2}}.$$

*Proof.* If  $\zeta = (\zeta_1, \zeta_2) \in \mathbb{R}^{2d}$ , then [32, Lemma 14.5.1] says that

$$|V_\Phi(W(g, f))(z, \zeta)| = |V_\varphi f(z + \frac{J\zeta}{2})| |V_\varphi g(z - \frac{J\zeta}{2})|.$$

Consequently

$$(14) \quad \|W(g, f)\|_{W(\mathcal{FL}_{1/v_J}^1, L^\infty)} \asymp \sup_{z \in \mathbb{R}^{2d}} \int_{\mathbb{R}^{2d}} |V_\varphi f(z + \frac{J\zeta}{2})| |V_\varphi g(z - \frac{J\zeta}{2})| \frac{1}{v(J\zeta)} d\zeta.$$

Making the change of variables  $u = J\zeta$  and observing that

$$\frac{1}{v(u)} \leq C \frac{m(z + \frac{u}{2})}{m(z - \frac{u}{2})},$$

$$\begin{aligned} \|W(g, f)\|_{W(\mathcal{FL}_{1/v_J}^1, L^\infty)} &\leq C \sup_{z \in \mathbb{R}^{2d}} \int_{\mathbb{R}^{2d}} |V_\varphi f(z + \frac{u}{2})| |V_\varphi g(z - \frac{u}{2})| \frac{m(z + \frac{u}{2})}{m(z - \frac{u}{2})} du \\ &= 2^{2d} C \sup_{z \in \mathbb{R}^{2d}} \int_{\mathbb{R}^{2d}} |V_\varphi f(z + u)| |V_\varphi g(z - u)| \frac{m(z + u)}{m(z - u)} du \\ &\leq \tilde{C} \|V_\varphi f m\|_{L^{p_1, p_2}} \|V_\varphi g \frac{1}{m}\|_{L^{p'_1, p'_2}} \\ &\lesssim \|f\|_{M_m^{p_1, p_2}} \|g\|_{M_{1/m}^{p'_1, p'_2}}. \end{aligned}$$

The claim is proved.  $\square$

**Lemma 3.2.** *Consider  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$ ,  $f \in M_m^2$ ,  $g \in M_{1/m}^2$ , then the Wigner distribution  $W(g, f) \in W(\mathcal{FL}_{1/v_J}^2, L^2)$ , with*

$$(15) \quad \|W(g, f)\|_{W(\mathcal{FL}_{1/v_J}^2, L^2)} \lesssim \|f\|_{M_m^2} \|g\|_{M_{1/m}^2}.$$

*Proof.* The technique is similar to the one in Lemma 3.1. Using (14) and the change of variables  $w = z + J\zeta/2$ ,  $u = J\zeta$ , we can write

$$\begin{aligned}
\|W(g, f)\|_{W(\mathcal{FL}_{1/v_J}^2, L^2)} &\asymp \left( \int_{\mathbb{R}^{2d}} \int_{\mathbb{R}^{2d}} |V_\varphi f(z + \frac{J\zeta}{2})|^2 |V_\varphi g(z - \frac{J\zeta}{2})|^2 \frac{1}{v^2(J\zeta)} d\zeta dz \right)^{\frac{1}{2}} \\
&= \left( \int_{\mathbb{R}^{2d}} \int_{\mathbb{R}^{2d}} |V_\varphi f(w)|^2 |V_\varphi g(w - u)|^2 \frac{1}{v^2(u)} dudw \right)^{\frac{1}{2}} \\
&\leq \tilde{C} \left( \int_{\mathbb{R}^{2d}} (|V_\varphi f|^2 m^2) * (|V_\varphi g|^2 \frac{1}{m^2}) du \right)^{\frac{1}{2}} \\
&\lesssim \| |V_\varphi f|^2 m^2 \|_1 \| |V_\varphi g|^2 \frac{1}{m^2} \|_1 \\
&\lesssim \|f\|_{M_m^2} \|g\|_{M_{1/m}^2},
\end{aligned}$$

where we have used Young's Inequality  $L^1 * L^1 \subset L^1$ . This concludes the proof.  $\square$

**3.1. Main result.** We address this section to the study of pseudodifferential operators acting on modulation spaces and having symbols in weighted Wiener amalgam spaces.

Here is our main result.

**Theorem 3.1.** *Assume that  $1 \leq p, q, r_1, r_2 \leq \infty$  satisfy*

$$(16) \quad q \leq p'$$

and

$$(17) \quad \max\{r_1, r_2, r'_1, r'_2\} \leq p.$$

*Consider  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$ . Then every Weyl operator  $\text{Op}_W(a)$  having symbol  $a \in W(\mathcal{FL}_{v_J}^p, L^q)$ , from  $\mathcal{S}(\mathbb{R}^d)$  to  $\mathcal{S}'(\mathbb{R}^d)$ , extends uniquely to a bounded operator on  $M_m^{r_1, r_2}(\mathbb{R}^d)$ , with the estimate*

$$(18) \quad \|\text{Op}_W(a)f\|_{M_m^{r_1, r_2}} \lesssim \|a\|_{W(\mathcal{FL}_{v_J}^p, L^q)} \|f\|_{M_m^{r_1, r_2}}.$$

The proof uses complex interpolation between Wiener amalgam spaces  $W(\mathcal{FL}_{v_J}^\infty, L^1)$  and  $W(\mathcal{FL}_{v_J}^2, L^2)$ , for which we first show the corresponding boundedness results.

**Proposition 3.2.** *Consider  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$  and  $a \in W(\mathcal{FL}_{v_J}^\infty, L^1)$ . Then the operator  $\text{Op}_W(a)$  is bounded on  $M_m^{r_1, r_2}$ , for every  $1 \leq r_1, r_2 \leq \infty$ , with*

$$(19) \quad \|\text{Op}_W(a)f\|_{M_m^{r_1, r_2}} \lesssim \|a\|_{W(\mathcal{FL}_{v_J}^\infty, L^1)} \|f\|_{M_m^{r_1, r_2}}.$$



*Proof.* For every  $f \in M_m^{r_1, r_2}$  and  $g \in M_{1/m}^{r'_1, r'_2}$ , we can write, for any fixed  $\Phi \in \mathcal{S}(\mathbb{R}^{2d}) \setminus \{0\}$ ,

$$|\langle \text{Op}_W(a)f, g \rangle| = |\langle a, W(g, f) \rangle| \leq \|V_\Phi a\|_{L_z^1(L_{v_J, \zeta}^\infty)} \|V_\Phi W(g, f)\|_{L_z^\infty(L_{1/v_J, \zeta}^1)}.$$

Observe that

$$\|W(g, f)\|_{W(\mathcal{F}L_{v_J}^1, L^\infty)} \asymp \|V_\Phi W(g, f)\|_{L_z^\infty(L_{1/v_J, \zeta}^1)} \lesssim \|f\|_{M_m^{r_1, r_2}} \|g\|_{M_{1/m}^{r'_1, r'_2}},$$

by Lemma 3.1. This concludes the proof.  $\square$

**Proposition 3.3.** *Consider  $m \in \mathcal{M}_v(\mathbb{R}^{2d})$  and  $a \in W(\mathcal{F}L_{v_J}^2, L^2)$ . Then the operator  $\text{Op}_W(a)$  is bounded on  $M_m^2$  with*

$$(20) \quad \|\text{Op}_W(a)f\|_{M_m^2} \lesssim \|a\|_{W(\mathcal{F}L_{v_J}^2, L^2)} \|f\|_{M_m^2}.$$

*Proof.* The arguments are the same as Proposition 3.2, with Lemma 3.1 replaced by 3.2. We leave the details to the interested reader.  $\square$

**Remark 3.4.** (i) *Observe that by (10),  $W(\mathcal{F}L_{v_J}^2, L^2) = \mathcal{F}M_{v_J \otimes 1}^2$  and a straightforward modification of [32, Theorem 11.3.5 (c)] gives*

$$\mathcal{F}M_{v_J \otimes 1}^2 = M_{1 \otimes v_{J-1}}^2 = M_{1 \otimes v_J}^2$$

*since by assumption  $v(-z) = v(z)$ .*

(ii) *Since  $v(-z) = v(z)$ , the weight  $v_J$  is even and the conclusion of the previous step (i) also follows by [29, Theorem 6], in the case  $p = 2$ .*

(iii) *Using (i) or (ii) we derive that the Wiener amalgam space  $W(\mathcal{F}L_{v_J}^2, L^2)$  coincides with the modulation space  $M_{1 \otimes v_J}^2$ . Then the conclusion of Proposition 3.3 also follows from [40, Theorem 4.3].*

*Proof of Theorem 3.1.* We make use of complex interpolation between Wiener amalgam and modulation spaces, using the boundedness results of Propositions 3.2 and 3.3. For  $\theta \in [0, 1]$ , we have

$$[W(\mathcal{F}L_{v_J}^\infty, L^1), W(\mathcal{F}L_{v_J}^2, L^2)]_\theta = W(\mathcal{F}L_{v_J}^p, L^{p'}),$$

with  $2 \leq p \leq \infty$ . As far as modulation spaces concern,  $[M_m^{s_1, s_2} M_m^2]_\theta = M_m^{r_1, r_2}$ , with

$$\frac{1}{r_1} = \frac{1-\theta}{s_1} + \frac{\theta}{2} = \frac{1-\theta}{s_1} + \frac{1}{p}$$

and

$$\frac{1}{r_2} = \frac{1-\theta}{s_2} + \frac{\theta}{2} = \frac{1-\theta}{s_2} + \frac{1}{p}$$

hence  $r_1, r_2 \leq p$ . Similarly we obtain  $r'_1, r'_2 \leq p$ , and the (17) follows. Finally, inclusion relations for Wiener amalgam spaces allow to consider symbols  $a \in W(\mathcal{F}L_{v_J}^p, L^q)$ , with  $q \leq p'$ , which gives (16) and concludes the proof.  $\square$

## ACKNOWLEDGEMENTS

The authors would like to thank Professors Elena Cordero and Fabio Nicola for fruitful conversations and comments.

## REFERENCES

- [1] I. Beltiță and D. Beltiță, Modulation Spaces of Symbols for Representations of Nilpotent Lie Groups, *J. Fourier Anal. Appl.*, (2011) 17:290.
- [2] A. Bényi, K. Gröchenig, K.A. Okoudjou and L.G. Rogers. Unimodular Fourier multipliers for modulation spaces. *J. Funct. Anal.*, 246(2): 366-384, 2007.
- [3] P. Boggiatto, G. De Donno, A. Oliaro, Time-frequency representations of Wigner type and pseudo-differential operators, *Trans. Amer. Math. Soc.*, 362(9) (2010), 4955–4981.
- [4] E. Cordero, M. de Gosson, F. Nicola. On the Invertibility of Born-Jordan Quantization. *J. Math. Pures Appl.*, 105(4):537–557, 2016.
- [5] E. Cordero, M. de Gosson and F. Nicola. Time-frequency Analysis of Born-Jordan Pseudo-differential Operators. *J. Funct. Anal.*, 272(2):577–598, 2017. DOI:10.1016/j.jfa.2016.10.004
- [6] E. Cordero and K. Gröchenig. Time-frequency analysis of localization operators. *J. Funct. Anal.*, 205 (1), 107–131, 2003.
- [7] E. Cordero, K. Gröchenig, F. Nicola and L. Rodino. Generalized Metaplectic Operators and the Schrödinger Equation with a Potential in the Sjöstrand Class, *J. Math. Phys.*, 55(8), art. no. 081506, 2014.
- [8] E. Cordero and F. Nicola. Metaplectic representation on Wiener amalgam spaces and applications to the Schrödinger equation. *J. Funct. Anal.*, 254:506–534, 2008.
- [9] E. Cordero and F. Nicola. Strichartz estimates in Wiener amalgam spaces for the Schrödinger equation. *Math. Nachr.*, 281(1):25–41, 2008.
- [10] E. Cordero and F. Nicola. Metaplectic representation on Wiener amalgam spaces and applications to the Schrödinger equation. *J. Funct. Anal.*, 254: 506-534, 2008.
- [11] E. Cordero and F. Nicola. Boundedness of Schrödinger type propagators on modulation spaces. *J. Fourier Anal. Appl.*, 16(3):311–339, 2010.
- [12] E. Cordero, F. Nicola, Sharp continuity results for the short-time Fourier transform and for localization operators, *Monatsh. Math.*, 162 (2011), 251–276.
- [13] E. Cordero, F. Nicola, Pseudodifferential operators on  $L^p$ , Wiener amalgam and modulation spaces, *Int. Math. Res. Notices*, 10 (2010), 1860–1893.
- [14] E. Cordero and F. Nicola. Sharp integral bounds for Wigner distributions. *International Mathematics Research Notices*, 2016(00), 1–29, 2016. DOI:10.1093/imrn/rnw250
- [15] E. Cordero and F. Nicola. On the Schrödinger equation with potential in modulation spaces. *J. Pseudo-Differ. Op. and Appl.*, 5(3):319–341, 2014.
- [16] E. Cordero, F. Nicola and L. Rodino. Time-frequency analysis of Fourier integral operators. *Commun. Pure Appl. Anal.*, 9(1):1–21, 2010.
- [17] E. Cordero, F. Nicola and L. Rodino, Sparsity of Gabor representation of Schrödinger propagators. *Appl. Comput. Harmon. Anal.*, 26(3):357–370, 2009.
- [18] E. Cordero, F. Nicola and L. Rodino, Gabor representations of evolution operators. *Trans. Amer. Math. Soc.*, 367(11):7639–7663, 2015.
- [19] E. Cordero, F. Nicola and L. Rodino. Schrödinger equations with rough Hamiltonians. *Journal Discrete and Continuous Dynamical System - A*, 35(10):4805–4821, 2015.

- [20] E. Cordero, F. Nicola and L. Rodino. Propagation of the Gabor Wave Front Set for Schrödinger Equations with non-smooth potentials. *Rev. Math. Phys.*, 27 (1) art. no. 1550001, 2015.
- [21] E. Cordero, F. Nicola and L. Rodino. Integral Representations for the Class of Generalized Metaplectic Operators. *J. Fourier Anal. Appl.*, 21:694–714, 2015.
- [22] E. Cordero, F. Nicola and L. Rodino. Wave packet analysis of Schrödinger equations in analytic function spaces. *Adv. Math.*, 278:182–209, 2015. DOI:10.1016/j.aim.2015.03.014
- [23] E. Cordero, F. Nicola and L. Rodino. Exponentially sparse representations of Fourier integral operators. *Rev. Math. Iberoamer.*, 31:461–476, 2015.
- [24] E. Cordero, A. Tabacco, P. Wahlberg, Schrödinger-type propagators, pseudodifferential operators and modulation spaces, *J. London Math. Soc.*, 88(2): 375–395, 2013.
- [25] E. Cordero, J. Toft and P. Wahlberg. Sharp results for the Weyl product on modulation spaces. *J. Funct. Anal.*, 267(8):3016–3057, 2014.
- [26] M. de Gosson, *Symplectic methods in Harmonic Analysis and in Mathematical Physics*, Birkhäuser, 2011.
- [27] H. G. Feichtinger. Banach convolution algebras of Wiener's type, In *Proc. Conf. "Function, Series, Operators"*, Budapest August 1980, Colloq. Math. Soc. János Bolyai, 35, 509–524, North-Holland, Amsterdam, 1983.
- [28] H. G. Feichtinger. Banach spaces of distributions of Wiener's type and interpolation. In *Proc. Conf. Functional Analysis and Approximation, Oberwolfach August 1980*, Internat. Ser. Numer. Math., 69:153–165. Birkhäuser, Boston, 1981.
- [29] H. G. Feichtinger. Generalized amalgams, with applications to Fourier transform. *Canad. J. Math.*, 42(3):395–409, 1990.
- [30] H. G. Feichtinger. Atomic characterizations of modulation spaces through Gabor-type representations. In *Proc. Conf. Constructive Function Theory, Rocky Mountain J. Math.*, 19:113–126, 1989.
- [31] K. Gröchenig. Time-Frequency Analysis of Sjöstrand's Class. *Rev. Mat. Iberoamericana*, 22(2):703–724, 2006.
- [32] K. Gröchenig. *Foundations of Time-Frequency Analysis*. Birkhäuser, Boston, 2001.
- [33] K. Gröchenig and C. Heil. Modulation spaces and pseudodifferential operators. *Integral Equations Operator Theory*, 34(4):439–457, 1999.
- [34] L. Hörmander. *The analysis of linear partial differential operators. III*, volume 274 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, Berlin, 1994.
- [35] D. Labate, Pseudodifferential operators on modulation spaces. *J. Math. Anal. Appl.*, 262:242–255, 2001.
- [36] L. H. Lieb. Integral bounds for radar ambiguity functions and Wigner distributions. *J. Math. Phys.*, 31, No.3, 1990.
- [37] J. Sjöstrand. An algebra of pseudodifferential operators. *Math. Res. Lett.*, 1:185–192, 1994.
- [38] M. Sugimoto and N. Tomita. Boundedness properties of pseudo-differential operators and Calderón-Zygmund operators on modulation spaces. *J. Fourier Anal. Appl.*, 14(1):124–143, 2008.
- [39] J. Toft. Continuity properties for modulation spaces, with applications to pseudo-differential calculus. I. *J. Funct. Anal.*, 207(2):399–429, 2004.
- [40] J. Toft. Continuity properties for modulation spaces, with applications to pseudo-differential calculus. II. *Ann. Global Anal. Geom.*, 26(1):73–106, 2004.
- [41] H. Weyl, *Quantenmechanik und Gruppentheories*. *Zeitschrift für Physik*, 46:1-46, 1927.

[42] M. W. Wong. Weyl Transforms, Springer, 1998.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123  
TORINO, ITALY

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123  
TORINO, ITALY

*Email address:* `lorenza.delia@edu.unito.it`

*Email address:* `salvatore.trapasso@edu.unito.it`