

Comment on “Groundwater Affects the Geomorphic and Hydrologic Properties of Coevolved Landscapes” by Litwin et al

Original

Comment on “Groundwater Affects the Geomorphic and Hydrologic Properties of Coevolved Landscapes” by Litwin et al / Kumar Anand, Shashank; Bonetti, Sara; Camporeale, CARLO VINCENZO; Hooshyar, Milad; Porporato, Amilcare. - In: JOURNAL OF GEOPHYSICAL RESEARCH. EARTH SURFACE. - ISSN 2169-9003. - 127:10(2022).
[10.1029/2022JF006669]

Availability:

This version is available at: 11583/2972959 since: 2022-11-10T21:27:18Z

Publisher:

AMER GEOPHYSICAL UNION

Published

DOI:10.1029/2022JF006669

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

This article is a comment on Litwin et al. (2022), <https://doi.org/10.1029/2021JF006239>.

Key Points:

- The specific contributing area is defined in the limit of an infinitesimal contour length
- Not all solutions of the minimalist landscape evolution model are rescaled replicas of each other
- Boundary conditions play an essential role in setting the landscape morphologies at the steady state

Correspondence to:

A. Porporato,
aporpora@princeton.edu

Citation:

Anand, S. K., Bonetti, S., Camporeale, C., Hooshyar, M., & Porporato, A. (2022). Comment on “Groundwater affects the geomorphic and hydrologic properties of coevolved landscapes” by Litwin et al. *Journal of Geophysical Research: Earth Surface*, 127, e2022JF006669. <https://doi.org/10.1029/2022JF006669>

Received 8 MAR 2022
Accepted 6 SEP 2022

Comment on “Groundwater Affects the Geomorphic and Hydrologic Properties of Coevolved Landscapes” by Litwin et al.

Shashank Kumar Anand¹, **Sara Bonetti²**, **Carlo Camporeale³**, **Milad Hooshyar⁴**, and **Amilcare Porporato^{1,5}**

¹Department of Civil and Environmental Engineering, Princeton University, Princeton, NJ, USA, ²School of Architecture, Civil and Environmental Engineering, École Polytechnique Fédérale de Lausanne, Lausanne, Switzerland, ³Department of Environment, Land and Infrastructure Engineering, Politecnico di Torino, Torino, Italy, ⁴Amazon, Seattle, WA, USA, ⁵High Meadows Environmental Institute, Princeton University, Princeton, NJ, USA

Abstract The objective of this comment is to correct two sets of statements in Litwin et al. (2022, <https://doi.org/10.1029/2021JF006239>), which consider our research work (Bonetti et al., 2018, <https://doi.org/10.1098/rspa.2017.0693>; Bonetti et al., 2020, <https://doi.org/10.1073/pnas.1911817117>). We clarify here that (a) the specific contributing area is defined in the limit of an infinitesimal contour length instead of the product of a reference contour width (Bonetti et al., 2018, <https://doi.org/10.1098/rspa.2017.0693>), and (b) not all solutions obtained from the minimalist landscape evolution model of Bonetti et al. (2020, <https://doi.org/10.1073/pnas.1911817117>) are rescaled copies of each other. We take this opportunity to demonstrate that the boundary conditions impact the obtained solutions, which has not been considered in the dimensional analysis of Litwin et al. (2022, <https://doi.org/10.1029/2021JF006239>). We clarify this point by using dimensional analysis and numerical simulations for a square domain, where only one horizontal length scale (the side length l) enters the physical law.

Plain Language Summary This comment is focused on rectifying the statements regarding the previous works of Bonetti et al. (2018, <https://doi.org/10.1098/rspa.2017.0693>) and Bonetti et al. (2020, <https://doi.org/10.1073/pnas.1911817117>) in Litwin et al. (2022, <https://doi.org/10.1029/2021JF006239>). We clarify here the definition of the specific contributing area and the role of boundary conditions in the obtained solutions of the landscape evolution model, which has been left unnoticed by Litwin et al. (2022, <https://doi.org/10.1029/2021JF006239>). Using dimensional analysis and simulations for a square domain, we show that only those solutions remain rescaled copies of each other where the non-dimensional group containing the influence of boundary conditions remains the same. This result was already incorporated in the dimensional analysis of Bonetti et al. (2020, <https://doi.org/10.1073/pnas.1911817117>), where a non-dimensional index C_l was derived explicitly.

Litwin et al. (2022) wrote a very interesting paper on the problem of the co-evolution of groundwater hydrology and land-surface dynamics. The authors are to be commended for this very nice research contribution, which quantifies the feedback between the spatial patterns in the emerging topography and sub-surface properties.

This comment is intended only to rectify some statements in this paper that regard our work (Bonetti et al., 2018, 2020). The first one considers the definition of the specific contributing area a and its relation to the total contributing area A . The second one is related to the dimensional analysis and the scaling properties of the landscape evolution model (LEM) solutions. We feel that these corrections are necessary to avoid misinterpretations of our results as well as to clarify important aspects regarding the scaling properties of the obtained solutions.

The statements in question concern the LEM with governing equations

$$\frac{\partial z}{\partial t} = D \nabla^2 z - K \sqrt{a} |\nabla z| + U, \tag{1}$$

$$-\nabla \cdot \left(a \frac{\nabla z}{|\nabla z|} \right) = 1, \tag{2}$$

where z is the topographic elevation, t is time, D is a diffusion coefficient, K is an erosion coefficient, a is the specific contributing area, and U is the uplift rate. For constant K , the model—referred to as the *NoHyd* model—is a specific version of the minimalist LEM of Bonetti et al. (2020).

Regarding the definition of the specific contributing area, Litwin et al. (2022) write on page 5: “To make the conversion between A and a , we represent A as the product of a and a characteristic contour width v_0 , which is a chosen constant value.” In reality, a is defined in the limit of an infinitesimal contour length (Bonetti et al., 2018; Gallant & Hutchinson, 2011), not as the product of a reference contour width. As a result, Equation 2 is only valid for the specific contributing area defined in this limit, as discussed in Bonetti et al. (2018).

With reference to similarity solutions and dimensional analysis, they write on page 4, “Additionally, the nondimensionalization generalizes our results and reconciles conflicting dimensional analyses provided by Theodoratos et al. (2018) and Bonetti et al. (2020).” Further, they mention on page 27, “We show that contrary to Bonetti et al. (2020) there is a single typology of the *NoHyd* model which can be rescaled to obtain all results the model may produce.” These conclusions need to be revised because the role of the boundary conditions is not included in the dimensional analysis. While the dimensional analysis of the LEM is facilitated by the fact that some of the main variables and parameters are clearly listed in the governing equations, the obtained solutions depend on the initial and boundary conditions as much as they do on the equations (Bursten, 2021).

Considering the domain geometries for which only one horizontal length scale enters the physical law (more complicated geometries will add further governing quantities), the LEM solution can be formally written as

$$z = f(K, D, U, x, y, t, l). \quad (3)$$

For concreteness, we consider here a square domain of side length l with boundaries at fixed zero elevation values that is,

$$z(0, y) = z(l, y) = 0, \quad 0 \leq y \leq l, \quad (4)$$

$$z(x, 0) = z(x, l) = 0, \quad 0 \leq x \leq l. \quad (5)$$

Taking L , H , and T as primary dimensions for horizontal directions, vertical direction, and time, respectively (Porporato, 2022), the dimensions of the quantities involved in Equation 3 are $[z] = H$, $[K] = L^{1/2}T^{-1}$, $[D] = L^2T^{-1}$, $[U] = HT^{-1}$, $[x] = L$, $[y] = L$, and $[l] = L$. Choosing K , D , U as the repeating variables and applying the Π theorem for the formulated physical law of Equation 3 yields

$$\Pi_z = \Phi(\Pi_x, \Pi_y, \Pi_t, \Pi_l), \quad (6)$$

with

$$\Pi_z = \frac{z}{\left(\frac{DU^3}{K^4}\right)^{1/3}} = \frac{z}{H} = \hat{z}, \quad (7)$$

$$\Pi_x = \frac{x}{\left(\frac{D^2}{K^2}\right)^{1/3}} = \frac{x}{L} = \hat{x}, \quad (8)$$

$$\Pi_y = \frac{y}{\left(\frac{D^2}{K^2}\right)^{1/3}} = \frac{y}{L} = \hat{y}, \quad (9)$$

$$\Pi_t = \frac{t}{\left(\frac{D}{K^4}\right)^{1/3}} = \frac{t}{T} = \hat{t}, \quad (10)$$

$$\Pi_l = \frac{l}{\left(\frac{D^2}{K^2}\right)^{1/3}} = \frac{l}{L} = C_l^{2/3}, \quad (11)$$

where C_l is the “channelization index” derived in Bonetti et al. (2020).

Using the above-defined scales \mathcal{H} , \mathcal{L} , and \mathcal{T} , the non-dimensionalized form of the LEM (governing equations: Equation 1 and Equation 2; boundary conditions: Equation 4 and Equation 5) can be written as

$$\frac{\partial \hat{z}}{\partial \hat{t}} = \hat{\nabla}^2 \hat{z} - \sqrt{\hat{a}} |\hat{\nabla} \hat{z}| + 1, \quad (12)$$

$$-\hat{\nabla} \cdot \left(\hat{a} \frac{\hat{\nabla} \hat{z}}{|\hat{\nabla} \hat{z}|} \right) = 1, \quad (13)$$

$$\hat{z}(0, \hat{y}) = \hat{z}(C_I^{2/3}, \hat{y}) = 0, \quad 0 \leq \hat{y} \leq C_I^{2/3}, \quad (14)$$

$$\hat{z}(\hat{x}, 0) = \hat{z}(\hat{x}, C_I^{2/3}) = 0, \quad 0 \leq \hat{x} \leq C_I^{2/3}. \quad (15)$$

The absence of $\Pi_l (= C_I^{2/3})$ in the non-dimensionalized governing Equations 12 and 13 does not mean that the solutions for this model do not depend on it. Π_l appears in the non-dimensionalized boundary conditions (Equation 14 and Equation 15) used to compute the steady-state solutions. While there is freedom in selecting the repeating variables, choosing one set over another for defining the relevant scales does not change the physics of the model at hand (Porporato, 2022). Selecting l , D , and U keeps Π_l in the dimensionless governing equations (Bonetti et al., 2020), while Π_l appears in the dimensionless boundary conditions when using K , D , and U for the non-dimensionalization.




Figure 1. Steady-state solutions for a square domain with fixed elevation boundary conditions, $D = 10^{-2} \text{m}^2 \text{yr}^{-1}$, and $U = 10^{-3} \text{m yr}^{-1}$. The dimensional form $z(x, y)$ is labeled in black, while the non-dimensional form $\hat{z}(\hat{x}, \hat{y})$ computed using scales \mathcal{H} and \mathcal{L} is written in blue and enclosed in parentheses. In panels a, b, and c (left to right), the domain length is fixed as $l = 100 \text{m}$ with increasing values of erosion coefficient K that lead to distinct topographies; also explained by the varied values of $\Pi_l = 2.92, 13.57$, and 63.0 . The non-dimensional solution $\hat{z}(\hat{x}, \hat{y})$ is also different in these cases. In panels d, b, and e (top to bottom), increasing values of K are offset by the decreasing values of domain length l so that the dimensionless number $\Pi_l = 13.57$ ($C_I = 50$) remains invariant and the topographies are rescaled version of each other (same non-dimensional form $\hat{z}(\hat{x}, \hat{y})$). The simulations were performed using the numerical algorithm developed in Anand et al. (2020).

The dependency of the non-dimensional solutions on the boundary conditions through Π_l signifies that not all the solutions that the model can produce are rescaled copies of each other. This is true only for those solutions in which the non-dimensional group Π_l remains invariant. A set of steady-state simulations for a square domain, represented in both dimensional and non-dimensional forms, demonstrates this point (Figure 1). In panels a, b, and c, the value of Π_l is changed by increasing the erosion coefficient K for a domain of fixed side length $l = 100$ m. This is the case discussed in Bonetti et al. (2020), which shows different solutions ranging from the unchannelized case (Figure 1a) to a case with multiple channels (Figure 1c). The non-dimensional solutions $\hat{z}(\hat{x}, \hat{y})$ computed by employing suitable scales \mathcal{H} and \mathcal{L} are also dissimilar in these cases. On the contrary, panels d, b, and e have different values of K , but they preserve the same Π_l because of the different length scale l , and therefore, are scaled versions of each other (similar non-dimensional forms labeled in the blue text). Theodoratos et al. (2018) consider this case only.

In reality, the reason why many solutions of LEMs look alike is that they have an interesting property of complete self-similarity that emerges at very large channelization index values ($C_I = \frac{Kl^{3/2}}{D}$) (Hooshyar, Anand, & Porporato, 2020; Hooshyar, Bonetti, et al., 2020; Hooshyar et al., 2021; Porporato, 2022). For these conditions, several spatially averaged quantities become invariant with respect to C_I , having reached a self-similar regime. This is similar to what happens to turbulent flows in the fully rough regime as they become independent of the Reynolds number or to regular polygons, which tends to circles as the number of sides tends to infinity (Barenblatt, 1996).

Data Availability Statement

The details of the numerical algorithm used for the simulations is described in Anand et al. (2020) and the well-commented Python code is available on GitHub <https://github.com/ShashankAnand1996/LEM> (Anand, 2022).

Acknowledgments

The authors acknowledge support from the US National Science Foundation (NSF) grants EAR-1331846 and EAR-1338694, Innovation Award - Moore Science-to-Action Fund, and BP through the Carbon Mitigation Initiative (CMI) at Princeton University. S.K.A. and A.P. also acknowledge the support from the High Meadows Environmental Institute (HMEI). The simulations in this article were performed on computational resources provided by Princeton Research Computing, a consortium of groups including the Princeton Institute for Computational Science and Engineering (PICSciE) and the Office of Information Technology's High Performance Computing Center and Visualization Laboratory at Princeton University.

References

- Anand, S. K. (2022). Shashankanand1996/lem: Version 1.0.1. [software]. Zenodo. <https://doi.org/10.5281/zenodo.6824822>
- Anand, S. K., Hooshyar, M., & Porporato, A. (2020). Linear layout of multiple flow-direction networks for landscape-evolution simulations. *Environmental Modelling & Software*, 133, 104804. <https://doi.org/10.1016/j.envsoft.2020.104804>
- Barenblatt, G. I. (1996). *Scaling, self-similarity, and intermediate asymptotics: Dimensional analysis and intermediate asymptotics*. Cambridge University Press. <https://doi.org/10.1017/CBO9781107050242>
- Bonetti, S., Bragg, A., & Porporato, A. (2018). On the theory of drainage area for regular and non-regular points. *Proceedings of the Royal Society A: Mathematical, Physical & Engineering Sciences*, 474(2211), 20170693. <https://doi.org/10.1098/rspa.2017.0693>
- Bonetti, S., Hooshyar, M., Camporeale, C., & Porporato, A. (2020). Channelization cascade in landscape evolution. *Proceedings of the National Academy of Sciences*, 117(3), 1375–1382. <https://doi.org/10.1073/pnas.1911817117>
- Bursten, J. R. (2021). The function of boundary conditions in the physical sciences. *Philosophy of Science*, 88(2), 234–257. <https://doi.org/10.1086/711502>
- Gallant, J. C., & Hutchinson, M. F. (2011). A differential equation for specific catchment area. *Water Resources Research*, 47(5). <https://doi.org/10.1029/2009WR008540>
- Hooshyar, M., Anand, S., & Porporato, A. (2020). Variational analysis of landscape elevation and drainage networks. *Proceedings of the Royal Society A*, 476(2239), 20190775. <https://doi.org/10.1098/rspa.2019.0775>
- Hooshyar, M., Bonetti, S., Singh, A., Fofoula-Georgiou, E., & Porporato, A. (2020). From turbulence to landscapes: Logarithmic mean profiles in bounded complex systems. *Physical Review E*, 102(3), 033107. <https://doi.org/10.1103/PhysRevE.102.033107>
- Hooshyar, M., Katul, G., & Porporato, A. (2021). Spectral signature of landscape channelization. *Geophysical Research Letters*, 48(8), e2020GL091015. <https://doi.org/10.1029/2020GL091015>
- Litwin, D. G., Tucker, G. E., Barnhart, K. R., & Harman, C. J. (2022). Groundwater affects the geomorphic and hydrologic properties of coevolved landscapes. *Journal of Geophysical Research: Earth Surface*, 127(1), e2021JF006239. <https://doi.org/10.1029/2021JF006239>
- Porporato, A. (2022). Hydrology without dimensions. *Hydrology and Earth System Sciences*, 26(2), 355–374. <https://doi.org/10.5194/hess-26-355-2022>
- Theodoratos, N., Seybold, H., & Kirchner, J. W. (2018). Scaling and similarity of a stream-power incision and linear diffusion landscape evolution model. *Earth Surface Dynamics*, 6(3), 779–808. <https://doi.org/10.5194/esurf-6-779-2018>