

Representing seesaw neutrino models and their motion in lepton flavour space

Original

Representing seesaw neutrino models and their motion in lepton flavour space / Di Bari, P; Re Fiorentin, M; Samanta, R.
- In: JOURNAL OF HIGH ENERGY PHYSICS. - ISSN 1029-8479. - ELETTRONICO. - 2019:5(2019).
[10.1007/JHEP05(2019)011]

Availability:

This version is available at: 11583/2974216 since: 2023-01-11T13:15:30Z

Publisher:

SPRINGER

Published

DOI:10.1007/JHEP05(2019)011

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Representing seesaw neutrino models and their motion in lepton flavour space

Pasquale Di Bari,^a Michele Re Fiorentin^b and Rome Samanta^a

^a*Physics and Astronomy, University of Southampton,
Southampton, SO17 1BJ, U.K.*

^b*Center for Sustainable Future Technologies, Istituto Italiano di Tecnologia,
corso Trento 21, 10129 Torino, Italy*

E-mail: P.Di-Bari@soton.ac.uk, michele.refioarentin@iit.it,
R.Samanta@soton.ac.uk

ABSTRACT: We discuss how seesaw neutrino models can be graphically represented in lepton flavour space. We examine various popular models and show how this representation helps understanding their properties and connection with experimental data showing in particular how certain texture zero models are ruled out. We also introduce a new matrix, the bridging matrix, that brings from the light to the heavy neutrino mass flavour basis, showing how this is related to the orthogonal matrix and how different quantities are easily expressed through it. We then show how one can randomly generate orthogonal and leptonic mixing matrices uniformly covering all flavour space in an unbiased way (Haar-distributed matrices). Using the isomorphism between the group of complex rotations and the Lorentz group, we also introduce the concept of Lorentz boost in flavour space for a seesaw model and how this has an insightful physical interpretation. Finally, as a significant application, we consider N_2 -leptogenesis. Using current experimental values of low energy neutrino parameters, we show that the probability that at least one flavoured decay parameter of the lightest right-handed neutrino is smaller than unity is about 49% (to be compared with the tiny probability that the total decay parameter is smaller than unity, $P(K_1 < 1) \sim 0.1\%$, confirming the crucial role played by flavour effects). On the other hand when $m_1 \gtrsim 0.1$ eV this probability reduces to less than 5%, showing how also N_2 -leptogenesis disfavors degenerate light neutrinos.

KEYWORDS: Beyond Standard Model, Cosmology of Theories beyond the SM, CP violation, Neutrino Physics

ARXIV EPRINT: [1812.07720](https://arxiv.org/abs/1812.07720)

Contents

1	Introduction	1
2	Representing seesaw models in lepton flavour space	3
3	Motion in lepton flavour space	10
3.1	Random generation of Haar-distributed U	11
3.2	Random generation of Haar-distributed Ω	12
3.3	An application: N_2 -leptogenesis	15
4	Conclusion	18
A	Excluding models with two heavy neutrino flavours coinciding with one charged lepton flavour	18
B	Orthogonal matrix for a generic boost in flavour space	19

1 Introduction

The possibility to identify the origin of neutrino masses and mixing clashes with the limited number of low energy neutrino parameters that we can access experimentally, those encoded in the light neutrino mass matrix (three mixing angles, three neutrino masses, one Dirac phase, two Majorana phases¹), in comparison with the large number of theoretical parameters typically introduced by models of new physics. Even within a minimal extension of the Standard Model explaining neutrino masses and mixing, the type I seesaw mechanism [1–6], there are far too many parameters to obtain definite predictions. This is true unless:

- (i) either this is embedded within a theoretical framework able to reduce the number of parameters (top-down approach);
- (ii) or an explanation of neutrino masses and mixing is linked to other observables (bottom-up approach) such as the matter-antimatter asymmetry of the universe with leptogenesis, parameters in the quark sector (as in grand-unified theories), lepton flavour violating processes (within different models), dark matter of the universe (from heavy-heavy neutrino mixing or from light-heavy neutrino mixing);
- (iii) or some combination of (i) and (ii) is realised; in this case the top-down and bottom-up approaches are complemented and both help to increase the predictive power.

¹The two Majorana phases are not fully measurable. However, a positive signal in neutrinoless double beta decay experiments would provide an experimental relation between them, placing constraints.

In the case of a pure bottom-up approach one would like to draw model independent conclusions based just on the experimental information. From this point of view a useful and widely used tool is the orthogonal parameterisation of the neutrino Dirac mass matrix within type-I seesaw mechanism since it allows to separate in a unambiguous way the light neutrino parameters, three light neutrino masses and six mixing parameters, from the heavy neutrino parameters (in the most attractive case of three heavy neutrinos one has three heavy neutrino masses and six parameters in the orthogonal matrix). Scans within this parameterisation within a particular model or imposing certain constraints such as successful leptogenesis can lead to interesting bounds on low energy neutrino parameters or even to specific predictions. They can also be used to study the impact of including specific effects in the calculation of the asymmetry or the validity of certain approximations.

In this paper we are interested in introducing new general tools for the study and understanding of seesaw models, in particular how these can be represented in lepton flavour space and randomly generated in an unbiased way. In section 2 we show how different models can be graphically represented in flavour space and how this helps understanding quite easily different properties or aspects of the model, for example whether it can reproduce successfully or not the experimental constraints. In particular we show how certain models with textures zeroes are now excluded by the experimental data. We also review how the parameters in the orthogonal matrix relate the light neutrino masses to the heavy neutrino masses and contain direct information on how fine tuned are the light neutrino masses from the seesaw formula. We introduce a new matrix, the *bridging matrix*, that relates in a simple way the light neutrino mass eigenstates to the lepton states produced by the decays of the heavy neutrino mass eigenstates. In section 3 we discuss a new parameterisation of the orthogonal matrix and of the leptonic mixing matrix such that if no experimental information is imposed, a random uniform generation of the parameters produces light and heavy neutrino flavours that cover uniformly all lepton flavour space without favouring any particular flavour direction or region. This new parameterisation is based on the isomorphism of the group of complex rotations with the restricted Lorentz group. In this way we introduce the concept of Lorentz boost in flavour space and, therefore, of motion of a model in flavour space with a specified velocity and along a certain direction in flavour space. This should be meant not as a continuous evolution in flavour space but rather as a property characterising each flavour model itself. In particular we show that models *at rest* in flavour space correspond to models with minimal fine-tuning.

We also apply this new parameterisation to leptogenesis, showing how in this way the distributions of all flavour decay parameters are identical if no experimental information on the low energy neutrino parameters is imposed and how these change when current experimental information is imposed. In particular, we consider the lightest right-handed (RH) neutrino flavoured decay parameters that play a special role in N_2 -leptogenesis. We are able to show, using latest measurements of neutrino mixing angles, how the probability that at least one of the lightest RH neutrino flavoured decay parameters is less than unity is $\sim 49\%$. Since this condition determines approximately whether the asymmetry produced by the next-to-lightest RH neutrino decays in that flavour can survive the lightest RH

neutrino wash-out, this result shows how successful N_2 -leptogenesis does not require special conditions at all. Finally, in section 4, we draw the conclusions.

2 Representing seesaw models in lepton flavour space

We consider a traditional extension of the standard model introducing N right-handed neutrinos N_{RJ} ($J = I, II, \dots, N$) with Yukawa couplings h^ν and, allowing for lepton number violation, with Majorana mass matrix M . In the flavour basis where the Majorana mass term and the charged lepton Yukawa matrices are both diagonal, the Yukawa interactions terms for neutrinos and charged leptons plus the Majorana mass term can be written as

$$-\mathcal{L}_{Y+M}^{\nu+\ell} = \overline{L}_\alpha h_{\alpha\alpha}^\ell \ell_{R\alpha} \Phi + \overline{L}_\alpha h_{\alpha J}^\nu N_{RJ} \tilde{\Phi} + \frac{1}{2} \overline{N_{RJ}^c} M_J N_{RJ} + \text{h.c.}, \quad (2.1)$$

where $L^T \equiv (\nu_L, \alpha_L)$ are the leptonic doublets, $M_I \leq \dots \leq M_N$ are the heavy neutrino masses and we indicated with Greek indexes the charged lepton flavours, $\alpha = e, \mu, \tau$, and with Roman indexes the heavy neutrino flavours, $J = I, II, \dots, N$. After spontaneous symmetry breaking the Higgs vev generates Dirac masses $m_D = v h^\nu$ and $m_\alpha = v h_{\alpha\alpha}^\ell$ respectively for neutrinos and charged leptons so that the total mass term of the Lagrangian for neutrinos and charged leptons can be written as

$$-\mathcal{L}_m^{\ell+\nu} = \overline{\alpha}_L m_\alpha \alpha_R + \overline{\nu}_{L\alpha} m_{D\alpha J} N_{RJ} + \frac{1}{2} \overline{N_{RJ}^c} M_J N_{RJ} + \text{h.c.} \quad (2.2)$$

In the limit $M \gg m_D$, the light neutrino mass matrix is given by the seesaw formula [1–6]

$$m_{\nu\alpha\beta} = -m_{D\alpha J} M_J^{-1} m_{D\beta J}. \quad (2.3)$$

This is diagonalised by the (unitary) leptonic mixing matrix U in a way that $m_{\nu\alpha\beta} = -U_{\alpha i} D_{m_{ij}} U_{\beta j}$, where $D_m \equiv \text{diag}(m_1, m_2, m_3)$. The light neutrino masses $m_1 \leq m_2 \leq m_3$ can then be expressed as

$$m_i = U_{i\alpha}^* m_{D\alpha J} M_J^{-1} (m_D^T)_{J\beta} U_{\beta i}^*. \quad (2.4)$$

This expression is equivalent to the orthogonality of the matrix [7]

$$\Omega_{iJ} = \frac{(U^\dagger m_D)_{iJ}}{\sqrt{m_i M_J}}, \quad (2.5)$$

that provides a useful (*orthogonal*) parameterisation of the neutrino Dirac mass matrix

$$m_{D\alpha J} = U_{\alpha i} \sqrt{m_i} \Omega_{iJ} \sqrt{M_J}. \quad (2.6)$$

The orthogonal matrix elements $\Omega_{iJ} = |\Omega_{iJ}| e^{i \frac{\varphi_{iJ}}{2}}$ have an important physical meaning [8]. They provide the fractional contribution to the light neutrino mass m_i from the term proportional to the inverse heavy neutrino mass M_J^{-1} and also, very importantly, they tell

how fine-tuned are phase cancellations in the seesaw formula to get each m_i as a sum of terms $\propto M_J^{-1}$. Indeed, it is simple to express each light neutrino mass m_i as²

$$m_i = \bar{m}_i \sum_J r_{iJ} e^{i\varphi_{iJ}}, \quad (2.7)$$

where each $r_{iJ} \equiv |\Omega_{iJ}^2| / \sum_J |\Omega_{iJ}^2| \propto 1/M_J$ is the fractional contribution to the neutrino mass m_i from the heavy neutrino inverse mass M_J^{-1} , and $\bar{m}_i \equiv m_i \sum_J |\Omega_{iJ}^2|$. In this way the quantities $\gamma_i \equiv \sum_J |\Omega_{iJ}^2| \geq 1$ can be regarded as a measure of the fine-tuning, from phase cancellations, that is required to reproduce the light neutrino masses m_i .

If we indicate with $|L_J\rangle$ the lepton quantum state produced (at tree level) in the decay of a RH neutrino N_J , its charged lepton flavour composition is determined by the neutrino Dirac mass matrix [9]

$$|L_J\rangle = \frac{m_{D\alpha J}}{\sqrt{(m_D^\dagger m_D)_{JJ}}} |L_\alpha\rangle. \quad (2.8)$$

If we use the leptonic mixing matrix U to express the charged lepton flavour eigenstates in terms of the neutrino mass eigenstates, $|L_\alpha\rangle = U_{\alpha i}^* |L_i\rangle$, we obtain

$$|L_J\rangle = \frac{m_{D\alpha J} U_{\alpha i}^*}{\sqrt{(m_D^\dagger m_D)_{JJ}}} |L_i\rangle = \frac{(U^\dagger m_D)_{iJ}}{\sqrt{(m_D^\dagger m_D)_{JJ}}} |L_i\rangle, \quad (2.9)$$

showing that the matrix

$$B_{iJ} \equiv \frac{(U^\dagger m_D)_{iJ}}{\sqrt{(m_D^\dagger m_D)_{JJ}}} \quad (2.10)$$

operates the transformation between the lepton flavour basis determined by the neutrino mass eigenstates to that one determined by heavy neutrino lepton flavour states.³ In terms of the orthogonal matrix one finds easily

$$B_{iJ} = \sqrt{\frac{m_i}{\tilde{m}_J}} \Omega_{iJ} = \frac{\sqrt{m_i} \Omega_{iJ}}{\sqrt{\sum_k m_k |\Omega_{kJ}|^2}}, \quad (2.11)$$

where we introduced the effective neutrino masses [10–12]

$$\tilde{m}_J \equiv \frac{(m_D^\dagger m_D)_{JJ}}{M_J} = \sum_k m_k |\Omega_{kJ}|^2. \quad (2.12)$$

This shows that for $N = 3$ the matrix B contains nine parameters: the three light neutrino masses and the six parameters in the orthogonal matrix. These are indeed the 3×3

²Notice that using the orthogonality of Ω one can write $m_i = m_i \sum_J \Omega_{iJ}^2$ and from this and the definition of φ_{iJ} one obtains (2.7). From eq. (2.5) one can see that $r_{iJ} \propto 1/M_J$.

³If one considers the lepton doublet fields, rather than the states, one has $L_J = B_{Ji}^* L_i$. Notice also that the matrix $U^\dagger m_D$ is the Dirac neutrino mass matrix in a flavour basis where both light neutrino and heavy neutrino mass matrices are diagonal [12]. The B matrix is obtained properly normalising $U^\dagger m_D$ and it basically bridges the energy gap between low and high energy states, more precisely bringing from low to high energy states. For this reason it could be also referred to as the *beanstalk matrix*, from the beanstalk narrated in the *Story of Jack and the Beanstalk*.

parameters necessary to determine the flavour compositions of the heavy neutrino flavour states $|L_J\rangle$.

The probability that a lepton L_J is measured as a lepton $L_{I \neq J}$ or, equivalently, the interference probability between a heavy neutrino N_J and a heavy neutrino $N_{I \neq J}$, can be simply expressed (at tree level) in terms of B as

$$p_{IJ}^0 \equiv |\langle L_J | L_I \rangle|^2 = \left| \sum_k B_{kJ}^* B_{kI} \right|^2, \quad (2.13)$$

and one can immediately verify from eq. (2.10) that $p_{JJ}^0 = 1$. The nought in the upper script indicates that they are calculated at tree level. On the other hand the probability that a lepton L_I is measured in a charged lepton flavour $\alpha = e, \mu, \tau$ is given by

$$p_{J\alpha}^0 \equiv |\langle L_\alpha | L_J \rangle|^2 = \frac{|m_{D\alpha J}|^2}{(m_D^\dagger m_D)_{JJ}} = \left| \sum_k U_{\alpha k} B_{kJ} \right|^2 = \frac{|\sum_k \sqrt{m_k} U_{\alpha k} \Omega_{kJ}|^2}{\sum_k m_k |\Omega_{kJ}|^2}. \quad (2.14)$$

These expressions for the probabilities clearly show the physical meaning of the B matrix as a transformation matrix between the light and the heavy neutrino flavour basis.

The seesaw formula is invariant under a generic unitary flavour transformation of the LH fields $\nu_{L\alpha'} = V_{L\alpha'\alpha} \nu_{L\alpha}$, so that one can write in the new flavour basis

$$m'_\nu = -m'_D \frac{1}{D_M} m_D'^T, \quad (2.15)$$

where $D_M \equiv \text{diag}(M_I, M_{II}, \dots, M_N)$ and $m'_{D\alpha'J} = V_{L\alpha'\alpha} m_{D\alpha J}$, while the transformed light neutrino mass matrix is given by $m'_{\nu\alpha'\beta'} = V_{L\alpha'\alpha} m_{\nu\alpha\beta} (V_L^T)_{\beta\beta'}$. In this new basis the charged lepton mass matrix is in general non-diagonal. The orthogonal matrix Ω and the bridging matrix B are of course invariant under this change of lepton flavour basis, since they are by definition transformations between the light and the heavy neutrino flavour bases and, therefore, are independent of which lepton flavour basis is chosen to represent the lepton fields and neutrino Dirac mass matrix. Therefore, in terms of the transformed Dirac mass matrix, they can be simply written as

$$\Omega_{iJ} = \frac{(W^\dagger m'_D)_{iJ}}{\sqrt{m_i M_J}}, \quad B_{iJ} = \frac{(W^\dagger m'_D)_{iJ}}{\sqrt{(m_D'^\dagger m'_D)_{JJ}}}, \quad (2.16)$$

where we introduced the unitary matrix $W_{\alpha'i} \equiv V_{\alpha'\alpha} U_{\alpha i}$ that brings from the light neutrino mass basis to the new generic primed flavour basis.

Neutrino Yukawa basis. A particularly important example of lepton flavour basis, useful especially to describe a model, is represented by the neutrino Yukawa basis. This is the basis where the neutrino Dirac mass matrix is diagonal. In general the change to this basis has to be done transforming simultaneously both the LH neutrino fields and the RH neutrino fields by means of a bi-unitary transformation, $\nu_{L\ell} = V_{L\ell\alpha}^Y \nu_{L\alpha}$ and $N_{R\ell}^Y = U_{R\ell I}^Y N_{RI}$ respectively ($\ell = a, b, c$), in a way that

$$m_D = V_L^{Y\dagger} D_{m_D} U_R^Y, \quad (2.17)$$

where $D_{m_D} \equiv \text{diag}(m_{Da}, m_{Db}, m_{Dc})$ and $m_{Da} \leq m_{Db} \leq m_{Dc}$ are the Dirac masses. The Yukawa basis has important physical properties. First of all whether the leptonic mixing matrix is generated either in the LH sector or in the RH sector is clearly something encoded by V_L^Y and U_R^Y respectively. If there is no right-right Majorana mass term (the Dirac neutrino case) then the light neutrino masses would be simply given by the Dirac masses (i.e., $m_1 = m_a, m_2 = m_b, m_3 = m_c$) and the leptonic mixing matrix would be simply given by $U = V_L^{Y\dagger}$: the Yukawa basis would simply coincide with the light neutrino mass basis. This would be still true when the Majorana mass term is turned on, the case of our interest, and $U_R^Y = I$, corresponding to say that the Majorana and the Dirac mass matrices are diagonalised in the same basis. The only difference would be that in this case one has seesawed neutrino masses $m_i = m_{D\ell}^2/M_I$ with $\ell = a, b, c$. Vice-versa, if $V_L^Y = I$, then leptonic mixing can only stem by a $U_R^Y \neq I$, as it can be immediately understood from the see-saw formula. Another important physical property of the Yukawa basis is that it sets the right basis where to describe medium effects in the description of RH-RH neutrino mixing in the early universe, proposed for example to be either the source of baryogenesis in the ARS mechanism [13] or of dark matter-genesis in [14], since the effective potential due to medium effects are diagonal in the Yukawa basis. Therefore, the RH neutrino mixing matrix should be identified with U_R^Y , at least in the absence of other (non-standard) RH neutrino interactions. Finally, notice that the neutrino Yukawa basis provides clearly the reference basis to compare the neutrino Yukawa interactions with those of other massive fermions and in case impose certain relations as in SO(10)-inspired models [11, 15–18] where the neutrino Dirac mass matrix is ‘not too different’ from the up quark Dirac mass matrix. Also, as we will see, often this is the right basis where to impose certain conditions rising from symmetries of the model, such as textures zeros or other relations on the mass matrices of other fermions.

Let us now consider a few interesting examples of lepton flavour bases associated to specific classes of models. These are graphically shown in the panels of figure 1 where we used the light neutrino flavour basis as a reference frame.

Charged lepton flavour basis. In panel (a) we show the usual charged lepton flavour basis and how this can be obtained, modulo the three phases, from the light neutrino flavour basis by means of three Euler rotations defining the three mixing angles in the leptonic mixing matrix [19].

Generic heavy neutrino flavour basis. In panel (b) we show a generic heavy lepton flavour basis that, in general, is not orthonormal. We have defined the angles θ_{IJ} simply in such a way that $p_{IJ}^0 = \cos \theta_{IJ}$. If the heavy neutrino flavour basis is orthonormal, then $p_{IJ}^0 = \delta_{IJ}$ and the equation (2.13) correctly shows that in this case B is unitary (however, we show in the following that in this case it has necessarily to coincide with the identity or any permutation matrix).

Coincident light and heavy neutrino flavour bases. A special case, shown graphically in panel (c), corresponds to have $B = P$ (where P here and elsewhere is the permutation matrix), so that the heavy neutrino flavour basis basically coincides with the light




Figure 1. Examples of lepton flavour bases with the light neutrino flavour basis as reference basis.

neutrino flavour basis. In this case one can easily see that necessarily also both $\Omega = P$ and $U_R^Y = P$. This corresponds to the situation described before when the neutrino masses are given simply by $m_i = m_{D\ell}^2/M_J$ and $U = V_L^{Y\dagger}$. Therefore, in this case necessarily the Yukawa basis also coincides (modulo axes permutations) with the light and heavy neutrino flavour bases. Indeed, it is correct to say that, since the heavy neutrino flavour basis is aligned with the Yukawa basis, then the resulting light neutrino basis, from the seesaw formula, is also coinciding. This situation corresponds to what has been called *limit of exact dominance* in [20] or *form dominance* in [21]. In this case heavy neutrinos do not mix and do not interfere in decays and indeed all CP asymmetries, both total [20] and flavoured [21], vanish. For this reason some departure from form dominance is necessary if one wants to realise leptogenesis. This class of models typically emerges when a non-Abelian flavour symmetry is imposed [23] in a way that $D_N(g) m_D^\dagger m_D D_N(g) = m_D^\dagger m_D$, where $D_N(g)$ is 3-dim irreducible representation of the non-Abelian flavour symmetry group G acting on the RH neutrinos and g is a generic group element. In this case the first Shur's lemma implies $m_D^\dagger m_D = \lambda_D^2 P$, where P is the permutation matrix and λ_D is the value of the degenerate Dirac neutrino masses, in a way that $m_i = \lambda_D^2/M_J$ realising form dominance corresponding indeed to $\Omega = P$. From eq. (2.5) one can see that the fact that $V_L^Y = U^\dagger$ is consistent with having $\Omega = P$ (and from eq. (2.10) that $B = P$). In order to have successful leptogenesis the flavour symmetry has to be broken and the CP asymmetries are related to the symmetry breaking parameter [22, 23].

One can wonder whether there can be models, generalising $\Omega = B = P$, characterised by a generic orthonormal heavy neutrino flavour basis that does not coincide with the light neutrino flavour basis. However, it is easy to show that this is impossible. The reason is that if the heavy neutrino flavour basis is orthonormal, then this has necessarily to coincide with the Yukawa basis since one can always find a matrix V_L that brings to a basis where m'_D is diagonal and, therefore, this necessarily implies $U_R = P$. However, in this case from the seesaw formula one immediately finds $V_L = U_L^\dagger$ and, therefore, the heavy neutrino flavour basis has necessarily to coincide with the light neutrino flavour basis, as confirmed also by the fact that one has $\Omega = B = P$.

Three coinciding heavy neutrino flavours. An opposite limit case, shown in panel (d), is realised when all three lepton heavy neutrino flavours coincide, i.e. $I = II = III = e'$, meaning that all three heavy neutrinos decay into leptons with the same flavour e' . It is easy to prove that this case is excluded by the experimental data since one can always perform a transformation, operated by a unitary matrix V'_L acting on the lepton doublets, from the (charged lepton) flavour basis (e, μ, τ) to a new orthonormal flavour basis (e', μ', τ') where e' coincides then with the common heavy neutrino flavour. In this new flavour basis the neutrino Dirac mass matrix takes the very simple form

$$m'_D = V'_L m_D = \begin{pmatrix} m_{De'I} & m_{De'II} & m_{De'III} \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}, \tag{2.18}$$

where V'_L is a unitary matrix that transforms the lepton doublets from the charged lepton flavour basis to the new flavour basis. From the seesaw formula one can see immediately that this form implies $m_1 = m_2 = 0$,⁴ and therefore this case is excluded since it cannot reproduce both solar and atmospheric neutrino mass scales.

Two coinciding heavy neutrino flavours. We can now consider a less special case where only two lepton heavy neutrino flavours coincide, while the third does not and is generic. For example, we can consider $I = II$. In this case we can always find a transformation, still operated by a unitary matrix V'_L , from (e, μ, τ) to a new orthonormal flavour basis (e', μ', τ') where $e' = I = II$. This case is shown graphically in panel (e) of figure 1. In this new flavour basis the Dirac mass matrix takes the form

$$m'_D = V'_L m_D = \begin{pmatrix} m_{De'I} & m_{De'II} & m_{De'III} \\ 0 & 0 & m_{D\mu'III} \\ 0 & 0 & m_{D\tau'III} \end{pmatrix}. \quad (2.19)$$

This form for m'_D can successfully reproduce all low energy neutrino data for a generic e' . However, if the flavour e' coincides with one of the charged lepton flavours (in this case $V'_L = P$), then the number of parameters gets considerably reduced and one has to verify for each case, whether it is possible to reproduce the low energy neutrino data. For example, if $e' = e$, then one obtains a seesaw model that implies a light neutrino mass matrix of the form respecting the so-called *strong scaling ansatz* [27, 28], leading necessarily to a vanishing θ_{13} now excluded by the data. This is only one out of nine cases corresponding to have $|L_I\rangle = |L_J\rangle = |L_\alpha\rangle$ with $I \neq J$ and $\alpha = e, \mu, \tau$. By inspection we have checked that also all the other eight cases, listed explicitly in appendix A, are excluded, since they give rise to a light neutrino mass matrix that is either again respecting the scaling ansatz made in [27] or has some similar scaling property also leading to unacceptable low energy neutrino data (see appendix A for more details).

A popular class of seesaw models where the number of parameters is considerably reduced is the two right-handed neutrino limit [29–32]. This can be obtained from the three RH neutrino case either in the limit of very large heaviest RH neutrino mass $M_3 \gg 10^{15}$ GeV or if one of the three RH neutrinos has negligible Yukawa couplings. In both cases one has $m_1 \rightarrow 0$ and effectively the heaviest RH neutrino decouples from the seesaw formula. In this case one effectively obtains a two RH neutrino formula with a 3×2 Dirac neutrino mass matrix. In this case the number of seesaw parameters reduces from eighteen to eleven. These are still too many to lead to predictions on the mixing parameters and usually one has to add some additional condition to this extent. For example, one could again consider a situation when both the two heavy neutrino flavours are aligned. However, analogously to the three RH neutrino case where all three heavy neutrino flavours are aligned, one would get a second vanishing light neutrino, so that one cannot reproduce both the solar and the atmospheric neutrino mass scales. Within these two RH neutrino models one can further reduce the number of parameters again imposing texture zeros in the neutrino Dirac mass matrix m_D , i.e., in the charged lepton flavour basis. In this case it has been shown that

⁴This is something expected since the matrix (2.18) has rank 1.

models with more than two texture zeros are all ruled out by the data and even among all possible models with two texture zeroes only one is still marginally allowed since it requires inverted hierarchy, now disfavoured at approximately 3σ [33], while all possibilities leading to normal hierarchical neutrino masses do not reproduce the measured values of the mixing angles [34, 35].

Lepton flavour basis leading to two texture zeros in the Dirac mass matrix.

Finally, let us conclude saying that of course one can always find a flavour basis (e', μ', τ') where m'_D has two textures zero, since one can always align one flavour along one of the heavy neutrino flavours, for example in a way that $e' = I$ as represented in panel (f) of figure 1.

3 Motion in lepton flavour space

The orthogonal parameterisation (see eq. (2.5)) is a useful tool that allows to scan over the (unknown) parameters in the orthogonal matrix and the RH neutrino masses taking into account the experimental information from low energy neutrino experiments also in combination with other phenomenological conditions (e.g., successful leptogenesis, reproducing the observed dark matter abundance, respecting constraints on rates of lepton flavour violating processes).

The scans are traditionally done using a parameterisation of the leptonic mixing matrix in terms of three Euler rotations (two real ones and one complex), defining the three mixing angles θ_{ij} , the CP violating Dirac phase δ and two CP violating Majorana phases ρ and σ , explicitly [19]

$$U = \begin{pmatrix} 1 & 0 & 0 \\ 0 & c_{23} & s_{23} \\ 0 & -s_{23} & c_{23} \end{pmatrix} \begin{pmatrix} c_{13} & 0 & s_{13} e^{-i\delta} \\ 0 & 1 & 0 \\ -s_{13} e^{i\delta} & 0 & c_{13} \end{pmatrix} \begin{pmatrix} c_{12} & s_{12} & 0 \\ -s_{12} & c_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix} \begin{pmatrix} e^{i\rho} & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & e^{i\sigma} \end{pmatrix} \quad (3.1)$$

where $s_{ij} \equiv \sin \theta_{ij}$ and $c_{ij} \equiv \cos \theta_{ij}$. The orthogonal matrix is usually analogously parameterised as the product of three complex rotations,

$$\Omega = \zeta \begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos z_{23} & \sin z_{23} \\ 0 & -\sin z_{23} & \cos z_{23} \end{pmatrix} \begin{pmatrix} \cos z_{13} & 0 & \sin z_{13} \\ 0 & 1 & 0 \\ -\sin z_{13} & 0 & \cos z_{13} \end{pmatrix} \begin{pmatrix} \cos z_{12} & \sin z_{12} & 0 \\ -\sin z_{12} & \cos z_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.2)$$

where the z_{ij} 's are three complex mixing angles and the overall sign, $\zeta = \pm 1$, takes into account two possible different options (branches), one with positive determinant and one with negative determinant. The three complex mixing angles can in turn be parameterised in terms of their real and imaginary parts writing $z_{ij} = x_{ij} + i y_{ij}$.

Points in flavour space, for a fixed set of light and heavy neutrino masses, can then be obtained generating random uniformly the values of the three mixing angles, the three phases, the three real and imaginary parts of the complex angles. The result of this random generation of points in flavour space is shown in the top-left panel of figure 2 for arbitrary values of the mixing angles and of the low energy phases, i.e., without imposing




Figure 2. Density distributions of flavour probabilities $p_{I\alpha}^0$ obtained by a random generation of U and Ω : in the top-left panel mixing angles have been generated randomly uniformly while in the top-right panel U and Ω have been generated according to the Haar measure as explained in the body text. In the bottom panels the mixing angles have been generated Gaussianly using the experimental results still for a uniform generation of Ω complex angles in the left panel or according to the Haar measure in the right panel.

any experimental constraint. More precisely we show, with triangular plots, the probability density distribution in the space of the charged lepton flavour probabilities $p_{I\alpha}^0$ ($\alpha = e, \mu, \tau$) for the lightest RH neutrino. As one can see, though we randomly uniformly generated the values of the parameters, the distribution exhibits a strong inhomogeneity toward large values of p_{Ie}^0 . This is of course an unpleasant feature if one wants to get unbiased flavour distributions where the flavour dependence originates only from the experimental data and/or from the properties of a model and is not an artefact of how the random generation of points is performed. Some basic results of group theory help explaining why this happens with the usual parameterisation and how the problem can be fixed but at the same time they will provide an insightful way to look at seesaw models in lepton flavour space. If one looks at the expression (2.14) in terms of U and Ω , then it is clear that the problem is that the usual parameterisations does not give a uniform distribution of the elements of U and Ω . In order to do that one has to generate random matrices in a way to cover uniformly the flavour space. Let us discuss separately how this can be done for U and Ω .

3.1 Random generation of Haar-distributed U

We want to generate U matrices in a way not to privilege any particular lepton flavour basis. Let us look again at the panel (a) in figure 1. Here the U matrix is regarded as a (proper) Euler rotation, an approximate picture that is valid only when phases are neglected. A




Figure 3. Distributions of real and imaginary parts of the entries of Haar-distributed U .

flavour unbiased random generation of weak lepton flavour bases, has to be such that given a certain flavour axis, for example the tauon axis, this points to any infinitesimal surface element on the unit sphere in lepton flavour space with equal probability. In this way, generating a large number of lepton flavour basis, each lepton axis will uniformly cover the unit sphere in lepton flavour space. This can be done using well known results of group theory that we briefly discuss [36].

Each (real) U matrix is an element of the group of real rotations $SO(3, \mathbb{R})$. When phases are taken into account each randomly generated U is an element of the group of unitary transformations $U(3)$. Therefore, in group theory language, a flavour unbiased random generation of U corresponds to generate randomly unitary matrices according to the Haar measure over the group $U(3)$ (*Haar-distributed random matrices*) that is given by

$$dV \equiv d(\sin^2 \theta_{12}) d(\sin^2 \theta_{23}) d(\cos^4 \theta_{13}) d\delta d\rho d\sigma. \quad (3.3)$$

In this way generating uniformly $\sin^2 \theta_{12}$, $\sin^2 \theta_{23}$ and $\cos^4 \theta_{13}$ in the interval $[0, 1]$, one obtains equal distributions for all $U_{\alpha i}$ elements, both for their real parts and for their imaginary parts, as shown in figure 3. Notice that the use of Haar-distributed U , and more generally light neutrino mass matrices, is the basis of anarchical prediction of low energy neutrino parameters [37, 38]. However, for us, more pragmatically, this is a way to generate flavoured unbiased scans over seesaw models.

3.2 Random generation of Haar-distributed Ω

If one uses the experimental information on U and, therefore, the experimental distribution for the mixing angles, the random generation of U is not actually necessary, except for the phases that however can be simply generated uniformly between 0 and 2π considering that the Haar measure is flat in the phases.

It is then actually more important to generate randomly Ω 's in a way not to introduce any bias in flavour space. Complex orthogonal matrices, as in eq. (3.2), provide a matrix representation of the Lie group $O(3, \mathbb{C})$ of complex rotations, the complex orthogonal group. It is well known that the special group of complex rotations $SO(3, \mathbb{C})$, i.e. those with

determinant +1, is isomorphic to the restricted (proper + orthochronous) Lorentz group $SO^+(3, 1)$. This can be seen showing that they have the same Lie algebra (see appendix B).⁵

For this reason a generic complex rotation matrix Ω with $\det(\Omega) = +1$ can be decomposed as

$$\Omega(z_{12}, z_{13}, z_{23}) = R(\alpha_{12}, \alpha_{13}, \alpha_{23}) \cdot \Omega_{\text{boost}}(\vec{\beta}), \quad (3.4)$$

where R is a real orthogonal matrix with $\det(R) = +1$ parameterised in terms of three Euler angles α_{ij} and Ω_{boost} is a pure Lorentz boost (in flavour space) parameterised in terms of a boost velocity vector $\vec{\beta} = \beta \hat{n}$ with an associated Lorentz factor $\gamma \equiv (1 - \beta^2)^{-1/2}$. For example, if one chooses a unit vector $\hat{n} = (0, 0, 1)$, then one simply has⁶

$$\Omega_{\text{boost}}(0, 0, \beta) = \begin{pmatrix} \cosh \psi & -i \sinh \psi & 0 \\ i \sinh \psi & \cosh \psi & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.5)$$

with $\beta = \tanh \psi$ and $\gamma = \cosh \psi$. This special case can be of course generalised for an arbitrary choice of \hat{n} (see appendix B). It is interesting to notice that for transformations with $\beta \neq 0$ there is a privileged direction in flavour space while transformations with $\beta = 0$ corresponds basically to a flavour invariant situation where $\Omega = B = P$ and the fine-tuning in the seesaw is minimum.⁷ Indeed, notice that the see-saw fine-tuning parameters associated to the light neutrino masses introduced in section 2 are in this case simply given by $\gamma_1 = \gamma_2 = \gamma^2 (1 + \beta^2)$, showing that the Lorentz factor of the transformation is related to the fine-tuning parameters. This is somehow another way to understand why imposing a flavour symmetry leads to $\Omega = B = P$: this is the case corresponding to vanishing velocity in flavour space, meaning that the model does not have any privileged flavour direction.

In the case of the special boost in eq. (3.5), the bridging matrix is given by

$$B_{\text{boost}} = \begin{pmatrix} \frac{\sqrt{m_1} \cosh \psi}{\sqrt{m_1 \cosh^2 \psi + m_2 \sinh^2 \psi}} & \frac{i \sqrt{m_1} \sinh \psi}{\sqrt{m_1 \sinh^2 \psi + m_2 \cosh^2 \psi}} & 0 \\ \frac{i \sqrt{m_2} \sinh \psi}{\sqrt{m_1 \sinh^2 \psi + m_2 \cosh^2 \psi}} & \frac{\sqrt{m_2} \cosh \psi}{\sqrt{m_1 \sinh^2 \psi + m_2 \cosh^2 \psi}} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.6)$$

an example confirming that the heavy lepton flavour basis is in general non-orthonormal.

If in the orthogonal matrix we turn on, in addition to a boost, a real rotation, then in the limit $\beta = 0$ one obtains $\Omega = R(\alpha_{12}, \alpha_{13}, \alpha_{23})$, i.e., in general one does not recover form dominance corresponding to $\Omega = P$. For example, let us consider a simple rotation around the third axis ($\alpha_{13} = \alpha_{23} = 0$), so that simply

$$\Omega = \begin{pmatrix} \cos \alpha_{12} & \sin \alpha_{12} & 0 \\ -\sin \alpha_{12} & \cos \alpha_{12} & 0 \\ 0 & 0 & 1 \end{pmatrix}. \quad (3.7)$$

⁵Notice that while the group $U(3)$ of unitary matrices is compact, the group $O(3, \mathbb{C})$ is not, that is why they have an intrinsically different parameterisation: in the case of a compact group parameters always vary within a finite interval, while in the case of a non compact group parameters can be arbitrarily large. This also leads to an intrinsic different Haar measure for complex orthogonal matrices compared to unitary matrices.

⁶See appendix B for details.

⁷We are barring the real rotation component $R(\alpha_{12}, \alpha_{13}, \alpha_{23})$.

This case still corresponds to a case of minimal fine tuning, since one clearly has $\gamma_1 = \gamma_2 = \gamma_3 = 1$. However, in this case one finds for the bridging matrix

$$B = \begin{pmatrix} \frac{\sqrt{m_1} \cos \alpha_{12}}{\sqrt{m_1 \cos^2 \alpha_{12} + m_2 \sin^2 \alpha_{12}}} & \frac{\sqrt{m_1} \sin \alpha_{12}}{\sqrt{m_1 \sin^2 \alpha_{12} + m_2 \cos^2 \alpha_{12}}} & 0 \\ -\frac{\sqrt{m_2} \sin \alpha_{12}}{\sqrt{m_1 \cos^2 \alpha_{12} + m_2 \sin^2 \alpha_{12}}} & \frac{\sqrt{m_2} \cos \alpha_{12}}{\sqrt{m_1 \sin^2 \alpha_{12} + m_2 \cos^2 \alpha_{12}}} & 0 \\ 0 & 0 & 1 \end{pmatrix}, \quad (3.8)$$

showing that this, in general, does not coincide with the orthogonal matrix (it is not a real rotation matrix) and also that it brings to an heavy lepton flavour basis that is not orthonormal. These kind of models, with a real orthogonal matrix coinciding with a rotation matrix, are interesting since they still correspond to minimal fine-tuning but, for the basis is not orthonormal, there can be in general interference among heavy neutrino flavours so that the flavoured CP asymmetries do not vanish in general. In this way in principle one could have leptogenesis stemming entirely from low energy neutrino phases [39–43]. However, unless one has a strong resonance enhancement, the observed asymmetry is usually not reproduced, implying that one needs to turn a boost on as well. Therefore, it seems that the explanation of the matter-antimatter asymmetry of the universe necessarily requires the existence of some privileged direction in lepton flavour space, corresponding to some mismatch between the bases where the Majorana and the Yukawa mass matrices are diagonal.

If we want again to generate flavour unbiased Ω matrices, it is then clear what we have to do now. First of all one has to generate Haar-distributed rotation matrices $R(\alpha_{12}, \alpha_{13}, \alpha_{23})$ as we did for U . For $SO(3, \mathbb{R})$, the Haar measure is quite simple

$$dV \equiv d(\sin \alpha_{13}) d\alpha_{23} d\alpha_{12}. \quad (3.9)$$

In the case of $\Omega_{\text{boost}}(\vec{\beta})$, it is clear that, for a fixed value of β , we need to generate isotropically unit vectors \hat{n} . For example, one can use polar coordinates and write

$$\hat{n} = (\cos \phi \sin \theta, \sin \phi \sin \theta, \cos \theta), \quad (3.10)$$

with $\theta \in [0, \pi]$ and $\phi \in [0, 2\pi]$. If one generates random uniformly θ and ϕ , one would obtain a clustered density of points near the poles. To generate isotropic orientation of \hat{n} , we use the standard technique of generating uniform points on a surface of a sphere by generating values of θ given by

$$\theta = \cos^{-1}(1 - 2a), \quad (3.11)$$

with the parameter a uniformly generated within the interval $[0, 1]$. Using this random generation procedure we have obtained distributions for the real and imaginary parts of the orthogonal matrix plotted in figure 4 Notice that this procedure can be easily extended to include also matrices Ω with negative determinant.

Finally, we have combined together the flavour unbiased procedure to random generate both U and Ω and again plotted the distribution of probabilities $p_{I\alpha}^0$ shown in the top right panel of figure 2 and, as one can see, this time, barring small statistical fluctuations,




Figure 4. Distribution of the real and imaginary parts of the entries $\Omega_{i,j}$.

we have obtained a perfectly flavour homogeneous distribution of points shown in the top-right panel.

In the bottom panels of figure, 2 we plotted the $p_{I\alpha}^0$ using the experimental information, generating the mixing angles in U random Gaussianly and using the following latest experimental results for the values of the mixing angles in the case of normal ordering [33]

$$\begin{aligned} \theta_{12} &= 33.82^\circ \pm 0.77^\circ, \\ \theta_{13} &= 8.61^\circ \pm 0.12^\circ, \\ \theta_{23} &= 49.7^\circ \pm 1.0^\circ. \end{aligned} \tag{3.12}$$

The inverted ordering case is now disfavoured at more than 3σ and we will not consider it in our following discussion. We have also compared again the case when mixing angles are random uniformly generated (left panel) with the case of Haar-distributed U (right panel). This time one can see that there is not a great difference since in any case the region that is biased is disfavoured by current data.

3.3 An application: N_2 -leptogenesis

The reason why we focused in figure 2 on $p_{I\alpha}^0$, is that the lightest RH neutrino plays a particular role in N_2 -leptogenesis [20, 44]. In this scenario of leptogenesis the current baryon asymmetry, expressed in terms of the baryon-to-photon number ratio at present η_{B0} , can be calculated as [45, 46]

$$\begin{aligned} \eta_{B0} \simeq 0.96 \times 10^{-2} & \left(\varepsilon_{IIe} \kappa(K_{IIe} + K_{II\mu}) e^{-\frac{3\pi}{8} K_{Ie}} \right. \\ & \left. + \varepsilon_{II\mu} \kappa(K_{IIe} + K_{II\mu}) e^{-\frac{3\pi}{8} K_{I\mu}} + \varepsilon_{II\tau} \kappa(K_{II\tau}) e^{-\frac{3\pi}{8} K_{I\tau}} \right). \end{aligned} \tag{3.13}$$

The lightest RH neutrino flavoured decay parameters $K_{I\alpha}$ play clearly a special role since they describe the exponential wash-out from lightest RH neutrino inverse decays and one needs that at least one of them is less than unity for the asymmetry produced by the N_2 -decays at a temperature $T \sim M_2$ to survive at present. The flavoured decay parameters are simply given by $K_{I\alpha} = p_{I\alpha}^0 K_I$, where $K_I = \tilde{m}_I/m_\star$ are the total decay parameters and




Figure 5. Distributions of the decay parameters for $m_1 = 0$ and generic values of the mixing from a random flavour blind generation of U and Ω .

m_\star is the equilibrium neutrino mass. Therefore, one can see the special role played by the $p_{I\alpha}^0$'s in N_2 -leptogenesis.

It is then particularly interesting to understand how special is the condition for the asymmetry produced by N_2 decays to survive at present. This is basically equivalent to understand how special is to have at least one $K_{I\alpha}$, for some lepton flavour α , less than unity.

The flavoured decay parameters are related to the orthogonal matrix through the eq. (2.12) for \tilde{m}_I and, therefore, for each choice of Ω and for a given value of m_1 , one has a corresponding set of values of $K_{I\alpha}$. We have therefore produced the distributions for the values of the $K_{I\alpha}$ for $\alpha = e, \mu, \tau$ adopting the flavoured unbiased procedure, based on the Haar measure, that we discussed. In figure 5 the distributions are shown without imposing any experimental information on the values of the mixing angles that, therefore, vary arbitrarily within $[0, 90^\circ]$ and in the hierarchical limit $m_1 = 0$. It can be seen how the distributions are identical independently of α as a result of the flavour blindness of the procedure we followed to generate randomly U and Ω .⁸ It is important to notice that the probability for each $K_{I\alpha}$ to be less than unity is about 12%, meaning that the probability that at least one $K_{I\alpha}$ is less than unity is approximately 36%.

How do these results change when the experimental information on the mixing angles is used? In figure 6 the distributions for the $K_{I\alpha}$ are now obtained using the experimental information on the mixing angles eq. (3.12). One can see how the fact that the experimental values favour small values of $p_{I\alpha}^0$, translates into a much higher probability, approximately 36%, for K_{Ie} to be less than unity compared to $K_{I\mu}$ and $K_{I\tau}$ whose probability to be less than unity drops to $\sim 6-7\%$. The probability that at least one $K_{I\alpha}$ is less than unity is therefore now about 49%. This result shows how the possibility to have a small lightest RH neutrino wash-out in one of the three flavours, a crucial condition to realise N_2 -leptogenesis, is not special at all (contrarily to some statements made in the literature). It should be also said that on the other hand the probability to have $K_I = \sum_\alpha K_{I\alpha} < 1$ is only 0.1% confirming and quantifying how accounting for flavour effects is crucial for N_2 -leptogenesis [44].

⁸We do not show the distributions for the $K_{II\alpha}$ and for the $K_{III\alpha}$ but they would also be identical since the procedure is flavour blind both to charged lepton flavour and to heavy neutrino flavour.




Figure 6. Same as in figure 5 but using the experimental values of the mixing angles in eq. (3.12).




Figure 7. Same as in figure 5 but for $m_1 = 0.01$ eV.




Figure 8. Same as in figure 6 but for $m_1 = 0.01$ eV.

In figure 7 and figure 8, again for arbitrary and experimental values of the mixing angles respectively, we also show how the distributions change departing from the hierarchical limit, for $m_1 = 0.01$ eV. It can be noticed how all probabilities drop and this is easily explained since $K_I = \tilde{m}_I/m_*$ and $\tilde{m}_I \geq m_1$ so that all decay parameters tend to increase. However, the probability that at least one of the three $K_{I\alpha} < 1$ is still quite large, approximately 23%. Notice also how for arbitrary mixing angles the distributions are still identical in the three flavours. For values $m_1 \gtrsim 0.1$ eV, the probability that at least one $K_{I\alpha} < 1$ drops below 5%. This can somehow be regarded as a kind of extension of the upper bound on neutrino masses holding in N_1 leptogenesis, also to the case of N_2 -leptogenesis, though it should be clear that in this case the upper bound should be interpreted more in a statistical way rather than as an absolute one.

4 Conclusion

We have seen how representing seesaw models in lepton flavour space allows a deeper understanding of different features of seesaw models. In particular different results have an easy graphical interpretation. The new bridging matrix, switching from the light to heavy neutrino lepton basis, is very useful for writing flavour probabilities and characterising seesaw models in an easy way. We have also seen how fine-tuning in seesaw models can be expressed in terms of a Lorentz boost in flavour space and sequential dominated models, characterised by minimal fine-tuning, are those for which the boost velocity is vanishing and therefore are flavour invariant, in agreement with previous results. A deviation from sequential dominated models, turning on some motion in flavour space, produces a deviation from orthonormality that is necessary to have non-vanishing CP decaying asymmetries and successful leptogenesis. We have seen also that one can deviate from sequential dominated models with a pure real rotation orthogonal matrix. This still corresponds to models at rest in flavour space, but again with some deviation from orthonormality producing non-vanishing CP violation and in principle allowing for successful leptogenesis. We have also seen how to generate randomly, in a flavour unbiased way, seesaw models. We have then applied these new tools to N_2 -leptogenesis and showed how it is actually very easy to realise the condition of no wash-out from the lightest RH neutrino, for $\sim 49\%$ of the points once the current values of the mixing angles are used for hierarchical light neutrinos. On the other hand, if $m_1 \gtrsim 0.1 \text{ eV}$, this probability drops to less than 5%, a result that confirms that the exclusion of quasi-degenerate neutrino from current cosmological observations, supports scenarios of minimal leptogenesis based on type-I seesaw and thermal production of RH neutrinos. The new tools and ideas introduced in this work will be very useful in different respects, both in the quest of models of new physics able to explain neutrino masses and mixing and, more pragmatically, in scanning seesaw models within different contexts such as leptogenesis.

Acknowledgments

We wish to thank Steve King for pointing out useful references. PDB acknowledges financial support from the STFC Consolidated Grant L000296/1. RS is supported by a Newton International Fellowship (NF 171202) from Royal Society (U.K.) and SERB (India). This project has received funding/support from the European Union Horizon 2020 research and innovation programme under the Marie Skłodowska-Curie grant agreements number 690575 and 674896. This work was performed in part at Aspen Center for Physics, which is supported by National Science Foundation grant PHY-1607611.

A Excluding models with two heavy neutrino flavours coinciding with one charged lepton flavour

In this appendix we list explicitly the forms of m_D corresponding to the nine cases discussed in section 2 with two heavy neutrino flavours coinciding with a charged lepton flavour, so

that $|L_I\rangle = |L_J\rangle = |L_\alpha\rangle$ with $I \neq J$ and $\alpha = e, \mu, \tau$. These are given by

$$m_D = \begin{pmatrix} m_{DeI} & m_{DeII} & m_{DeIII} \\ 0 & 0 & m_{D\mu III} \\ 0 & 0 & m_{D\tau III} \end{pmatrix}, \begin{pmatrix} m_{DeI} & m_{DeII} & m_{DeIII} \\ 0 & m_{D\mu II} & 0 \\ 0 & m_{D\tau II} & 0 \end{pmatrix}, \begin{pmatrix} m_{DeI} & m_{DeII} & m_{DeIII} \\ m_{D\mu I} & 0 & 0 \\ m_{D\tau I} & 0 & 0 \end{pmatrix}, \\
 \begin{pmatrix} 0 & 0 & m_{D\tau I} \\ m_{D\mu I} & m_{D\mu II} & m_{D\mu III} \\ 0 & 0 & m_{D\tau III} \end{pmatrix}, \begin{pmatrix} 0 & m_{D\mu II} & 0 \\ m_{D\mu I} & m_{D\mu II} & m_{D\mu III} \\ 0 & m_{D\tau II} & 0 \end{pmatrix}, \begin{pmatrix} m_{DeI} & 0 & 0 \\ m_{D\mu I} & m_{D\mu II} & m_{D\mu III} \\ m_{D\tau I} & 0 & 0 \end{pmatrix}, \\
 \begin{pmatrix} 0 & 0 & m_{DeIII} \\ 0 & 0 & m_{D\mu III} \\ m_{D\tau I} & m_{D\tau II} & m_{D\mu III} \end{pmatrix}, \begin{pmatrix} 0 & m_{DeII} & 0 \\ 0 & m_{D\mu II} & 0 \\ m_{D\tau I} & m_{D\tau II} & m_{D\tau III} \end{pmatrix}, \begin{pmatrix} m_{DeI} & 0 & 0 \\ m_{D\mu I} & 0 & 0 \\ m_{D\tau I} & m_{D\tau II} & m_{D\tau III} \end{pmatrix}. \tag{A.1}$$

All these cases are excluded, since they give rise to a light neutrino mass matrix that is either again respecting the scaling ansatz made in [27] or has some similar scaling property also leading to unacceptable low energy neutrino data. Let us give here a few more details. If, for example, $\alpha = e$, then the resulting light neutrino Majorana mass matrix (m_ν) is of a special form obeying *strong scaling ansatz* that is ruled out [27]. Basically this corresponds to a situation when in the light neutrino Majorana matrix one of the rows is c times of an other row, where, ‘ c ’ is a common scale factor which can be expressed as a function of the elements of m_D . For example, if $I = I$ and $J = II$, then the second row is c times the third row. This leads to a vanishing eigenvalue and the corresponding eigenvector has one vanishing entry. This results in vanishing U_{e3} or U_{e1} , depending whether one has inverted or the normal mass ordering. It can be checked that the other two cases, i.e., $(I, J) = (I, III)$ and $(I, J) = (II, III)$, also lead to a form obeying the strong scaling ansatz (though with different scale factors) and, as in the previous case, this results into vanishing U_{e3} or U_{e1} . In the remaining six cases, for $\alpha = \mu, \tau$, we found analogously that either $U_{\alpha 3}$ or $U_{\alpha 1}$ vanishes, again for inverted and normal mass ordering respectively. Since a zero entry in U is excluded by the experimental data, we conclude that all nine cases corresponding to have two coinciding heavy neutrino flavours are ruled out.

B Orthogonal matrix for a generic boost in flavour space

In this appendix we generalise the parameterisation of the orthogonal matrix in terms of three real angles and the three components of the boost velocity. As well known, a generic proper orthochronous Lorentz transformation can be written as $\Lambda = e^{-i(\vec{\alpha} \cdot \vec{J} + \vec{\xi} \cdot \vec{K})}$ where \vec{J} and \vec{K} are respectively the rotation and boost generators of $SO^+(3, 1)$ and obey the Lie algebra⁹

$$[J_i, J_j] = i \epsilon_{ijk} J_k, \tag{B.1}$$

$$[K_i, K_j] = -i \epsilon_{ijk} K_k, \tag{B.2}$$

$$[J_i, K_j] = i \epsilon_{ijk} K_k, \tag{B.3}$$

⁹Both \vec{J} and \vec{K} are represented by 4×4 matrices.

where ϵ_{ijk} is the totally antisymmetric tensor. If we now consider a generic complex rotation Ω belonging to $SO(3, \mathbb{C})$, this can be written as $\Omega = e^{-i(\vec{\alpha} \cdot \vec{L} + \vec{\xi} \cdot \vec{\Sigma})}$, where \vec{L} are the generators of the real rotations and $\vec{\Sigma} = i\vec{L}$ are the generators of the imaginary rotations (hyperbolic rotations), both represented by 3×3 matrices. The generators of $SO(3, \mathbb{C})$ satisfy the Lie algebra

$$[L_i, L_j] = i\epsilon_{ijk} L_k, \tag{B.4}$$

$$[\Sigma_i, \Sigma_j] = -i\epsilon_{ijk} \Sigma_k, \tag{B.5}$$

$$[L_i, \Sigma_j] = i\epsilon_{ijk} \Sigma_k, \tag{B.6}$$

clearly coinciding with that one of $SO(3, 1)^+$, with the identification $\vec{J} \leftrightarrow \vec{L}$ and $\vec{K} \leftrightarrow \vec{\Sigma}$. Therefore, $\vec{\Sigma}$ correspond to the boost generators in $SO(3, \mathbb{C})$. This shows that the two groups are indeed isomorphic and one can find a map between each other. We can then decompose the 3×3 complex orthogonal matrix $\Omega \in SO(3, \mathbb{C})$ as

$$\Omega = R(\vec{\alpha}) \cdot \Omega_{\text{boost}}(\vec{\beta}), \tag{B.7}$$

where the 3×3 matrices $R(\vec{\alpha}) = e^{-i\vec{\alpha} \cdot \vec{L}} \in SO(3, \mathbb{R})$ and $\Omega_{\text{boost}} = e^{-i\vec{\xi} \cdot \vec{\Sigma}}$ are generated by \vec{L} and $\vec{\Sigma}$ respectively and we defined $\vec{\alpha} \equiv (\alpha_{12}, \alpha_{13}, \alpha_{23})$. Since $\vec{\Sigma} = i\vec{L}$, then one has simply $\Omega_{\text{boost}}(\vec{\beta}) = e^{-i(i\vec{\xi}) \cdot \vec{L}}$, showing that boosts are rotations with complex angles $i\vec{\xi}$. In the special case $\vec{\beta} = (0, 0, \beta)$, considered in the body text, one obtains easily eq. (3.5). In general, for $\vec{\beta} = \beta \hat{n}$ and $\hat{n} = (n_1, n_2, n_3)$, one obtains

$$\Omega_{\text{boost}}(\vec{\beta}) = \begin{pmatrix} \cosh \xi + n_1^2(1 - \cosh \xi) & n_1 n_2(1 - \cosh \xi) - i n_3 \sinh \xi & n_1 n_3(1 - \cosh \xi) + i n_2 \sinh \xi \\ n_1 n_2(1 - \cosh \xi) + i n_3 \sinh \xi & \cosh \xi + n_2^2(1 - \cosh \xi) & n_2 n_3(1 - \cosh \xi) - i n_1 \sinh \xi \\ n_1 n_3(1 - \cosh \xi) - i n_2 \sinh \xi & n_2 n_3(1 - \cosh \xi) + i n_1 \sinh \xi & \cosh \xi + n_3^2(1 - \cosh \xi) \end{pmatrix}, \tag{B.8}$$

with $\beta = \tanh \xi$.

Open Access. This article is distributed under the terms of the Creative Commons Attribution License ([CC-BY 4.0](https://creativecommons.org/licenses/by/4.0/)), which permits any use, distribution and reproduction in any medium, provided the original author(s) and source are credited.

References

- [1] P. Minkowski, $\mu \rightarrow e\gamma$ at a Rate of One Out of 10^9 Muon Decays?, *Phys. Lett.* **67B** (1977) 421 [[INSPIRE](#)].
- [2] T. Yanagida, *Horizontal gauge symmetry and masses of neutrinos*, in *Proceedings of the Workshop on Unified Theory and Baryon Number of the Universe*, O. Sawada and A. Sugamoto eds., KEK, (1979), p. 95.
- [3] P. Ramond, *The Family Group in Grand Unified Theories*, in *International Symposium on Fundamentals of Quantum Theory and Quantum Field Theory Palm Coast, Florida, February 25 – March 2, 1979*, pp. 265–280, 1979, [hep-ph/9809459](#) [[INSPIRE](#)].
- [4] M. Gell-Mann, P. Ramond and R. Slansky, *Complex Spinors and Unified Theories*, in *Supergravity*, P. van Nieuwenhuizen and D. Freedman eds., North Holland, Amsterdam, (1979), *Conf. Proc.* C790927 p. 315, PRINT-80-0576.

- [5] R. Barbieri, D.V. Nanopoulos, G. Morchio and F. Strocchi, *Neutrino Masses in Grand Unified Theories*, *Phys. Lett.* **90B** (1980) 91 [INSPIRE].
- [6] R.N. Mohapatra and G. Senjanović, *Neutrino Mass and Spontaneous Parity Nonconservation*, *Phys. Rev. Lett.* **44** (1980) 912 [INSPIRE].
- [7] J.A. Casas and A. Ibarra, *Oscillating neutrinos and $\mu \rightarrow e, \gamma$* , *Nucl. Phys.* **B 618** (2001) 171 [hep-ph/0103065] [INSPIRE].
- [8] S. Lavignac, I. Masina and C.A. Savoy, *Large solar angle and seesaw mechanism: A bottom up perspective*, *Nucl. Phys.* **B 633** (2002) 139 [hep-ph/0202086] [INSPIRE].
- [9] S. Blanchet, P. Di Bari, D.A. Jones and L. Marzola, *Leptogenesis with heavy neutrino flavours: from density matrix to Boltzmann equations*, *JCAP* **01** (2013) 041 [arXiv:1112.4528] [INSPIRE].
- [10] M. Plümacher, *Baryogenesis and lepton number violation*, *Z. Phys.* **C 74** (1997) 549 [hep-ph/9604229] [INSPIRE].
- [11] E. Nezri and J. Orloff, *Neutrino oscillations versus leptogenesis in SO(10) models*, *JHEP* **04** (2003) 020 [hep-ph/0004227] [INSPIRE].
- [12] M. Fujii, K. Hamaguchi and T. Yanagida, *Leptogenesis with almost degenerate Majorana neutrinos*, *Phys. Rev.* **D 65** (2002) 115012 [hep-ph/0202210] [INSPIRE].
- [13] E.K. Akhmedov, V.A. Rubakov and A. Yu. Smirnov, *Baryogenesis via neutrino oscillations*, *Phys. Rev. Lett.* **81** (1998) 1359 [hep-ph/9803255] [INSPIRE].
- [14] A. Anisimov and P. Di Bari, *Cold Dark Matter from heavy Right-Handed neutrino mixing*, *Phys. Rev.* **D 80** (2009) 073017 [arXiv:0812.5085] [INSPIRE].
- [15] A. Yu. Smirnov, *Seesaw enhancement of lepton mixing*, *Phys. Rev.* **D 48** (1993) 3264 [hep-ph/9304205] [INSPIRE].
- [16] W. Buchmüller and M. Plümacher, *Baryon asymmetry and neutrino mixing*, *Phys. Lett.* **B 389** (1996) 73 [hep-ph/9608308] [INSPIRE].
- [17] F. Buccella, D. Falcone and F. Tramontano, *Baryogenesis via leptogenesis in SO(10) models*, *Phys. Lett.* **B 524** (2002) 241 [hep-ph/0108172] [INSPIRE].
- [18] G.C. Branco, R. Gonzalez Felipe, F.R. Joaquim and M.N. Rebelo, *Leptogenesis, CP-violation and neutrino data: What can we learn?*, *Nucl. Phys.* **B 640** (2002) 202 [hep-ph/0202030] [INSPIRE].
- [19] S.F. King, *Neutrino mass models*, *Rept. Prog. Phys.* **67** (2004) 107 [hep-ph/0310204] [INSPIRE].
- [20] P. Di Bari, *Seesaw geometry and leptogenesis*, *Nucl. Phys.* **B 727** (2005) 318 [hep-ph/0502082] [INSPIRE].
- [21] M.-C. Chen and S.F. King, *A_4 See-Saw Models and Form Dominance*, *JHEP* **06** (2009) 072 [arXiv:0903.0125] [INSPIRE].
- [22] E.E. Jenkins and A.V. Manohar, *Tribimaximal Mixing, Leptogenesis and θ_{13}* , *Phys. Lett.* **B 668** (2008) 210 [arXiv:0807.4176] [INSPIRE].
- [23] E. Bertuzzo, P. Di Bari, F. Feruglio and E. Nardi, *Flavor symmetries, leptogenesis and the absolute neutrino mass scale*, *JHEP* **11** (2009) 036 [arXiv:0908.0161] [INSPIRE].

- [24] P.F. Harrison, D.H. Perkins and W.G. Scott, *Tri-bimaximal mixing and the neutrino oscillation data*, *Phys. Lett. B* **530** (2002) 167 [[hep-ph/0202074](#)] [[INSPIRE](#)].
- [25] G. Altarelli and F. Feruglio, *Discrete Flavor Symmetries and Models of Neutrino Mixing*, *Rev. Mod. Phys.* **82** (2010) 2701 [[arXiv:1002.0211](#)] [[INSPIRE](#)].
- [26] S.F. King and C. Luhn, *Neutrino Mass and Mixing with Discrete Symmetry*, *Rept. Prog. Phys.* **76** (2013) 056201 [[arXiv:1301.1340](#)] [[INSPIRE](#)].
- [27] R.N. Mohapatra and W. Rodejohann, *Scaling in the neutrino mass matrix*, *Phys. Lett. B* **644** (2007) 59 [[hep-ph/0608111](#)] [[INSPIRE](#)].
- [28] R. Sinha, R. Samanta and A. Ghosal, *Generalized $\mathbb{Z}_2 \times \mathbb{Z}_2$ in scaling neutrino Majorana mass matrix and baryogenesis via flavored leptogenesis*, *JHEP* **12** (2017) 030 [[arXiv:1706.00946](#)] [[INSPIRE](#)].
- [29] S.F. King, *Large mixing angle MSW and atmospheric neutrinos from single right-handed neutrino dominance and U(1) family symmetry*, *Nucl. Phys. B* **576** (2000) 85 [[hep-ph/9912492](#)] [[INSPIRE](#)].
- [30] P.H. Frampton, S.L. Glashow and T. Yanagida, *Cosmological sign of neutrino CP-violation*, *Phys. Lett. B* **548** (2002) 119 [[hep-ph/0208157](#)] [[INSPIRE](#)].
- [31] P.H. Chankowski and K. Turzyski, *Limits on T_{reh} for thermal leptogenesis with hierarchical neutrino masses*, *Phys. Lett. B* **570** (2003) 198 [[hep-ph/0306059](#)] [[INSPIRE](#)].
- [32] A. Ibarra and G.G. Ross, *Neutrino phenomenology: The case of two right-handed neutrinos*, *Phys. Lett. B* **591** (2004) 285 [[hep-ph/0312138](#)] [[INSPIRE](#)].
- [33] I. Esteban, M.C. Gonzalez-Garcia, A. Hernandez-Cabezudo, M. Maltoni and T. Schwetz, *Global analysis of three-flavour neutrino oscillations: synergies and tensions in the determination of θ_{23} , δ_{CP} and the mass ordering*, *JHEP* **01** (2019) 106 [[arXiv:1811.05487](#)] [[INSPIRE](#)].
- [34] K. Harigaya, M. Ibe and T.T. Yanagida, *Seesaw Mechanism with Occam's Razor*, *Phys. Rev. D* **86** (2012) 013002 [[arXiv:1205.2198](#)] [[INSPIRE](#)].
- [35] D.M. Barreiros, R.G. Felipe and F.R. Joaquim, *Minimal type-I seesaw model with maximally restricted texture zeros*, *Phys. Rev. D* **97** (2018) 115016 [[arXiv:1802.04563](#)] [[INSPIRE](#)].
- [36] K Zyczkowski and M Kus, *Random unitary matrices*, *J. Phys. A* **27** (1994) 4235.
- [37] L.J. Hall, H. Murayama and N. Weiner, *Neutrino mass anarchy*, *Phys. Rev. Lett.* **84** (2000) 2572 [[hep-ph/9911341](#)] [[INSPIRE](#)].
- [38] N. Haba and H. Murayama, *Anarchy and hierarchy*, *Phys. Rev. D* **63** (2001) 053010 [[hep-ph/0009174](#)] [[INSPIRE](#)].
- [39] E. Nardi, Y. Nir, E. Roulet and J. Racker, *The importance of flavor in leptogenesis*, *JHEP* **01** (2006) 164 [[hep-ph/0601084](#)] [[INSPIRE](#)].
- [40] S. Blanchet and P. Di Bari, *Flavor effects on leptogenesis predictions*, *JCAP* **03** (2007) 018 [[hep-ph/0607330](#)] [[INSPIRE](#)].
- [41] S. Pascoli, S.T. Petcov and A. Riotto, *Connecting low energy leptonic CP-violation to leptogenesis*, *Phys. Rev. D* **75** (2007) 083511 [[hep-ph/0609125](#)] [[INSPIRE](#)].
- [42] A. Anisimov, S. Blanchet and P. Di Bari, *Viability of Dirac phase leptogenesis*, *JCAP* **04** (2008) 033 [[arXiv:0707.3024](#)] [[INSPIRE](#)].

- [43] K. Moffat, S. Pascoli, S.T. Petcov and J. Turner, *Leptogenesis from Low Energy CP Violation*, *JHEP* **03** (2019) 034 [[arXiv:1809.08251](#)] [[INSPIRE](#)].
- [44] O. Vives, *Flavor dependence of CP asymmetries and thermal leptogenesis with strong right-handed neutrino mass hierarchy*, *Phys. Rev. D* **73** (2006) 073006 [[hep-ph/0512160](#)] [[INSPIRE](#)].
- [45] P. Di Bari and A. Riotto, *Successful type-I Leptogenesis with SO(10)-inspired mass relations*, *Phys. Lett. B* **671** (2009) 462 [[arXiv:0809.2285](#)] [[INSPIRE](#)].
- [46] P. Di Bari and A. Riotto, *Testing SO(10)-inspired leptogenesis with low energy neutrino experiments*, *JCAP* **04** (2011) 037 [[arXiv:1012.2343](#)] [[INSPIRE](#)].