

Anisotropic global microlocal analysis for tempered distributions

*Original*

Anisotropic global microlocal analysis for tempered distributions / Rodino, L.; Wahlberg, P.. - In: MONATSHFTE FÜR MATHEMATIK. - ISSN 0026-9255. - 202:2(2023), pp. 397-434. [10.1007/s00605-022-01812-z]

*Availability:*

This version is available at: 11583/2974734 since: 2023-10-04T06:16:24Z

*Publisher:*

Springer

*Published*

DOI:10.1007/s00605-022-01812-z

*Terms of use:*

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

*Publisher copyright*

Springer postprint/Author's Accepted Manuscript

This version of the article has been accepted for publication, after peer review (when applicable) and is subject to Springer Nature's AM terms of use, but is not the Version of Record and does not reflect post-acceptance improvements, or any corrections. The Version of Record is available online at: <http://dx.doi.org/10.1007/s00605-022-01812-z>

(Article begins on next page)

# ANISOTROPIC GLOBAL MICROLOCAL ANALYSIS FOR TEMPERED DISTRIBUTIONS

LUIGI RODINO AND PATRIK WAHLBERG

**ABSTRACT.** We study an anisotropic version of the Shubin calculus of pseudodifferential operators on  $\mathbf{R}^d$ . Anisotropic symbols and Gabor wave front sets are defined in terms of decay or growth along curves in phase space of power type parametrized by one positive parameter that distinguishes space and frequency variables. We show that this gives subcalculi of Shubin's isotropic calculus, and we show a microlocal as well as a microelliptic inclusion in the framework. Finally we prove an inclusion for the anisotropic Gabor wave front set of chirp type oscillatory functions with a real polynomial phase function.

## 1. INTRODUCTION

In this paper we study an anisotropic version of Shubin's calculus of pseudodifferential operators on  $\mathbf{R}^d$  [26] and a naturally appearing anisotropic Gabor wave front set.

Shubin symbols for pseudodifferential operators satisfy estimates involving  $1 + |x| + |\xi|$ , and they are thus isotropic on the phase space  $(x, \xi) \in T^*\mathbf{R}^d$ . In particular they behave in a way that does not distinguish between  $x \in \mathbf{R}^d$  and  $\xi \in \mathbf{R}^d$ .

Otherwise expressed, the symbols satisfy growth or decay restrictions on straight lines in phase space of the form  $\mathbf{R}_+ \ni \lambda \mapsto (\lambda x, \lambda \xi)$  for  $(x, \xi) \in T^*\mathbf{R}^d \setminus 0$ . For Shubin operators there are results concerning global microlocal analysis involving the Gabor wave front set, introduced by Hörmander in [14] and elaborated in several recent works [2, 4, 5, 7, 8, 21–23, 25, 27]. The Gabor wave front set detects the lack of superpolynomial decay along straight lines in phase space of the short-time Fourier transform of a tempered distribution. It is global in the sense that it measures smoothness and decay at infinity of the distribution comprehensively. It is empty exactly when a tempered distribution is a Schwartz function.

In this paper we replace the weight  $1 + |x| + |\xi|$  by  $1 + |x| + |\xi|^{\frac{1}{s}}$  where  $s > 0$ . We introduce Shubin type symbols with anisotropic behaviour, with decay or growth along power type curves in phase space of the form

$$(1.1) \quad \mathbf{R}_+ \ni \lambda \mapsto (\lambda x, \lambda^s \xi)$$

for  $(x, \xi) \in T^*\mathbf{R}^d \setminus 0$ .

The idea of anisotropy in pseudodifferential calculus has been around for a long time, cf. [16, 20], with recent contributions exemplified by [10]. These works treat mainly anisotropic behavior in the frequency variable  $\xi \in \mathbf{R}^d$  with  $d$  parameters, for fixed

---

2010 *Mathematics Subject Classification.* 46F05, 46F12, 35A27, 35S05, 35A18, 81S30, 58J47.

*Key words and phrases.* Tempered distributions, global wave front sets, pseudodifferential operators, Shubin calculus, microlocality, microellipticity, phase space, anisotropy.

$x \in \mathbf{R}^d$ . Our idea is to study global anisotropy comprehensively in the phase space  $T^*\mathbf{R}^d$ . For simplicity we use only one parameter for the relation between the space and the frequency variables. Even the idea of anisotropic pseudodifferential calculus on  $T^*\mathbf{R}^d$  is not new, cf. [3, 6, 18, 20], but as far as we know a systematic microlocal analysis has not yet been fully developed. The aim of our paper is to contribute to such a calculus and adapted microlocal analysis.

For  $s > 0$  and  $m \in \mathbf{R}$  we study symbols that are smooth and satisfy estimates of the form

$$|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \lesssim (1 + |x| + |\xi|^{1/s})^{m - |\alpha| - s|\beta|}, \quad (x, \xi) \in T^*\mathbf{R}^d, \quad \alpha, \beta \in \mathbf{N}^d.$$

This is a generalization of the isotropic Shubin symbols that satisfy the estimates with  $s = 1$ . When  $s \neq 1$  an anisotropic symbol is still embedded in an isotropic Shubin symbol space of possibly higher order. These symbol classes were introduced in [20, Definition 3.1], and the corresponding basic calculus is briefly stated there without proofs. In this paper we provide detailed proofs of the calculus from scratch, and extend the analysis to an adapted anisotropic Gabor microlocal analysis.

For fixed  $s > 0$  we show that the anisotropic symbols give rise to a subcalculus of the isotropic Shubin calculus. More precisely the anisotropic symbol classes are independent of the quantization parameter that admits transfer between Weyl and Kohn–Nirenberg quantization. They are also stable with respect to operator composition as well as formal adjoint.

Then we introduce the corresponding notion of anisotropic Gabor wave front set  $\text{WF}_g^s(u)$  of a tempered distribution  $u$ . This means the complement of curves of the form (1.1) in a neighborhood of which the short-time Fourier transform decays super-polynomially. The neighborhoods are  $s$ -conic, that is if a point  $(x, \xi) \in T^*\mathbf{R}^d \setminus 0$  belongs to the neighborhood then it contains the whole curve (1.1), and so is the anisotropic Gabor wave front set.

The first main result that we present is the microlocal inclusion

$$\text{WF}_g^s(a^w(x, D)u) \subseteq \text{WF}_g^s(u),$$

where  $u$  is a tempered distribution,  $a$  is an isotropic Shubin symbol, and  $a^w(x, D)$  denotes the Weyl quantization.

The second main result is the microelliptic inclusion

$$\text{WF}_g^s(u) \subseteq \text{WF}_g^s(a^w(x, D)u) \bigcup \text{char}_{s, m_1}(a)$$

where again  $u$  is a tempered distribution,  $a$  is an anisotropic Shubin symbol with parameter  $s > 0$  and order  $m$ ,  $m_1 \leq m$  and  $\text{char}_{s, m_1}(a)$  is a notion of microlocal characteristic set adapted to the anisotropic Shubin calculus (see Definition 3.8).

Taken together these results imply

$$\text{WF}_g^s(a^w(x, D)u) = \text{WF}_g^s(u)$$

if  $\text{char}_{s, m_1}(a) = \emptyset$  for some  $m_1 \leq m$ .

The paper is organized as follows. Section 2 sets the stage in terms of notations and some definitions, and a background on pseudodifferential operators in the Weyl quantization with isotropic Shubin symbols. In Section 3 we introduce the anisotropic Shubin symbols for a fixed parameter  $s > 0$ . We show adapted asymptotic expansions,

and invariance under a commonly used family of quantizations parametrized by a real parameter. This family includes the Weyl as well as the Kohn–Nirenberg quantization. We also show the continuity of the Weyl product acting on the anisotropic symbol Fréchet spaces, and we discuss  $s$ -conic cutoff functions.

Section 4 is devoted to the anisotropic Gabor wave front set. We state the definition, discuss a few properties and show that it does not depend of the chosen nonzero Schwartz window function in the short-time Fourier transform. The full metaplectic invariance of the isotropic ( $s = 1$ ) Gabor wave front set does not hold when  $s \neq 1$  but we show a few partial such invariances.

In Section 5 we show that pseudodifferential operators with isotropic Shubin symbols are microlocal with respect to all anisotropic Gabor wave front sets. In particular microlocality holds for anisotropic Shubin symbols. Another consequence is the invariance of anisotropic Gabor wave front sets with respect to translation and modulation.

Section 6 treats a microelliptic inclusion for the anisotropic Gabor wave front set and anisotropic Shubin symbols with  $s > 0$  fixed. Finally in Section 7 we show inclusions and equalities for the anisotropic Gabor wave front set of oscillatory functions with phase functions that are real polynomials on  $\mathbf{R}^d$  of order  $m \geq 2$ . The anisotropy parameter is  $s = m - 1$ .

## 2. PRELIMINARIES

The unit sphere in  $\mathbf{R}^d$  is denoted by  $\mathbf{S}^{d-1} \subseteq \mathbf{R}^d$ . A ball of radius  $r > 0$  in  $\mathbf{R}^d$  is denoted by  $B_r$ , and  $e_j \in \mathbf{R}^d$  is the vector of zeros except for position  $j$ ,  $1 \leq j \leq d$ , where it is one. The transpose of a matrix  $A \in \mathbf{R}^{d \times d}$  is denoted by  $A^T$ . We write  $f(x) \lesssim g(x)$  provided there exists  $C > 0$  such that  $f(x) \leq C g(x)$  for all  $x$  in the domain of  $f$  and of  $g$ . If  $f(x) \lesssim g(x) \lesssim f(x)$  then we write  $f \asymp g$ . We use the bracket  $\langle x \rangle = (1 + |x|^2)^{\frac{1}{2}}$  for  $x \in \mathbf{R}^d$ . Peetre's inequality with optimal constant [24, Lemma 2.1] is

$$\langle x + y \rangle^s \leq \left( \frac{2}{\sqrt{3}} \right)^{|s|} \langle x \rangle^s \langle y \rangle^{|s|} \quad x, y \in \mathbf{R}^d, \quad s \in \mathbf{R}.$$

The normalization of the Fourier transform is

$$\mathcal{F}f(\xi) = \widehat{f}(\xi) = (2\pi)^{-\frac{d}{2}} \int_{\mathbf{R}^d} f(x) e^{-i\langle x, \xi \rangle} dx, \quad \xi \in \mathbf{R}^d,$$

for  $f \in \mathcal{S}(\mathbf{R}^d)$  (the Schwartz space), where  $\langle \cdot, \cdot \rangle$  denotes the scalar product on  $\mathbf{R}^d$ . The conjugate linear action of a tempered distribution  $u \in \mathcal{S}'(\mathbf{R}^d)$  on a test function  $\phi \in \mathcal{S}(\mathbf{R}^d)$  is written  $(u, \phi)$ , consistent with the  $L^2$  inner product  $(\cdot, \cdot) = (\cdot, \cdot)_{L^2}$  which is conjugate linear in the second argument.

Denote translation by  $T_x f(y) = f(y - x)$  and modulation by  $M_\xi f(y) = e^{i\langle y, \xi \rangle} f(y)$  for  $x, y, \xi \in \mathbf{R}^d$  where  $f$  is a function or distribution defined on  $\mathbf{R}^d$ . The composed operator is denoted by  $\Pi(x, \xi) = M_\xi T_x$ . Let  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus \{0\}$ . The short-time Fourier transform (STFT) of a tempered distribution  $u \in \mathcal{S}'(\mathbf{R}^d)$  is defined by

$$V_\varphi u(x, \xi) = (2\pi)^{-\frac{d}{2}} (u, M_\xi T_x \varphi) = \mathcal{F}(u T_x \overline{\varphi})(\xi), \quad x, \xi \in \mathbf{R}^d.$$

The function  $V_\varphi u$  is smooth and polynomially bounded [11, Theorem 11.2.3], that is there exists  $k \geq 0$  such that

$$(2.1) \quad |V_\varphi u(x, \xi)| \lesssim \langle (x, \xi) \rangle^k, \quad (x, \xi) \in T^*\mathbf{R}^d.$$

We have  $u \in \mathcal{S}(\mathbf{R}^d)$  if and only if

$$(2.2) \quad |V_\varphi u(x, \xi)| \lesssim \langle (x, \xi) \rangle^{-N}, \quad (x, \xi) \in T^*\mathbf{R}^d, \quad \forall N \geq 0.$$

The inverse transform is given by

$$(2.3) \quad u = (2\pi)^{-\frac{d}{2}} \iint_{\mathbf{R}^{2d}} V_\varphi u(x, \xi) M_\xi T_x \varphi \, dx \, d\xi$$

provided  $\|\varphi\|_{L^2} = 1$ , with action under the integral understood, that is

$$(2.4) \quad (u, f) = (V_\varphi u, V_\varphi f)_{L^2(\mathbf{R}^{2d})}$$

for  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $f \in \mathcal{S}(\mathbf{R}^d)$ , cf. [11, Theorem 11.2.5].

We will use

$$|x + y|^{\frac{1}{s}} \leq \kappa(s^{-1}) \left( |x|^{\frac{1}{s}} + |y|^{\frac{1}{s}} \right), \quad x, y \in \mathbf{R}^d, \quad s > 0,$$

where

$$\kappa(t) = \begin{cases} 1 & \text{if } 0 < t \leq 1 \\ 2^{t-1} & \text{if } t > 1 \end{cases}.$$

Let  $s > 0$ . We use the weight function on  $(x, \xi) \in T^*\mathbf{R}^d$

$$(2.5) \quad \mu_s(x, \xi) = 1 + |x| + |\xi|^{\frac{1}{s}}.$$

The following inequality of Peetre type holds.

**Lemma 2.1.** *If  $t \in \mathbf{R}$  then*

$$\mu_s(x + y, \xi + \eta)^t \leq C_{s,t} \mu_s(x, \xi)^{|t|} \mu_s(y, \eta)^t, \quad x, y, \xi, \eta \in \mathbf{R}^d.$$

*Proof.* We may assume  $t = 1$ . We have

$$\begin{aligned} \mu_s(x + y, \xi + \eta) &= 1 + |x + y| + |\xi + \eta|^{\frac{1}{s}} \\ &\leq 1 + |x| + |y| + \kappa(s^{-1})|\xi|^{\frac{1}{s}} + \kappa(s^{-1})|\eta|^{\frac{1}{s}} \\ &\leq \left( 1 + |x| + \kappa(s^{-1})|\xi|^{\frac{1}{s}} \right) \left( 1 + |y| + \kappa(s^{-1})|\eta|^{\frac{1}{s}} \right) \\ &\leq \kappa(s^{-1})^2 \mu_s(x, \xi) \mu_s(y, \eta). \end{aligned}$$

□

For  $s > 0$  we will use subsets of  $T^*\mathbf{R}^d \setminus 0$  that are  $s$ -conic, that is subsets closed under the operation  $T^*\mathbf{R}^d \setminus 0 \ni (x, \xi) \mapsto (\lambda x, \lambda^s \xi)$  for all  $\lambda > 0$ .

**2.1. Pseudodifferential operators.** We need some elements from the calculus of pseudodifferential operators [9, 13, 19, 26]. Let  $a \in C^\infty(\mathbf{R}^{2d})$ ,  $m \in \mathbf{R}$  and  $0 \leq \rho \leq 1$ . Then  $a$  is a *Shubin symbol* of order  $m$  and parameter  $\rho$ , denoted  $a \in G_\rho^m$ , if for all  $\alpha, \beta \in \mathbf{N}^d$  there exists a constant  $C_{\alpha, \beta} > 0$  such that

$$(2.6) \quad |\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \leq C_{\alpha, \beta} \langle (x, \xi) \rangle^{m - \rho|\alpha + \beta|}, \quad x, \xi \in \mathbf{R}^d.$$

The Shubin symbols  $G_\rho^m$  form a Fréchet space where the seminorms are given by the smallest possible constants in (2.6). We write  $G_1^m = G^m$ .

For  $a \in G_\rho^m$  and  $t \in \mathbf{R}$  a pseudodifferential operator in the  $t$ -quantization is defined by

$$(2.7) \quad a_t(x, D)f(x) = (2\pi)^{-d} \int_{\mathbf{R}^{2d}} e^{i\langle x-y, \xi \rangle} a((1-t)x + ty, \xi) f(y) dy d\xi, \quad f \in \mathcal{S}(\mathbf{R}^d),$$

when  $m < -d$ . The definition extends to  $m \in \mathbf{R}$  if the integral is viewed as an oscillatory integral. If  $t = 0$  we get the Kohn–Nirenberg quantization  $a_0(x, D)$  and if  $t = \frac{1}{2}$  we get the Weyl quantization  $a_{1/2}(x, D) = a^w(x, D)$ . The relation between symbols in different quantizations is [13]

$$e^{it\langle D_x, D_\xi \rangle} a_t(x, \xi) = e^{is\langle D_x, D_\xi \rangle} a_s(x, \xi), \quad t, s \in \mathbf{R}$$

where  $e^{it\langle D_x, D_\xi \rangle}$  is the Fourier multiplier operator with symbol  $e^{it\langle x, \xi \rangle}$ . Using [13, Theorem 7.6.1] we may write for  $t \in \mathbf{R} \setminus 0$  and  $a \in \mathcal{S}(\mathbf{R}^{2d})$

$$(2.8) \quad e^{it\langle D_x, D_\xi \rangle} a(x, \xi) = (2\pi|t|)^{-d} \iint_{\mathbf{R}^{2d}} a(y, \eta) e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle} dy d\eta.$$

If  $0 < \rho \leq 1$  then the Shubin symbols are invariant with respect to the parameter  $t$  in the sense of  $a_t \in G_\rho^m$  if and only if  $a_s = e^{i(t-s)\langle D_x, D_\xi \rangle} a_t \in G_\rho^m$  for any  $t, s \in \mathbf{R}$  [26, Theorem 23.2]. If  $t \in \mathbf{R}$  then for the formal adjoint we have  $a_t(x, D)^* = \bar{a}_{1-t}(x, D)$ . Thus if  $a_t \in G_\rho^m$  then  $a_t(x, D)^* = b_t(x, D)$  where  $b_t \in G_\rho^m$  [26, Theorem 23.5].

We will use exclusively the Weyl quantization which has several particular features. One important such feature is the simplicity of the formal adjoint:  $a^w(x, D)^* = \bar{a}^w(x, D)$ . As for the Shubin symbols, we will see that also the anisotropic symbol classes that we will use in this paper give pseudodifferential calculi that are invariant with respect to the quantization parameter  $t \in \mathbf{R}$  (see Proposition 3.3).

If  $0 < \rho \leq 1$  and  $a \in G_\rho^m$  then the operator  $a^w(x, D)$  acts continuously on  $\mathcal{S}(\mathbf{R}^d)$  and extends uniquely by duality to a continuous operator on  $\mathcal{S}'(\mathbf{R}^d)$ . By Schwartz's kernel theorem the Weyl quantization may be extended to a weak formulation which yields continuous linear operators  $a^w(x, D) : \mathcal{S}(\mathbf{R}^d) \rightarrow \mathcal{S}'(\mathbf{R}^d)$ , even if  $a$  is only an element of  $\mathcal{S}'(\mathbf{R}^{2d})$ .

If  $a \in \mathcal{S}'(\mathbf{R}^{2d})$  then

$$(2.9) \quad (a^w(x, D)f, g) = (2\pi)^{-d} (a, W(g, f)), \quad f, g \in \mathcal{S}(\mathbf{R}^d),$$

where the cross-Wigner distribution [9, 11] is defined as

$$W(g, f)(x, \xi) = \int_{\mathbf{R}^d} g(x + y/2) \overline{f(x - y/2)} e^{-i\langle y, \xi \rangle} dy, \quad (x, \xi) \in \mathbf{R}^{2d}.$$

We have  $W(g, f) \in \mathcal{S}(\mathbf{R}^{2d})$  when  $f, g \in \mathcal{S}(\mathbf{R}^d)$ .

The real phase space  $T^*\mathbf{R}^d \simeq \mathbf{R}^d \oplus \mathbf{R}^d$  is a real symplectic vector space equipped with the canonical symplectic form

$$\sigma((x, \xi), (x', \xi')) = \langle x', \xi \rangle - \langle x, \xi' \rangle, \quad (x, \xi), (x', \xi') \in T^*\mathbf{R}^d.$$

This form can be expressed with the inner product as  $\sigma(X, Y) = \langle \mathcal{J}X, Y \rangle$  for  $X, Y \in T^*\mathbf{R}^d$  where

$$\mathcal{J} = \begin{pmatrix} 0 & I_d \\ -I_d & 0 \end{pmatrix} \in \mathbf{R}^{2d \times 2d}.$$

The real symplectic group  $\mathrm{Sp}(d, \mathbf{R})$  is the set of matrices in  $\mathrm{GL}(2d, \mathbf{R})$  that leaves  $\sigma$  invariant. Hence  $\mathcal{J} \in \mathrm{Sp}(d, \mathbf{R})$ .

To each symplectic matrix  $\chi \in \mathrm{Sp}(d, \mathbf{R})$  is associated an operator  $\mu(\chi)$  that is unitary on  $L^2(\mathbf{R}^d)$ , and determined up to a complex factor of modulus one, such that

$$\mu(\chi)^{-1}a^w(x, D)\mu(\chi) = (a \circ \chi)^w(x, D), \quad a \in \mathcal{S}'(\mathbf{R}^{2d})$$

(cf. [9, 13]). The operator  $\mu(\chi)$  is a homeomorphism on  $\mathcal{S}$  and on  $\mathcal{S}'$ .

The mapping  $\mathrm{Sp}(d, \mathbf{R}) \ni \chi \rightarrow \mu(\chi)$  is called the metaplectic representation [9]. It is in fact a representation of the so called 2-fold covering group of  $\mathrm{Sp}(d, \mathbf{R})$ , which is called the metaplectic group. The metaplectic representation satisfies the homomorphism relation modulo a change of sign:

$$\mu(\chi\chi') = \pm\mu(\chi)\mu(\chi'), \quad \chi, \chi' \in \mathrm{Sp}(d, \mathbf{R}).$$

We do not enter into the geometric subtleties of this construction since they are not needed in this paper.

Let  $0 < \rho \leq 1$ . The Weyl product  $a\# b$  of two symbols  $a \in G_\rho^m$  and  $b \in G_\rho^n$  is defined as the product of symbols corresponding to operator composition:  $(a\# b)^w(x, D) = a^w(x, D)b^w(x, D)$ . According to [26, Theorem 23.6]  $a\# b \in G_\rho^{m+n}$  if  $a \in G_\rho^m$  and  $b \in G_\rho^n$ , and the bilinear map  $(a, b) \mapsto a\# b$  is continuous  $G_\rho^m \times G_\rho^n \rightarrow G_\rho^{m+n}$ . When  $a, b \in \mathcal{S}(\mathbf{R}^{2d})$  we have the formula [13, Eq. (18.5.6)]

$$(2.10) \quad a\# b(x, \xi) = e^{\frac{i}{2}\sigma(D_x, D_\xi; D_y, D_\eta)} a(x, \xi) b(y, \eta) \Big|_{(y, \eta) = (x, \xi)}.$$

Using [13, Vol. 3 p. 152] we may write for  $a, b \in \mathcal{S}(\mathbf{R}^{2d})$

$$(2.11) \quad a\# b(z) = \pi^{-2d} \iint_{\mathbf{R}^{4d}} a(w) b(u) e^{2i\sigma(z-u, z-w)} dw du, \quad z \in T^*\mathbf{R}^d.$$

### 3. ANISOTROPIC SHUBIN CALCULUS

Let  $s > 0$  be fixed. We need a simplified version of a tool taken from [16, 20] and their references. Given  $(x, \xi) \in \mathbf{R}^{2d} \setminus 0$  there is a unique  $\lambda = \lambda(x, \xi) = \lambda_s(x, \xi) > 0$  such that

$$\lambda(x, \xi)^{-2}|x|^2 + \lambda(x, \xi)^{-2s}|\xi|^2 = 1.$$

Then  $(x, \xi) \in \mathbf{S}^{2d-1}$  if and only if  $\lambda(x, \xi) = 1$ . By the implicit function theorem the function  $\lambda : \mathbf{R}^{2d} \setminus 0 \rightarrow \mathbf{R}_+$  is smooth [17].

If  $\mu > 0$  and  $(x, \xi) \in \mathbf{S}^{2d-1}$  then  $\lambda(\mu x, \mu^s \xi) = \mu = \mu \lambda(x, \xi)$ . In fact

$$(3.1) \quad \lambda(\mu x, \mu^s \xi) = \mu \lambda(x, \xi)$$

holds for any  $(x, \xi) \in \mathbf{R}^{2d} \setminus 0$  and  $\mu > 0$  by the following argument. Given  $(x, \xi) \in \mathbf{R}^{2d} \setminus 0$  set  $\mu_1 = \lambda(x, \xi)$  so that  $(x/\mu_1, \xi/\mu_1^s) \in \mathbf{S}^{2d-1}$ . Then for  $\mu > 0$

$$\lambda(\mu x, \mu^s \xi) = \lambda(\mu \mu_1 x / \mu_1, (\mu \mu_1)^s \xi / \mu_1^s) = \mu \mu_1 = \mu \lambda(x, \xi).$$

We may define the projection  $p(x, \xi) = p_s(x, \xi)$  of  $(x, \xi) \in \mathbf{R}^{2d} \setminus 0$  along the curve  $\mathbf{R}_+ \ni \mu \mapsto (\mu x, \mu^s \xi)$  onto  $\mathbf{S}^{2d-1}$ . This means

$$(3.2) \quad p(x, \xi) = (\lambda(x, \xi)^{-1} x, \lambda(x, \xi)^{-s} \xi), \quad (x, \xi) \in \mathbf{R}^{2d} \setminus 0.$$

Due to (3.1)  $p(\mu x, \mu^s \xi) = p(x, \xi)$  does not depend on  $\mu > 0$ . The function  $p : \mathbf{R}^{2d} \setminus 0 \rightarrow \mathbf{S}^{2d-1}$  is smooth since  $\lambda \in C^\infty(\mathbf{R}^{2d} \setminus 0)$  and  $\lambda(x, \xi) > 0$  for all  $(x, \xi) \in \mathbf{R}^{2d} \setminus 0$ .

From [20], or by straightforward arguments, we have the bounds

$$(3.3) \quad |x| + |\xi|^{\frac{1}{s}} \lesssim \lambda(x, \xi) \lesssim |x| + |\xi|^{\frac{1}{s}}, \quad (x, \xi) \in \mathbf{R}^{2d} \setminus 0$$

and

$$(3.4) \quad \langle (x, \xi) \rangle^{\min(1, \frac{1}{s})} \lesssim 1 + \lambda(x, \xi) \lesssim \langle (x, \xi) \rangle^{\max(1, \frac{1}{s})}, \quad (x, \xi) \in \mathbf{R}^{2d} \setminus 0.$$

Hörmander type symbol classes with anisotropic behavior in the frequency domain can be found in [16, Définition 1.3] and in [20, Definition 1.4]. Now we define symbol classes that are adaptations of this concept to the Shubin calculus.

**Definition 3.1.** Let  $s > 0$  and  $m \in \mathbf{R}$ . The space of  $(s)$ -anisotropic Shubin symbols  $G^{m,s}$  of order  $m$  consists of functions  $a \in C^\infty(\mathbf{R}^{2d})$  that satisfy the estimates

$$|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \lesssim (1 + |x| + |\xi|^{\frac{1}{s}})^{m - |\alpha| - s|\beta|}, \quad (x, \xi) \in T^*\mathbf{R}^d, \quad \alpha, \beta \in \mathbf{N}^d.$$

The symbols  $G^{m,s}$  enjoy the following symmetry: If  $b(x, \xi) = a(\xi, x)$  then  $a \in G^{m,s}$  if and only if  $b \in G^{m/s, 1/s}$ . It is clear that

$$\bigcap_{m \in \mathbf{R}} G^{m,s} = \mathcal{S}(\mathbf{R}^{2d}).$$

Referring to the weight (2.5) we use the seminorms on  $a \in G^{m,s}$  indexed by  $j \in \mathbf{N}$

$$(3.5) \quad \|a\|_j = \max_{|\alpha|+|\beta| \leq j} \sup_{(x, \xi) \in \mathbf{R}^{2d}} \mu_s(x, \xi)^{-m+|\alpha|+s|\beta|} \left| \partial_x^\alpha \partial_\xi^\beta a(x, \xi) \right|.$$

The symbol classes  $G^{m,s}$  with  $s \in \mathbf{Q}_+$  (positive rationals) were introduced in [20, Definition 3.1] as a tool in order to construct parametrices for pseudodifferential operators. Here we generalize to  $s \in \mathbf{R}_+$  which is a straightforward extension concerning the calculus. In [20, Section 3] results for a calculus for the symbol classes  $G^{m,s}$  are briefly stated without proofs. In this section we prove in detail the basic calculus results for the anisotropic Shubin symbols  $G^{m,s}$ .

We have  $G^{m,1} = G^m = G_1^m$ , that is the usual Shubin class, and we cannot embed  $G_\rho^m$  in a space  $G^{n,s}$  unless  $\rho = s = 1$ . Using (3.3) and (3.4) the embedding

$$(3.6) \quad G^{m,s} \subseteq G_\rho^{m_0},$$

where  $m_0 = \max(m, m/s)$  and  $\rho = \min(s, 1/s)$ , can be confirmed. Thus the Shubin calculus [26] applies to the anisotropic Shubin symbols.

We also note that the more general pseudodifferential calculus in [19] is not directly applicable to the symbol classes  $G^{m,s}$  unless  $s = 1$ . In fact if  $s \neq 1$  then either the space weight function  $\Phi(x, \xi) = 1 + |x| + |\xi|^{\frac{1}{s}}$  or the frequency weight function  $\Psi(x, \xi) = 1 + |x|^s + |\xi|$  is not sublinear. Nevertheless from (3.4) it follows that  $G^{m,s} \subseteq S(M; \Phi, \Psi)$  as defined in [19, Definition 1.1.1] with  $M(x, \xi) = \langle (x, \xi) \rangle^{\max(m, m/s)}$ ,  $\Phi(x, \xi) = \langle (x, \xi) \rangle^{\min(1, \frac{1}{s})}$  and  $\Psi(x, \xi) = \langle (x, \xi) \rangle^{\min(1, s)}$ . Thus the pseudodifferential calculus in [19, Chapter 1.2] applies to  $G^{m,s}$ , but the anisotropy is again lost.

There is a more subtle anisotropic subcalculus adapted to the anisotropic Shubin symbols  $G^{m,s}$ , for each fixed  $s > 0$ , which preserves the anisotropy. We deduce a minimal such calculus and start with asymptotic expansions.

Given a sequence of symbols  $a_j \in G^{m_j, s}$ ,  $j = 1, 2, \dots$ , such that  $m_j \rightarrow -\infty$  as  $j \rightarrow +\infty$  we write

$$a \sim \sum_{j=1}^{\infty} a_j$$

provided that for any  $n \geq 2$

$$a - \sum_{j=1}^{n-1} a_j \in G^{\mu_n, s}$$

where  $\mu_n = \max_{j \geq n} m_j$ .

**Lemma 3.2.** *Let  $s > 0$ . Given a sequence of symbols  $a_j \in G^{m_j, s}$ ,  $j = 1, 2, \dots$ , such that  $m_j \rightarrow -\infty$  as  $j \rightarrow +\infty$ , there exists a symbol  $a \in G^{m, s}$  where  $m = \max_{j \geq 1} m_j$  such that  $a \sim \sum_{j=1}^{\infty} a_j$ . The symbol  $a$  is unique modulo addition with a function in  $\mathcal{S}(\mathbf{R}^{2d})$ .*

*Proof.* Let  $\varphi \in C^\infty(\mathbf{R}^{2d})$  satisfy  $0 \leq \varphi \leq 1$ ,  $\varphi(z) = 0$  if  $|z| \leq \frac{1}{2}$  and  $\varphi(z) = 1$  if  $|z| \geq 1$ . Set for  $t \geq 1$

$$\psi(x, \xi) = \varphi(t^{-1}x, t^{-s}\xi), \quad (x, \xi) \in T^*\mathbf{R}^d.$$

Then for all  $t \geq 1$  we have

$$\left| \partial_x^\alpha \partial_\xi^\beta \psi(x, \xi) \right| \leq C_{\alpha, \beta} \mu_s(x, \xi)^{-|\alpha| - s|\beta|}.$$

If fact this is trivial if  $\alpha = \beta = 0$ . If instead  $(\alpha, \beta) \in \mathbf{N}^{2d} \setminus 0$  then

$$\frac{1}{4} \leq t^{-2}|x|^2 + t^{-2s}|\xi|^2 \leq 1$$

in the support of  $\partial_x^\alpha \partial_\xi^\beta \varphi(t^{-1}x, t^{-s}\xi)$ . Thus  $|x| + |\xi|^{\frac{1}{s}} \lesssim t$  in said support. This gives

$$\begin{aligned} (3.7) \quad \left| \partial_x^\alpha \partial_\xi^\beta \psi(x, \xi) \right| &= t^{-|\alpha| - s|\beta|} \left| (\partial_x^\alpha \partial_\xi^\beta \varphi)(t^{-1}x, t^{-s}\xi) \right| \\ &\leq C_{\alpha, \beta} \mu_s(x, \xi)^{-|\alpha| - s|\beta|}. \end{aligned}$$

The symbol  $a$  is constructed as

$$a(x, \xi) = \sum_{j=1}^{\infty} \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi)$$

for a sufficiently rapidly increasing sequence  $(t_j) \subseteq \mathbf{R}_+$ . Given  $n \geq 2$  we must show  $a - \sum_{j=1}^{n-1} a_j \in G^{\mu_n, s}$ . We have

$$a(x, \xi) - \sum_{j=1}^{n-1} a_j(x, \xi) = \sum_{j=1}^{n-1} \left( \varphi(t_j^{-1}x, t_j^{-s}\xi) - 1 \right) a_j(x, \xi) + \sum_{j=n}^{\infty} \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi).$$

The first sum is compactly supported and hence belongs to  $G^{\mu_n, s}$  trivially so it suffices to prove

$$(3.8) \quad \sum_{j=n}^{\infty} \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi) \in G^{\mu_n, s}.$$

First we show

$$(3.9) \quad \left| \partial_x^\alpha \partial_\xi^\beta \left( \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi) \right) \right| \leq 2^{-j} \mu_s(x, \xi)^{m_j+1-|\alpha|-s|\beta|}$$

for all  $j \geq 1$  and  $|\alpha| + |\beta| \leq j$ , provided  $t_j > 0$  is sufficiently large. In fact this estimate is a consequence of  $a_j \in G^{m_j, s}$ , (3.7), Leibniz' rule, and the support properties of  $\varphi(t_j^{-1}x, t_j^{-s}\xi)$ , if  $t_j > 0$  is sufficiently large.

Let  $\alpha, \beta \in \mathbf{N}^d$  and pick  $N \geq \max(n+1, |\alpha| + |\beta|)$  such that  $\mu_N \leq \mu_n - 1$ . Then for all  $j \geq N$  it holds  $m_j \leq \mu_j \leq \mu_N \leq \mu_n - 1$ . Combined with (3.9) this gives

$$\sum_{j=N}^{\infty} \left| \partial_x^\alpha \partial_\xi^\beta \left( \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi) \right) \right| \leq 2^{1-N} \mu_s(x, \xi)^{\mu_n-|\alpha|-s|\beta|}.$$

Since  $\sum_{j=N}^{N-1} \varphi(t_j^{-1}x, t_j^{-s}\xi) a_j(x, \xi) \in G^{\mu_n, s}$  we have proved (3.8).  $\square$

We have the following asymptotic expansion for the Weyl product of  $a \in G^{m, s}$  and  $b \in G^{n, s}$ ,  $m, n \in \mathbf{R}$  [26]:

$$(3.10) \quad a \# b(x, \xi) \sim \sum_{\alpha, \beta \geq 0} \frac{(-1)^{|\beta|}}{\alpha! \beta!} 2^{-|\alpha+\beta|} D_x^\beta \partial_\xi^\alpha a(x, \xi) D_x^\alpha \partial_\xi^\beta b(x, \xi).$$

Each term in the sum belongs to  $G^{m+n-(1+s)|\alpha+\beta|, s}$ .

In the next result we show that the symbol classes  $G^{m, s}$  are invariant with respect to the parameter  $t \in \mathbf{R}$  in (2.7). In other words if one changes quantization one gets a new symbol in the same class. Combined with  $a^w(x, D)^* = \overline{a^w}(x, D)$ , an immediate consequence is that for each  $t \in \mathbf{R}$  the symbol class  $G^{m, s}$  is closed with respect to formal adjoint: If  $a_t \in G^{m, s}$  and  $a_t(x, D)^* = b_t(x, D)$  then  $b_t \in G^{m, s}$ .

We also show the continuity of the bilinear Weyl product on the symbol classes  $G^{m, s}$ . Again by the first result the continuity extends to the symbol product in the  $t$ -quantization for any  $t \in \mathbf{R}$ .

**Proposition 3.3.** *Let  $s > 0$  and  $m, n \in \mathbf{R}$ .*

- (i) *If  $t \in \mathbf{R}$  and  $a \in G^{m, s}$  then  $b(x, \xi) = e^{it\langle D_x, D_\xi \rangle} a(x, \xi) \in G^{m, s}$ , and the map  $a \mapsto b$  is continuous on  $G^{m, s}$ .*
- (ii) *If  $a \in G^{m, s}$  and  $b \in G^{n, s}$  then  $a \# b \in G^{m+n, s}$ , and the Weyl product is continuous*

$$\# : G^{m, s} \times G^{n, s} \rightarrow G^{m+n, s}.$$

*Proof.* (i) We may assume  $t \neq 0$  since the claim is trivial otherwise. Let  $\alpha, \beta \in \mathbf{N}^d$ . The operator  $e^{it\langle D_x, D_\xi \rangle}$  commutes with differential operators  $\partial_x^\alpha \partial_\xi^\beta$ . The distribution  $\partial_x^\alpha \partial_\xi^\beta b = \mathcal{F}^{-1} \left( e^{it\langle \cdot, \cdot \rangle} \widehat{\partial_x^\alpha \partial_\xi^\beta a} \right)$  is well defined in  $\mathcal{S}'(\mathbf{R}^{2d})$ .

Let  $\chi \in C_c^\infty(\mathbf{R}^{2d})$  satisfy  $0 \leq \chi \leq 1$ ,  $\chi(z) = 1$  when  $|z| \leq 1$  and  $\chi(z) = 0$  when  $|z| \geq 2$ . Set  $\chi_\varepsilon(z) = \chi(\varepsilon z)$  for  $\varepsilon > 0$ . Then  $\chi_\varepsilon(\partial_x^\alpha \partial_\xi^\beta a) \rightarrow \partial_x^\alpha \partial_\xi^\beta a$  in  $\mathcal{S}'(\mathbf{R}^{2d})$  as  $\varepsilon \rightarrow 0^+$ . Hence we obtain from (2.8)

$$\begin{aligned} \partial_x^\alpha \partial_\xi^\beta b(x, \xi) &= \mathcal{F}^{-1} \left( e^{it\langle \cdot, \cdot \rangle} \widehat{\partial_x^\alpha \partial_\xi^\beta a} \right) (x, \xi) = \lim_{\varepsilon \rightarrow 0^+} \mathcal{F}^{-1} \left( e^{it\langle \cdot, \cdot \rangle} \mathcal{F} \left( \chi_\varepsilon \partial_x^\alpha \partial_\xi^\beta a \right) \right) (x, \xi) \\ &= (2\pi|t|)^{-d} \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbf{R}^{2d}} e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle} \chi_\varepsilon(y, \eta) \partial_x^\alpha \partial_\xi^\beta a(y, \eta) dy d\eta \end{aligned}$$

in  $\mathcal{S}'(\mathbf{R}^{2d})$ .

Define the operator

$$(Sf)(y, \eta) = (1 - \Delta_{y, \eta}) \left( \langle t^{-1}(x - y, \xi - \eta) \rangle^{-2} f(y, \eta) \right)$$

acting on  $f \in C^\infty(\mathbf{R}^{2d})$ . From

$$(1 - \Delta_{y, \eta}) e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle} = \langle t^{-1}(x - y, \xi - \eta) \rangle^2 e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle}$$

we obtain from integration by parts for  $N \in \mathbf{N}$

$$\begin{aligned} (2\pi|t|)^d \partial_x^\alpha \partial_\xi^\beta b(x, \xi) &= \lim_{\varepsilon \rightarrow 0^+} \int_{\mathbf{R}^{2d}} e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle} S^N \left( \chi_\varepsilon(y, \eta) \partial_x^\alpha \partial_\xi^\beta a(y, \eta) \right) dy d\eta \\ &= \int_{\mathbf{R}^{2d}} e^{-\frac{i}{t}\langle x-y, \xi-\eta \rangle} S^N \left( \partial_x^\alpha \partial_\xi^\beta a(y, \eta) \right) dy d\eta \end{aligned}$$

by dominated convergence, since  $S^N \partial_x^\alpha \partial_\xi^\beta a \in L^1(\mathbf{R}^{2d})$  provided  $N$  is large enough.

This gives using (3.4), (3.5) and Lemma 2.1

$$\begin{aligned} |\partial_x^\alpha \partial_\xi^\beta b(x, \xi)| &\lesssim \int_{\mathbf{R}^{2d}} \left| S^N \left( \partial_x^\alpha \partial_\xi^\beta a(y, \eta) \right) \right| dy d\eta \\ &\leq C_{t,N} \|a\|_{2N+|\alpha+\beta|} \int_{\mathbf{R}^{2d}} \langle (x - y, \xi - \eta) \rangle^{-2N} \mu_s(y, \eta)^{m-|\alpha|-s|\beta|} dy d\eta \\ &= C_{t,N} \|a\|_{2N+|\alpha+\beta|} \int_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{-2N} \mu_s(x - y, \xi - \eta)^{m-|\alpha|-s|\beta|} dy d\eta \\ &\lesssim C_{t,N} \|a\|_{2N+|\alpha+\beta|} \mu_s(x, \xi)^{m-|\alpha|-s|\beta|} \int_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{-2N} \mu_s(y, \eta)^{|m|+|\alpha|+s|\beta|} dy d\eta \\ &\leq C_{t,N} \|a\|_{2N+|\alpha+\beta|} \mu_s(x, \xi)^{m-|\alpha|-s|\beta|} \int_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{-2N+(|m|+|\alpha|+s|\beta|)\max(1, \frac{1}{s})} dy d\eta \\ &\leq C_{t,N} \|a\|_{2N+|\alpha+\beta|} \mu_s(x, \xi)^{m-|\alpha|-s|\beta|} \end{aligned}$$

after possibly increasing  $N$  (which may depend on  $|\alpha + \beta|$ ). In view of (3.5) we obtain for any  $j \in \mathbf{N}$

$$\|b\|_j \leq C_{t,N} \|a\|_{2N_j+j}$$

for some  $N_j \in \mathbf{N}$ , which proves claim (i).

(ii) Due to (3.6) we may use results for the calculus of Shubin symbols  $G_\rho^m$ . When  $a, b \in \mathcal{S}(\mathbf{R}^{2d})$  we have by (2.10)  $a \# b(z) = f(z, z)$  where

$$f(z, w) = e^{\frac{i}{2}\sigma(D_z, D_w)}(a \otimes b)(z, w), \quad z, w \in \mathbf{R}^{2d}.$$

Suppose  $a \in G^{m,s}$  and  $b \in G^{n,s}$ . Set  $a_\varepsilon = \chi_\varepsilon a$  and  $b_\varepsilon = \chi_\varepsilon b$  where  $\chi \in C_c^\infty(\mathbf{R}^{2d})$  and  $\chi_\varepsilon$  is defined as above. Then  $a_\varepsilon \otimes b_\varepsilon \rightarrow a \otimes b$  in  $\mathcal{S}'(\mathbf{R}^{4d})$  as  $\varepsilon \rightarrow 0^+$ . Since  $e^{\frac{i}{2}\sigma(D_z, D_w)}$  is continuous on  $\mathcal{S}'(\mathbf{R}^{4d})$  it follows that

$$(3.11) \quad f(z, w) = \lim_{\varepsilon \rightarrow 0^+} e^{\frac{i}{2}\sigma(D_z, D_w)}(a_\varepsilon \otimes b_\varepsilon)(z, w)$$

in  $\mathcal{S}'(\mathbf{R}^{4d})$ .

From the argument in the proof of [25, Theorem A.5] it follows that the limit (3.11) is actually pointwise for all  $z, w \in \mathbf{R}^{2d}$ . The Fourier multiplier operator  $e^{\frac{i}{2}\sigma(D_z, D_w)}$  commutes with differential operators so for any  $\alpha, \beta \in \mathbf{N}^{2d}$  we have the pointwise limit

$$(3.12) \quad \partial_z^\alpha \partial_w^\beta f(z, w) = \lim_{\varepsilon \rightarrow 0^+} e^{\frac{i}{2}\sigma(D_z, D_w)}(\partial_z^\alpha a_\varepsilon \otimes \partial_w^\beta b_\varepsilon)(z, w)$$

which yields using (2.11)

$$(3.13) \quad \begin{aligned} \partial^\alpha(a \# b)(z) &= \partial^\alpha(f(z, z)) = \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} (\partial_z^\beta \partial_w^{\alpha-\beta} f)(z, z) \\ &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \lim_{\varepsilon \rightarrow 0^+} e^{\frac{i}{2}\sigma(D_z, D_w)}(\partial^\beta a_\varepsilon \otimes \partial^{\alpha-\beta} b_\varepsilon)(z, z) \\ &= \pi^{-2d} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \lim_{\varepsilon \rightarrow 0^+} \iint_{\mathbf{R}^{4d}} e^{2i\sigma(z-v, z-u)} \partial^\beta a_\varepsilon(u) \partial^{\alpha-\beta} b_\varepsilon(v) \, du \, dv. \end{aligned}$$

Next we note

$$(1 - \Delta_{u,v})e^{2i\sigma(z-v, z-u)} = \langle 2(z-u, z-v) \rangle^2 e^{2i\sigma(z-v, z-u)}.$$

If we define the operator

$$(Sf)(u, v) = (1 - \Delta_{u,v}) \left( \langle 2(z-u, z-v) \rangle^{-2} f(u, v) \right), \quad u, v \in \mathbf{R}^{2d},$$

acting on  $f \in C^\infty(\mathbf{R}^{4d})$ , then we obtain for  $N \in \mathbf{N}$  using integration by parts and dominated convergence

$$\begin{aligned} &\lim_{\varepsilon \rightarrow 0^+} \iint_{\mathbf{R}^{4d}} e^{2i\sigma(z-v, z-u)} \partial^\beta a_\varepsilon(u) \partial^{\alpha-\beta} b_\varepsilon(v) \, du \, dv \\ &= \lim_{\varepsilon \rightarrow 0^+} \iint_{\mathbf{R}^{4d}} e^{2i\sigma(z-v, z-u)} S^N \left( \partial^\beta a_\varepsilon(u) \partial^{\alpha-\beta} b_\varepsilon(v) \right) \, du \, dv \\ &= \iint_{\mathbf{R}^{4d}} e^{2i\sigma(z-v, z-u)} S^N \left( \partial^\beta a(u) \partial^{\alpha-\beta} b(v) \right) \, du \, dv \end{aligned}$$

since  $S^N(\partial^\beta a \otimes \partial^{\alpha-\beta} b) \in L^1(\mathbf{R}^{4d})$  provided  $N$  is sufficiently large.

We denote  $\alpha = (\alpha_1, \alpha_2) \in \mathbf{N}^{2d}$  with  $\alpha_1, \alpha_2 \in \mathbf{N}^d$ . Combining with (3.13) and using (3.4), (3.5) and Lemma 2.1 we obtain

$$\begin{aligned}
& |\partial^\alpha(a \# b)(z)| \\
& \lesssim \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \iint_{\mathbf{R}^{4d}} \left| S^N \left( \partial^\beta a(u) \partial^{\alpha-\beta} b(v) \right) \right| du dv \\
& \lesssim \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \|a\|_{2N+|\beta|} \|b\|_{2N+|\alpha-\beta|} \\
& \quad \times \iint_{\mathbf{R}^{4d}} \langle (z-u, z-v) \rangle^{-2N} \mu_s(u)^{m-|\beta_1|-s|\beta_2|} \mu_s(v)^{n-|\alpha_1-\beta_1|-s|\alpha_2-\beta_2|} du dv \\
& \leq \|a\|_{2N+|\alpha|} \|b\|_{2N+|\alpha|} \\
& \quad \times \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \iint_{\mathbf{R}^{4d}} \langle (u, v) \rangle^{-2N} \mu_s(z-u)^{m-|\beta_1|-s|\beta_2|} \mu_s(z-v)^{n-|\alpha_1-\beta_1|-s|\alpha_2-\beta_2|} du dv \\
& \lesssim \|a\|_{2N+|\alpha|} \|b\|_{2N+|\alpha|} \mu_s(z)^{m+n-|\alpha_1|-s|\alpha_2|} \\
& \quad \times \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \iint_{\mathbf{R}^{4d}} \langle (u, v) \rangle^{-2N+(|m|+|n|+2|\alpha_1|+2s|\alpha_2|)\max(1, \frac{1}{s})} du dv \\
& \lesssim \|a\|_{2N+|\alpha|} \|b\|_{2N+|\alpha|} \mu_s(z)^{m+n-|\alpha_1|-s|\alpha_2|}
\end{aligned}$$

if  $N$  is sufficiently large. This shows that for any  $\alpha \in \mathbf{N}^{2d}$  we have

$$\sup_{z \in \mathbf{R}^{2d}} \mu_s(z)^{-m-n+|\alpha_1|+s|\alpha_2|} |\partial^\alpha(a \# b)(z)| \lesssim \|a\|_{2N+|\alpha|} \|b\|_{2N+|\alpha|}$$

and the claimed continuity follows in view of (3.5).  $\square$

**3.1.  $s$ -conic cutoff functions.** A family of open  $s$ -conic subsets are defined and denoted as follows. Recall the projection function (3.2)  $p: \mathbf{R}^{2d} \setminus 0 \rightarrow \mathbf{S}^{2d-1}$ .

**Definition 3.4.** Suppose  $s, \varepsilon > 0$  and  $z_0 \in \mathbf{S}^{2d-1}$ . Then

$$\Gamma_{s, z_0, \varepsilon} = \{(x, \xi) \in \mathbf{R}^{2d} \setminus 0, |z_0 - p(x, \xi)| < \varepsilon\} \subseteq T^*\mathbf{R}^d \setminus 0.$$

For simplicity we write  $\Gamma_{z_0, \varepsilon} = \Gamma_{s, z_0, \varepsilon}$  when  $s$  is fixed and understood from the context. If  $\varepsilon > 2$  then  $\Gamma_{z_0, \varepsilon} = T^*\mathbf{R}^d \setminus 0$  so we usually restrict to  $\varepsilon \leq 2$ .

Next we construct cutoff functions  $\chi \in G^{0, s}$  such that  $0 \leq \chi \leq 1$ ,  $\text{supp } \chi \subseteq \Gamma_{z_0, 2\varepsilon} \setminus B_{r/2}$ ,  $\chi|_{\Gamma_{z_0, \varepsilon} \setminus \overline{B}_r} \equiv 1$  for given  $\varepsilon, r > 0$ , and  $z_0 \in \mathbf{S}^{2d-1}$ . They will be needed in Section 6.

**Lemma 3.5.** Let  $s > 0$ . If  $r > 0$ ,  $0 < \varepsilon \leq 1$  and  $z_0 \in \mathbf{S}^{2d-1}$  then there exists  $\chi \in G^{0, s}$  such that  $0 \leq \chi \leq 1$ ,  $\text{supp } \chi \subseteq \Gamma_{z_0, 2\varepsilon} \setminus B_{r/2}$  and  $\chi|_{\Gamma_{z_0, \varepsilon} \setminus \overline{B}_r} \equiv 1$ .

*Proof.* Let  $\varphi \in C_c^\infty(\mathbf{R}^{2d})$  satisfy  $0 \leq \varphi \leq 1$ ,  $\text{supp } \varphi \subseteq z_0 + B_{2\varepsilon}$  and  $\varphi|_{z_0 + B_\varepsilon} \equiv 1$ . Let  $g \in C^\infty(\mathbf{R})$  satisfy  $0 \leq g \leq 1$ ,  $g(x) = 0$  if  $x \leq \frac{1}{2}$  and  $g(x) = 1$  if  $x \geq 1$ . Set

$$(3.14) \quad \psi(\lambda x, \lambda^s \xi) = \varphi(x, \xi), \quad (x, \xi) \in \mathbf{S}^{2d-1}, \quad \lambda > 0,$$

and

$$(3.15) \quad \chi(z) = g(r^{-1}|z|)\psi(z), \quad z \in \mathbf{R}^{2d}.$$

Note that (3.14) can be written

$$\psi(x, \xi) = \varphi(p(x, \xi)), \quad (x, \xi) \in \mathbf{R}^{2d} \setminus 0,$$

and it follows that  $\psi \in C^\infty(\mathbf{R}^{2d} \setminus 0)$ , and thus  $\chi \in C^\infty(\mathbf{R}^{2d})$ . The properties  $\chi|_{\Gamma_{z_0, \varepsilon} \setminus \bar{B}_r} \equiv 1$  and  $\text{supp } \chi \subseteq \Gamma_{z_0, 2\varepsilon} \setminus B_{r/2}$  follow.

From (3.14) we obtain

$$(3.16) \quad \partial_x^\alpha \partial_\xi^\beta \varphi(x, \xi) = \lambda^{|\alpha|+s|\beta|} (\partial_x^\alpha \partial_\xi^\beta \psi)(\lambda x, \lambda^s \xi), \quad (x, \xi) \in \mathbf{S}^{2d-1}, \quad \lambda > 0.$$

Let  $(y, \eta) \in \mathbf{R}^{2d}$  satisfy  $|(y, \eta)| > \frac{r}{2}$ . Then  $(y, \eta) = (\lambda x, \lambda^s \xi)$  for a unique  $(x, \xi) \in \mathbf{S}^{2d-1}$  and a unique

$$\lambda > \delta := \min \left( \frac{r}{2}, \left( \frac{r}{2} \right)^{\frac{1}{s}} \right) > 0.$$

We have

$$1 + |y| + |\eta|^{\frac{1}{s}} = 1 + \lambda(|x| + |\xi|^{\frac{1}{s}}) \leq 2(1 + \lambda).$$

Thus we obtain from (3.16) for any  $\alpha, \beta \in \mathbf{N}^d$

$$\left| \partial_y^\alpha \partial_\eta^\beta \psi(y, \eta) \right| \leq C_{\alpha, \beta} (1 + \lambda)^{-|\alpha| - s|\beta|} \lesssim (1 + |y| + |\eta|^{\frac{1}{s}})^{-|\alpha| - s|\beta|}.$$

From (3.15) we may conclude that  $\chi \in G^{0, s}$ .  $\square$

Sometimes it is useful to have the following alternative to the  $s$ -conic neighborhoods of Definition 3.4.

**Definition 3.6.** Suppose  $s, \varepsilon > 0$  and  $(x_0, \xi_0) \in \mathbf{S}^{2d-1}$ . Then

$$\begin{aligned} \tilde{\Gamma}_{s, (x_0, \xi_0), \varepsilon} &= \tilde{\Gamma}_{(x_0, \xi_0), \varepsilon} = \{(x, \xi) \in \mathbf{R}^{2d} \setminus 0 : (x, \xi) = (\lambda(x_0 + y), \lambda^s(\xi_0 + \eta)), \lambda > 0, (y, \eta) \in B_\varepsilon\} \\ &= \{(x, \xi) \in \mathbf{R}^{2d} \setminus 0 : \exists \lambda > 0 : (\lambda x, \lambda^s \xi) \in (x_0, \xi_0) + B_\varepsilon\} \subseteq T^* \mathbf{R}^d \setminus 0. \end{aligned}$$

Again  $\tilde{\Gamma}_{(x_0, \xi_0), \varepsilon}$  is  $s$ -conic.

The neighborhoods  $\Gamma_{s, (x_0, \xi_0), \varepsilon}$  and  $\tilde{\Gamma}_{s, (x_0, \xi_0), \varepsilon}$  are not identical, even if  $s = 1$  in which case  $p(x, \xi) = (x, \xi)/|(x, \xi)|$ . But by the following result the  $s$ -conic neighborhoods of the form  $\Gamma_{s, z_0, \varepsilon}$  and  $\tilde{\Gamma}_{s, z_0, \varepsilon}$  are equivalent topologically.

**Lemma 3.7.** Let  $z_0 \in \mathbf{S}^{2d-1}$ . For each  $\varepsilon > 0$  there exists  $\delta > 0$  such that

$$(3.17) \quad \Gamma_{z_0, \delta} \subseteq \tilde{\Gamma}_{z_0, \varepsilon}$$

and

$$(3.18) \quad \tilde{\Gamma}_{z_0, \delta} \subseteq \Gamma_{z_0, \varepsilon}.$$

*Proof.* Let  $z_0 = (x_0, \xi_0)$ . If  $\varepsilon > 0$  and  $(x, \xi) \in \Gamma_{z_0, \varepsilon} \cap \mathbf{S}^{2d-1}$  then  $(x, \xi) \in (x_0, \xi_0) + B_\varepsilon$  so  $(x, \xi) \in \tilde{\Gamma}_{z_0, \varepsilon}$ . Since both  $\Gamma_{z_0, \varepsilon}$  and  $\tilde{\Gamma}_{z_0, \varepsilon}$  are  $s$ -conic, this shows

$$\Gamma_{z_0, \varepsilon} \subseteq \tilde{\Gamma}_{z_0, \varepsilon}$$

for any  $\varepsilon > 0$ . Thus (3.17) follows with  $\delta = \varepsilon$ .

In order to show (3.18) let  $\varepsilon > 0$ , and suppose  $0 < \delta < 1$ . If  $(x, \xi) \in \tilde{\Gamma}_{z_0, \delta} \cap \mathbf{S}^{2d-1}$  then there exists  $\mu = \mu(x, \xi) > 0$  such that  $|(\mu x, \mu^s \xi) - (x_0, \xi_0)| < \delta$ . We have

$$\begin{aligned} \min(\mu, \mu^s) &\leq |(\mu x, \mu^s \xi)| < 1 + \delta, \\ \max(\mu, \mu^s) &\geq |(\mu x, \mu^s \xi)| > 1 - \delta \end{aligned}$$

which gives

$$(1 - \delta)^{\max(1, \frac{1}{s})} < \mu(x, \xi) < (1 + \delta)^{\max(1, \frac{1}{s})} \quad \forall (x, \xi) \in \tilde{\Gamma}_{z_0, \delta} \cap \mathbf{S}^{2d-1}.$$

Thus we may pick  $\delta < \varepsilon/2$  such that

$$\max(|1 - \mu(x, \xi)|, |1 - \mu(x, \xi)^s|) < \varepsilon/2 \quad \forall (x, \xi) \in \tilde{\Gamma}_{z_0, \delta} \cap \mathbf{S}^{2d-1}.$$

If  $(x, \xi) \in \tilde{\Gamma}_{z_0, \delta} \cap \mathbf{S}^{2d-1}$  then  $p(x, \xi) = (x, \xi)$  so we obtain

$$\begin{aligned} |p(x, \xi) - (x_0, \xi_0)| &= |(\mu(x, \xi)x, \mu(x, \xi)^s \xi) - (x_0, \xi_0) + ((1 - \mu(x, \xi))x, (1 - \mu(x, \xi)^s)\xi)| \\ &< \delta + \max(|1 - \mu(x, \xi)|, |1 - \mu(x, \xi)^s|) < \varepsilon. \end{aligned}$$

Again due to  $s$ -conic property of  $\tilde{\Gamma}_{z_0, \delta}$  and  $\Gamma_{z_0, \varepsilon}$ , this shows  $\tilde{\Gamma}_{z_0, \delta} \subseteq \Gamma_{z_0, \varepsilon}$ , that is (3.18).  $\square$

In Example 3.9 and in Section 6 we will use the following definition which is a natural anisotropic microlocal version of [26, Definition 25.1] as well as of [3, Eq. (1.11)] (cf. [6]).

**Definition 3.8.** Let  $s > 0$ ,  $z_0 \in \mathbf{R}^{2d} \setminus 0$ , and  $a \in G^{m, s}$ . Then  $z_0$  is called non-characteristic of order  $m_1 \leq m$ ,  $z_0 \notin \text{char}_{s, m_1}(a)$ , if there exists  $\varepsilon > 0$  such that, with  $\Gamma = \Gamma_{s, p(z_0), \varepsilon}$ ,

$$(3.19) \quad |a(x, \xi)| \geq C \mu_s(x, \xi)^{m_1}, \quad (x, \xi) \in \Gamma, \quad |x| + |\xi|^{\frac{1}{s}} \geq R,$$

$$(3.20)$$

$$|\partial_x^\alpha \partial_\xi^\beta a(x, \xi)| \lesssim |a(x, \xi)| \mu_s(x, \xi)^{-|\alpha| - s|\beta|}, \quad \alpha, \beta \in \mathbf{N}^d, \quad (x, \xi) \in \Gamma, \quad |x| + |\xi|^{\frac{1}{s}} \geq R,$$

for suitable  $C, R > 0$ .

If  $m_1 = m$  we write  $\text{char}_{s, m}(a) = \text{char}_s(a)$ , and then the condition (3.20) is then redundant. Note that  $\text{char}_{s, m_1}(a)$  is a closed  $s$ -conic subset of  $T^*\mathbf{R}^d \setminus 0$ , and  $\text{char}_{s, m_1}(a) \subseteq \text{char}_{s, m_2}(a)$  if  $m_1 \leq m_2 \leq m$ .

**Example 3.9.** In [3, 6] polynomial symbols of the form

$$(3.21) \quad a(x, \xi) = \sum_{\frac{|\alpha|}{k} + \frac{|\beta|}{m} \leq 1} c_{\alpha\beta} x^\alpha \xi^\beta, \quad x, \xi \in \mathbf{R}^d, \quad c_{\alpha\beta} \in \mathbf{C},$$

are studied for  $k, m \in \mathbf{N}$ . Then  $a \in G^{\max(k, m)}$  and  $a \in G^{k, \frac{k}{m}}$ . In fact we have for  $(x, \xi) \in \mathbf{S}^{2d-1}$  and  $\lambda > 0$

$$(\partial_x^\gamma \partial_\xi^\kappa a)(\lambda x, \lambda^{\frac{k}{m}} \xi) = \sum_{\frac{|\alpha|}{k} + \frac{|\beta|}{m} \leq 1} c_{\alpha\beta\gamma\kappa} \lambda^{|\alpha - \gamma| + \frac{k}{m}|\beta - \kappa|} x^{\alpha - \gamma} \xi^{\beta - \kappa}.$$

If  $(y, \eta) \in \mathbf{R}^{2d}$  and  $|(y, \eta)| \geq 1$  then we write  $(y, \eta) = (\lambda x, \lambda^{\frac{k}{m}} \xi)$  for  $(x, \xi) \in \mathbf{S}^{2d-1}$  and  $\lambda \geq 1$ . Since

$$|y| + |\eta|^{\frac{m}{k}} = \lambda \left( |x| + |\xi|^{\frac{m}{k}} \right) \asymp \lambda$$

we obtain

$$\begin{aligned} |\partial_x^\gamma \partial_\xi^\kappa a(y, \eta)| &\lesssim \sum_{\frac{|\alpha|}{k} + \frac{|\beta|}{m} \leq 1} (1 + |y| + |\eta|^{\frac{m}{k}})^k \left( \frac{|\alpha|}{k} + \frac{|\beta|}{m} \right) - |\gamma| - \frac{k}{m} |\kappa| \\ &\lesssim (1 + |y| + |\eta|^{\frac{m}{k}})^{k - |\gamma| - \frac{k}{m} |\kappa|} \end{aligned}$$

which proves that  $a \in G^{k, \frac{k}{m}}$ .

In [3, Eq. (1.11)] the symbol  $a$  given by (3.21) is called  $(k, m)$ -globally elliptic if

$$|a(x, \xi)| \geq C \left( |x| + |\xi|^{\frac{m}{k}} \right)^k, \quad |x| + |\xi|^{\frac{m}{k}} \geq R$$

for some  $C, R > 0$ . Thus Definition 3.8 can be viewed as a microlocalization of  $(k, m)$ -global ellipticity. A  $(k, m)$ -globally elliptic symbol as above satisfies  $\text{char}_{k/m}(a) = \text{char}_{k/m, k}(a) = \emptyset$ .

#### 4. ANISOTROPIC GABOR WAVE FRONT SETS

The following definition is inspired by H. Zhu's [28, Definition 1.5] of a quasi-homogeneous wave front set defined by two non-negative parameters. Zhu uses a semiclassical formulation whereas we use the STFT. As far as we know it is an open question to determine if the concepts coincide.

Given positive parameters  $t, s > 0$  we define the  $t, s$ -Gabor wave front set  $\text{WF}_g^{t,s}(u) \subseteq T^*\mathbf{R}^d \setminus 0$  of  $u \in \mathcal{S}'(\mathbf{R}^d)$ .

**Definition 4.1.** Suppose  $u \in \mathcal{S}'(\mathbf{R}^d)$ ,  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$ , and  $t, s > 0$ . A point  $z_0 = (x_0, \xi_0) \in T^*\mathbf{R}^d \setminus 0$  satisfies  $z_0 \notin \text{WF}_g^{t,s}(u)$  if there exists an open set  $U \subseteq T^*\mathbf{R}^d$  such that  $z_0 \in U$  and

$$(4.1) \quad \sup_{(x, \xi) \in U, \lambda > 0} \lambda^N |V_\varphi u(\lambda^t x, \lambda^s \xi)| < +\infty \quad \forall N \geq 0.$$

If  $s = t$  we have  $\text{WF}_g^{t,t}(u) = \text{WF}_g(u)$  which denotes the usual Gabor wave front set [14, 22]. In the definition of  $\text{WF}_g^{t,s}(u)$  only the fraction  $s/t$  matters. Therefore we may assume in the sequel that  $t = 1$ , and we write  $\text{WF}_g^{1,s}(u) = \text{WF}_g^s(u)$  for simplicity. We call  $\text{WF}_g^s(u)$  the anisotropic  $s$ -Gabor wave front set. It is clear that  $\text{WF}_g^s(u)$  is  $s$ -conic.

Referring to (2.1) and (2.2) we see that  $\text{WF}_g^s(u)$  records  $s$ -conic curves  $0 < \lambda \mapsto (\lambda x, \lambda^s \xi)$  where  $V_\varphi u$  does not behave like the STFT of a Schwartz function. From (2.1) it also follows that it suffices to check (4.1) for  $\lambda \geq L$  where  $L > 0$  may be arbitrarily large.

From (2.2) it follows that  $\text{WF}_g^s(u) = \emptyset$  if  $u \in \mathcal{S}(\mathbf{R}^d)$ . Conversely, if  $\text{WF}_g^s(u) = \emptyset$  then

$$\sup_{(x, \xi) \in \mathbf{S}^{2d-1}, \lambda > 0} \lambda^N |V_\varphi u(\lambda x, \lambda^s \xi)| < +\infty \quad \forall N \geq 0$$

due to the compactness of the unit sphere  $\mathbf{S}^{2d-1}$ . Given  $(y, \eta) \in T^*\mathbf{R}^d \setminus 0$  there is a unique  $\lambda > 0$  such that  $(y, \eta) = (\lambda x, \lambda^s \xi)$  and  $(x, \xi) \in \mathbf{S}^{2d-1}$ , and  $|(y, \eta)|^2 = \lambda^2|x|^2 + \lambda^{2s}|\xi|^2 \leq \lambda^2 + \lambda^{2s}$ . This implies that (2.2) is satisfied, and thus  $u \in \mathcal{S}(\mathbf{R}^d)$ . We have now shown that  $\text{WF}_g^s(u) = \emptyset$  if and only if  $u \in \mathcal{S}(\mathbf{R}^d)$ , for any  $s > 0$ .

**4.1. Window invariance and consequences.** First we show that  $\text{WF}_g^s(u)$  does not depend on the window function  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$ .

**Proposition 4.2.** *Let  $s > 0$ ,  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $z_0 \in T^*\mathbf{R}^d \setminus 0$ . If  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$  and (4.1) holds with  $t = 1$  for an open set  $U \subseteq T^*\mathbf{R}^d \setminus 0$  containing  $z_0$ , and  $\psi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$ , then there exists an open set  $V \subseteq U$  such that  $z_0 \in V$  and*

$$(4.2) \quad \sup_{(x, \xi) \in V, \lambda > 0} \lambda^N |V_\psi u(\lambda x, \lambda^s \xi)| < \infty, \quad \forall N \geq 0.$$

*Proof.* Since  $z_0 \in U \subseteq \mathbf{R}^{2d}$  where  $U$  is open we may pick an open set  $V \subseteq U$  such that  $z_0 \in V$  and  $V + B_\varepsilon \subseteq U$  for some  $0 < \varepsilon \leq 1$ , and we may assume

$$(4.3) \quad \sup_{z \in V} |z| \leq |z_0| + 1 := \mu.$$

By [11, Lemma 11.3.3] we have

$$|V_\psi u(z)| \leq (2\pi)^{-\frac{d}{2}} \|\varphi\|_{L^2}^{-2} |V_\varphi u| * |V_\psi \varphi|(z), \quad z \in \mathbf{R}^{2d}.$$

Let  $\lambda \geq 1$  and  $N \in \mathbf{N}$ . We have

$$\begin{aligned} & \lambda^N |V_\psi u(\lambda x, \lambda^s \xi)| \\ & \lesssim \iint_{\mathbf{R}^{2d}} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| |V_\psi \varphi(y, \eta)| \, dy \, d\eta \\ & = I_1 + I_2 \end{aligned}$$

where we split the integral into the two terms

$$\begin{aligned} I_1 &= \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| |V_\psi \varphi(y, \eta)| \, dy \, d\eta, \\ I_2 &= \iint_{\Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| |V_\psi \varphi(y, \eta)| \, dy \, d\eta \end{aligned}$$

where

$$\Omega_\lambda = \{(y, \eta) \in \mathbf{R}^{2d} : |(y, \eta)| < 2^{-\frac{1}{2}} \varepsilon \lambda^{\min(1, s)}\}.$$

First we estimate  $I_1$  when  $(x, \xi) \in V$ . From (2.1), (2.2) and (4.3) we obtain for some  $k \geq 0$  and any  $L \geq k$

$$\begin{aligned}
(4.4) \quad I_1 &\lesssim \lambda^N \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (\lambda x - y, \lambda^s \xi - \eta) \rangle^k |V_\psi \varphi(y, \eta)| \, dy \, d\eta \\
&\lesssim (1 + \mu^2)^{\frac{k}{2}} \lambda^{N+k \max(1,s)} \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (y, \eta) \rangle^k |V_\psi \varphi(y, \eta)| \, dy \, d\eta \\
&\lesssim (1 + \mu^2)^{\frac{k}{2}} \lambda^{N+k \max(1,s)} \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (y, \eta) \rangle^{k-L-2d-1} \, dy \, d\eta \\
&\lesssim \lambda^{N+k \max(1,s)} \left(1 + \frac{1}{2} \varepsilon^2 \lambda^{2 \min(1,s)}\right)^{\frac{1}{2}(k-L)} \iint_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{-2d-1} \, dy \, d\eta \\
&\lesssim \lambda^{N+k \max(1,s) + \min(1,s)(k-L)} \\
&\leq C_{N,L,\mu,\varepsilon}
\end{aligned}$$

for any  $\lambda \geq 1$ , provided we pick  $L \geq k + \min(1, s)^{-1} (N + k \max(1, s))$ . Here  $C_{N,L,\mu,\varepsilon} > 0$  is a constant that depends on  $N, L, \mu, \varepsilon$  but not on  $\lambda > 0$ . Thus we have obtained the required estimate for  $I_1$ .

It remains to estimate  $I_2$ . If  $(y, \eta) \in \Omega_\lambda$  then  $|y|^2 < \frac{1}{2} \varepsilon^2 \lambda^2$  and  $|\eta|^2 < \frac{1}{2} \varepsilon^2 \lambda^{2s}$  which implies  $(\lambda^{-1}y, \lambda^{-s}\eta) \in B_\varepsilon$ . Hence if  $(x, \xi) \in V$  then  $(x - \lambda^{-1}y, \xi - \lambda^{-s}\eta) \in U$  and we may use the estimate (4.1) with  $t = 1$ . This gives

$$\begin{aligned}
(4.5) \quad I_2 &= \iint_{\Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| |V_\psi \varphi(y, \eta)| \, dy \, d\eta \\
&\leq C_N \iint_{\mathbf{R}^{2d}} |V_\psi \varphi(y, \eta)| \, dy \, d\eta \\
&\lesssim C_N
\end{aligned}$$

for all  $\lambda \geq 1$ . Thus we have obtained the required estimate for  $I_2$ . Combining (4.4) and (4.5), we have proved (4.2).  $\square$

If  $\tilde{u}(x) = u(-x)$  then

$$(4.6) \quad V_{\tilde{\psi}} \tilde{u}(x, \xi) = V_\psi u(-x, -\xi).$$

Using Proposition 4.2 it follows that we have the following symmetry:

$$(4.7) \quad \tilde{u} = \pm u \implies \text{WF}_g^s(u) = -\text{WF}_g^s(u).$$

We also have

$$(4.8) \quad V_\psi \bar{u}(x, \xi) = \overline{V_{\tilde{\psi}} u(x, -\xi)}.$$

Referring to [24, Definition 3.2] we observe that

$$(4.9) \quad \text{WF}_g^s(u) \subseteq \text{WF}^{1,s}(u), \quad s > 0, \quad u \in \mathcal{S}'(\mathbf{R}^d),$$

where  $\text{WF}^{1,s}(u)$  is a particular case of a  $t, s$ -Gelfand–Shilov wave front set, a concept that requires super-exponential rather than super-polynomial decay along curves in phase space.

**4.2. Metaplectic properties.** The Gabor wave front set is symplectically invariant as (cf. [14, Proposition 2.2])

$$(4.10) \quad \text{WF}_g(\mu(\chi)u) = \chi \text{WF}_g(u), \quad \chi \in \text{Sp}(d, \mathbf{R}), \quad u \in \mathcal{S}'(\mathbf{R}^d).$$

When  $s \neq 1$  the  $s$ -Gabor wave front set  $\text{WF}_g^s(u)$  is no longer symplectically invariant. Nevertheless, two of the generators of the symplectic group behave invariantly in certain individual senses which we now describe. By [9, Proposition 4.10] each matrix  $\chi \in \text{Sp}(d, \mathbf{R})$  is a finite product of matrices in  $\text{Sp}(d, \mathbf{R})$  of the form

$$\mathcal{J}, \quad \begin{pmatrix} A^{-1} & 0 \\ 0 & A^T \end{pmatrix}, \quad \begin{pmatrix} I & 0 \\ B & I \end{pmatrix},$$

for  $A \in \text{GL}(d, \mathbf{R})$  and  $B \in \mathbf{R}^{d \times d}$  symmetric. The corresponding metaplectic operators are  $\mu(\mathcal{J}) = \mathcal{F}$ ,

$$\mu \begin{pmatrix} A^{-1} & 0 \\ 0 & A^T \end{pmatrix} f(x) = |A|^{\frac{1}{2}} f(Ax),$$

if  $A \in \text{GL}(d, \mathbf{R})$ , and

$$\mu \begin{pmatrix} I & 0 \\ B & I \end{pmatrix} f(x) = e^{\frac{i}{2}\langle Bx, x \rangle} f(x),$$

if  $B \in \mathbf{R}^{d \times d}$  is symmetric.

**Proposition 4.3.** *Let  $s > 0$  and  $u \in \mathcal{S}'(\mathbf{R}^d)$ . Then we have*

(i)

$$\text{WF}_g^s(\widehat{u}) = \mathcal{J} \text{WF}_g^{\frac{1}{s}}(u).$$

(ii) *If  $A \in \text{GL}(d, \mathbf{R})$  and  $u_A(x) = |A|^{\frac{1}{2}} u(Ax)$  then*

$$\text{WF}_g^s(u_A) = \begin{pmatrix} A^{-1} & 0 \\ 0 & A^T \end{pmatrix} \text{WF}_g^s(u).$$

(iii) *If  $B \in \mathbf{R}^{d \times d}$  is symmetric and  $v(x) = e^{\frac{i}{2}\langle Bx, x \rangle} u(x)$  then if  $s = 1$*

$$(4.11) \quad \text{WF}_g^s(v) = \begin{pmatrix} I & 0 \\ B & I \end{pmatrix} \text{WF}_g^s(u),$$

*if  $s > 1$  then*

$$(4.12) \quad \text{WF}_g^s(v) = \text{WF}_g^s(u),$$

*and finally if  $0 < s < 1$  then*

$$(4.13) \quad (x, \xi) \in \text{WF}_g^s(u) \quad \text{for some } \xi \in \mathbf{R}^d \implies (x, Bx) \in \text{WF}_g(v).$$

*Proof.* Let  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$ . We have from the proof of [7, Corollary 4.5]

$$(4.14) \quad |V_{\mu(\chi)\varphi}(\mu(\chi)u)(\chi(x, \xi))| = |V_\varphi u(x, \xi)|$$

for all  $\chi \in \text{Sp}(d, \mathbf{R})$ .

(i) If  $\chi = \mathcal{J}$  we obtain

$$|V_{\widehat{\varphi}} \widehat{u}(\mathcal{J}(x, \xi))| = |V_{\widehat{\varphi}} \widehat{u}(\xi, -x)| = |V_\varphi u(x, \xi)|.$$

From this and Proposition 4.2 it follows that  $(x, \xi) \notin \text{WF}_g^{\frac{1}{s}}(u)$  if and only if  $\mathcal{J}(x, \xi) \notin \text{WF}_g^s(\widehat{u})$  which proves claim (i).

(ii) Next we insert  $u_A$  for  $A \in \text{GL}(d, \mathbf{R})$  into (4.14) which gives

$$|V_{\psi_A} u_A(A^{-1}x, A^T \xi)| = |V_\psi u(x, \xi)|.$$

Note that  $\psi_A \in \mathcal{S}(\mathbf{R}^d) \setminus 0$ . We obtain  $(x, \xi) \notin \text{WF}_g^s(u)$  if and only if  $(A^{-1}x, A^T \xi) \notin \text{WF}_g^s(u_A)$  which shows claim (ii).

(iii) When  $s = 1$  (4.11) is a particular case of (4.10).

Suppose  $s \neq 1$ . With  $\psi(x) = e^{\frac{i}{2}\langle Bx, x \rangle} \varphi(x) \in \mathcal{S}(\mathbf{R}^d) \setminus 0$  we obtain from (4.14)

$$|V_\varphi u(x, \xi)| = |V_\psi v(x, Bx + \xi)|$$

or equivalently

$$|V_\varphi u(x, -Bx + \xi)| = |V_\psi v(x, \xi)|.$$

If  $\lambda > 0$  then

$$(4.15) \quad |V_\varphi u(\lambda x, \lambda^s \xi)| = |V_\psi v(\lambda x, \lambda^s(\lambda^{1-s} Bx + \xi))| = |V_\psi v(\lambda x, \lambda(Bx + \lambda^{s-1} \xi))|$$

and

$$(4.16) \quad |V_\psi v(\lambda x, \lambda^s \xi)| = |V_\varphi u(\lambda x, \lambda^s(-\lambda^{1-s} Bx + \xi))| = |V_\varphi u(\lambda x, \lambda(-Bx + \lambda^{s-1} \xi))|.$$

Suppose  $s > 1$  and  $0 \neq (x_0, \xi_0) \notin \text{WF}_g^s(u)$ . Then for some  $\varepsilon > 0$  we have

$$(4.17) \quad \sup_{x \in x_0 + B_\varepsilon, \xi \in \xi_0 + B_{2\varepsilon}, \lambda > 0} \lambda^N |V_\varphi u(\lambda x, \lambda^s \xi)| < +\infty \quad \forall N \geq 0.$$

We have  $\lambda^{1-s}|Bx| < \varepsilon$  when  $x \in x_0 + B_\varepsilon$  if  $\lambda \geq L$  for  $L \geq 1$  sufficiently large. Thus  $\xi - \lambda^{1-s} Bx \in \xi_0 + B_{2\varepsilon}$  if  $\xi \in \xi_0 + B_\varepsilon$  and  $\lambda \geq L$ . From (4.16) and (4.17) we obtain

$$\sup_{x \in x_0 + B_\varepsilon, \xi \in \xi_0 + B_\varepsilon, \lambda > 0} \lambda^N |V_\psi v(\lambda x, \lambda^s \xi)| < +\infty \quad \forall N \geq 0$$

which shows that  $(x_0, \xi_0) \notin \text{WF}_g^s(v)$ . Thus  $\text{WF}_g^s(v) \subseteq \text{WF}_g^s(u)$ . Likewise one shows the opposite inclusion using (4.15). We have now proved (4.12).

Suppose  $0 < s < 1$  and  $0 \neq (x_0, Bx_0) \notin \text{WF}_g(v)$ . Then for some  $\varepsilon > 0$  we have

$$(4.18) \quad \sup_{x \in x_0 + B_\varepsilon, \xi \in Bx_0 + B_{2|B|\varepsilon}, \lambda > 0} \lambda^N |V_\psi v(\lambda x, \lambda \xi)| < +\infty \quad \forall N \geq 0.$$

Let  $\eta_0 \in \mathbf{R}^d$ . We have  $Bx + \lambda^{s-1} \xi \in Bx_0 + B_{2|B|\varepsilon}$  when  $x \in x_0 + B_\varepsilon$  and  $\xi \in \eta_0 + B_\varepsilon$  if  $\lambda \geq L$  for  $L \geq 1$  sufficiently large. From (4.15) we obtain

$$\sup_{x \in x_0 + B_\varepsilon, \xi \in \eta_0 + B_\varepsilon, \lambda > 0} \lambda^N |V_\varphi u(\lambda x, \lambda^s \xi)| < +\infty \quad \forall N \geq 0$$

and it follows that  $(x_0, \eta_0) \notin \text{WF}_g^s(u)$ . We have shown (4.13). □

## 5. MICROLOCALITY FOR ANISOTROPIC GABOR WAVE FRONT SETS

Let  $m \in \mathbf{R}$ ,  $0 \leq \rho \leq 1$ ,  $a \in G_\rho^m$  and  $\varphi \in \mathcal{S}(\mathbf{R}^{2d}) \setminus 0$ . According to [5, Proposition 3.2] the estimates

$$(5.1) \quad |V_\varphi a(z, \zeta)| \lesssim \langle z \rangle^m \langle \zeta \rangle^{-L}, \quad (z, \zeta) \in T^*\mathbf{R}^{2d},$$

hold for any  $L \geq 0$ . Note that the case  $\rho = 0$  is included, so (5.1) is valid under the assumption

$$\left| \partial_x^\alpha \partial_\xi^\beta a(x, \xi) \right| \lesssim \langle (x, \xi) \rangle^m, \quad \alpha, \beta \in \mathbf{N}^d.$$

The next result concerns microlocality with respect to the  $s$ -Gabor wave front set for pseudodifferential operators in the isotropic Shubin calculus. Due to (3.6) the result is also true for the anisotropic Shubin symbols  $G^{m,s}$ .

**Proposition 5.1.** *Let  $s > 0$ ,  $m \in \mathbf{R}$  and  $0 \leq \rho \leq 1$ . If  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $a \in G_\rho^m$  then*

$$(5.2) \quad \text{WF}_g^s(a^w(x, D)u) \subseteq \text{WF}_g^s(u).$$

*Proof.* Pick  $\varphi \in \mathcal{S}(\mathbf{R}^d)$  such that  $\|\varphi\|_{L^2} = 1$ . Denoting the formal adjoint of  $a^w(x, D)$  by  $a^w(x, D)^*$ , (2.4) gives for  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $z \in \mathbf{R}^{2d}$

$$\begin{aligned} (2\pi)^{\frac{d}{2}} V_\varphi(a^w(x, D)u)(z) &= (a^w(x, D)u, \Pi(z)\varphi) \\ &= (u, a^w(x, D)^* \Pi(z)\varphi) \\ &= \int_{\mathbf{R}^{2d}} V_\varphi u(w) (\Pi(w)\varphi, a^w(x, D)^* \Pi(z)\varphi) dw \\ &= \int_{\mathbf{R}^{2d}} V_\varphi u(w) (a^w(x, D) \Pi(w)\varphi, \Pi(z)\varphi) dw \\ &= \int_{\mathbf{R}^{2d}} V_\varphi u(z - w) (a^w(x, D) \Pi(z - w)\varphi, \Pi(z)\varphi) dw. \end{aligned}$$

By e.g. [12, Lemma 3.1], or a computation using (2.9), we have

$$|(a^w(x, D) \Pi(z - w)\varphi, \Pi(z)\varphi)| = \left| V_\Phi a \left( z - \frac{w}{2}, \mathcal{J}w \right) \right|$$

where  $\Phi$  is the Wigner distribution  $\Phi = W(\varphi, \varphi) \in \mathcal{S}(\mathbf{R}^{2d})$ .

Combining the preceding identities we deduce

$$(5.3) \quad |V_\varphi(a^w(x, D)u)(z)| \lesssim \int_{\mathbf{R}^{2d}} |V_\varphi u(z - w)| \left| V_\Phi a \left( z - \frac{w}{2}, \mathcal{J}w \right) \right| dw.$$

Suppose  $0 \neq z_0 \notin \text{WF}_g^s(u)$ . Then there exists an open set  $U$  such that  $z_0 \in U$  and (4.1) holds with  $t = 1$ . We pick an open set  $V$  such that  $z_0 \in V$  and  $V + B_\varepsilon \subseteq U$  for some  $0 < \varepsilon \leq 1$ , and we may assume that (4.3) holds.

Let  $\lambda \geq 1$  and  $N \in \mathbf{N}$ . We have

$$\begin{aligned} &\lambda^N |V_\varphi(a^w(x, D)u)(\lambda x, \lambda^s \xi)| \\ &\lesssim \iint_{\mathbf{R}^{2d}} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\ &= I_1 + I_2 \end{aligned}$$

where the integral is decomposed into the two terms

$$I_1 = \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta,$$

$$I_2 = \iint_{\Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta$$

where

$$\Omega_\lambda = \{(y, \eta) \in \mathbf{R}^{2d} : |(y, \eta)| < 2^{-\frac{1}{2}} \varepsilon \lambda^{\min(1, s)}\}.$$

First we estimate  $I_1$  when  $(x, \xi) \in V$ . From (2.1), (4.3) and (5.1) we obtain for some  $k \geq 0$  and any  $L \geq k + |m|$

(5.4)

$$\begin{aligned} I_1 &\lesssim \lambda^N \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (\lambda x - y, \lambda^s \xi - \eta) \rangle^k \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\ &\lesssim (1 + \mu^2)^{\frac{k}{2}} \lambda^{N+k \max(1, s)} \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (y, \eta) \rangle^k \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\ &\lesssim (1 + \mu^2)^{\frac{k+|m|}{2}} \lambda^{N+(k+|m|) \max(1, s)} \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (y, \eta) \rangle^{k+|m|-L-2d-1} dy d\eta \\ &\lesssim \lambda^{N+(k+|m|) \max(1, s)} \iint_{\mathbf{R}^{2d} \setminus \Omega_\lambda} \langle (y, \eta) \rangle^{k+|m|-L} \langle (y, \eta) \rangle^{-2d-1} dy d\eta \\ &\leq \lambda^{N+(k+|m|) \max(1, s)} \left( 1 + \frac{1}{2} \varepsilon^2 \lambda^{2 \min(1, s)} \right)^{\frac{1}{2}(k+|m|-L)} \iint_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{-2d-1} dy d\eta \\ &\lesssim \lambda^{N+(k+|m|) \max(1, s) + \min(1, s)(k+|m|-L)} \\ &\leq C_{N, L, a, \mu, \varepsilon} \end{aligned}$$

for any  $\lambda \geq 1$ , provided we pick  $L \geq k + |m| + \min(1, s)^{-1} (N + (k + |m|) \max(1, s))$ . Here  $C_{N, L, a, \mu, \varepsilon} > 0$  is a constant that depends on  $N, L, a, \mu, \varepsilon$  but not on  $\lambda > 0$ . Thus we have obtained the required estimate for  $I_1$ .

It remains to estimate  $I_2$ . If  $(y, \eta) \in \Omega_\lambda$  then  $|y|^2 < \frac{1}{2} \varepsilon^2 \lambda^2$  and  $|\eta|^2 < \frac{1}{2} \varepsilon^2 \lambda^{2s}$  which implies  $(\lambda^{-1}y, \lambda^{-s}\eta) \in B_\varepsilon$ . Hence if  $(x, \xi) \in V$  then  $(x - \lambda^{-1}y, \xi - \lambda^{-s}\eta) \in U$  and we may use the estimate (4.1) with  $t = 1$ .

This gives for any  $L \geq 0$  and a constant  $C_{N,s,m} > 0$ , using (5.1) and (4.3)  
(5.5)

$$\begin{aligned}
I_2 &= \iint_{\Omega_\lambda} \lambda^N |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\
&= \lambda^{-|m| \max(1,s)} \iint_{\Omega_\lambda} \lambda^{N+|m| \max(1,s)} |V_\varphi u(\lambda(x - \lambda^{-1}y), \lambda^s(\xi - \lambda^{-s}\eta))| \\
&\quad \times \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\
&\leq C_{N,s,m} \lambda^{-|m| \max(1,s)} \iint_{\Omega_\lambda} \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\
&\leq C_{N,s,m} \lambda^{-|m| \max(1,s) + |m| \max(1,s)} \iint_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^{|m|-L} dy d\eta \\
&\lesssim C_{N,s,m}
\end{aligned}$$

provided  $L > |m| + 2d$ , for all  $\lambda \geq 1$ . Thus we have obtained the required estimate for  $I_2$ . Combining (5.4) and (5.5), referring to Definition 4.1, we may conclude that  $z_0 \notin \text{WF}_g^s(a^w(x, D)u)$  and hence we have proved (5.2).  $\square$

A consequence of Proposition 5.1 is the invariance of the anisotropic Gabor wave front set under translations and modulations, as well as time-frequency shifts [11].

**Corollary 5.2.** *Suppose  $s > 0$ . For any  $z \in \mathbf{R}^{2d}$  and any  $u \in \mathcal{S}'(\mathbf{R}^d)$  we have*

$$\text{WF}_g^s(\Pi(z)u) = \text{WF}_g^s(u).$$

*Proof.* Let  $z = (x, \xi) \in \mathbf{R}^{2d}$ . By a calculation it is verified that  $\Pi(x, \xi) = a_{x,\xi}^w(x, D)$  where

$$a_{x,\xi}(y, \eta) = e^{\frac{i}{2}\langle x, \xi \rangle + i(\langle y, \xi \rangle - \langle x, \eta \rangle)}, \quad (y, \eta) \in \mathbf{R}^{2d}.$$

For any  $\alpha, \beta \in \mathbf{N}^d$  we have

$$\left| \partial_y^\alpha \partial_\eta^\beta a_{x,\xi}(y, \eta) \right| = |\xi^\alpha x^\beta| := C_{\alpha,\beta}$$

where we may consider  $|\xi^\alpha x^\beta| \geq 0$  as a constant as a function of  $(y, \eta) \in \mathbf{R}^{2d}$ . This implies that  $a_{x,\xi} \in G_0^0$ . Thus we may apply Proposition 5.1 which gives

$$\text{WF}_g^s(\Pi(z)u) \subseteq \text{WF}_g^s(u).$$

The opposite inclusion follows from  $u = e^{-i\langle x, \xi \rangle} \Pi(-(x, \xi)) \Pi(x, \xi) u$ .  $\square$

We finish this section with the anisotropic Gabor wave front sets for a few important tempered distributions.

**Proposition 5.3.** *If  $s > 0$  then:*

- (i) *for any  $x \in \mathbf{R}^d$  and any  $\alpha \in \mathbf{N}^d$*   
(5.6)  $\text{WF}_g^s(D^\alpha \delta_x) = \{0\} \times (\mathbf{R}^d \setminus 0);$
- (ii) *for any  $\alpha \in \mathbf{N}^d$*   
 $\text{WF}_g^s(x^\alpha) = (\mathbf{R}^d \setminus 0) \times \{0\};$

(iii) for any  $\xi \in \mathbf{R}^d$

$$\mathrm{WF}_g^s(e^{i\langle \cdot, \xi \rangle}) = (\mathbf{R}^d \setminus 0) \times \{0\}.$$

*Proof.* Due to Corollary 5.2 we may assume  $x = 0$  in (i) and  $\xi = 0$  in (iii). By Proposition 4.3 (i) it suffices to show (i), since  $\widehat{D^\alpha \delta_0}(\xi) = (2\pi)^{-\frac{d}{2}} \xi^\alpha$ .

Let  $\varphi \in \mathcal{S}(\mathbf{R}^d)$  satisfy  $\varphi \equiv 1$  in a neighborhood of the origin. We have

$$V_\varphi D^\alpha \delta_0(x, \xi) = (2\pi)^{-\frac{d}{2}} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \xi^\beta \overline{D^{\alpha-\beta} \varphi(-x)}.$$

If  $\xi \neq 0$  we obtain for  $\lambda > 0$

$$V_\varphi D^\alpha \delta_0(0, \lambda^s \xi) = (2\pi)^{-\frac{d}{2}} \lambda^{s|\alpha|} \xi^\alpha$$

which does not decay as a function of  $\lambda$ . Thus

$$(5.7) \quad \{0\} \times (\mathbf{R}^d \setminus 0) \subseteq \mathrm{WF}_g^s(D^\alpha \delta_0).$$

Suppose on the other hand  $(x_0, \xi_0) \in T^*\mathbf{R}^d$  and  $x_0 \neq 0$ . If  $0 < \varepsilon < |x_0|/2$ ,  $x \in x_0 + B_\varepsilon$ ,  $\xi \in \xi_0 + B_\varepsilon$  and  $\lambda \geq 1$  then for any  $n \in \mathbf{N}$  we have

$$\begin{aligned} |V_\varphi D^\alpha \delta_0(\lambda x, \lambda^s \xi)| &= (2\pi)^{-\frac{d}{2}} \left| \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \lambda^{s|\beta|} \xi^\beta \overline{D^{\alpha-\beta} \varphi(-\lambda x)} \right| \\ &\lesssim \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \lambda^{s|\beta|} |\xi|^\beta \langle \lambda x \rangle^{-n} \\ &\lesssim \lambda^{s|\alpha| - n}. \end{aligned}$$

This shows

$$\mathrm{WF}_g^s(D^\alpha \delta_0) \subseteq \{0\} \times (\mathbf{R}^d \setminus 0)$$

so combining with (5.7) we have shown (5.6) when  $x = 0$ .  $\square$

## 6. MICROELLIPTICITY FOR ANISOTROPIC GABOR WAVE FRONT SETS

The main result in this section is the microelliptic inclusion expressed in Theorem 6.4. To get there we need a definition and several auxiliary results.

**Definition 6.1.** Suppose  $s > 0$ ,  $a \in G^{m,s}$  and let  $p$  be the projection (3.2). The  $s$ -conical support  $\mathrm{conesupp}_s(a) \subseteq T^*\mathbf{R}^d \setminus 0$  of  $a$  is defined as follows. A point  $z_0 \in T^*\mathbf{R}^d \setminus 0$  satisfies  $z_0 \notin \mathrm{conesupp}_s(a)$  if there exists  $\varepsilon > 0$  such that

$$\begin{aligned} &\mathrm{supp}(a) \cap \overline{\{z \in \mathbf{R}^{2d} \setminus 0, |p(z) - p(z_0)| < \varepsilon\}} \\ &= \mathrm{supp}(a) \cap \overline{\Gamma}_{p(z_0), \varepsilon} \quad \text{is compact in } \mathbf{R}^{2d}. \end{aligned}$$

Clearly  $\mathrm{conesupp}_s(a) \subseteq T^*\mathbf{R}^d \setminus 0$  is  $s$ -conic.

**Proposition 6.2.** Let  $s > 0$ . If  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $a \in G_0^m$  then

$$(6.1) \quad \mathrm{WF}_g^s(a^w(x, D)u) \subseteq \mathrm{conesupp}_s(a).$$

*Proof.* We have  $|\partial^\beta a(w)| \lesssim \langle w \rangle^m$  for any  $\beta \in \mathbf{N}^{2d}$ . We may assume that  $\text{conesupp}_s(a) \neq T^*\mathbf{R}^d \setminus 0$  since the inclusion is trivial otherwise. Let  $0 \neq z_0 \notin \text{conesupp}_s(a)$ . We may assume  $|z_0| = 1$ .

By Lemma 3.7 we may assume that

$$(6.2) \quad \text{supp}(a) \subseteq B_R \cup \left( \mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon} \right)$$

for some  $R > 0$  and  $0 < \varepsilon < 1$ .

Let  $\varphi \in \mathcal{S}(\mathbf{R}^d) \setminus 0$  and set  $\Phi = W(\varphi, \varphi) \in \mathcal{S}(\mathbf{R}^{2d})$ . We start by proving the following estimate for any  $\alpha, \beta \in \mathbf{N}^{2d}$  such that  $\beta \leq \alpha$ , any  $\lambda \geq 1$ ,  $(x, \xi) \in z_0 + B_\varepsilon$ , and any  $L \geq |m| + 2d + 1$ . We have

$$(6.3) \quad \int_{\mathbf{R}^{2d}} \left| \partial^\beta a(w) \right| \left| \partial^{\alpha-\beta} \Phi \left( w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right) \right| dw \lesssim \lambda^{-L \min(1,s) + |m| \max(1,s)} \langle (y, \eta) \rangle^{2L}.$$

In fact using Peetre's inequality we obtain on the one hand for any  $L \geq 0$

$$(6.4) \quad \begin{aligned} & \int_{B_R} \left| \partial^\beta a(w) \right| \left| \partial^{\alpha-\beta} \Phi \left( w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right) \right| dw \\ & \lesssim \int_{B_R} \left\langle w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right\rangle^{-L} dw \lesssim \int_{B_R} \langle w \rangle^L \langle (y, \eta) \rangle^L \langle (\lambda x, \lambda^s \xi) \rangle^{-L} dw \\ & \lesssim \lambda^{-L \min(1,s)} \langle (y, \eta) \rangle^L. \end{aligned}$$

On the other hand, since

$$|(\lambda^{-1}u, \lambda^{-s}\theta) - z_0| \geq 2\varepsilon \quad \forall \lambda > 0 \quad \forall (u, \theta) \in \mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}$$

we have for  $(x, \xi) \in z_0 + B_\varepsilon$

$$|(\lambda^{-1}u, \lambda^{-s}\theta) - (x, \xi)| \geq \varepsilon \quad \forall \lambda > 0 \quad \forall (u, \theta) \in \mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}.$$

It follows that for  $\lambda \geq 1$ ,  $(x, \xi) \in z_0 + B_\varepsilon$  and  $w = (u, \theta) \in \mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}$  we have

$$\begin{aligned} |w - (\lambda x, \lambda^s \xi)|^2 &= \lambda^2 |\lambda^{-1}u - x|^2 + \lambda^{2s} |\lambda^{-s}\theta - \xi|^2 \\ &\geq \lambda^{2 \min(1,s)} \varepsilon^2. \end{aligned}$$

This gives for  $(x, \xi) \in z_0 + B_\varepsilon$  and any  $L \geq |m| + 2d + 1$

$$(6.5) \quad \begin{aligned} & \int_{\mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}} \left| \partial^\beta a(w) \right| \left| \partial^{\alpha-\beta} \Phi \left( w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right) \right| dw \\ & \lesssim \int_{\mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}} \langle w \rangle^{|m|} \left\langle w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right\rangle^{-2L} dw \\ & \lesssim \langle (y, \eta) \rangle^{2L} \int_{\mathbf{R}^{2d} \setminus \tilde{\Gamma}_{z_0, 2\varepsilon}} \langle w \rangle^{|m|} \langle w - (\lambda x, \lambda^s \xi) \rangle^{-L} \langle w - (\lambda x, \lambda^s \xi) \rangle^{-L} dw \\ & \lesssim \lambda^{-L \min(1,s)} \langle (y, \eta) \rangle^{2L} \int_{\mathbf{R}^{2d}} \langle w + (\lambda x, \lambda^s \xi) \rangle^{|m|} \langle w \rangle^{-L} dw \\ & \lesssim \lambda^{-L \min(1,s)} \langle (y, \eta) \rangle^{2L} \langle (\lambda x, \lambda^s \xi) \rangle^{|m|} \int_{\mathbf{R}^{2d}} \langle w \rangle^{|m|-L} dw \\ & \lesssim \lambda^{-L \min(1,s) + |m| \max(1,s)} \langle (y, \eta) \rangle^{2L}. \end{aligned}$$

Combining (6.2), (6.4) and (6.5) we have now shown (6.3).

Next we observe that integration by parts gives for any  $\alpha \in \mathbf{N}^{2d}$  and  $z, \zeta \in \mathbf{R}^{2d}$

$$\begin{aligned} |\zeta^\alpha V_\Phi a(z, \zeta)| &= (2\pi)^{-d} \left| \int_{\mathbf{R}^{2d}} a(w) \partial_w^\alpha \left( e^{-i\langle w, \zeta \rangle} \right) \overline{\Phi(w-z)} dw \right| \\ &\lesssim \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbf{R}^{2d}} |\partial^\beta a(w)| |\partial^{\alpha-\beta} \Phi(w-z)| dw. \end{aligned}$$

Combining this with (6.3) we obtain for  $(x, \xi) \in z_0 + B_\varepsilon$ , and any  $M \in \mathbf{N}$ ,  $L \geq |m| + 2d + 1$  and  $\lambda \geq 1$

$$\begin{aligned} &\langle (y, \eta) \rangle^{2(M+L)} \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| \\ &\lesssim \max_{|\alpha| \leq 2(M+L)} \left| (y, \eta)^\alpha V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| \\ &\lesssim \max_{|\alpha| \leq 2(M+L)} \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} \int_{\mathbf{R}^{2d}} |\partial^\beta a(w)| |\partial^{\alpha-\beta} \Phi \left( w - \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2} \right) \right)| dw \\ &\lesssim \lambda^{-L \min(1, s) + |m| \max(1, s)} \langle (y, \eta) \rangle^{2L}. \end{aligned}$$

Given any  $N, M \geq 0$  we may pick  $L \geq 0$  such that  $L \min(1, s) - |m| \max(1, s) \geq N$ . We thus have

$$(6.6) \quad \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| \lesssim \lambda^{-N} \langle (y, \eta) \rangle^{-M}$$

for any  $N, M \geq 0$ ,  $\lambda \geq 1$  and  $(x, \xi) \in z_0 + B_\varepsilon$ .

Finally we prove that  $z_0 \notin \text{WF}_g^s(a^w(x, D)u)$ . We use (5.3) from the proof of Proposition 5.1 and (6.6). This gives for  $(x, \xi) \in z_0 + B_\varepsilon$ , using (2.1) for some  $k \geq 0$ , for any  $N, M \geq 0$ ,  $\lambda \geq 1$

$$\begin{aligned} &|V_\varphi(a^w(x, D)u)(\lambda x, \lambda^s \xi)| \\ &\lesssim \int_{\mathbf{R}^{2d}} |V_\varphi u((\lambda x, \lambda^s \xi) - (y, \eta))| \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\ &\lesssim \langle (\lambda x, \lambda^s \xi) \rangle^k \int_{\mathbf{R}^{2d}} \langle (y, \eta) \rangle^k \left| V_\Phi a \left( \lambda x - \frac{y}{2}, \lambda^s \xi - \frac{\eta}{2}, \eta, -y \right) \right| dy d\eta \\ &\lesssim \lambda^{k \max(1, s) - N} \end{aligned}$$

provided  $M \geq k + 2d + 1$ . Since  $N \geq 0$  is arbitrary we have shown  $z_0 \notin \text{WF}_g^s(a^w(x, D)u)$ , and thus (6.1).  $\square$

As another tool for the microellipticity result Theorem 6.4 we need the following lemma where we use Definition 3.8.

**Lemma 6.3.** *Suppose  $s > 0$ ,  $a \in G^{m, s}$  and  $\text{char}_{s, m_1}(a) \neq T^*\mathbf{R}^d \setminus 0$  for some  $m_1 \leq m$ . Let  $\Gamma \subseteq T^*\mathbf{R}^d \setminus 0$  be a closed  $s$ -conic set such that  $\text{char}_{s, m_1}(a) \cap \Gamma = \emptyset$ . Then there exists  $\rho > 0$  such that for any  $\chi \in G^{0, s}$  with  $\text{supp}(\chi) \subseteq \Gamma \setminus B_\rho$ , there exists  $b \in G^{-m_1, s}$  such that*

$$b \# a = \chi + r$$

where  $r \in \mathcal{S}(\mathbf{R}^{2d})$ .

*Proof.* The proof follows established principles in pseudodifferential calculus. Therefore we content ourselves with a sketch of the main steps of the construction of the microlocal parametrix  $b$ .

As a first approximation set  $b_0 := a^{-1}\chi$ . The estimates

$$|\partial_x^\alpha \partial_\xi^\beta (a^{-1})(x, \xi)| \leq C_{\alpha\beta} |a(x, \xi)|^{-1} \mu_s(x, \xi)^{-|\alpha|-s|\beta|}, \quad \alpha, \beta \in \mathbf{N}^d, \quad (x, \xi) \in \Gamma, \quad |x| + |\xi|^{\frac{1}{s}} \geq R,$$

are consequences of the non-characteristic estimates (3.19), (3.20) and induction.

By Leibniz' rule they imply the estimates

$$|\partial_x^\alpha \partial_\xi^\beta b_0(x, \xi)| \leq C_{\alpha\beta} |a(x, \xi)|^{-1} \mu_s(x, \xi)^{-|\alpha|-s|\beta|}, \quad \alpha, \beta \in \mathbf{N}^d, \quad |x| + |\xi|^{\frac{1}{s}} \geq R,$$

and consequently  $b_0 \in G^{-m_1, s}$  if  $\rho > 0$  is sufficiently large.

Then, by (3.10) and again the non-characteristic estimates (3.19) and (3.20) it follows that  $b_0 \# a = \chi + r_0 + r_{0, \mathcal{S}}$  with  $r_0 \in G^{-(1+s), s}$  satisfying  $\text{supp}(r_0) \subseteq \text{supp}(\chi)$  and  $r_{0, \mathcal{S}} \in \mathcal{S}(\mathbf{R}^{2d})$ . Subsequently, setting  $b_1 := -a^{-1}r_0$ , we notice that we obtain the estimates

$$|\partial_x^\alpha \partial_\xi^\beta b_1(x, \xi)| \leq C_{\alpha\beta} |a(x, \xi)|^{-1} \mu_s(x, \xi)^{-(1+s)-|\alpha|-s|\beta|}, \quad \alpha, \beta \in \mathbf{N}^d, \quad |x| + |\xi|^{\frac{1}{s}} \geq R,$$

and consequently  $b_1 \in G^{-m_1-(1+s), s}$ .

This gives

$$(b_0 + b_1) \# a = \chi + r_0 + r_{0, \mathcal{S}} - r_0 + r_1 + r_{1, \mathcal{S}} = \chi + r_1 + r_{0, \mathcal{S}} + r_{1, \mathcal{S}}$$

with  $r_1 \in G^{-2(1+s), s}$ ,  $\text{supp}(r_1) \subseteq \text{supp}(\chi)$  and  $r_{1, \mathcal{S}} \in \mathcal{S}(\mathbf{R}^{2d})$ . Constructing in this way recursively  $b_{j+1} := -a^{-1}r_j \in G^{-m_1-(s+1)(j+1), s}$  and  $r_{j+1} \in G^{-(s+1)(j+2), s}$  with  $\text{supp}(r_{j+1}) \subseteq \text{supp}(\chi)$ ,  $j = 1, 2, \dots$ , one obtains a sequence of symbols  $(b_j)_{j \geq 0}$ .

Finally set  $b \sim \sum_{j=0}^{\infty} b_j \in G^{-m_1, s}$ . The symbol  $b$  satisfies  $b \# a = \chi + r$  with  $r \in \mathcal{S}(\mathbf{R}^{2d})$ .  $\square$

Finally we are in a position to state and prove the main result on microellipticity in the anisotropic Shubin calculus. The proof is short due to the long preparation. Note that we require that the symbol is anisotropic, as opposed to Proposition 5.1 where the symbol is allowed to be isotropic.

**Theorem 6.4.** *Let  $s > 0$ . If  $u \in \mathcal{S}'(\mathbf{R}^d)$  and  $a \in G^{m, s}$  then for any  $m_1 \leq m$*

$$\text{WF}_g^s(u) \subseteq \text{WF}_g^s(a^w(x, D)u) \bigcup \text{char}_{s, m_1}(a).$$

*Proof.* We may assume that  $\text{WF}_g^s(a^w(x, D)u) \neq T^*\mathbf{R}^d \setminus 0$  and  $\text{char}_{s, m_1}(a) \neq T^*\mathbf{R}^d \setminus 0$ , since the inclusion is trivial otherwise. Let  $0 \neq z_0 \notin \text{WF}_g^s(a^w(x, D)u)$  and  $z_0 \notin \text{char}_{s, m_1}(a)$ . Due to  $s$ -conic invariance we may assume  $|z_0| = 1$ .

Pick  $\varepsilon > 0$  such that  $\bar{\Gamma}_{z_0, 2\varepsilon} \cap \text{char}_{s, m_1}(a) = \emptyset$ , and pick  $\chi \in G^{0, s}$  such that  $\text{supp } \chi \subseteq \Gamma_{z_0, 2\varepsilon} \setminus B_R$  and  $\chi|_{\Gamma_{z_0, \varepsilon} \setminus \bar{B}_{2R}} \equiv 1$ , for  $R > 0$  to be chosen. This is possible due to Lemma 3.5. Then  $z_0 \notin \text{conesupp}_s(1 - \chi)$ , and due to (3.6) we have  $\chi \in G_0^0$ . By Proposition 6.2 we may thus conclude

$$z_0 \notin \text{WF}_g^s((1 - \chi)^w(x, D)u).$$

According to Lemma 6.3 we may pick  $R > 0$  such that there exists  $b \in G^{-m_1, s}$  and  $r \in \mathcal{S}(\mathbf{R}^{2d})$  such that  $1 = b \# a + r + 1 - \chi$ , so we have

$$u = b^w(c, D)a^w(x, D)u + r^w(x, D)u + (1 - \chi)^w(x, D)u.$$

Here  $r^w(x, D)u \in \mathcal{S}(\mathbf{R}^d)$  which means that  $z_0 \notin \text{WF}_g^s(r^w(x, D)u)$  trivially. By Proposition 5.1 we have  $z_0 \notin \text{WF}_g^s(b^w(x, D)a^w(x, D)u)$ . Thus we may conclude that  $z_0 \notin \text{WF}_g^s(u)$ .  $\square$

**Corollary 6.5.** *Let  $s > 0$ . If  $u \in \mathcal{S}'(\mathbf{R}^d)$ ,  $a \in G^{m, s}$  and  $\text{char}_{s, m_1}(a) = \emptyset$  for some  $m_1 \leq m$  then*

$$\text{WF}_g^s(a^w(x, D)u) = \text{WF}_g^s(u).$$

## 7. THE $s$ -GABOR WAVE FRONT SET OF OSCILLATORY FUNCTIONS

An important reason for the introduction of the anisotropic Gabor wave front set  $\text{WF}_g^s(u)$  is that it describes accurately the phase space singularities of oscillatory functions known generically as chirp signals.

Let  $\varphi : \mathbf{R}^d \rightarrow \mathbf{R}$  be a real polynomial of order  $m \geq 2$

$$(7.1) \quad \varphi(x) = \varphi_m(x) + p(x)$$

where

$$(7.2) \quad p(x) = \sum_{0 \leq |\alpha| < m} c_\alpha x^\alpha, \quad c_\alpha \in \mathbf{R},$$

and

$$(7.3) \quad \varphi_m(x) = \sum_{|\alpha|=m} c_\alpha x^\alpha, \quad c_\alpha \in \mathbf{R}, \quad \exists \alpha \in \mathbf{N}^d : |\alpha| = m, \quad c_\alpha \in \mathbf{R} \setminus 0,$$

is the principal part.

We will study chirp functions of the form

$$(7.4) \quad u(x) = e^{i\varphi(x)}, \quad x \in \mathbf{R}^d.$$

First we note that for any  $\lambda > 0$  and any  $1 \leq j \leq d$  we have

$$(7.5) \quad \lambda^{-m} \partial_j (\varphi(\lambda y)) = \partial_j \varphi_m(y) + \lambda^{1-m} \partial_j p(\lambda y)$$

and if  $|y| \leq R$  and  $\lambda \geq 1$  then

$$(7.6) \quad \lambda^{1-m} |\partial_j p(\lambda y)| = \left| \sum_{0 \leq |\alpha| \leq m-1} \alpha_j c_\alpha y^{\alpha - e_j} \lambda^{|\alpha| - m} \right| \leq C_R \lambda^{-1}.$$

The following result shows that only the principal part  $\varphi_m(x)$  of  $\varphi$  is recorded in  $\text{WF}_g^{m-1}(u)$ , and the  $(m-1)$ -Gabor wave front set is contained in the  $(m-1)$ -conic set in phase space which is the graph of its gradient, that is  $0 \neq x \mapsto (x, \nabla \varphi_m(x))$ . The gradient of the phase function is known as the instantaneous frequency [1].

**Theorem 7.1.** *If  $m \geq 2$  and  $\varphi$  is a real polynomial defined by (7.1), (7.2) and (7.3), and  $u$  is defined by (7.4), then*

$$(7.7) \quad \text{WF}_g^{m-1}(u) \subseteq \{(x, \nabla \varphi_m(x)) \in \mathbf{R}^{2d} : x \neq 0\}.$$

*If  $d = 1$  and  $\varphi$  is even or odd then*

$$(7.8) \quad \text{WF}_g^{m-1}(u) = \{(x, \varphi'_m(x)) \in \mathbf{R}^2 : x \neq 0\}.$$

*Proof.* Set

$$W = \{(x, \nabla \varphi_m(x)) \in \mathbf{R}^{2d} : x \in \mathbf{R}^d \setminus 0\} \subseteq T^*\mathbf{R}^d \setminus 0.$$

Then  $W$  is an  $(m-1)$ -conic set in  $T^*\mathbf{R}^d \setminus 0$ .

Suppose  $(x_0, \xi_0) \in \mathbf{R}^{2d} \setminus 0$  and  $(x_0, \xi_0) \notin W$ . Then there exists  $1 \leq j \leq d$  such that  $\xi_{0,j} \neq \partial_j \varphi_m(x_0)$ . Thus there exists an open set  $U$  such that  $(x_0, \xi_0) \in U$ , and  $0 < \varepsilon \leq 1$ ,  $\delta > 0$ , such that

$$(x, \xi) \in U, \quad |x - y| \leq \delta\sqrt{2} \implies |\xi_j - \partial_j \varphi_m(x)| \geq 2\varepsilon, \quad |\partial_j(\varphi_m(x) - \varphi_m(y))| \leq \frac{\varepsilon}{2}.$$

By (7.6) we have

$$\lambda^{1-m} |\partial_j p(\lambda y)| \leq \frac{\varepsilon}{2}$$

if  $(x, \xi) \in U$ ,  $|x - y| \leq \delta\sqrt{2}$  and  $\lambda \geq L$  where  $L \geq 1$  is sufficiently large.

Using (7.5) we obtain if  $(x, \xi) \in U$ ,  $|x - y| \leq \delta\sqrt{2}$  and  $\lambda \geq L$

$$(7.9) \quad |\xi_j - \lambda^{-m} \partial_j(\varphi(\lambda y))| \geq |\xi_j - \partial_j \varphi_m(x)| - (|\partial_j(\varphi_m(y) - \varphi_m(x))| + \lambda^{1-m} |\partial_j p(\lambda y)|) \geq \varepsilon.$$

Let  $\psi \in C_c^\infty(\mathbf{R}^d) \setminus 0$  have  $\text{supp } \psi \subseteq B_\delta$ . We denote by  $y' \in \mathbf{R}^{d-1}$  the vector  $y \in \mathbf{R}^d$  except coordinate  $j$ . The stationary phase theorem [13, Theorem 7.7.1] gives, for any  $k \in \mathbf{N}$ , and any  $\lambda \geq L$ , if  $(x, \xi) \in U$ , using (7.9),

$$\begin{aligned} |V_\psi u(\lambda x, \lambda^{m-1} \xi)| &= (2\pi)^{-\frac{d}{2}} \left| \int_{\mathbf{R}^d} e^{i(\varphi(y) - \lambda^{m-1} \langle y, \xi \rangle)} \overline{\psi(\lambda(\lambda^{-1}y - x))} dy \right| \\ &= (2\pi)^{-\frac{d}{2}} \lambda^d \left| \int_{|x-y| \leq \delta} e^{i\lambda^m(\lambda^{-m}\varphi(\lambda y) - \langle y, \xi \rangle)} \overline{\psi(\lambda(y - x))} dy \right| \\ &\leq C\lambda^d \int_{|x'-y'| \leq \delta} \sum_{n=0}^k \lambda^n \sup_{|x_j - y_j| \leq \delta} |(\partial_j^n \psi)(\lambda(y - x))| |\xi_j - \lambda^{-m} \partial_j(\varphi(\lambda y))|^{n-2k} \\ &\quad \times \lambda^{m(n-2k)} dy' \\ &\leq C_k \varepsilon^{-2k} \sum_{n=0}^k \lambda^{d+n+m(n-2k)} \\ &\leq C_{k,\varepsilon} \lambda^{d-k(m-1)}. \end{aligned}$$

This shows that  $(x_0, \xi_0) \notin \text{WF}_g^{m-1}(u)$  and the inclusion (7.7) follows.

Next let  $d = 1$ . If  $\varphi$  is even then  $u$  is even, and  $W = -W$  since  $m$  is even, so by (4.7) we have either  $\text{WF}_g^{m-1}(u) = \emptyset$  or  $\text{WF}_g^{m-1}(u) = W$ . The former is not true since  $u \notin \mathcal{S}(\mathbf{R})$ . Thus we have proved (7.8) when  $\varphi$  is even.

If  $\varphi$  is odd then  $m$  is odd and  $\tilde{u}(x) = \overline{u(x)} = e^{-i\varphi(x)}$ . Again  $\text{WF}_g^{m-1}(u) = \emptyset$  cannot hold since  $u \notin \mathcal{S}(\mathbf{R})$ . If we assume that the inclusion (7.7) is strict we get a contradiction from (4.6) and (4.8). Indeed suppose e.g.

$$\text{WF}_g^{m-1}(u) = \{(x, \varphi'_m(x)) \in \mathbf{R}^2 : x > 0\}.$$

By (4.6) and (4.8) we then get the contradiction

$$\begin{aligned} \text{WF}_g^{m-1}(\tilde{u}) &= \{(x, -\varphi'_m(x)) \in \mathbf{R}^2 : x < 0\} \\ &= \{(x, -\varphi'_m(x)) \in \mathbf{R}^2 : x > 0\} = \text{WF}_g^{m-1}(\bar{u}). \end{aligned}$$

This proves (7.8) when  $\varphi$  is odd.  $\square$

We would also like to determine  $\text{WF}_g^s(u)$  when  $s \neq m-1$ . The following two results treat this question.

**Proposition 7.2.** *If  $m \geq 2$ ,  $s > m-1$ , and  $\varphi$  is a real polynomial defined by (7.1), (7.2) and (7.3), and  $u$  is defined by (7.4), then*

$$(7.10) \quad \text{WF}_g^s(u) \subseteq (\mathbf{R}^d \setminus 0) \times \{0\}.$$

*If  $d = 1$  and  $\varphi$  is even or odd then*

$$(7.11) \quad \text{WF}_g^s(u) = (\mathbf{R} \setminus 0) \times \{0\}.$$

*Proof.* Suppose  $(x_0, \xi_0) \in T^*\mathbf{R}^d$  and  $\xi_0 \neq 0$ , that is  $\xi_{0,j} \neq 0$  for some  $1 \leq j \leq d$ . From (7.5) we obtain

$$\lambda^{-1-s} \partial_j (\varphi(\lambda y)) = \lambda^{m-1-s} (\partial_j \varphi_m(y) + \lambda^{1-m} \partial_j p(\lambda y)).$$

Thus from  $s > m-1$ , using (7.6), it follows that there exists  $U \subseteq \mathbf{R}^{2d}$  such that  $(x_0, \xi_0) \in U$ , and  $0 < \varepsilon \leq 1$ ,  $L \geq 1$  such that

$$|\xi_j - \lambda^{-1-s} \partial_j (\varphi(\lambda y))| \geq \varepsilon$$

when  $(x, \xi) \in U$ ,  $|x - y| \leq \sqrt{2}$  and  $\lambda \geq L$ .

Let  $\psi \in C_c^\infty(\mathbf{R}) \setminus 0$  be such that  $\text{supp } \psi \subseteq B_1$ . The stationary phase theorem [13, Theorem 7.7.1] yields, for any  $k \in \mathbf{N}$ , and any  $\lambda \geq L$ , if  $(x, \xi) \in U$ ,

$$\begin{aligned} |V_\psi u(\lambda x, \lambda^s \xi)| &= (2\pi)^{-\frac{d}{2}} \left| \int_{\mathbf{R}^d} e^{i(\varphi(y) - \lambda^s \langle y, \xi \rangle)} \overline{\psi(\lambda(\lambda^{-1}y - x))} dy \right| \\ &= (2\pi)^{-\frac{d}{2}} \lambda^d \left| \int_{\mathbf{R}^d} e^{i\lambda^{1+s}(\lambda^{-1-s}\varphi(\lambda y) - \langle y, \xi \rangle)} \overline{\psi(\lambda(y - x))} dy \right| \\ &\leq C \lambda^d \int_{|x' - y'| \leq 1} \sum_{n=0}^k \lambda^n \sup_{|x_j - y_j| \leq 1} |(\partial_j^n \psi)(\lambda(y - x))| |\xi_j - \lambda^{-1-s} \partial_j (\varphi(\lambda y))|^{n-2k} \\ &\quad \times \lambda^{(1+s)(n-2k)} dy' \\ &\leq C_k \lambda^{d-ks} \varepsilon^{-2k}. \end{aligned}$$

This shows that  $(x_0, \xi_0) \notin \text{WF}_g^s(u)$  and (7.10) follows.

When  $d = 1$  and  $\varphi$  is either even or odd then (7.11) follows as in the proof of Theorem 7.1.  $\square$

**Proposition 7.3.** *Let  $m \geq 2$ ,  $0 < s < m - 1$ , and  $\varphi$  be a real polynomial defined by (7.1), (7.2) and (7.3). Suppose  $\varphi_m(x) \neq 0$  for all  $x \in \mathbf{R}^d \setminus 0$ . If  $u$  is defined by (7.4) then*

$$(7.12) \quad \text{WF}_g^s(u) \subseteq \{0\} \times (\mathbf{R}^d \setminus 0).$$

If  $d = 1$  and  $\varphi$  is even then

$$(7.13) \quad \text{WF}_g^s(u) = \{0\} \times (\mathbf{R} \setminus 0).$$

*Proof.* Suppose  $(x_0, \xi_0) \in T^*\mathbf{R}^d$  and  $x_0 \neq 0$ . The assumption  $\varphi_m(x) \neq 0$  for all  $x \in \mathbf{R}^d \setminus 0$  and Euler's homogeneous function theorem imply that  $\nabla \varphi_m(x_0) \neq 0$ , that is  $\partial_j \varphi_m(x_0) \neq 0$  for some  $1 \leq j \leq d$ . From (7.5) and (7.6) and  $s < m - 1$  it follows that there exists  $U \subseteq \mathbf{R}^{2d}$  such that  $(x_0, \xi_0) \in U$ ,  $1 \leq j \leq d$  and  $0 < \varepsilon \leq 1$ ,  $L \geq 1$  such that

$$|\lambda^{1+s-m} \xi_j - \lambda^{-m} \partial_j (\varphi(\lambda y))| \geq \varepsilon$$

when  $(x, \xi) \in U$ ,  $|x - y| \leq \varepsilon \sqrt{2}$  and  $\lambda \geq L$ .

Let  $\psi \in C_c^\infty(\mathbf{R}) \setminus 0$  be such that  $\text{supp } \psi \subseteq B_\varepsilon$ . Again by the stationary phase theorem [13, Theorem 7.7.1] we obtain, for any  $k \in \mathbf{N}$ , and any  $\lambda \geq L$ , if  $(x, \xi) \in U$ ,

$$\begin{aligned} |V_\psi u(\lambda x, \lambda^s \xi)| &= (2\pi)^{-\frac{d}{2}} \left| \int_{\mathbf{R}^d} e^{i(\varphi(y) - \lambda^s \langle y, \xi \rangle)} \overline{\psi(\lambda(\lambda^{-1}y - x))} dy \right| \\ &= (2\pi)^{-\frac{d}{2}} \lambda^d \left| \int_{\mathbf{R}^d} e^{i\lambda^m (\lambda^{-m} \varphi(\lambda y) - \lambda^{1+s-m} \langle y, \xi \rangle)} \overline{\psi(\lambda(y - x))} dy \right| \\ &\leq C \lambda^d \int_{|x' - y'| \leq \varepsilon} \sum_{n=0}^k \lambda^n \sup_{|x_j - y_j| \leq \varepsilon} |(\partial_j^n \psi)(\lambda(y - x))| |\lambda^{1+s-m} \xi_j - \lambda^{-m} \partial_j (\varphi(\lambda y))|^{n-2k} \\ &\quad \times \lambda^{m(n-2k)} dy' \\ &\leq C_k \lambda^{d-k(m-1)} \varepsilon^{-2k}. \end{aligned}$$

This shows that  $(x_0, \xi_0) \notin \text{WF}_g^s(u)$  and (7.10) follows.

When  $d = 1$  and  $\varphi$  is even then (7.13) follows as in the proof of Theorem 7.1.  $\square$

**Example 7.4.** Let  $k \in \mathbf{N} \setminus 0$  and consider the differential equation

$$u^{(k)} - xu = 0.$$

for  $u \in \mathcal{S}'(\mathbf{R})$ . When  $k = 2$  this is the Airy equation. Fourier transformation gives

$$(7.14) \quad i^k \xi^k \widehat{u} + D \widehat{u} = 0$$

which is solved by

$$\widehat{u}(\xi) = C \exp \left( -i^{k+1} \frac{\xi^{k+1}}{k+1} \right), \quad C \in \mathbf{C}.$$

This function belongs to  $\mathcal{S}'(\mathbf{R})$  provided  $k \notin 1 + 4\mathbf{N}$ .

The equation (7.14) can be written  $a^w(x, D) \widehat{u} = 0$  where

$$a(x, \xi) = i^k x^k + \xi.$$

By Example 3.9 we know that  $a \in G^k \cap G^{k,k}$ . Suppose  $k = 2n$  with  $n \in \mathbf{N} \setminus 0$ . Since  $a(x, \xi) = 0$  when  $\xi = (-1)^{n+1}x^{2n}$ , it follows from Definition 3.8 that  $(x, (-1)^{n+1}x^{2n}) \in \text{char}_{2n}(a)$  for any  $x \neq 0$ . It holds

$$(7.15) \quad \text{char}_{2n}(a) = \{(x, (-1)^{n+1}x^{2n}) \in \mathbf{R}^2, x \neq 0\}.$$

In fact it suffices to show

$$(7.16) \quad \text{char}_{2n}(a) \subseteq \{(x, (-1)^{n+1}x^{2n}) \in \mathbf{R}^2, x \neq 0\}.$$

Suppose  $(x_0, \xi_0) \in \mathbf{S}^1$  with  $\xi_0 \neq (-1)^{n+1}x_0^{2n}$ . In order to show (7.16) we must show  $(x_0, \xi_0) \notin \text{char}_{2n}(a)$ . There exist  $\varepsilon, \delta > 0$  such that

$$|\xi - (-1)^{n+1}x^{2n}| \geq \delta(|x| + |\xi|^{\frac{1}{2n}})^{2n}$$

if  $(x, \xi) \in (x_0, \xi_0) + B_\varepsilon$ . This inequality is  $2n$ -conic, that is invariant to the transformation  $T^*\mathbf{R} \setminus 0 \ni (x, \xi) \mapsto (\lambda x, \lambda^{2n}\xi)$  for  $\lambda > 0$ . It follows that

$$|a(x, \xi)| \geq \delta(|x| + |\xi|^{\frac{1}{2n}})^{2n}$$

when  $(x, \xi) \in \tilde{\Gamma}_{2n, (x_0, \xi_0), \varepsilon}$ . Hence from Lemma 3.7 it follows  $(x_0, \xi_0) \notin \text{char}_{2n}(a)$  and we have shown (7.16) and thereby (7.15).

Invoking Theorem 6.4 we obtain

$$\begin{aligned} \text{WF}_g^{2n}(\hat{u}) &\subseteq \text{WF}_g^{2n}(a^w(x, D)\hat{u}) \bigcup \text{char}_{2n}(a) \\ &= \text{char}_{2n}(a) = \{(x, (-1)^{n+1}x^{2n}) \in \mathbf{R}^2, x \neq 0\}. \end{aligned}$$

Thus we have found an alternative proof of a particular case of the inclusion (7.7) in Theorem 7.1 when  $m$  is odd. From (7.8) we know that the inclusion is actually an equality.

Adding this information and applying Proposition 4.3 (i) we obtain

$$\text{WF}_g^{\frac{1}{2n}}(u) = -\mathcal{J}\text{WF}_g^{2n}(\hat{u}) = \{((-1)^n x^{2n}, x) \in \mathbf{R}^2, x \neq 0\}.$$

If  $n = 1$  then  $u$  is the Airy function (multiplied by  $C$ ) [13], and thus

$$\text{WF}_g^{\frac{1}{2}}(u) = \{(-x^2, x) \in \mathbf{R}^2, x \neq 0\}.$$

This can be compared to [21, Example 8.5] which says that

$$\text{WF}_g(u) = \text{WF}_g^1(u) = \{(x, 0) \in \mathbf{R}^2, x < 0\}.$$

#### ACKNOWLEDGMENT

Work partially supported by the MIUR project “Dipartimenti di Eccellenza 2018-2022” (CUP E11G18000350001).

## REFERENCES

- [1] P. Boggiatto, A. Oliaro and P. Wahlberg, *The wave front set of the Wigner distribution and instantaneous frequency*, J. Fourier Anal. Appl. **18** (2012), 410–438.
- [2] C. Boiti, D. Jornet and A. Oliaro, *The Gabor wave front set in spaces of ultradifferentiable functions*, Monatsh. Math. **188** (2019), 199–246.
- [3] M. Cappiello, T. Gramchev and L. Rodino, *Entire extensions and exponential decay for semilinear elliptic equations*, J. Anal. Math. **111** (2010), 339–367.
- [4] M. Cappiello and R. Schulz, *Microlocal analysis of quasianalytic Gelfand–Shilov type ultradistributions*, Compl. Var. Elliptic Equ. **61** (4) (2016), 538–561.
- [5] M. Cappiello, R. Schulz and P. Wahlberg, *Conormal distributions in the Shubin calculus of pseudodifferential operators*, J. Math. Phys. **59** 021502 (2018).
- [6] M. Cappiello, T. Gramchev, S. Pilipović and L. Rodino, *Anisotropic Shubin operators and eigenfunction expansions in Gelfand–Shilov spaces*, J. Anal. Math. **138** (2) (2019), 857–870.
- [7] E. Carylps and P. Wahlberg, *Propagation of exponential phase space singularities for Schrödinger equations with quadratic Hamiltonians*, J. Fourier Anal. Appl. **23** (3) (2017), 530–571. Correction: **27**:35 (2021).
- [8] E. Cordero and L. Rodino, *Time-Frequency Analysis of Operators*, De Gruyter Studies in Mathematics **75**, De Gruyter, Berlin, 2020.
- [9] G. B. Folland, *Harmonic Analysis in Phase Space*, Princeton University Press, 1989.
- [10] G. Garello and A. Morando, *m-Microlocal elliptic pseudodifferential operators acting on  $L^p_{\text{loc}}(\Omega)$* , Math. Nachr. **289** (2016), 1820–1837.
- [11] K. Gröchenig, *Foundations of Time-Frequency Analysis*, Birkhäuser, Boston, 2001.
- [12] ———, *Time-frequency analysis of Sjöstrand’s class*, Rev. Mat. Iberoamer. **22** (2), 703–724, 2006.
- [13] L. Hörmander, *The Analysis of Linear Partial Differential Operators*, Vol. I, III, IV, Springer, Berlin, 1990.
- [14] L. Hörmander, *Quadratic hyperbolic operators*, Microlocal Analysis and Applications, Lecture Notes in Math. **1495**, Eds. L. Cattabriga, L. Rodino, pp. 118–160, Springer, 1991.
- [15] L. Hörmander, *Symplectic classification of quadratic forms, and general Mehler formulas*, Math. Z. **219** (1995), 413–449.
- [16] R. Lascar, *Propagation des singularités d’équations pseudodifférentielles quasi homogènes*, Ann. Inst. Fourier Grenoble **27** (1977), 79–123.
- [17] S. G. Krantz and H. R. Parks, *The Implicit Function Theorem. History, Theory, and Applications*, Birkhäuser, Boston, 2003.
- [18] J. Martin, *Spectral inequalities for anisotropic Shubin operators*, arXiv:2205.11868 [math.AP] (2022).
- [19] F. Nicola and L. Rodino, *Global Pseudo-Differential Calculus on Euclidean Spaces*, Birkhäuser, Basel, 2010.
- [20] C. Parenti and L. Rodino, *Parametrices for a class of pseudo differential operators I,II*, Ann. Mat. Pura Appl. **125** (4) (1980), 221–254 and 255–278.
- [21] K. Pravda-Starov, L. Rodino and P. Wahlberg, *Propagation of Gabor singularities for Schrödinger equations with quadratic Hamiltonians*, Math. Nachr. **291** (1) (2018), 128–159.
- [22] L. Rodino and P. Wahlberg, *The Gabor wave front set*, Monats. Math. **173** (4) (2014), 625–655.
- [23] L. Rodino and S. I. Trapasso, *An introduction to the Gabor wave front set*, Anomalies in partial differential equations, Springer INdAM Ser. **43**, Springer, Cham, 369–393, 2021.
- [24] L. Rodino and P. Wahlberg, *Microlocal analysis of Gelfand–Shilov spaces*, arXiv:2202.05543 [math.AP] (2022).
- [25] R. Schulz and P. Wahlberg, *Equality of the homogeneous and the Gabor wave front set*, Comm. PDE **42** (5) (2017), 703–730.
- [26] M. A. Shubin, *Pseudodifferential Operators and Spectral Theory*, Springer, 2001.
- [27] P. Wahlberg, *The Gabor wave front set of compactly supported distributions*, Advances in Microlocal and Time-Frequency Analysis, P. Boggiatto, M. Cappiello, E. Cordero, S. Coriasco, G. Garello, A. Oliaro, J. Seiler (Eds.) Birkhäuser Verlag, pp. 507–520, 2020.

- [28] H. Zhu, *Propagation of singularities for gravity-capillary water waves*, arXiv:1810.09339 [math.AP] (2020).

DEPARTMENT OF MATHEMATICS, UNIVERSITÀ DI TORINO, VIA CARLO ALBERTO 10, 10123 TORINO, ITALY

*Email address:* `luigi.rodino[AT]unito.it`

DIPARTIMENTO DI SCIENZE MATEMATICHE, POLITECNICO DI TORINO, CORSO DUCA DEGLI ABRUZZI 24, 10129 TORINO, ITALY

*Email address:* `patrik.wahlberg[AT]polito.it`