

EGU22-9190, updated on 31 Mar 2023

<https://doi.org/10.5194/egusphere-egu22-9190>

EGU General Assembly 2022

© Author(s) 2023. This work is distributed under the Creative Commons Attribution 4.0 License.



Seasonal compartmentalisation of water in a grassland at 2600 m a.s.l.

Alessio Gentile¹, Ivan Bevilacqua¹, Davide Canone¹, Natalie Ceperley^{2,3,4}, Davide Gisolo¹, Mesmer N'Sassila¹, Maurizio Previati¹, Giulia Zuecco⁵, Bettina Schaeffli^{2,3,4}, and Stefano Ferraris¹

¹Interuniversity Department of Regional and Urban Studies and Planning, Politecnico and Università degli Studi di Torino, Torino, Italy (alessio.gentile@polito.it)

²Institute of Earth Surface Dynamics, Faculty of Geosciences and Environment, University of Lausanne, Lausanne, Switzerland

³Institute of Geography, Faculty of Science, University of Bern, Bern, Switzerland

⁴Oeschger Centre for Climate Change Research, University of Bern, Bern, Switzerland

⁵Department of Land, Environment, Agriculture and Forestry, University of Padova, Legnaro, Italy

High Alpine catchments are precious water-resources since they act as natural storage reservoirs, storing water in the snow cover and in the subsurface and thereby providing water during the dry seasons. Thus, a deeper knowledge of the hydrological functioning of these systems is necessary, in particular to make climate change projections. The role of seasonality is crucial in these catchments that generally exhibit a snow-dominated hydro-climatic regime.

Here we use high-frequency observations of stable isotopes of water to identify the seasonal origin of streamwater in a high-elevation Alpine catchment located in the Valle d'Aosta Region, Italy. We quantify the relative contribution of winter and summer precipitation reaching the stream through the Seasonal Origin Index (SOI_Q), calculated using the $\delta^{18}O$ values and the volumes of precipitation and streamflow. Highly negative SOI_Q values are obtained suggesting that streamwater is mainly composed of winter precipitation. Conversely, the Seasonal Origin Index for evapotranspiration (SOI_{ET}), which can be directly inferred from SOI_Q , returns a positive value reflecting that plants preferentially take up water deriving from summer precipitation.

These findings allow us to develop a conceptual model of this Alpine system. This conceptual model suggests:

- a deep infiltration component, mainly composed by snowmelt water, reaching the stream through a preferential flow.
- a shallow infiltration component, predominantly represented by summer rainfall, that dominates the shallow soils and that is used by plants.

Therefore, we presume a seasonal compartmentalisation of water in this high-elevation catchment.

Nevertheless, a previous study in Switzerland revealed $SOI_Q \approx 0$ for the Allenbach and Dischmabach snow-dominated catchments, indicating that similar fractions of summer and winter precipitation become streamflow. This different result achieved in systems with an apparently similar functioning highlights the need for a deep insight into the flow paths governing high-elevation catchments and it opens the way for new challenges to understand the hydrological processes hidden behind this difference.