

A note on very ample Terracini loci

Original

A note on very ample Terracini loci / Ballico, E.; Ventura, E.. - In: RENDICONTI DEL CIRCOLO MATEMATICO DI PALERMO. - ISSN 0009-725X. - (2023). [10.1007/s12215-023-00886-4]

Availability:

This version is available at: 11583/2978104 since: 2023-04-22T06:27:50Z

Publisher:

Springer

Published

DOI:10.1007/s12215-023-00886-4

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



A note on very ample Terracini loci

Edoardo Ballico¹ · Emanuele Ventura²

Received: 27 December 2022 / Accepted: 22 March 2023
© The Author(s) 2023

Abstract

In this short note we show that, for any ample embedding of a variety of dimension at least two in a projective space, all high enough degree Veronese re-embeddings have non-empty Terracini loci.

Keywords Secant variety · Terracini locus

Mathematics Subject Classification 14N05

1 Introduction

Terracini loci were introduced by the first author and Chiantini in [2]. Their emptiness implies non-defectivity of secant varieties due to the celebrated Terracini's lemma, whereas the converse is not true: there exist non-empty Terracini loci even in the presence of non-defective secants. This triggered the interest for this geometric notion, leading to the results in the aforementioned article. The Terracini locus has been the subject of recent investigations [3, 4], especially for Segre and Veronese varieties, that are crucial in the context of tensors. We start off by defining set-theoretically these loci.

Definition Let $X \subset \mathbb{P}^N$ be a non-degenerate projective variety of dimension $n \geq 1$ over an algebraically closed field \mathbb{K} . Let $S \subset X_{\text{reg}}$ be a finite subset of smooth points of X whose cardinality is k . Let $(2S, X)$ be the union of the corresponding 2-fat points $(2p, X)$ supported at the points $p \in S$. Then S is in the k th *Terracini locus* $\mathbb{T}_k(X)$ if and only if $h^0(\mathcal{I}_{(2S, X)}(1)) > 0$ and $h^1(\mathcal{I}_{(2S, X)}(1)) > 0$. Equivalently, S is in $\mathbb{T}_k(X)$ whenever the n -dimensional tangent spaces $T_p X$, for $p \in S$, are linearly dependent and their projective linear span is not the ambient space \mathbb{P}^N .

✉ Emanuele Ventura
emanuele.ventura@polito.it
Edoardo Ballico
edoardo.ballico@unitn.it

¹ Università di Trento, Via Sommarive 14, 38123 Povo, Trento, Italy

² Politecnico di Torino, Dipartimento di Scienze Matematiche "G. L. Lagrange", Corso Duca degli Abruzzi 24, 10129 Torino, Italy

A consequence of a deep result of Alexander and Hirschowitz [1, Theorem 1.1 and Corollary 1.2] (where in their notation one chooses $m = 2$) states that for any projective variety X there exists a very ample embedding such that all the secant varieties of X under this embedding are non-defective. The aim of this note is to point out that, even in this very ample regime, the emptiness of the corresponding Terracini locus *does not* generally hold. Thus we answer in the negative the question whether a statement similar to the one by Alexander and Hirschowitz works for Terracini loci.

2 Very ample regime

Let \mathbb{K} be an algebraically closed field and let X be a projective variety of dimension n over \mathbb{K} . We say that an embedding $X \subset \mathbb{P}^r$ of X is not secant defective if for each positive integer k the k -secant variety of X has dimension $\min\{r, k(n+1) - 1\}$. For a very ample line bundle L on X , let $\nu_L : X \rightarrow |L|^\vee$ denote the associated embedding. The k th secant variety and the k th Terracini locus of $\nu_L(X)$ are denoted $\sigma_k(\nu_L(X))$ and $\mathbb{T}_k(\nu_L(X))$, respectively. We say that $\nu_L(X)$ is *secant non-defective* if $\sigma_k(\nu_L(X))$ is non-defective for every $k \geq 1$.

Theorem 1 *Let $n \geq 2$ and X be as above. Let $F, L \in \text{Pic}(X)$, where L is an ample line bundle. Then there exists an integer m_0 (depending only on X, F, L) such that for all $m \geq m_0$ the line bundle $F + mL$ is very ample, $\nu_{F+mL}(X)$ is secant non-defective, and there exists $k > 0$ such that $\sigma_k(\nu_{F+mL}(X)) \neq |F + mL|^\vee$ and $\mathbb{T}_k(\nu_{F+mL}(X)) \neq \emptyset$.*

Proof Let $L = \mathcal{L}(D)$ and define $\alpha = D \cdots D > 0$, the n times self-intersection of the Cartier divisor D . Fix an integral curve $Y \subset X$ such that $Y \cap X_{\text{reg}} \neq \emptyset$, where Y is possibly singular. Let $\beta = Y \cdot D \cdots D$, the intersection of Y with $n - 1$ copies of D , i.e. $\beta = \deg(L|_Y)$ and $\beta > 0$ because L is ample. Fix a real number ε such that $\alpha > \varepsilon > 0$. By the result of Alexander and Hirschowitz [1, Theorem 1.1], by the asymptotic Riemann-Roch and by the ampleness of L , we find an integer m_1 such that for all $m \geq m_1$ we have that: $F + mL$ is very ample, $\nu_{F+mL}(X)$ is secant non-defective, and $h^0(F + mL) \geq \frac{\alpha - \varepsilon}{n!} m^n$.

Thus, for $1 \leq k < \left\lfloor \frac{\alpha - \varepsilon}{(n+1)!} m^n \right\rfloor$, we have $\sigma_k(\nu_{F+mL}(X)) \subsetneq |F + mL|^\vee$. By the asymptotic Riemann-Roch, $h^0(Y, (F + mL)|_Y)$ grows like a linear function of the form βm . Therefore there exists $m_0 \geq m_1$ such that for all $m \geq m_0$ one has $1 \leq h^0(Y, (F + mL)|_Y)/2 < \left\lfloor \frac{\alpha - \varepsilon}{(n+1)!} m^n \right\rfloor$.

Define $k - 1 = \lceil h^0(Y, (F + mL)|_Y)/2 \rceil$. Note that the projective linear span of the curve Y has dimension $\dim(Y) \leq 2k - 3$. Fix a set $S \subset Y \cap X_{\text{reg}}$ with cardinality k . The zero-dimensional scheme $(2S, X) \cap Y \subset Y$ has degree at least $2k$. Hence, if $(2S, X) \cap Y \subset Y$ was linearly independent, then its projective linear span would be at least $(2k - 1)$ -dimensional. Therefore $(2S, X) \cap Y$ is linearly dependent, i.e. $h^1(\mathcal{I}_{(2S, X) \cap Y}(1)) > 0$. Moreover, since $k < \left\lfloor \frac{\alpha - \varepsilon}{(n+1)!} m^n \right\rfloor$ and $\deg((2S, X)) = k(n + 1)$, the projective linear span of this scheme cannot fill the ambient space, i.e. one has $h^0(\mathcal{I}_{(2S, X)}(1)) > 0$.

Now, let $Z \subset W$ be two zero-dimensional schemes. Then one has the exact sequence of sheaves

$$0 \longrightarrow \mathcal{I}_W(1) \longrightarrow \mathcal{I}_Z(1) \longrightarrow \mathcal{I}_Z(1)/\mathcal{I}_W(1) \longrightarrow 0.$$

Here the cokernel sheaf is either zero or supported on a zero-dimensional scheme. Taking the long exact sequence in cohomology, we then find a surjective map in cohomology

$H^1(\mathcal{I}_W(1)) \rightarrow H^1(\mathcal{I}_Z(1))$. The zero-dimensional scheme $(2S, X) \cap Y$ is a closed subscheme of $(2S, X)$ and so we likewise have a surjection

$$H^1(\mathcal{I}_{(2S, X)}(1)) \rightarrow H^1(\mathcal{I}_{(2S, X) \cap Y}(1)).$$

Therefore $h^1(\mathcal{I}_{(2S, X)}(1)) > 0$ too. So any collection of k smooth points of $Y \cap X_{\text{reg}}$ are in the k th Terracini locus of $v_{F+mL}(X)$. □

Remark 2 Let $X \subset \mathbb{P}^N$ be a projective variety with $\dim X = n \geq 2$ and consider $v_d(X)$. For any integer $k > 0$, the set $S^k v_d(X_{\text{reg}})$ of all subsets of $v_d(X_{\text{reg}})$ with cardinality k is a variety of dimension kn . For $d \gg 0$, the families of $S \in \mathbb{T}_k(v_d(X))$ we found in the proof of Theorem 1 on a fixed curve Y have codimension k in $S^k v_d(X_{\text{reg}})$. Varying Y , we do not decrease significantly the codimension of $\mathbb{T}_k(v_d(X))$ in $S^k v_d(X_{\text{reg}})$: the magnitude of this is $O(k)$. We do not have examples for which, when k is increasing with d , $\mathbb{T}_k(v_d(X))$ has codimension 1 in $S^k v_d(X_{\text{reg}})$, which is the least codimension allowed in view of the secant non-defectivity result in [1].

Proposition 3 Let $N \geq 1$ and let $C \subset \mathbb{P}^N$ be a smooth and non-degenerate rational curve of degree d . For all $d' \geq d + 1 - N$, the curve $v_{d'}(C) \subset \langle v_{d'}(C) \rangle$ has empty Terracini loci.

Proof Suppose $N = d = 1$ so that $C = \mathbb{P}^1$. For $d' \geq 1$, consider the rational normal curve $v_{d'}(\mathbb{P}^1)$. Its k th Terracini locus consists of those subsets $S \subset \mathbb{P}^1$ such that $(2S, v_{d'}(\mathbb{P}^1))$ does not span $\langle v_{d'}(C) \rangle$, i.e. $h^0(\mathcal{I}_{2S}(d')) > 0$, and such that $h^1(\mathcal{I}_{2S}(d')) > 0$. Since $C = \mathbb{P}^1$, for any zero-dimensional scheme $Z \subset C$ either $h^0(\mathcal{I}_Z(d')) = 0$ or $h^1(\mathcal{I}_Z(d')) = 0$. Hence any Terracini locus of the rational normal curve $v_{d'}(C)$ is empty.

For the general case, let $d \geq 2$ and $d' \geq d + 1 - N$. One has $h^1(\mathcal{I}_C(d')) = 0$ [5, Theorem p. 492]. Hence $v_{d'}(C)$ is an embedding of \mathbb{P}^1 by the complete linear system $|\mathcal{O}_{\mathbb{P}^1}(d \cdot d')|$. So this has empty Terracini loci by the first part. □

The case of curves with positive arithmetic genus is treated in the following proposition. Here different behaviours appear according to the parity of the degree.

Proposition 4 Let C be an integral projective curve over \mathbb{K} , with $\text{char}(\mathbb{K}) \neq 2$, whose arithmetic genus is $g > 0$. Let F and L be line bundles on C , where L is ample, of degrees $\alpha = \text{deg}(L)$ and $\beta = \text{deg}(F)$. For each integer $m > 0$, consider the complete linear system $|F + mL|$. Assume that $\beta + m\alpha \geq 4g + 2$ and assume that $\beta + m\alpha$ is even. Then $v_{F+mL}(C)$ has a non-empty Terracini locus.

Proof Recall that a line bundle E on C is very ample if $\text{deg}(E) \geq 2g + 1$ [6, Corollary 3.2, Chapter IV]. Since the Picard group $\text{Pic}^0(C)$ is a quasi-projective irreducible group and $\text{char}(\mathbb{K}) \neq 2$, the kernel of the multiplication morphism $\otimes 2 : \text{Pic}^0(C) \rightarrow \text{Pic}^0(C)$ is finite. So this morphism is surjective. Since $\text{deg}(F + mL)$ is even and $\otimes 2$ is surjective, there is a line bundle R_m such that $R_m^{\otimes 2} \cong F + mL$. Thus $\text{deg}(R_m) = (\beta + m\alpha)/2$. Since $\beta + m\alpha \geq 4g + 2$, the line bundle R_m is very ample. Thus $|R_m| \neq \emptyset$ and a general $S \in |R_m|$ consists of k distinct reduced points and $S \subset C_{\text{reg}}$. Note that $2S \in |F + mL|$ and hence $\langle 2v_{F+mL}(S) \rangle \subsetneq |F + mL|^\vee$ is a hyperplane. Since $\text{deg}(F + mL) > 2g - 1$, one has $h^0(F + mL) = \text{deg}(F + mL) + 1 - g = \text{deg}(2S) + 1 - g$. Since $g > 0$, $2S$ does not give $\text{deg}(2S)$ independent conditions to $|F + mL|$. Then, by definition, $v_{F+mL}(S)$ is in the k th Terracini locus of $v_{F+mL}(C)$. □

Corollary 5 Let $C \subset \mathbb{P}^N$ be an integral and non-degenerate projective curve with arithmetic genus $g = 1$ of degree d over \mathbb{K} , with $\text{char}(\mathbb{K}) \neq 2$. If $d' \geq d + 1 - N$ and $d \cdot d'$ is even, then $\mathbb{T}_{d \cdot d'/2}(v_d(C)) \neq \emptyset$. If $d \cdot d'$ is odd, then all Terracini loci of $v_{d'}(C)$ are empty.

Proof Since $d' \geq d + 1 - N$, we have $h^1(\mathcal{I}_C(d')) = 0$ [5, Theorem p. 492]. Hence $v_{d'}(C)$ is an embedding of C by a complete linear system. By Proposition 4, if $d \cdot d'$ is even, then $\mathbb{T}_{d \cdot d' / 2}(v_{d'}(C)) \neq \emptyset$.

Suppose a line bundle L on C has $\deg(L) = 2m + 1$; let $S \subset C_{\text{reg}}$ have cardinality k . Then $\deg(L(-2S)) = 2(m - k) + 1 \neq 0$. If $\deg(L(-2S)) < 0$, then $h^0(L(-2S)) = 0$. If $\deg(L(-2S)) > 0$, by Serre duality, we find $h^1(L(-2S)) = 0$. Therefore any Terracini locus is empty. \square

Acknowledgements The open-ended problem of looking at very ample Terracini loci was proposed by Luca Chiantini and Ciro Ciliberto (along with many other interesting problems) during the *Geometry of Secants Workshop* held in October 2022, within the AGATES semester at University of Warsaw and IMPAN. We warmly thank Chiantini, Ciliberto, and the organizers of the semester and of the workshop. This work is supported by the Thematic Research Programme “Tensors: geometry, complexity and quantum entanglement”, University of Warsaw, Excellence Initiative—Research University and the Simons Foundation Award No. 663281 granted to the Institute of Mathematics of the Polish Academy of Sciences for the years 2021–2023. We thank an anonymous referee for useful comments and corrections.

Funding Open access funding provided by Politecnico di Torino within the CRUI-CARE Agreement.

Declarations

Conflict of Interest All authors certify that they have no affiliations with or involvement in any organization or entity with any financial interest or non-financial interest in the subject matter or materials discussed in this manuscript.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Alexander, J., Hirschowitz, A.: An asymptotic vanishing theorem for generic unions of multiple points. *Invent. Math.* **140**, 303–325 (2000)
2. Ballico, E., Chiantini, L.: On the Terracini Locus of Projective Varieties. *Milan J. Math.* **89**, 1–17 (2021)
3. Ballico, E., Bernardi, A., Santarsiero, P.: Terracini locus for three points on a Segre variety, preprint at [arXiv:2012.00574](https://arxiv.org/abs/2012.00574), (2020)
4. Chiantini, L., Gesmundo, F.: Decompositions and Terracini loci of cubic forms of low rank, preprint at [arXiv:2302.03715](https://arxiv.org/abs/2302.03715), (2023)
5. Gruson, L., Lazarsfeld, R., Peskine, C.: On a theorem of Castelnuovo and the equations defining space curves. *Invent. Math.* **72**, 491–506 (1983)
6. Hartshorne, R.: Algebraic geometry. Graduate Texts in Mathematics, No. 52, pp. 16–496. Springer-Verlag, New York-Heidelberg (1977)

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.