

Surface decontamination protocols for surgical treatment of peri-implantitis: A systematic review with meta-analysis

Original

Surface decontamination protocols for surgical treatment of peri-implantitis: A systematic review with meta-analysis / Baima, G.; Citterio, F.; Romandini, M.; Romano, F.; Mariani, G. M.; Buduneli, N.; Aimetti, M.. - In: CLINICAL ORAL IMPLANTS RESEARCH. - ISSN 0905-7161. - 33:11(2022), pp. 1069-1086. [10.1111/clr.13992]

Availability:

This version is available at: 11583/2979502 since: 2023-06-22T14:14:26Z

Publisher:

John Wiley and Sons Inc

Published

DOI:10.1111/clr.13992

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Surface decontamination protocols for surgical treatment of peri-implantitis: A systematic review with meta-analysis

Giacomo Baima¹  | Filippo Citterio¹  | Mario Romandini²  | Federica Romano¹  |
Giulia Maria Mariani¹  | Nurcan Buduneli³  | Mario Aimetti¹ 

¹Department of Surgical Sciences, C.I.R. Dental School, University of Turin, Turin, Italy

²Section of Graduate Periodontology, Faculty of Odontology, University Complutense, Madrid, Spain

³Department of Periodontology, School of Dentistry, Ege University, İzmir, Turkey

Correspondence

Giacomo Baima, Department of Surgical Sciences, C.I.R. Dental School, University of Turin, Turin, Italy.

Email: giacomo.baima@unito.it

Mario Romandini, Giovanni XXIII Square, 24; 74123 Taranto, Italy.

Email: mario.romandini@gmail.com

Funding information

This study was funded solely by the institutions of the authors.

Abstract

Objective: To answer the following PICO question: “In patients requiring surgical treatment of peri-implantitis (P), is any implant surface decontamination protocol (I) superior to others (C) in terms of clinical and radiographic parameters (O)?”

Methods: Randomized clinical trials (RCTs) comparing two or more decontamination protocols as part of the surgical treatment of peri-implantitis were included. Two authors independently searched for eligible studies, screened titles and abstracts, did full-text analysis, extracted data, and performed the risk-of-bias assessment. Whenever possible, results were summarized through random effects meta-analyses.

Results: Twenty-two manuscripts reporting on 16 RCTs were included, testing mechanical, chemical and physical decontamination protocols. All of them resulted in an improvement in clinical parameters; however, the superiority of specific protocols over others is mainly based on single RCTs. The use of titanium brushes and implantoplasty showed favorable results as single decontamination methods. Meta-analyses indicated a lack of added effect of Er:Yag laser on probing pocket depth (PPD) reduction ($n = 2$, WMD = -0.24 mm, 95% confidence interval [CI] $[-1.10; 0.63]$, $p = .59$); while systemic antimicrobials (amoxicillin or azithromycin) showed an added effect on treatment success ([PPD ≤ 5 mm, no bleeding or suppuration, no progressive bone loss]; $n = 2$, RR = 1.84, 95% CI $[1.17; 2.91]$, $p = .008$), but not in terms of PPD reduction ($n = 2$, WMD = 0.93 mm, 95% CI $[-0.69; 2.55]$, $p = .26$), even if with substantial heterogeneity.

Conclusions: No single decontamination method demonstrated clear evidence of superiority compared to the others. Systemic antibiotics, but not Er:Yag laser, may provide short-term clinical benefits in terms of treatment success (CRD42020182303).

KEYWORDS

air-powder devices, curettes, decontamination, dental implants, implantoplasty, peri-implant diseases, surgical therapy, titanium brushes

1 | INTRODUCTION

Peri-implantitis is a plaque-associated pathological condition affecting tissues around dental implants characterized by the inflammation of the peri-implant mucosa and the progressive resorption of supporting bone (Berglundh, Armitage, et al., 2018). Its estimated prevalence has been reported between 1 and 47%, depending on the employed case definitions (Derks & Tomasi, 2015; Romandini, Berglundh, et al., 2021; Romandini, Lima, et al., 2021). Peri-implantitis has a bacterial etiology, and therefore the success of treatment mostly depends on arresting the inflammatory process through efficient control of infection and removal of dysbiotic biofilm from the implant surface (Lindhe & Meyle, 2008).

Despite no clinical guidelines being available yet, a stepwise therapeutic approach similar to the one used for periodontitis is employed in the management of peri-implantitis (Sanz et al., 2020). After an initial phase, including oral hygiene instructions, risk factor control and supra-mucosal instrumentation, implants affected by moderate-severe peri-implantitis undergo surgical treatment, which comprehends access, resective or reconstructive procedures (Heitz-Mayfield & Mombelli, 2014). Although those surgical approaches demonstrated favorable treatment outcomes in terms of probing pocket depth (PPD) reduction (Roccuzzo et al., 2018), in most cases the composite criteria employed for defining treatment success are not achieved (Carcuac et al., 2016; Khoury et al., 2019).

Incomplete implant decontamination represents the main reason for this limited predictability (Meyle, 2012), due to the complex micro- and macro-topography of titanium interfaces and bony defects anatomies (Koo et al., 2019). Several mechanical (curettes, ultrasonic, irrigations with saline, air powder abrasion, titanium brushes, implantoplasty), chemical (citric acid, chlorhexidine - CHX, enamel matrix derivatives - EMD, topical or systemic antimicrobials) and physical (laser, photodynamic therapy) decontamination methods have been proposed either alone or in combination (Carcuac et al., 2016; Klinge et al., 2002; Louropoulou et al., 2014). Despite some reviews having tried to comprehensively assess the efficacy of adjunctive measures for the treatment of peri-implantitis (Ramanauskaite et al., 2021; Schwarz et al., 2015), a focused synthesis of the effect of the proposed decontamination protocols during surgical treatment of peri-implantitis, as well as the identification of the eventual superiority of specific methods over others is currently lacking.

Therefore, the primary aim of the present systematic review was to answer the following PICOS question: 'In patients requiring surgical treatment of peri-implantitis (P), is any implant surface decontamination protocol (I) superior to others (C) in terms of clinical [changes in probing pocket depth (PPD) - primary outcome] and radiographic parameters (O) in randomized clinical trials (RCTs) (S)?' Moreover, the present systematic review aimed at comprehensively analyzing the longitudinal effects of the implant surface decontamination protocols tested in RCTs.

2 | MATERIALS AND METHODS

This systematic review is reported according to the PRISMA statement (Moher et al., 2009), and the protocol was registered on PROSPERO (CRD42020182303).

2.1 | Eligibility criteria

Inclusion criteria were determined a priori and organized according to the PICOS acronym:

(P) Population. Patients in good general health requiring surgical treatment of peri-implantitis.

(I) Interventions. Any type of local or systemic (i.e., systemic antimicrobials) implant surface decontamination protocol used during surgical treatment of peri-implantitis.

(C) Comparisons. Any possible comparison between different protocols for intra-surgical decontamination (including placebo).

(O) Outcomes of interest:

- Primary outcome: changes in PPD.

- Other considered outcomes: changes in marginal bone level (MBL), treatment success (possibly adhering to the definition to Carcuac et al., 2016 - residual PPD ≤5 mm, no bleeding/suppurative on probing [BoP/SoP], no progressive marginal bone loss after treatment), BoP/SoP, soft tissue level changes, clinical/relative attachment level (CAL/RAL), implant survival, need of retreatment, patient-reported outcome measures (PROMs) and adverse events.

(S) Types of studies. RCTs with at least 6-months follow-up and a minimum of 10 patients (5 per group). RCTs not directly comparing different decontamination protocols were excluded.

2.2 | Search methods for the identification of studies

Four electronic databases were independently searched for relevant articles using the following search algorithms by two authors (GB and FC).

2.2.1 | [MEDLINE] (via PUBMED) (2021-12-20)

("peri-implantitis"[MeSH Terms] OR "peri-implantitis"[All Fields] OR "peri implantitis"[All Fields]) AND ("therapy"[All Fields] OR "treatment"[All Fields] OR "Decontamination"[Mesh] OR Antiinfective agents OR antimicrobials OR antibiotics OR "Therapeutics"[Mesh] OR "therapeutics"[All Fields]) NOT (retrospective OR review OR in vitro OR case report OR orthopedic OR animal OR experimental)

Filter: English.

Search 2: [EMBASE] (via ELSEVIER) (2021-12-20)

('periimplantitis'/exp OR periimplantitis OR (perimplant AND ['disease'/exp OR disease])) AND ('therapy'/exp OR 'decontamination'/exp OR 'debridement'/exp OR antimicrobials OR antibiotics) NOT (([retrospective OR review OR in] AND vitro OR case) AND report OR orthopedic OR animal OR experimental)

Filter: English.

Search 3: [SCOPUS] (2021-12-20)

periimplantitis AND (therapy OR decontamination) AND NOT (review OR in AND vitro OR animal).

Search 4: [CENTRAL (Cochrane central register of controlled trials)] (2021-12-20)

(peri-implantitis OR periimplantitis OR peri implantitis) AND (surgical treatment OR surgery OR surgical)

In addition, duplicate (GB and NB) hand-searching was performed from January 2010 to June 2021 on the following journals: Journal of Clinical Periodontology, Journal of Periodontology and Clinical Oral Implants Research. Reference lists and previous systematic reviews were also screened.

2.3 | Study selection

The titles and abstracts of all identified studies were screened independently and in duplicate by two calibrated reviewers (NB and GB). Initial calibration of investigators was achieved by online discussion sessions. Any disagreement was resolved by discussion with a third reviewer (FC). Full text of studies of possible relevance were assessed in duplicate by two reviewers (NB and FC), in order to make a final decision about their inclusion. Percentage of agreement and kappa statistics were employed to score inter-rater agreement (yes/no) of the screening and full-text analysis processes. Disagreements were again resolved by a joint discussion with a third review author (GB). The reasons for study exclusion after full text analysis were recorded.

2.4 | Data extraction and management

Data from included studies were extracted in duplicate by two reviewers (GB and FC) using predefined data extraction forms. If necessary, corresponding authors of the included studies were contacted for clarification of any missing information. Data on general information (first author, year of publication and setting); methods (study design, diagnostic criteria for peri-implantitis, follow-up period); participants (inclusion criteria, number of randomized participants and implants, drop-outs, number of analyzed participants and implants, age, gender, smoking, history of periodontitis, implant surface), interventions and controls (pre-surgical procedures, type of surgery, decontamination protocols,

biomaterials, post-surgical care, frequency of supportive peri-implant care) and outcome/results of interest (for each outcome considered: collected or not, definition, time-points, results). The type of surgery was categorized as access flap, resective, reconstructive or combined (Appendix S1).

2.5 | Assessment of risk of bias in the included studies

The risk of bias in the included studies was assessed independently and in duplicate by two review authors (GB and FC) according to the RoB2 tool, considering PPD reduction as the main outcome of interest (Sterne et al., 2019).

2.6 | Data synthesis

In the presence of at least 2 studies, random-effects meta-analyses were carried out using specific softwares (OpenMeta [Analyst], Brown University, RI, USA; RevMan v5.4, The Cochrane Collaboration, 2020), using the Mantel-Haenszel method for dichotomous data, and the inverse of variance method for continuous data. In order to account for within-patient correlation in studies which failed to adjust for it, an intracluster correlation coefficient of 0.07 was assumed for the calculation of the effective sample size and CIs (Campbell et al., 2012; see Appendix S2). Statistical significance was set in advance to $p < 0.05$. Only subgroup analyses according to the employed surgical approach (access, resective or reconstructive) were performed.

Two different sets of analyses were conducted. First, the effect of each implant surface decontamination protocol was assessed by comparing baseline values with values at follow-up. Continuous data were combined in weighted mean effects (WME) and 95% confidence intervals (CIs), while binary data were pooled as weighted mean percentage (WMP) and 95% CIs. Second, when possible, pairwise comparisons were carried out to compare different decontamination protocols. The estimates of the effect were expressed as weighted mean differences (WMD) and 95% CIs for continuous outcomes and as risk ratio (RR) and 95% CIs for dichotomous outcomes.

Statistical heterogeneity among studies was explored using the I^2 index and the Cochrane's Q statistic ($p < 0.1$). Network meta-analysis was not possible due to the lack of common comparators.

2.7 | Certainty of evidence

The Grades of Recommendation, Assessment, Development and Evaluation (GRADE) tool has been used to summarize the overall quality of the evidence for the questions for which pairwise meta-analyses were available (Guyatt et al., 2011). The certainty of the

TABLE 1 Included studies: Decontamination protocols employed

Study	Study design	Groups	Randomized patients	Randomized implants	Follow-up (mo)	Diagnosis of peri-implantitis	Pre-treatment phase at affected implants
Lasserre et al. (2020)	RCT parallel (test/ control)	Implantoplasty	16	22	6	MBL ≥ 2 mm, PPD ≥ 5 mm, BOP and/or SoP	OHI and supragingival instrumentation with scalers, polishing paste, and rubber cups 4 w before surgery
		Glicine air polishing	15	20			
Cha et al. (2019)	RCT parallel (test/control)	Minocycline ointments	25	25	12	MBL ≥ 3 mm, PPD ≥ 6 mm, and BOP	Supragingival instrumentation, and standardized OHI
		Placebo	25	25			
Toma et al. (2019)	RCT parallel (2 test/1 control)	Perio Flow	16	22	6	MBL ≥ 2 mm, PPD ≥ 5 mm, BOP and/ or SoP	2 w before surgery, OHI and professional supragingival instrumentation, using a rubber cup with polishing paste
		Titanium brushes	16	23			
		Plastic curettes	15	25			
Albaker et al. (2018)	RCT parallel (test/control)	Photodynamic Therapy	11	11	12	MBL ≥ 2 mm, PPD ≥ 5 mm, BOP and/ or SoP	Full mouth SRP using ultrasonic scaler and hand instruments
		Curettes	13	13			
Hallström et al. (2017)	RCT parallel (test/control)	Access flap + Systemic antimicrobials	20	20	12	MBL ≥ 3 mm, PPD ≥ 5 mm and BOP/SoP	NR
		Access flap + placebo	19	19			
Carcuac et al. (2016) also reported in: Carcuac et al. (2017)	RCT parallel (2 test/2 control)	Antibiotic + Antiseptic +	27	47	12	MBL ≥ 2 mm, PPD ≥ 6 mm, BOP and/or SoP	Supragingival instrumentation using rubber cups, polishing paste, and OHI
		Antibiotic + Antiseptic -	25	46			
		Antibiotic Antiseptic +	24	49			
		Antibiotic -Antiseptic -	24	37			
Isehmed et al. (2016) also reported in: Isehmed et al. (2018)	RCT parallel (test/control)	Access flap + EMD	15	15	12	MBL ≥ 3 mm, PPD ≥ 5 mm BOP and/or SoP	Periodontitis was treated with mechanical debridement and OHI
		Access flap	14	14			

Systemic ABX	Mechanical decontamination	Chemical decontamination	Physical decontamination	Biomaterials	Post-op	Frequency of SPT
No	Plastic curettes, diamond burs Plastic curettes, air powder device	Sterile saline	NA	NA	0.2% CHX x 10 d; Ibuprofen 3 x 600 mg for 2 d, paracetamol 1 g	1w, 3 m, 6 m
Amoxicillin 3 x 500 mg for 3 d	Titanium curettes, ultrasonic scaler, titanium brush, air-powder device	Minocycline ointment Placebo ointment	NA	NA	Ibuprofen 3 x 600 for 3 d	1w, 1 m, 3 m, 6 m (Minocycline or placebo administered at 1 and 3 mo)
No	Air powder device Titanium brushes Plastic curettes	Sterile saline	NA	NA	0.2% CHX and paracetamol 3 g for 10 d	1w, 3 m, 6 m
Amoxicillin + Clavulanic Acid 3 x 625 mg for 7 d	Curettes plus saline soaked cotton gauzes Curettes plus saline soaked cotton gauzes	Sterile saline Sterile saline	Photodynamic therapy (methylene blue + diode laser) NA	NA	Ibuprofen 3 x 600 mg for 7 d, 0.2% CHX for 2 weeks	1w, 3 m, 6 m, 9 m, 12 m
Azithromycin (2 x 250 mg at surgery, and 1 x 250 mg for 4 d)	Titanium curettes plus gauze soaked in saline	NA	NA	NA	CHX 0.12% twice x 10 d	2w, 6w, 3 m, 6 m, 12 m
No	Titanium curettes plus gauze soaked in saline	NA	NA			
Amoxicillin 2 x 750 mg for 10 d commenced 3 d prior to surgery	Titanium curettes Titanium curettes	Gauze soaked in 0.2% CHX Sterile saline	NA	NA	1 min 0.2% CHX twice daily for 14 d	2w, 3 m, 6 m, 9 m, 12 m
No	Titanium curettes Titanium curettes	Gauze soaked in 0.2% CHX Sterile saline				
No	Ultrasonic scaler, titanium curettes	Sterile saline plus EMD Sterile saline	NA	NA	2 x 10 ml CHX for 6 w and not chew or brush on the treated side for 2 weeks	2w, 6w, 3 m, 6 m, 9 m, 12 m

(Continues)

TABLE 1 (Continued)

Study	Study design	Groups	Randomized patients	Randomized implants	Follow-up (mo)	Diagnosis of peri-implantitis	Pre-treatment phase at affected implants
Papadopoulos et al. (2015)	RCT parallel (test/control)	Access flap + Diode laser	9	9	6	MBL ≥ 2 mm, PPD ≥ 5 mm BOP and/or SoP	Mechanical debridement using ultrasonics and hand instruments to the whole dentition
		Access flap	10	10			
Wang et al. (2020)	RCT parallel (test/control)	Er:Yag Laser	12	12	6	MBL > 2 mm, PPD ≥ 5 mm BOP and/or SoP	Full mouth prophylaxis performed with piezo-instruments and stainless-steel hand scalers
		Placebo	12	12			
deTapia et al (2019)	RCT parallel (test/control)	Titanium brushes	15	15	12	MBL $> 30\%$, PPD ≥ 6 mm and BOP and/or SoP	Subgingival scaling with plastic curettes and irrigation with 0.12% CHX
		Ultrasonic scalers	15	15			
Schwarz et al., 2011 also reported in: Schwarz et al. (2012, 2013, 2017)	RCT parallel, single blind (test/control)	Er:Yag Laser	16	16	6	PPD of ≥ 5 mm and an intrabony component of > 3 mm as estimated clinically	Non-surgical instrumentation using plastic curettes, combined with 0.2% CHX solution and CHX gel 0.2%.
		Plastic curettes	16	16			
Schlee et al. (2019)	RCT parallel (test/control)	Electrolytic method (EC)	12	12	6	MBL ≥ 3 mm, PPD ≥ 6 mm, and BOP and/ or SoP	Suprastructures removed 14 days before surgery, implants cleaned by powder spray and CHX. Cover screw was placed
		Powder spray plus EC	12	12			
Isler et al. (2018)	RCT parallel (test/control)	Ozone therapy	23	42	12	MBL ≥ 2 mm, deepening PPD, and BOP and/or SoP	Non-surgical treatment provided. In test group, ozone therapy was initiated
		Saline irrigation	23	NR			
de Waal et al. (2015)	RCT parallel, double blind (test/control)	2% CHX	22	49	12	MBL ≥ 2 mm, PPD ≥ 5 mm and BOP and/or SoP	Mechanical debridement of implants, suprastructures, and remaining dentition
		0.12% CHX + 0.05% CPC	22	59			
de Waal et al. (2013)	RCT parallel, single blind (test/control)	0.12% CHX + 0.05% CPC	15	31	12	MBL ≥ 2 mm, PPD ≥ 5 mm and BOP and/or SoP	Mechanical debridement of implants, suprastructures, and remaining dentition
		Gauze soaked in saline	15	48			

Systemic ABX	Mechanical decontamination	Chemical decontamination	Physical decontamination	Biomaterials	Post-op	Frequency of SPT
No	Plastic curettes, plus gauzes soaked in saline	NA	Diode laser NA	NA	2 x 0.12% CHX for 2 w and a careful tooth brushing with a soft toothbrush	2w, 3 m, 6 m
Amoxicillin 3 x 500 mg for 10 d	Ultrasonic scaler, stainless-steel curettes	NA	Er:Yag Laser Sham laser application	Alloplastic bone graft + acellular dermal matrix	2 x CHX for 1 w and Ibuprofen 600 mg as needed	2w, 1 m, 3 m, 6 m
Amoxicillin 3 x 500 mg and Metronidazole 3 x 500 mg for 7 d	Plastic ultrasonic scaler, titanium brushes (intraosseous) Plastic ultrasonic scaler (intraosseous)	3% H2O2	NA	Alloplastic bone graft + collagen membrane	2 x 0.12% CHX for 2 w	1w, 2w, 3w, 1 m, 3 m, 6 m, 9 m, 12 m
No	Plastic curettes + cotton pellets soaked in saline (intraosseous part)	NA	Er:Yag Laser (intraosseous part) NA	Xenogenic bone graft + collagen membrane	2 x 0.12% CHX for 2 w	2w, 4w, 6w, 8w, 4 m, 6 m
No	Curettes and/or ultrasonic devices Curettes and/or ultrasonic devices plus powder spray		Pilot electrolytic approach for 120 s, then sterile saline.	Autogenous bone graft and xenograft 50:50 + collagen membrane	NR	2w, 6w, 6 m
Amoxicillin 3 x 500 mg and Metronidazole 3 x 500 mg for 1 week	Titanium curettes	Sterile saline plus ozone delivery Sterile saline	NA	Xenogenic bone graft + growth factors	2 x 0.12% CHX for 2 weeks, anti-inflammatory and analgesic drugs for the first 3 d	1w, 1 m, 3 m, 6 m, 9 m, 12 m
No	Gauze soaked in saline	2% CHX 0.12% CHX + 0.05% CPC x 1 min	NA	NA	0.12% CHX + 0.05% CPC without alcohol (two times daily x 2 weeks)	2w, 3 m, 6 m, 9 m, 12 m
No	Gauze soaked in saline	2% CHX x 1 min NA	NA	NA	0.12% CHX + 0.05% CPC without alcohol (two times daily x 2 weeks)	2w, 3 m, 6 m, 9 m, 12 m

TABLE 1 (Continued)

Study	Study design	Groups	Randomized patients	Randomized implants	Follow-up (mo)	Diagnosis of peri-implantitis	Pre-treatment phase at affected implants
Romeo et al (2005) also reported in: Romeo et al. (2007)	RCT parallel, single blind (test/control)	Implantoplasty	NR	19	12	PPD ≥ 5 mm, radiographic evidence of horizontal peri-implant radiolucency and BOP and/or SoP	NR
		No implantoplasty	NR	16			

Abbreviations: BOP, bleeding on probing; CHX, chlorhexidine; CPC, cetylpyridinium chloride; EMD, enamel matrix derivatives; MBL, marginal bone level; NR, not reported; PPD, probing pocket depth; SoP, suppuration on probing.

	Randomization	Deviations from intended intervention	Missing outcome data	Measurement of the outcome	Selection of reported results	Overall risk of bias		
Romeo et al. 2005	+	+	+	+	+	+	Low risk	+
Schwartz et al. 2011	+	+	+	+	+	+	Some concerns	?
De Waal et al. 2013	+	+	+	+	+	+	High risk	-
Papadopoulos et al. 2015	+	+	?	+	+	?		
De Waal et al. 2015	+	+	+	+	+	+		
Isehede et al. 2016	+	+	+	+	+	+		
Carcuac et al. 2016	?	+	+	+	?	?		
Hallström et al. 2017	+	?	+	+	?	?		
Isler et al. 2018	?	+	+	+	+	?		
Albaker et al. 2018	+	?	+	+	+	?		
Thoma et al. 2019	?	+	+	+	+	?		
Schlee et al. 2019	+	+	+	+	-	-		
De Tapia et al. 2019	+	+	+	+	+	+		
Cha et al. 2019	+	+	+	+	+	+		
Lassere et al. 2020	?	+	+	+	+	?		
Wang et al. 2020	+	+	+	+	+	+		

FIGURE 1 Risk of bias assessment

body of evidence was not evaluated for comparisons for which meta-analyses were not possible.

Schlee et al., 2019; Schwarz et al., 2011; Schwarz et al., 2012; Schwarz et al., 2013; Schwarz et al., 2017; Tapia et al., 2019; Toma et al., 2019; Wang et al., 2020).

3 | RESULTS

3.1 | Study selection

The electronic search yielded 1835 records and hand searching identified 6 additional studies (Figure S1). After removal of duplicates, the total number of screened articles was 1497. Twenty-six records were selected for full-text analysis (agreement: 98.6%; Kappa = 0.68, 95% CI: 0.55–0.81), which resulted in the exclusion of 4 of them (agreement: 96.1%; Kappa = 0.83, 95% CI: 0.52–1.14 - reasons for exclusion reported in Table S1) and the final inclusion of 22 manuscripts reporting on 16 RCTs (Albaker et al., 2018; Carcuac et al., 2016; Cha et al., 2019; de Waal et al., 2013; de Waal et al., 2015; Hallström et al., 2017; Isehede et al., 2016; Isehede et al., 2018; Isler et al., 2018; Lasserre et al., 2020; Papadopoulos et al., 2015; Romeo et al., 2004; Romeo et al., 2007;

3.2 | Characteristics of the included studies

Table 1 shows details about the characteristics of the included studies.

3.2.1 | Design and settings

All RCTs had a parallel arm design, and all of them except 3 were carried out in Europe. Fourteen studies included one experimental and one control group, while 2 studies adopted multiple arms. Six included studies had a 6-months follow-up, while the remaining 12-months. Four trials were also reported at longer follow-up periods, such as 2 years (Schwarz et al., 2012), 3 years (Carcuac

Systemic ABX	Mechanical decontamination	Chemical decontamination	Physical decontamination	Biomaterials	Post-op	Frequency of SPT
Amoxicillin 50 mg/kg/die for 8 d per os	Implantoplasty (burs) None	Gel of metronidazole plus a solution of tetracycline hydrochloride	NA	NA	0.2% CHX (10 mL for 1 min at interval of 8 h for 2 w)	NR

et al., 2017; Isehmed et al., 2018; Romeo et al., 2007), 4 years (Schwarz et al., 2013), 5 years (Isehmed et al., 2018) and 7 years (Schwarz et al., 2017).

3.2.2 | Patients and implants

Diagnostic criteria for peri-implantitis were consistent across most included studies except for minor differences (either bone loss/level ≥ 2 or 3 mm, together with PPD ≥ 5 or 6 mm, and BoP/SoP). In total, 849 implants were treated in 604 patients. Percentage of current smokers ranged from 14.2% to 50%, whereas in 5 studies smokers were excluded. Only 5 studies included implants with a machined surface, with proportions ranging from 1.3 to 35%.

3.2.3 | Interventions

Eight studies performed a non-surgical sub-mucosal instrumentation prior to the surgical treatment, while the remaining 8 only performed supra-mucosal instrumentation or did not report this information. Eight trials reported decontamination protocols during access surgery, 3 in resective surgery, 2 in reconstructive and 3 in combined procedures.

Curettes were the most used method for local mechanical decontamination (25 arms of 12 RCTs), either alone or in combination with other devices. Plastic curettes were employed in 6 arms of 4 RCTs, titanium curettes in 12 arms of 5 RCTs, only one study employed stainless steel curettes in 2 arms, while two studies did not report this information (Albaker et al., 2018; Schlee et al., 2019). Gauzes soaked in saline were employed in 16 arms of 7 studies; ultrasonic scalers were adopted in 10 arms of 5 studies; air-powder devices in 5 arms of 4 studies; titanium brushes in 4 arms of 3 studies; while implantoplasty was carried out in 6 arms of 4 studies. Among local chemical decontamination agents, CHX-based formulations were applied in 4 arms of 3 studies, either in 2% gel formulation or in solution (at 0.2% concentration or at 0.12% in combination with 0.05% cetylpyridinium chloride - CPC). Other local chemical decontamination employed were a gel of metronidazole, a solution of tetracycline hydrochloride, minocycline ointments, 3% hydrogen

peroxide (H₂O₂), and EMD (each one in 1 arm of 1 study). Among local physical decontamination, lasers were used in 3 arms of 3 different studies (2 studies employed a Er:Yag laser, while the remaining one a diode laser); electrolytic current was employed in 2 arms of one study; while ozone therapy and photodynamic therapy were each one part of 1 arm of 1 study.

Six studies utilized systemic chemical decontamination agents (i.e., antibiotics) in all treatment groups, 2 only in test groups as part of the studied comparisons, while the remaining 8 did not administer them. Of the 8 studies which utilized systemic antibiotics, 4 trials employed amoxicillin alone, 3 amoxicillin in combination with either clavulanic acid or metronidazole, while the remaining one azithromycin. Systemic antibiotics were administered prior to the surgical treatment in two RCTs, while in 6 studies they were prescribed on the day of the surgery.

3.3 | Risk of bias within studies

The risk of bias assessment for the included RCTs is summarized in Figure 1. Seven trials resulted at low risk of bias, eight trials with some concerns and the remaining one at high risk.

3.4 | Effect of different decontamination protocols

3.4.1 | Results of individual studies

The results of the included studies in terms of effect of different decontamination protocols are reported in the Appendix. Moreover, a summary of the results on the main outcomes considered in the present systematic review is reported in Table 2, while on the other outcomes is reported in Table S2 ("Effect of studied decontamination protocols" columns).

3.4.2 | Meta-analyses

Data were pooled using WMEs and WMPs (comparisons against baseline). With regards to access flap procedures, the single use of

TABLE 2 Included studies: Summary of the results for the main outcomes of interest (PPD, MBL and treatment success)

Study	Type of surgery	Tested methods	Common methods	Analyzed patients	Analyzed implants	Success criteria	Effect of studied decontamination protocols	Comparison of the studied protocols (effect of test procedure)
Lasserre et al. (2020)	Access flap	Test: Implantoplasty (diamond burs) Control: Glycine air polishing	Plastic curettes, sterile saline	15	20	PPD ≤5 mm, no BoP/SoP, and no further bone loss ≥0.5 mm from baseline	ΔPPD: -4.00 ± 1.83 mm ΔMBL: -0.26 ± 1.08 mm Success: 15.0% ΔPPD: -3.29 ± 1.73 mm ΔMBL: -0.53 ± 0.94 mm Success: 26.0%	ΔPPD: 0.61 mm ΔMBL: 0.27 mm (in favour of control group)
Cha et al. (2019)	Access flap	Test: Repeated application of minocycline ointments Control: Repeated application of placebo ointments	Titanium curettes, ultrasonic scaler, titanium brush, air-powder device, systemic amoxicillin	24	24	PPD <5 mm, no BoP, and no further bone loss	ΔmPPD: -2.68 ± 1.73 mm ΔwPPD: -3.58 ± 2.32 mm ΔMBL: -0.72 ± 0.56 mm Success: 66.7% ΔmPPD: -1.55 ± 1.86 mm ΔwPPD: -2.45 ± 2.13 mm ΔMBL: -0.31 ± 0.49 mm Success: 36.3%	ΔmPPD: 1.13 mm ^a ΔMBL: 0.41 mm ^a Success: RR = 2 ^a
Toma et al. (2019)	Access flap	Test A: Air powder device Test B: Titanium brushes Control: Plastic curettes	Sterile saline	16	22	PPD ≤5 mm, no BoP/SoP, and no further bone loss ≥0.5 mm from baseline	ΔPPD: -2.39 ± 1.12 mm ΔMBL: -0.95 ± 0.24 mm Success: 29.0% ΔPPD: -2.41 ± 0.48 mm ΔMBL: -1.12 ± 0.24 mm Success: 33.0% ΔPPD: -1.37 ± 0.87 mm ΔMBL: -0.51 ± 0.17 mm Success: 22.0%	A versus B ΔPPD: 0.2 mm ^a ΔMBL: 0.17 mm ^a A versus control ΔPPD: 1.02 mm ΔMBL: 0.44 mm B versus control ΔPPD: 1.04 mm ΔMBL: 0.61 mm
Albaker et al. (2018)	Access flap	Test: Photodynamic Therapy Control: NA	Curettes, saline soaked cotton gauze, systemic amoxicillin plus clavulanic acid	11	11	NR	NR	NR
Hallström et al. (2017)	Access flap	Test: Systemic azithromycin Control: Placebo	Titanium curettes plus gauze soaked in saline	15	15	PPD ≤5 mm, no BoP/SoP, and no further bone loss ≥0.5 mm from baseline to 1y	ΔPPD: -1.7 ± 1.1 mm Success: 46.6% ΔPPD: -1.6 ± 1.5 mm Success: 25.0%	ΔPPD: 0.1 mm ΔMBL: 0.5 mm (in favour of control group) Success: RR = 1.8

TABLE 2 (Continued)

Study	Type of surgery	Tested methods	Common methods	Analyzed patients	Analyzed implants	Success criteria	Effect of studied decontamination protocols	Comparison of the studied protocols (effect of test procedure)
Carcuac et al. (2016) also reported in: Carcuac et al. (2017)	Access flap	Test A: Gauze soaked in 0.2% CHX +systemic amoxicillin	Titanium curettes	27	47	PPD ≤5mm, no BoP/SoP, and no further bone loss ≥0.5 mm	ΔwPPD: -2.80 ± 1.87 mm ΔMBL: -0.18 ± 1.15 mm Success: 40.4%	Test A vs Control A ΔwPPD: 0.64 mm ΔMBL: 0.87 mm
		Test B: Gauze soaked in saline + systemic amoxicillin		25	46	from baseline	ΔwPPD: -3.44 ± 1.66 mm ΔMBL: -0.51 ± 0.84 mm Success: 65.2%	Test B vs Control B ΔwPPD: 1.75 mm ΔMBL: 1.47 mm
	Control A: Gauze soaked in 0.2% CHX Control B: Gauze soaked in saline		23	48		ΔwPPD: -2.16 ± 1.79 mm ΔMBL: +0.69 ± 1.32 mm Success: 37.5%	Test A vs Test B ΔwPPD: 0.64 mm ^a ΔMBL: 0.33 mm ^a ^a : in favour of B	Control A vs Control B ΔwPPD: 0.47 mm ΔMBL: 0.3 mm
			24	37		ΔwPPD: -1.69 ± 2.22 mm ΔMBL: +0.96 ± 1.42 mm Success: 35.1%	ΔPPD: 1.5 mm ΔMBL: 0.5 mm (in favour of control group)	
Ished et al. (2016) also reported in: Isehmed et al. (2018)	Access flap	Test: EMD	Ultrasonic scaler, titanium curettes	12	12	NR	ΔPPD: -2.5 ± 2.0 mm ΔMBL: -0.7 ± 1.1 mm	
		Control: NA		13	13		ΔPPD: -4.0 ± 2.9 mm ΔMBL: -0.2 ± 1.1 mm	
Papadopoulos et al. (2015)	Access flap	Test: Diode laser Control: NA	Plastic curettes	8 8	8 8	NR NR	NR NR	NR
Wang et al. (2020)	Combined	Test: Er:Yag laser	Ultrasonic scaler, stainless-steel	12	12	NR	ΔPPD: -2.65 ± 2.14 mm ΔMBL: -1.27 ± 1.14 mm	ΔPPD: 0.80 mm ^a ΔMBL: 0.19 mm
		Control: Sham laser application	curettes, systemic amoxicillin	12	12		ΔPPD: -1.85 ± 1.71 mm ΔMBL: -1.08 ± 1.04 mm	
deTapia et al (2019)	Combined	Test: Titanium brushes	Ultrasonic scalers, 3% H2O2, systemic amoxicillin plus metronidazole	15	15	Absence of PPD ≥5mm, with no BOP/SOP and no additional peri-implant bone loss	ΔmPPD: -2.84 ± 0.93 mm ΔwPPD: -4.87 ± 1.55 mm ΔMBL: -2.51 ± 1.21 mm Success: 66.7%	ΔmPPD: 1.29 mm ^a ΔmMBL: 1.78 mm ^a ΔwMBL: 1.68 mm ^a Success: RR = 2.6 ^a
		Control: NA		12	12		ΔmPPD: -1.55 ± 1.86 mm ΔwPPD: -2.85 ± 1.91 mm ΔMBL: -0.73 ± 1.26 mm Success: 23.0%	

TABLE 2 (Continued)

Study	Type of surgery	Tested methods	Common methods	Analyzed patients	Analyzed implants	Success criteria	Effect of studied decontamination protocols	Comparison of the studied protocols (effect of test procedure)
Schwarz et al., 2011 also reported in: Schwarz et al. (2012, 2013, 2017)	Combined	Test: Er:Yag Laser Control: NA	Plastic curettes + cotton pellets soaked in saline	15 15	15 15	NR	ΔPPD: -1.7 ± 1.4 mm ΔPPD: -2.4 ± 1.5 mm	ΔPPD: 0.7 mm (in favour of control group)
Schlee et al. (2019)	Rec	Test: NA Control: Powder spray	Curettes and/or ultrasonic devices, pilot electrolytic method	12 11	12 11	NR	ΔMBL: -2.71 ± 1.70 mm ΔMBL: -2.81 ± 2.15 mm	ΔMBL: 0.10 mm (in favour of control group)
Isler et al. (2018)	Rec	Test: Ozone therapy Control: NA	Titanium curettes, saline irrigation, systemic amoxicillin plus metronidazole	20 21	30 30	PPD < 5 mm without BOP and/or SoP, no further BL, and DF ≥ 1 mm	ΔPPD: -3.5 ± 1.31 mm ΔMBL: -2.32 ± 1.28 mm Success: 50.0% ΔPPD: -2.42 ± 1.23 mm ΔMBL: -1.17 ± 0.77 mm Success: 36.6%	ΔPPD: 1.1 mm ΔMBL: 1.15 mm ^a
de Waal et al. (2015)	Res	Test: 2% CHX Control: 0.12% CHX + 0.05% CPC	Gauze soaked in saline	21 20	48 54	PPD ≤ 5 mm, no BoP/SoP, and no further bone loss ≥ 0.5 mm from baseline	ΔPPD: -1.68 ± 1.06 mm ΔMBL: -0.24 ± 0.72 mm Success: 14.3% ΔPPD: -2.01 ± 1.26 mm ΔMBL: -0.14 ± 0.49 mm Success: 28.8%	ΔPPD: 0.33 mm (in favour of control group) ΔMBL: 0.10 mm
de Waal et al. (2013)	Res	Test: 0.12% CHX + 0.05% CPC Control: NA	Gauze soaked in saline	15 12	31 38	PPD ≤ 5 mm, no BoP/SoP, and no further bone loss ≥ 0.5 mm from baseline	ΔPPD: -2.21 ± 2.01 mm ΔMBL: -0.78 mm ± 0.93 mm Success: 3.2% ΔPPD: -1.64 ± 1.03 mm ΔMBL: -0.58 mm ± 0.86 mm Success: 2.1%	ΔPPD: 0.57 mm ΔMBL: 0.18 mm
Romeo et al (2005) also reported in: Romeo et al. (2007)	Res	Test: Implantoplasty (diamond burs) Control: No implantoplasty	Gel of metronidazole, solution of tetracycline hydrochloride, systemic amoxicillin	19 16	19 16	NR	ΔmMBL: 0 ± 0.14 mm ΔmMBL: -0.01 ± 0.13 mm ΔmMBL: +0.51 ± 0.49 mm ΔmMBL: +0.56 ± 0.43 mm	ΔPPD: 2.48 mm ΔmMBL: 0.51 mm ΔmMBL: 0.57 mm

Abbreviations: BoP, bleeding on probing; CHX, chlorhexidine; CPC, cetylpyridinium chloride; MBL, mean bone level (mMBL: mesial, dMBL: distal); PPD, probing pocket depth (mPPD: mean, wPPD: worst); Rec, reconstructive surgery; Res, restorative surgery; RR, relative risk; SoP, suppuration on probing; wPPD, worst probing pocket depth.
^aSignificant difference between the test and control group.




FIGURE 2 Meta-analyses for the effect of different decontamination protocols tested in 2 or more RCTs. A-C: effect of the single use of curettes in access flap surgery on PPD reduction, MBL change and treatment success. D-F: effect of systemic antibiotics (AB) in addition to mechanical decontamination in access flap surgery on PPD reduction, MBL change and treatment success; in parenthesis are reported other eventual chemical decontaminants used in association with AB. G-I: effect of 0.12% chlorhexidine (CHX) and 0.05% cetylpyridinium chloride (CPC) in addition to mechanical decontamination in resective surgery on PPD reduction, MBL change and treatment success. L: effect of Er:Yag laser in addition to mechanical and/or chemical decontamination in combined surgery on PPD reduction.

curettes for surface decontamination resulted in a WME of 1.46 mm for PPD reduction (3 arms of 3 trials; 95% CI:1.17/1.74; $I^2=0.00\%$), in a negligible MBL change (2 arms of 2 trials; WME = -0.21 mm; 95% CI:-1.65/1.23; $I^2=97.35\%$), while the WMP for treatment success amounted to 28.6% (3 arms of 3 trials; 95% CI:18.5/38.7; $I^2=0.00\%$) (Figure 2a,b,c). The use of systemic antimicrobials in combination with local decontamination methods resulted in a WME of 2.46 mm for PPD reduction (5 arms of 3 trials; 95% CI:1.74/3.18; $I^2=86.01\%$), in a WME of 0.44 mm for MBL changes (4 arms of 2 trials; 95% CI:0.22/0.67; $I^2=70.16\%$) and in a WMP of 51% for treatment success (3 arms of 2 trials; 95% CI:33/69; $I^2=69.95\%$) (Figure 2d,e,f).

With regards to resective surgery, the use of 0.12% CHX+0.05% CPC resulted in a WME of 2.05 mm for PPD reduction (2 arms of 2 trials; 95% CI:1.72/2.38; $I^2=0$), a WME of -0.49 mm for MBL changes (2 arms of 2 trials; 95% CI:-1.02/0.04; $I^2=85.73$) and a WMP of 7.8% for treatment success (2 arms of 2 trials; 95% CI:2.1/17.7; $I^2=63.15$) (Figure 2g,h,i).

With regards to reconstructive surgery, no meta-analysis was carried out since the studied decontamination protocols were only tested once.


With regards to combined surgery, the use of the Er:Yag laser resulted in a WME of 2.04 mm for PPD reduction (2 arms of 2 trials; 95% CI:1.15/2.94; $I^2=43.24$) (Figure 2l).

3.5 | Comparison among different decontamination protocols


3.5.1 | Results of individual studies

The results of the included studies in terms of comparison between different decontamination protocols on the main outcomes considered in the present systematic review are reported in Table 2, while the results on the other considered outcomes are reported in

(a) Systemic AB vs placebo: PPD reduction



(b) Systemic AB vs placebo: Treatment success



(c) Er:Yag laser vs no laser: PPD reduction




FIGURE 3 Meta-analyses for the comparison among different decontamination protocols reported in 2 or more RCTs. A-B: added effect of systemic antibiotics (AB) over mechanical decontamination in access flap surgery on PPD reduction and treatment success. C: added effect of Er:Yag laser in addition to mechanical and/or chemical decontamination in combined surgery on PPD reduction.

Table S2 ("Comparison of the studied protocols" columns). Briefly, only 7 studies reported better clinical and/or radiographic results at 6- or 12-months examinations for one decontamination protocol over the others.

With regards to access flap surgery, Cha et al. (2019) reported 1.13 mm more PPD reduction, as well as 0.41 mm better MBL changes and 30.4% more treatment success using repeated local delivery of minocycline ointment over placebo. Toma et al. (2019) reported a 1.04 mm more mean PPD reduction and 0.61 more bone gain at the 6-months examination with the use of titanium brushes vs. the use of plastic curettes, as well as a 1.02 mm more mean PPD reduction and 0.43 mm more bone gain with the use of a glycine air-powder device versus the use of plastic curettes. Carcuac et al. (2016) reported a better efficacy of amoxicillin alone versus CHX soaked gauzes in terms of deepest PPD reduction (1.28 mm) and of bone level changes (1.20 mm). Moreover, amoxicillin alone showed 1.75 mm more deepest PPD reduction and 1.47 mm better MBL changes compared to the use of neither amoxicillin nor CHX soaked gauzes. In addition, amoxicillin combined with CHX soaked gauzes resulted in better MBL changes than the use of CHX alone (0.87 mm) and saline soaked gauzes (1.14 mm). Finally, the use of amoxicillin, but not of CHX soaked gauzes, has shown an added

effect on treatment success, but only in implants with modified surfaces.

With regards to resective surgery, Romeo et al. (2004) studied the added effect of implantoplasty over the use of a 25% metronidazole gel followed by the use of a solution of 50 mg/mL tetracycline hydrochloride for 3 minutes. At the 12-months examination, 2.48 mm less PPD values were found at the site-level analysis in the implantoplasty group.

With regards to reconstructive surgery, Isler et al. (2018) reported an added effect of ozone therapy over the combined use of titanium curettes and sterile saline irrigation in terms of bone gain (1.15 mm) at the 12-months examination.

With regards to combined surgery, Wang et al. (2020) showed that Er:Yag laser resulted in an added effect of 0.8 mm in mean PPD reduction at 6-months follow-up than sham laser application, when both used in addition to ultrasonic scalers and steel curettes. De Tapia et al. (2019) reported an added effect of titanium brushes over the combined use of plastic ultrasonic scaler and 3% H₂O₂ at 12-months follow-up in terms of mean and deepest PPD reduction (added effect: 1.29 mm and 2.02 mm, respectively), mean and deepest bone level change (added effect: 1.78 mm and 1.68 mm, respectively) and treatment success (added effect: 33.6%).

Conversely, the remaining 9 studies showed no differences in clinical and radiographic parameters comparing two or more decontamination approaches (Albaker et al., 2018; de Waal et al., 2013, 2015; Hallström et al., 2017; Isehmed et al., 2016; Lasserre et al., 2020; Papadopoulos et al., 2015; Schlee et al., 2019; Schwarz et al., 2011).

Considering the studies with longer follow-up periods, the 6–12 months results were confirmed in the majority of the reports (Romeo et al., 2007; Schwarz et al., 2012, 2013, 2017; Isehmed et al., 2018), except for the study by Carcuac et al. (2017) in which the short-term benefits of systemic antibiotics at implants with modified surfaces were not sustained over 3 years.

3.5.2 | Meta-analyses

Meta-analyses comparing clinical and radiographic outcomes of different decontamination protocols were only possible for the added effect of systemic antimicrobials in access flap surgery and for the added effect of Er:Yag laser in combined surgery.

The use of systemic antimicrobials in access surgery resulted in no added effect in terms of PPD reduction (2 trials; WMD = 0.63 mm; 95% CI: -0.69/2.55; $I^2 = 84\%$), whereas it provided added effect in terms of treatment success (2 trials; RR = 1.84; 95% CI: 1.17/2.91; $I^2 = 0\%$) (Figure 3a,b).

The use of the Er:Yag laser in combined surgery resulted in no added effect in terms of PPD reduction (2 trials; WMD = -0.24 mm; 95% CI: -1.10/0.63; $I^2 = 60\%$) (Figure 3c).

The “summary of findings” tables for the effect of systemic antimicrobials and Er:Yag laser are reported in Figure S2 and S3, respectively.

4 | DISCUSSION

The findings from the present systematic review indicate that several decontamination protocols resulted in improved clinical and radiographic outcomes after surgical treatment of peri-implantitis. Owing to paucity of available trials, evidence regarding the superiority of some protocols over the others is mainly based on single RCTs. However, meta-analyses indicated a short-term added effect of systemic antimicrobials on treatment success but not on PPD reduction in access surgery, and a lack of added effect of Er:Yag laser on PPD reduction in combined surgery.

Ideally, implant surface decontamination should remove biofilm without causing surface damage not to render surfaces more conducive to bacterial colonization (Louropoulou et al., 2014). For such a purpose, mechanical, chemical and physical decontamination protocols have been tested in RCTs so far.

With regards to mechanical methods, *in vitro* studies indicated that non-metal curettes and rubber cups were minimally traumatic but ineffective to clean contaminated titanium surfaces; while ultrasonic scalers, metal curettes and rotating titanium brushes were effective particularly on modified titanium surfaces (John et al., 2014).

The air abrasive system resulted effective in all types of implant surfaces, with glycine/erythritol powder causing less alterations than sodium bicarbonate (Cochis et al., 2013; Pranno et al., 2020). The included trials indicate a better clinical performance of titanium brushes over plastic curettes, ultrasonic scalers and air powder devices (Toma et al., 2019). Moreover, there is controversy regarding the role of implantoplasty. No clinical benefit of implantoplasty on implant survival rate was observed in a recent long-term retrospective study (Ravidà et al., 2020). The removal of threads and of the superficial portion of the implant surface may enhance intra-surgical decontamination, and the long-term recontamination may be potentially prevented, thanks to the reduced plaque-retention of the smoothed implant surface. The trials included in the present systematic review reported an added benefit of implantoplasty in terms of PPD reduction in both access flap and resective surgery (Lasserre et al., 2020; Romeo et al., 2004).

Coupling mechanical instruments with chemical/physical agents may improve the overall cleaning ability as the chemical agent may reach niches inaccessible mechanically (Carcuac et al., 2017). Different concentrations of CHX showed limited benefits both in *in vitro* models and in clinical trials, and a cytotoxic effect was reported (Schwarz et al., 2005). The available trials confirm that CHX alone and in combination with CPC does not provide clinical benefits when employed in access flap or resective surgery. EMD application on fixture surfaces switched subgingival microbiota to Gram+ aerobic populations (Isehmed et al., 2016), and this ecological shift was linked in the only identified RCT with an increase in bone levels as compared with non-EMD controls. Moreover, in an exploratory analysis performed at 5 years, EMD application was associated with higher survival rates (Isehmed et al., 2018). Being peri-implantitis an infection-driven disease, local and systemic antimicrobials have been proposed as an adjunctive method of decontamination. The only identified RCT on the added effect of local intra-surgical application of minocycline (and its subsequent sub-mucosal application 1- and 3-months after) resulted in a statistically significant bone gain compared to placebo, and in the highest rate of treatment success among all the included trials (66.7%) (Cha et al., 2019). With regards to their systemic administration, the meta-analyses reported in the present systematic review identified a greater probability of treatment success when they are employed (RR = 1.84). Nevertheless, a careful risk/benefit evaluation should be performed before systemic administration of systemic antimicrobials, in light of the potential onset of side effects and of the growing issue of antibiotic resistances (WHO, 2020).

Many different types of lasers have been proposed to decontaminate the implant surface and enhance the healing potential during treatment of peri-implantitis, with the Er:YAG laser having demonstrated a high degree of bactericidal effect at low-power intensity (Lin et al., 2018). However, the meta-analysis performed in the present systematic review showed that the adjunctive application of Er:Yag laser in combined surgery had no significant effect on PPD reduction over other mechanical and chemical decontaminating agents. The effect of photodynamic therapy was only evaluated in

one RCT, showing also no additional benefits on clinical and radiographic results (Albaker et al., 2018; Chambrone et al., 2018).

To the best of the authors' knowledge, this systematic review is the first providing a comprehensive qualitative and quantitative analysis on the decontamination protocols tested in RCTs as part of the surgical treatment of peri-implantitis. Limitations worth mentioning are mainly related to the available trials, since most of the evidence on decontamination protocols is based on single RCTs, and that half of the included trials were not considered at low risk of bias. Moreover, several factors such as implant surface characteristics (Berglundh, Wennström, & Lindhe, 2018) configurations of peri-implant defect (Schwarz et al., 2010) frequency and quality of supportive therapy (Heitz-Mayfield et al., 2018; Rocuzzo et al., 2018) can affect the results of surgical treatment of peri-implantitis, but their impact could not be analyzed due to the high heterogeneity among the included trials.

5 | CONCLUSIONS

The present systematic review highlighted the absence of consistent evidence of superiority of any decontamination protocol over the others in the surgical treatment of peri-implantitis. Meta-analyses indicated an added benefit of systemic antimicrobials, but not of Er:Yag laser, in increasing treatment success rates. However, this effect was not present on PPD reduction, and it was only detected in the short-term.

Well-designed RCTs are needed to definitely identify the most effective mechanical decontamination method and to verify the added effect of adjunctive chemical/physical measures. When multiple implants per patient are included, the use of mixed models analysis should be implemented. In addition to implant survival and to PPD, BoP/SoP and MBL changes, researchers are encouraged to analyze also additional relevant outcomes which have been sparsely reported so far, including treatment success, soft tissue level changes, PROMs and rates of re-interventions and adverse events (e.g., implant fracture, emphysemas, side effects, etc.). Researchers are also suggested to employ a common comparator (i.e., titanium brushes) in order to favor future analyses of the literature, until clear evidence of superiority of a different protocol is identified.

AUTHOR CONTRIBUTIONS

GB, FC, NB and MA made substantial contributions to study conception. GB, FC, NB, FR and MA contributed to the study design. GB, FC and NB searched and collected the data. GB, FC, MR and FR performed data analysis and interpretation. GB, MR, FC, GMM and FR prepared the first draft of the manuscript. All authors have read, revised critically, and approved the final manuscript.

ACKNOWLEDGMENTS

The authors kindly thank S. Isler, J. Lasserre, E. Romeo, S. Toma, and Y. de Waal, for providing more information and estimates about their studies, regardless of the final decision to include or

not include them in the present systematic review. The authors received no financial support and declare no potential conflicts of interest with respect to the authorship and/or publication of this article. Open Access Funding provided by Università degli Studi di Torino within the CRUI-CARE Agreement. Open Access Funding provided by Università degli Studi di Torino within the CRUI-CARE Agreement.

FUNDING INFORMATION

This study was funded solely by the institutions of the authors.

CONFLICT OF INTEREST

The authors declare no conflict of interest.

DATA AVAILABILITY STATEMENT

All data generated or analyzed during this study are included in this published article [and its supplementary information files].

ETHICAL APPROVAL

Ethics approval was not required for this systematic review.

ORCID

Giacomo Baima  <https://orcid.org/0000-0002-9395-4967>

Filippo Citterio  <https://orcid.org/0000-0002-4513-7479>

Mario Romandini  <https://orcid.org/0000-0001-5646-083X>

Federica Romano  <https://orcid.org/0000-0002-5172-299X>

Nurcan Buduneli  <https://orcid.org/0000-0002-1590-5801>

Mario Aimetti  <https://orcid.org/0000-0003-0657-0787>

REFERENCES

- Albaker, A. M., ArRejaie, A. S., Alrabiah, M., Al-Aali, K. A., Mokeem, S., Alasqah, M. N., Vohra, F., & Abduljabbar, T. (2018). Effect of antimicrobial photodynamic therapy in open flap debridement in the treatment of peri-implantitis: A randomized controlled trial. *Photodiagnosis and Photodynamic Therapy*, 23, 71–74. <https://doi.org/10.1016/j.pdpdt.2018.05.003>
- Berglundh, T., Armitage, G., Araujo, M. G., Avila-Ortiz, G., Blanco, J., Camargo, P. M., Chen, S., Cochran, D., Derks, J., Figuero, E., Hämmerle, C. H. F., Heitz-Mayfield, L. J. A., Huynh-Ba, G., Iacono, V., Koo, K.-T., Lambert, F., McCauley, L., Quirynen, M., Renvert, S., ... Zitzmann, N. (2018). Peri-implant diseases and conditions: Consensus report of workgroup 4 of the 2017 world workshop on the classification of periodontal and peri-implant diseases and conditions. *Journal of Clinical Periodontology*, 45(S20), S286–S291. <https://doi.org/10.1111/jcpe.12957>
- Berglundh, T., Wennström, J. L., & Lindhe, J. (2018). Long-term outcome of surgical treatment of peri-implantitis: A 2-11-year retrospective study. *Clinical Oral Implants Research*, 29(4), 404–410. <https://doi.org/10.1111/clr.13138>
- Campbell, M. K., Piaggio, G., Elbourne, D. R., Altman, D. G., & CONSORT Group. (2012). Consort 2010 statement: Extension to cluster randomised trials. *BMJ (Clinical Research Ed.)*, 345, e5661. <https://doi.org/10.1136/bmj.e5661>
- Carcuac, O., Derks, J., Abrahamsson, I., Wennström, J. L., Petzold, M., & Berglundh, T. (2017). Surgical treatment of peri-implantitis: 3-Year results from a randomized controlled clinical trial. *Journal of Clinical Periodontology*, 44(12), 1294–1303. <https://doi.org/10.1111/jcpe.12813>

- Carcuac, O., Derks, J., Charalampakis, G., Abrahamsson, I., Wennström, J., & Berglundh, T. (2016). Adjunctive systemic and local antimicrobial therapy in the surgical treatment of peri-implantitis: A randomized controlled clinical trial. *Journal of Dental Research*, 95(1), 50–57. <https://doi.org/10.1177/0022034515601961>
- Cha, J. K., Lee, J. S., & Kim, C. S. (2019). Surgical therapy of peri-Implantitis with local minocycline: A 6-month randomized controlled clinical trial. *Journal of Dental Research*, 98(3), 288–295. <https://doi.org/10.1177/0022034518818479>
- Chambrone, L., Wang, H.-L., & Romanos, G. E. (2018). Antimicrobial photodynamic therapy for the treatment of periodontitis and peri-implantitis: An American Academy of periodontology best evidence review. *Journal of Periodontology*, 89(7), 783–803. <https://doi.org/10.1902/jop.2017.170172>
- Cochis, A., Fini, M., Carrassi, A., Migliario, M., Visai, L., & Rimondini, L. (2013). Effect of air polishing with glycine powder on titanium abutment surfaces. *Clinical Oral Implants Research*, 24(8), 904–909. <https://doi.org/10.1111/j.1600-0501.2012.02490.x>
- de Waal, Y. C. M., Raghoobar, G. M., Huddleston Slater, J. J. R., Meijer, H. J. A., Winkel, E. G., & van Winkelhoff, A. J. (2013). Implant decontamination during surgical peri-implantitis treatment: A randomized, double-blind, placebo-controlled trial. *Journal of Clinical Periodontology*, 40(2), 186–195. <https://doi.org/10.1111/jcpe.12034>
- de Waal, Y. C. M., Raghoobar, G. M., Meijer, H. J. A., Winkel, E. G., & van Winkelhoff, A. J. (2015). Implant decontamination with 2% chlorhexidine during surgical peri-implantitis treatment: A randomized, double-blind, controlled trial. *Clinical Oral Implants Research*, 26(9), 1015–1023. <https://doi.org/10.1111/clr.12419>
- Derks, J., & Tomasi, C. (2015). Peri-implant health and disease. A systematic review of current epidemiology. *Journal of Clinical Periodontology*, 42 Suppl 16, S158–S171. <https://doi.org/10.1111/jcpe.12334>
- Guyatt, G., Oxman, A. D., Akl, E. A., Kunz, R., Vist, G., Brozek, J., Norris, S., Falck-Ytter, Y., Glasziou, P., DeBeer, H., Jaeschke, R., Rind, D., Meerpohl, J., Dahm, P., & Schünemann, H. J. (2011). GRADE guidelines: 1. Introduction-GRADE evidence profiles and summary of findings tables. *Journal of Clinical Epidemiology*, 64(4), 383–394. <https://doi.org/10.1016/j.jclinepi.2010.04.026>
- Hallström, H., Persson, G. R., Lindgren, S., & Renvert, S. (2017). Open flap debridement of peri-implantitis with or without adjunctive systemic antibiotics: A randomized clinical trial. *Journal of Clinical Periodontology*, 44(12), 1285–1293. <https://doi.org/10.1111/jcpe.12805>
- Heitz-Mayfield, L. J. A., & Mombelli, A. (2014). The therapy of peri-implantitis: A systematic review. *The International Journal of Oral & Maxillofacial Implants*, 29(Suppl), 325–345. <https://doi.org/10.11607/jomi.2014suppl.g5.3>
- Heitz-Mayfield, L. J. A., Salvi, G. E., Mombelli, A., Loup, P.-J., Heitz, F., Kruger, E., & Lang, N. P. (2018). Supportive peri-implant therapy following anti-infective surgical peri-implantitis treatment: 5-year survival and success. *Clinical Oral Implants Research*, 29(1), 1–6. <https://doi.org/10.1111/clr.12910>
- Ished, C., Holmlund, A., Renvert, S., Svenson, B., Johansson, I., & Lundberg, P. (2016). Effectiveness of enamel matrix derivative on the clinical and microbiological outcomes following surgical regenerative treatment of peri-implantitis. A randomized controlled trial. *Journal of Clinical Periodontology*, 43(10), 863–873. <https://doi.org/10.1111/jcpe.12583>
- Ished, C., Svenson, B., Lundberg, P., & Holmlund, A. (2018). Surgical treatment of peri-implantitis using enamel matrix derivative, an RCT: 3- And 5-year follow-up. *Journal of Clinical Periodontology*, 45(6), 744–753. <https://doi.org/10.1111/jcpe.12894>
- Isler, S. C., Unsal, B., Soysal, F., Ozcan, G., Peker, E., & Karaca, I. R. (2018). The effects of ozone therapy as an adjunct to the surgical treatment of peri-implantitis. *Journal of Periodontal & Implant Science*, 48(3), 136–151. <https://doi.org/10.5051/jpis.2018.48.3.136>
- John, G., Becker, J., & Schwarz, F. (2014). Rotating titanium brush for plaque removal from rough titanium surfaces—An in vitro study. *Clinical Oral Implants Research*, 25(7), 838–842. <https://doi.org/10.1111/clr.12147>
- Khoury, F., Keeve, P. L., Ramanauskaitė, A., Schwarz, F., Koo, K., Sculean, A., & Romanos, G. (2019). Surgical treatment of peri-implantitis – Consensus report of working group 4. *International Dental Journal*, 69(S2), 18–22. <https://doi.org/10.1111/idj.12505>
- Klinge, B., Gustafsson, A., & Berglundh, T. (2002). A systematic review of the effect of anti-infective therapy in the treatment of peri-implantitis. *Journal of Clinical Periodontology*, 29(Suppl 3), 213–225; discussion 232–233. <https://doi.org/10.1034/j.1600-051x.29.s3.13.x>
- Koo, K.-T., Khoury, F., Keeve, P. L., Schwarz, F., Ramanauskaitė, A., Sculean, A., & Romanos, G. (2019). Implant surface decontamination by surgical treatment of periimplantitis: A literature review. *Implant Dentistry*, 28(2), 173–176. <https://doi.org/10.1097/ID.0000000000000840>
- Lasserre, J., Brex, M., & Toma, S. (2020). Implantoplasty versus glycine air abrasion for the surgical treatment of peri-implantitis: A randomized clinical trial. *The International Journal of Oral & Maxillofacial Implants*, 35(1), 197–206. <https://doi.org/10.11607/jomi.6677>
- Lin, G. H., Suárez López Del Amo, F., & Wang, H. L. (2018). Laser therapy for treatment of peri-implant mucositis and peri-implantitis: An American Academy of periodontology best evidence review. *Journal of Periodontology*, 89(7), 766–782. <https://doi.org/10.1902/jop.2017.160483>
- Lindhe, J., Meyle, J., & on behalf of Group D of the European Workshop on Periodontology. (2008). Peri-implant diseases: Consensus report of the sixth European workshop on periodontology. *Journal of Clinical Periodontology*, 35, 282–285. <https://doi.org/10.1111/j.1600-051X.2008.01283.x>
- Louropoulou, A., Slot, D. E., & Weijden, F. (2014). The effects of mechanical instruments on contaminated titanium dental implant surfaces: A systematic review. *Clinical Oral Implants Research*, 25(10), 1149–1160. <https://doi.org/10.1111/clr.12224>
- Meyle, J. (2012). Mechanical, chemical and laser treatments of the implant surface in the presence of marginal bone loss around implants. *European Journal of Oral Implantology*, 5(Suppl), S71–S81.
- Moher, D., Liberati, A., Tetzlaff, J., Altman, D. G., & PRISMA Group. (2009). Preferred reporting items for systematic reviews and meta-analyses: The PRISMA statement. *Journal of Clinical Epidemiology*, 62(10), 1006–1012. <https://doi.org/10.1016/j.jclinepi.2009.06.005>
- Papadopoulos, C. A., Vouros, I., Menexes, G., & Konstantinidis, A. (2015). The utilization of a diode laser in the surgical treatment of peri-implantitis. A randomized clinical trial. *Clinical Oral Investigations*, 19(8), 1851–1860. <https://doi.org/10.1007/s00784-014-1397-9>
- Pranno, N., Cristalli, M. P., Mengoni, F., Sauzullo, I., Annibali, S., Polimeni, A., & La Monaca, G. (2020). Comparison of the effects of air-powder abrasion, chemical decontamination, or their combination in open-flap surface decontamination of implants failed for peri-implantitis: An ex vivo study. *Clinical Oral Investigations*, 25, 2667–2676. <https://doi.org/10.1007/s00784-020-03578-w>
- Ramanauskaitė, A., Fretwurst, T., & Schwarz, F. (2021). Efficacy of alternative or adjunctive measures to conventional non-surgical and surgical treatment of peri-implant mucositis and peri-implantitis: A systematic review and meta-analysis. *International Journal of Implant Dentistry*, 7(1), 112. <https://doi.org/10.1186/s40729-021-00388-x>
- Ravidà, A., Siqueira, R., Saleh, I., Saleh, M. H. A., Giannobile, A., & Wang, H. L. (2020). Lack of clinical benefit of Implantoplasty to improve implant survival rate. *Journal of Dental Research*, 99(12), 1348–1355. <https://doi.org/10.1177/0022034520944158>

- Roccuzzo, M., Layton, D. M., Roccuzzo, A., & Heitz-Mayfield, L. J. (2018). Clinical outcomes of peri-implantitis treatment and supportive care: A systematic review. *Clinical Oral Implants Research*, 29(S16), 331–350. <https://doi.org/10.1111/clr.13287>
- Romandini, M., Berglundh, J., Derks, J., Sanz, M., & Berglundh, T. (2021). Diagnosis of peri-implantitis in the absence of baseline data: A diagnostic accuracy study. *Clinical Oral Implants Research*, 32, 297–313. <https://doi.org/10.1111/clr.13700>
- Romandini, M., Lima, C., Pedrinaci, I., Araoz, A., Soldini, M. C., & Sanz, M. (2021). Prevalence and risk/protective indicators of peri-implant diseases: A university-representative cross-sectional study. *Clinical Oral Implants Research*, 32(1), 112–122. <https://doi.org/10.1111/clr.13684>
- Romeo, E., Ghisolfi, M., Murgolo, N., Chiapasco, M., Lops, D., & Vogel, G. (2004). Therapy of peri-implantitis with resective surgery: A 3-year clinical trial on rough screw-shaped oral implants. Part I: Clinical outcome. *Clinical Oral Implants Research*, 16(1), 9–18. <https://doi.org/10.1111/j.1600-0501.2004.01084.x>
- Romeo, E., Lops, D., Chiapasco, M., Ghisolfi, M., & Vogel, G. (2007). Therapy of peri-implantitis with resective surgery. A 3-year clinical trial on rough screw-shaped oral implants. Part II: Radiographic outcome. *Clinical Oral Implants Research*, 18(2), 179–187. <https://doi.org/10.1111/j.1600-0501.2006.01318.x>
- Sanz, M., Herrera, D., Kerschull, M., Chapple, I., Jepsen, S., Berglundh, T., Sculean, A., & Tonetti, M. S. (2020). Treatment of stage I–III periodontitis—the EFP S3 level clinical practice guideline. *Journal of Clinical Periodontology*, 47(S22), 4–60. <https://doi.org/10.1111/jcpe.13290>
- Schlee, R., Rathe, F., Brodbeck, R., Ratka, C., Weigl, P., & Zipprich, H. (2019). Treatment of peri-implantitis—Electrolytic cleaning versus mechanical and electrolytic cleaning—A randomized controlled clinical trial—Six-month results. *Journal of Clinical Medicine*, 8(11), 1909. <https://doi.org/10.3390/jcm8111909>
- Schwarz, F., Hegewald, A., John, G., Sahm, N., & Becker, J. (2013). Four-year follow-up of combined surgical therapy of advanced peri-implantitis evaluating two methods of surface decontamination. *Journal of Clinical Periodontology*, 40(10), 962–967. <https://doi.org/10.1111/jcpe.12143>
- Schwarz, F., John, G., Mainusch, S., Sahm, N., & Becker, J. (2012). Combined surgical therapy of peri-implantitis evaluating two methods of surface debridement and decontamination. A two-year clinical follow up report. *Journal of Clinical Periodontology*, 39(8), 789–797. <https://doi.org/10.1111/j.1600-051X.2012.01867.x>
- Schwarz, F., John, G., Schmucker, A., Sahm, N., & Becker, J. (2017). Combined surgical therapy of advanced peri-implantitis evaluating two methods of surface decontamination: A 7-year follow-up observation. *Journal of Clinical Periodontology*, 44(3), 337–342. <https://doi.org/10.1111/jcpe.12648>
- Schwarz, F., Sahm, N., Iglhaut, G., & Becker, J. (2011). Impact of the method of surface debridement and decontamination on the clinical outcome following combined surgical therapy of peri-implantitis: A randomized controlled clinical study: Surgical therapy of peri-implantitis. *Journal of Clinical Periodontology*, 38(3), 276–284. <https://doi.org/10.1111/j.1600-051X.2010.01690.x>
- Schwarz, F., Sahm, N., Schwarz, K., & Becker, J. (2010). Impact of defect configuration on the clinical outcome following surgical regenerative therapy of peri-implantitis. *Journal of Clinical Periodontology*, 37(5), 449–455. <https://doi.org/10.1111/j.1600-051X.2010.01540.x>
- Schwarz, F., Schmucker, A., & Becker, J. (2015). Efficacy of alternative or adjunctive measures to conventional treatment of peri-implant mucositis and peri-implantitis: A systematic review and meta-analysis. *International Journal of Implant Dentistry*, 1(1), 22. <https://doi.org/10.1186/s40729-015-0023-1>
- Schwarz, F., Sculean, A., Romanos, G., Herten, M., Horn, N., Scherbaum, W., & Becker, J. (2005). Influence of different treatment approaches on the removal of early plaque biofilms and the viability of SAOS2 osteoblasts grown on titanium implants. *Clinical Oral Investigations*, 9(2), 111–117. <https://doi.org/10.1007/s00784-005-0305-8>
- Sterne, J. A. C., Savović, J., Page, M. J., Elbers, R. G., Blencowe, N. S., Boutron, I., Cates, C. J., Cheng, H.-Y., Corbett, M. S., Eldridge, S. M., Emberson, J. R., Hernán, M. A., Hopewell, S., Hróbjartsson, A., Junqueira, D. R., Jüni, P., Kirkham, J. J., Lasserson, T., Li, T., ... Higgins, J. P. T. (2019). RoB 2: A revised tool for assessing risk of bias in randomised trials. *BMJ (Clinical Research Ed.)*, 366, l4898. <https://doi.org/10.1136/bmj.l4898>
- Tapia, B., Valles, C., Ribeiro-Amaral, T., Mor, C., Herrera, D., Sanz, M., & Nart, J. (2019). The adjunctive effect of a titanium brush in implant surface decontamination at peri-implantitis surgical regenerative interventions: A randomized controlled clinical trial. *Journal of Clinical Periodontology*, 46(5), 586–596. <https://doi.org/10.1111/jcpe.13095>
- Toma, S., Brex, M. C., & Lasserre, J. F. (2019). Clinical evaluation of three surgical modalities in the treatment of peri-implantitis: A randomized controlled clinical trial. *Journal of Clinical Medicine*, 8(7), 966. <https://doi.org/10.3390/jcm8070966>
- Wang, C., Ashnagar, S., Gianfilippo, R. D., Arnett, M., Kinney, J., & Wang, H. (2020). Laser-assisted regenerative surgical therapy for peri-implantitis: A randomized controlled clinical trial. *Journal of Periodontology*, 92(3), 378–388. <https://doi.org/10.1002/JPER.20-0040>
- World Health Organization (2020). WHO library cataloguing-in-publication data. Antimicrobial resistance: Global report on surveillance. Retrieved from <https://www.who.int/news-room/fact-sheet/s/detail/antibiotic-resistance>. Accessed 11 July 2022.

SUPPORTING INFORMATION

Additional supporting information can be found online in the Supporting Information section at the end of this article.

How to cite this article: Baima, G., Citterio, F., Romandini, M., Romano, F., Mariani, G. M., Buduneli, N., & Aimetti, M. (2022). Surface decontamination protocols for surgical treatment of peri-implantitis: A systematic review with meta-analysis. *Clinical Oral Implants Research*, 00, 1–18. <https://doi.org/10.1111/clr.13992>