

Complete intersections on Veronese surfaces

Original

Complete intersections on Veronese surfaces / Canino, Stefano; Carlini, Enrico. - In: JOURNAL OF ALGEBRA. - ISSN 0021-8693. - 622:(2023), pp. 328-350. [10.1016/j.jalgebra.2023.01.025]

Availability:

This version is available at: 11583/2979587 since: 2023-07-07T12:42:00Z

Publisher:

Elsevier

Published

DOI:10.1016/j.jalgebra.2023.01.025

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)

Complete intersections on Veronese surfaces

Stefano Canino^{*1} and Enrico Carlini ^{†1}

¹DISMA-Department of Mathematics, Politenico di Torino, Corso Duca degli Abruzzi 24,
10129, Torino

June 14, 2022

Abstract

In this paper we describe all possible reduced complete intersection sets of points on Veronese surfaces. We formulate a conjecture for the general case of complete intersection subvarieties of any dimension and we prove it in the case of the quadratic Veronese threefold. Our main tool is an effective characterization of all possible Hilbert functions of reduced subvarieties of Veronese surfaces.

1 Introduction

Complete intersection subvarieties are both a classical both a modern topic of study in Algebraic Geometry.

In [Eul53] Euler asked when a sets of points in the plane is the intersection of two curves, that is, using the modern terminology, when a set of points in the plane is a complete intersection. In the same period, Cramer asked similar questions so that this type of questions is presently known as the *Cramer-Euler problem*. Euler solution in the case of nine points in the plane gave rise to what are now known as the Cayley-Bacharach Theorems, see [EGH96].

Complete intersections and their algebraic counterpart, regular sequences, play a central role in Commutative Algebra and in Algebraic geometry. Consider, for example, the well known Hartshorne conjecture, stated for the first time by Hartshorne in [Har70] and still open, which is probably one of the most studied problems regarding complete intersection. More recently, complete intersections have shown to have unexpected applications. For example, in [BBOV21] and [BO21], the strength and the slice rank of polynomials are studied using complete intersections. See also [CN17] for an application in proving the existence of special families of vector bundles on quartic surfaces of \mathbb{P}^3 . For a a more exhaustive overview on complete intersections we advise to see [GS84].

In this paper, we consider a generalization of the Cramer-Euler problem: characterize the possible complete intersections lying on a Veronese surface $V_{2,d}$, and more generally on a Veronese variety. We recall that, given non-negative integers n and d , a Veronese variety of \mathbb{P}^N is projectively

^{*}stefano.canino@polito.it

[†]enrico.carlini@polito.it

equivalent to $V_{n,d} \subseteq \mathbb{P}^N$, that is the image of \mathbb{P}^n via the usual Veronese map:

$$\begin{aligned} \nu_{n,d} : \quad \mathbb{P}^n &\rightarrow \mathbb{P}^N \\ [x_0, \dots, x_n] &\mapsto [x_0^d, x_0^{d-1}x_1, \dots, x_n^d] \end{aligned}$$

where $N = \binom{n+d}{d} - 1$. Notice that for $d = 1$ the Veronese surface $V_{2,1}$ is the plane \mathbb{P}^2 , so that our problem in this special case is exactly the Cramer-Euler problem. In Theorem 4.5 we completely solve the problem showing that for $d > 2$ the only reduced complete intersections of \mathbb{P}^N lying on $V_{2,d}$ are finite sets of either one or two points. While, for the Veronese surface $V_{2,2} \subseteq \mathbb{P}^5$, one also has plane conics and their intersections with suitable hypersurfaces. Moreover, in Theorem 4.3 we show that, except for the case $d = 2$, the only complete intersections lying on rational normal curves $V_{1,d}$ are the trivial ones, that is one single point or the set of two points. The case $V_{1,2}$, that is of a plane conic, is different. In fact, by cutting with any properly chosen curve, one will produce a complete intersection set of points.

Inspired by these evidences we formulate Conjecture 5.2: the only reduced complete intersections of $V_{n,d}$, $d \geq 3$, are finite sets of either one or two points while for $d = 2$ one also has plane conics and their intersections with suitable hypersurfaces. We also checked the validity of the conjecture for $V_{3,2}$, see Proposition 5.1.

In order to prove our main result, Theorem 4.5, we characterize the possible Hilbert functions of reduced subvarieties of Veronese varieties in Theorem 2.16. In other words, we characterize all possible Hilbert functions of radical ideals in the Veronese ring $\mathbb{C}[x_0, \dots, x_N]/\mathcal{I}(V_{n,d})$, where $\mathcal{I}(V_{n,d})$ is the defining ideal of $V_{n,d}$. Beyond their application to the proof of our theorem, Hilbert functions play a central role in Commutative Algebra and in Algebraic Geometry, for example see [Sta78], [Mig07], and [BGM94]. Indeed, in recent times Hilbert functions have also been used as tools in other fields. For example, in the study of Waring rank, that is the tensor rank for symmetric tensors, in the paper [CCG12]. Another example is the study of Strassen's Conjecture, a crucial conjecture in complexity theory, now proved to be false in general [Shi19], but still open in the relevant case of symmetric tensors, see [CCC⁺18] and [CMM18]. As a last example, we also mention the study of the identifiability of tensors, which plays a crucial role in Algebraic Statistic, see [ABC18],[AC20] and [AGMO18].

In this paper, we first characterize the Hilbert functions of reduced subvarieties of $V_{n,d}$. Thus, we generalise 0-sequences and differentiable 0-sequences introduced in [GMR83], see Definition 2.10 and Definition 2.11. Successively, we give more effective characterization for the case of the rational normal curves $V_{1,d}$ in Theorem 2.9, so recovering a classical result, and for the case of the surfaces $V_{2,d}$ in Theorem 3.4. In [GMP11, Theorem 4.5] a similar characterization is given in the case of subschemes of $V_{n,d}$, that is in the case of any ideal in a Veronese ring. However, we consider these characterization not to be enough effective for our purposes. In fact, given a candidate Hilbert function, one has to solve a kind of interpolation problem to decide whether the given function is the Hilbert function of a reduced subvariety, or subscheme, of $V_{n,d}$, see Remark 3.7.

More precisely, the paper is structured as follows. In §2, we recall some basic notions needed in the paper and we study the relationship between $\mathcal{I}(\nu_{n,d}^{-1}(\mathbb{X}))$ and $\mathcal{I}(\mathbb{X})$, where $\mathbb{X} \subseteq V_{n,d}$ is a reduced subvariety of $V_{n,d}$. Using this we characterize the Hilbert functions of subvarieties lying on $V_{n,d}$ and introduce d -sequences and differentiable d -sequences. Moreover, we use these results to study the Hilbert functions of divisors of $V_{n,d}$ and the Hilbert functions of finite sets of reduced points lying on a rational normal curve $V_{1,d}$, recovering the classical results. In §3, we apply Theorem 2.6 to the special case of Veronese surfaces getting a more explicit characterization for the Hilbert

functions of reduced subvarieties of Veronese surfaces in Theorem 3.4. In §4, we use the previously introduced tools to prove our main theorem characterizing all possible complete intersections lying on Veronese surfaces. In §5, we study the reduced complete intersection lying on $V_{3,2}$ and state Conjecture 5.2 about reduced complete intersections lying on Veronese varieties.

Acknowledgments: the authors wish to thank Ciro Ciliberto, Luca Chiantini, Alessandro Gimigliano, and Giorgio Ottaviani for some useful discussions. Moreover, the authors acknowledge the kindness of Marvi Catalisano and Juan Migliore in parting from a similar project which was never completed. The authors are members of GNSAGA of INDAM and were supported by MIUR grant Dipartimenti di Eccellenza 2018-2022 (E11G18000350001).

2 Basic notions and first results

In this section we introduce the needed basic notions and some preliminary results including a complete characterization of the possible Hilbert functions of reduced subvarieties of a Veronese variety. Let \mathbb{K} be an algebraic closed field with $\text{char } \mathbb{K} = 0$ and $R = \mathbb{K}[x_0, \dots, x_n]$ the coordinate ring of \mathbb{P}^n . If d is a positive integer, we set $N = \binom{n+d}{d} - 1$ and we denote the coordinate ring of \mathbb{P}^N by $S = \mathbb{K}[y_0, \dots, y_N]$.

If M is a graded module, we denote by M_p the p -th graded component of M , so that $M = \bigoplus_{p=0}^{\infty} M_p$. In particular, since an ideal $I \subseteq R$ is an R -module, we use the same notation I_p for the p -th graded component of I .

For a graded module M , we indicate its Hilbert function by $H_M(t) = \dim M_t$ and its first difference function by $\Delta H_M(t) = H_M(t) - H_M(t-1)$, setting $H_M(-1) = 0$. If $\mathbb{X} \subseteq \mathbb{P}^n$ is a reduced projective scheme we denote by $\mathcal{I}(\mathbb{X})$ its associated ideal and by $H_{\mathbb{X}}(t)$, its Hilbert function, that is the Hilbert function of the module $R/\mathcal{I}(\mathbb{X})$.

Notation 2.1 We will denote by $\alpha_0, \dots, \alpha_N$ the multi-indices of $\{0, 1, \dots, d\}^{n+1}$ such that $|\alpha_j| = d$ ordered by the usual lexicographic order. Moreover, if $\alpha_i = (\alpha_{i0}, \dots, \alpha_{in})$, we set

$$\underline{x}^{\alpha_i} = x_0^{\alpha_{i0}} \dots x_n^{\alpha_{in}}.$$

Definition 2.2 For each $n, d \in \mathbb{N}_{>0}$ we define the (n, d) -Veronese embedding

$$\begin{aligned} \nu_{n,d} : \quad \mathbb{P}^n &\rightarrow \mathbb{P}^N \\ [x_0, \dots, x_n] &\mapsto [\underline{x}^{\alpha_0}, \dots, \underline{x}^{\alpha_N}] \end{aligned}$$

and the (n, d) -Veronese variety $V_{n,d} := \nu_{n,d}(\mathbb{P}^n)$.

The following lemma will be very useful.

Lemma 2.3 Let us consider the graded morphism

$$\begin{aligned} \varphi_d : \quad S &\rightarrow R \\ a &\mapsto a \\ y_i &\mapsto \underline{x}^{\alpha_i} \end{aligned}$$

for all $a \in \mathbb{K}$, and for $i \in \{0, \dots, N\}$. Then the following hold:

1. $\ker \varphi_d = \mathcal{I}(V_{n,d})$.

2. $\text{Im } \varphi_d = \bigoplus_{\ell=0}^{\infty} R_{\ell d}$ and, in particular, $\varphi_d(S_t) = R_{td}$.
3. If $\mathbb{X} \subseteq V_{n,d}$, then $(\mathcal{I}(\nu_{n,d}^{-1}(\mathbb{X})))_{td} = \varphi_d(\mathcal{I}(\mathbb{X})_t)$. In particular

$$\varphi_d(\mathcal{I}(\mathbb{X})) = \bigoplus_{s=0}^{\infty} (\mathcal{I}(\nu_{n,d}^{-1}(\mathbb{X})))_{sd}.$$

4. If \mathbb{X} is a subvariety of $V_{n,d}$ and we set $\mathbb{Y} = \nu_{n,d}^{-1}(\mathbb{X})$, then

$$H_{\mathbb{X}}(t) = H_{\mathbb{Y}}(td) \quad \forall t \geq 0.$$

Proof The proof of (1) is trivial. To prove (2) it is enough to note that any monomial of degree td can be written as product of monomials of degree d . The proof of (3) is a straightforward check of a double inclusion. The proof of (4) follows from the chain of graded isomorphisms

$$S/\mathcal{I}(\mathbb{X}) \cong \frac{S/\mathcal{I}(V_{n,d})}{\mathcal{I}(\mathbb{X})/\mathcal{I}(V_{n,d})} \cong \frac{\psi_d(S/\mathcal{I}(V_{n,d}))}{\psi_d(\mathcal{I}(\mathbb{X})/\mathcal{I}(V_{n,d}))} = \frac{\bigoplus_{s=0}^{\infty} R_{sd}}{\varphi_d(\mathcal{I}(\mathbb{X}))}$$

where $\psi_d : S/\mathcal{I}(V_{n,d}) \rightarrow \bigoplus_{s=0}^{\infty} R_{sd}$ is the canonical isomorphism induced by φ_d . ■

Remark 2.4 Since φ_d is a ring homomorphism, $\varphi_d(\mathcal{I}(\mathbb{X}))$ is an ideal of $\text{Im } \varphi_d = \bigoplus_{s=0}^{\infty} R_{sd}$, but it is not an ideal of R . Nevertheless, one has

$$(\varphi_d(\mathcal{I}(\mathbb{X}))R)_{td} = (\varphi_d(\mathcal{I}(\mathbb{X})))_{td}.$$

Remark 2.5 In the notations of Lemma 2.3, if we choose $\mathbb{X} = V_{n,d}$ then we have $\mathbb{Y} = \mathbb{P}^n$ and thus we get that the Hilbert Function of $V_{n,d}$ is $H_{V_{n,d}}(t) = \binom{n+td}{n}$.

The following theorem is an immediate consequence of Lemma 2.3.

Theorem 2.6 *Let $h(t) : \mathbb{N} \rightarrow \mathbb{N}$ be the Hilbert function of a projective variety in \mathbb{P}^N . Then there exists $\mathbb{X} \subseteq V_{n,d} \subseteq \mathbb{P}^N$ such that $H_{\mathbb{X}}(t) = h(t)$ if and only there exists $k(t) : \mathbb{N} \rightarrow \mathbb{N}$ Hilbert function of a projective variety in \mathbb{P}^n such that $h(t) = k(dt)$.*

Remark 2.7 We note that, if a subvariety \mathbb{Y} has Hilbert function satisfying the conditions of Theorem 2.6, this does not mean that \mathbb{Y} lies on a Veronese variety. The theorem only guarantees that there exists *some* subvariety \mathbb{X} of a Veronese variety having the same Hilbert function of \mathbb{Y} . Consider, for example, seven generic points in \mathbb{P}^3 . By genericity they do not lie on a $V_{1,d}$, that is they do not lie on rational normal curve, but their Hilbert function satisfies the hypothesis of the theorem.

In the case of divisors of $V_{n,d}$ we can be more explicit.

Proposition 2.8 *If \mathbb{X} is a divisor of $V_{n,d}$, let $\deg \mathbb{X} = de$, then*

$$H_{\mathbb{X}}(t) = \begin{cases} \binom{n+dt}{n}, & \text{if } t \leq \lfloor \frac{e-1}{d} \rfloor \\ \binom{n+dt}{n} - \binom{n+dt-e}{n}, & \text{if } t \geq \lfloor \frac{e-1}{d} \rfloor + 1 \end{cases}$$

Proof Since \mathbb{X} is a divisor and $\deg \mathbb{X} = de$, there exists a (unique) hypersurface $\mathbb{Y} : F = 0$ of degree e in \mathbb{P}^n such that $\nu_{n,d}(\mathbb{Y}) = \mathbb{X}$. For each $t \in \mathbb{N}$ we have the following short exact sequence

$$0 \longrightarrow R(-e)_t \xrightarrow{F} R_t \xrightarrow{\pi} (R/(F))_t \longrightarrow 0$$

and as a consequence, we get $H_{\mathbb{Y}}(t) = \dim(R/(F))_t = \dim R_t - \dim R(-e)_t$ and this ends the proof.

Remark 2.9 As special case of divisor we can consider $\mathbb{X} \subseteq V_{1,d} \subseteq \mathbb{P}^d$ a finite set of s reduced points on the rational normal curve of degree d . Using Proposition 2.8 we get the well known result:

$$H_{\mathbb{X}}(t) = \begin{cases} dt + 1, & \text{if } t \leq \lfloor \frac{s-2}{d} \rfloor \\ s, & \text{if } t \geq \lfloor \frac{s-2}{d} \rfloor + 1 \end{cases}.$$

We conclude recalling the following definitions from [GMR83].

Definition 2.10 A sequence of non-negative integers $(c_t)_{t \in \mathbb{N}}$ is called a *0-sequence* if $c_0 = 1$ and $c_{t+1} \leq c^{(t)}$ for all $t \geq 1$.

Definition 2.11 Let $(b_t)_{t \in \mathbb{N}}$ be a 0-sequence. Then $(b_t)_{t \in \mathbb{N}}$ is *differentiable* if the difference sequence $(c_t)_{t \in \mathbb{N}}$, $c_t = b_t - b_{t-1}$ is again a 0-sequence (where $b_{-1} = 0$).

In [GMR83] the authors completely characterized the Hilbert functions of reduced varieties with the following theorem.

Theorem 2.12 ([GMR83], Theorem 3.3) *Let \mathbb{K} be an infinite field and let $(b_t)_{t \in \mathbb{N}}$ be a differentiable 0-sequence with $b_1 = n + 1$. There is a radical ideal I in $\mathbb{K}[x_0, \dots, x_n]$ such that $(b_t)_{t \in \mathbb{N}}$ is the Hilbert function of $\mathbb{K}[x_0, \dots, x_n]/I$.*

The complete characterization of Theorem 2.12 can be rephrased: if S is a sequence of non-negative integers, then there exists a reduced \mathbb{K} -algebra A such that S is the Hilbert function of A if and only if S is a differentiable 0-sequence.

Theorem 2.6 suggests us to extend Definition 2.10 and 2.11 as follows.

Definition 2.13 A 0-sequence $(b_t)_{t \in \mathbb{N}}$ is called *d-sequence* if there exists a 0-sequence $(c_t)_{t \in \mathbb{N}}$ such that $b_t = c_{(d+1)t}$.

Definition 2.14 A 0-sequence $(b_t)_{t \in \mathbb{N}}$ is called *differentiable d-sequence* if there exists a differentiable 0-sequence $(c_t)_{t \in \mathbb{N}}$ such that $b_t = c_{(d+1)t}$.

Remark 2.15 We note that that a differentiable d -sequence is necessarily a differentiable 0-sequence.

We can now rephrase Theorem 2.6 as follows:

Theorem 2.16 *Let $(h_t)_{t \in \mathbb{N}}$ be a sequence of non-negative integers such that $h_0 = 1$ and $h_1 = N + 1$. There exists a projective variety $\mathbb{X} \subseteq V_{n,d} \subseteq \mathbb{P}^N$ such that $H_{\mathbb{X}}(t) = h_t$ if and only if $(h_t)_{t \in \mathbb{N}}$ is a differentiable $(d - 1)$ -sequence.*

It is natural to ask for an effective characterization of d -sequences similar to the one of Theorem 2.12. The question does not have an answer in general yet, nevertheless one can give an answer in a special case using our results. In the case $(d - 1)$ -sequences with $h_1 = \binom{d+2}{2}$, such a characterization can be easily produced using Theorem 3.4 in the next section.

3 Hilbert functions of points on Veronese surfaces

In this section we focus our attention on the case of Veronese surfaces $V_{2,d}$. In particular, we give an effective characterization of the Hilbert function of any reduced subvariety of $V_{2,d}$ in Theorem 3.4.

Notation 3.1 Given $d, t, s \in \mathbb{N}$ such that $s \geq d^2t + \frac{d(d+3)}{2}$ we define the following two functions:

$$\mu_1(d, t, s) := d^2t + \frac{d(d+3)}{2} - s$$

$$\mu_2(d, t, s) := \begin{cases} \left\lfloor \frac{2d(t+1)+3-\sqrt{1+8\mu_1(d,t,s)}}{2} \right\rfloor, & \text{if } 1 \leq \mu_1(d, t, s) \leq \binom{d+1}{2} \\ dt - n, & \text{if } \binom{d+1}{2} + dn < \mu_1(d, t, s) \leq \binom{d+1}{2} + d(n+1), 0 \leq n \leq dt \end{cases}$$

We begin with a technical result.

Lemma 3.2 *Let $d, t \in \mathbb{N}$. Consider a function $h : \{1, 2, \dots, d\} \rightarrow \mathbb{N}$ such that there exists $i_0 \in \{1, 2, \dots, d\}$ with the properties*

1. $h(i) = dt + i + 1$ for each $1 \leq i \leq i_0 - 1$;
2. $h(i) \geq h(i+1)$ for each $i_0 \leq i \leq d - 1$.

Then $h(d) \leq \mu_2(d, t, \sum_{i=1}^d h(i))$ and moreover the inequality is sharp.

Proof See Appendix A. ■

In the following proposition we characterise Hilbert functions of reduced points in $\mathbb{P}^{\frac{d(d+3)}{2}}$ which arise from Hilbert functions of reduced points in \mathbb{P}^2 by sampling with steps of length d .

Proposition 3.3 *Let us consider a finite set of reduced points $\mathbb{X} \subseteq \mathbb{P}^{\frac{d(d+3)}{2}}$ and set*

$$t_1 = \max \{t \mid H_{\mathbb{X}}(t) = H_{V_{2,d}}(t)\}, \quad t_2 = \min \{t \mid H_{\mathbb{X}}(t) = |\mathbb{X}|\}.$$

Then $H_{\mathbb{X}}(t)$ is a $(d-1)$ -sequence if and only if the following conditions hold

i.

$$\mu_2(d, t_1, \Delta H_{\mathbb{X}}(t_1 + 1)) \geq \left\lceil \frac{\Delta H_{\mathbb{X}}(t_1 + 2)}{d} \right\rceil;$$

ii. For all $t_1 + 2 \leq t \leq t_2 - 1$

$$\left\lfloor \frac{\Delta H_{\mathbb{X}}(t)}{d} \right\rfloor \geq \left\lfloor \frac{\Delta H_{\mathbb{X}}(t+1)}{d} \right\rfloor.$$

Proof First, we assume that there exists $\mathbb{Y} \subseteq \mathbb{P}^2$ such that $H_{\mathbb{Y}}(dt) = H_{\mathbb{X}}(t)$. Let us set

$$H_{\mathbb{X}}(t) = h_t, \quad \Delta H_{\mathbb{X}}(t) = \Delta h_t,$$

and

$$H_{\mathbb{Y}}(t) = k_t, \quad \Delta H_{\mathbb{Y}}(t) = \Delta k_t.$$

Since $(k_t)_{t \in \mathbb{N}}$ is the Hilbert function of a finite set of reduced points of \mathbb{P}^2 , then (see [KK98], Proposition 1.1) there exists $t' \in \mathbb{N}$ such that the following conditions hold

- $k_t = \binom{2+t}{t}$ for all $t \leq t'$ and $k_t < \binom{2+t}{t}$ for all $t > t'$, and thus $\Delta k_t = t + 1$ for all $t \leq t'$;

- $\Delta k_t \geq \Delta k_{t+1}$ for all $t > t'$, and Δk_t is eventually equal to 0.

Note that

$$\Delta h_t = h_t - h_{t-1} = k_{dt} - k_{d(t-1)} = k_{dt} + \sum_{i=1}^{d-1} (k_{dt-i} - k_{dt-i}) - k_{dt-d} = \sum_{i=0}^{d-1} (k_{dt-i} - k_{dt-(i+1)}) = \sum_{i=0}^{d-1} \Delta k_{dt-i}$$

and thus

$$\Delta h_{t+1} = \sum_{i=0}^{d-1} \Delta k_{d(t+1)-i} = \sum_{i=1}^d \Delta k_{dt+i}. \quad (1)$$

Thus, Remark 2.5 yields that

$$\sum_{i=1}^d \Delta k_{dt+i} = \Delta h_{t+1} = \binom{2+d(t+1)}{2} - \binom{2+dt}{2} = d^2t + \frac{d(d+3)}{2} \text{ for all } t \leq t_1 - 1.$$

Since $\Delta k_t \leq t+1$ for all t and

$$\sum_{i=1}^d (dt+i+1) = d^2t + \frac{d(d+3)}{2}$$

it follows that

$$\sum_{i=1}^d \Delta k_{dt+i} = d^2t + \frac{d(d+3)}{2} \text{ for all } t \leq t_1 - 1$$

if and only if

$$\Delta k_t = t+1 \text{ for all } t \leq dt_1$$

and thus $t' \geq dt_1$. Moreover, since $h_t < \binom{2+dt}{2}$ for $t > t_1$, the same argument shows that

$$\sum_{i=1}^d \Delta k_{dt+i} < d^2t + \frac{d(d+3)}{2} \text{ for all } t \geq t_1$$

and for $t = t_1$ we have

$$\sum_{i=1}^d \Delta k_{dt_1+i} < d^2t_1 + \frac{d(d+3)}{2}.$$

As a consequence, there exists a minimum $i_0 \in \{1, 2, \dots, d\}$ such that $\Delta k_{dt_1+i_0} < dt_1 + i_0 + 1$ and therefore $t' \leq dt_1 + d - 1 = d(t_1 + 1) - 1$. It follows that

$$\Delta k_{dt_1+i} = dt_1 + i + 1 \quad (2)$$

for $1 \leq i \leq i_0 - 1$ and

$$\Delta k_{dt_1+i_0} \geq \Delta k_{dt_1+i_0+1} \geq \dots \geq \Delta k_{dt_1+i_0+a} = 0$$

for some $a \in \mathbb{N}$. Moreover

$$\Delta k_{dt_1+i_0} \geq \Delta k_{dt_1+i_0+1} \geq \dots \geq \Delta k_{dt_1+d} \quad (3)$$

and

$$\Delta k_{dt+1} \geq \Delta k_{dt+2} \geq \cdots \geq \underbrace{\Delta k_{dt+d+1}}_{=\Delta k_{d(t+1)+1}} \quad (4)$$

for each $t \geq t_1 + 1$. Now, for a fixed $t \geq t_1 + 1$, by (4) we have

$$(d-1)\Delta k_{dt+1} \geq \sum_{i=2}^d \Delta k_{dt+i},$$

and

$$(d-1)\Delta k_{dt+d} \leq \sum_{i=1}^{d-1} \Delta k_{dt+i}.$$

Using formula (1), we obtain

$$\Delta k_{dt+1} = \Delta h_{t+1} - \sum_{i=2}^d \Delta k_{dt+i} \geq \Delta h_{t+1} - (d-1)\Delta k_{dt+1}$$

and hence

$$\Delta k_{dt+1} \geq \frac{\Delta h_{t+1}}{d}$$

and similarly

$$\Delta k_{dt+d} = \Delta h_{t+1} - \sum_{i=1}^{d-1} \Delta k_{dt+i} \leq \Delta h_{t+1} - (d-1)\Delta k_{dt+d}$$

and hence

$$\Delta k_{dt+d} \leq \frac{\Delta h_{t+1}}{d}.$$

Moreover, since $(h_t)_{t \in \mathbb{N}}$ and $(k_t)_{t \in \mathbb{N}}$ are integer valued, we have that

$$\Delta k_{dt+1} \geq \left\lceil \frac{\Delta h_{t+1}}{d} \right\rceil, \quad \Delta k_{dt+d} \leq \left\lfloor \frac{\Delta h_{t+1}}{d} \right\rfloor$$

for each $t \geq t_1 + 1$. By (2) and (3) it follows that the function

$$\begin{array}{ccc} \Delta k_{dt_1+i} : & \{1, 2, \dots, d\} & \rightarrow \mathbb{N} \\ & i & \mapsto \Delta k_{dt_1+i} \end{array}$$

satisfies the hypothesis of Lemma 3.2. Hence, we get that

$$\mu_2(d, t_1 \Delta h_{t_1+1}) \geq \Delta k_{dt_1+d} \geq \Delta k_{d(t_1+1)+1} \geq \left\lceil \frac{\Delta h_{t_1+2}}{d} \right\rceil$$

thus proving condition (i.). Finally, using (4) we get that

$$\Delta k_{dt+d} \geq \Delta k_{dt+d+1} = \Delta k_{d(t+1)+1}$$

for each $t \geq t_1 + 1$, and hence

$$\left\lceil \frac{\Delta h_{t+1}}{d} \right\rceil \geq \left\lceil \frac{\Delta h_{t+2}}{d} \right\rceil$$

for each $t \geq t_1 + 1$. We notice that this inequality is always verified for $t \geq t_2$ since

$$\left\lceil \frac{\Delta h_{t+1}}{d} \right\rceil = 0$$

for each $t \geq t_2$. Hence also condition (ii.) is proved.

Now we assume that conditions (i.) and (ii.) hold and we prove that there exists $\mathbb{Y} \subseteq \mathbb{P}^2$ such that $H_{\mathbb{Y}}(dt) = h_t$. Since

$$k_t = \sum_{i=0}^t \Delta k_t,$$

we can construct the Hilbert function $(k_t)_{t \in \mathbb{N}}$ by its first difference Δk_t . For each t let e_t be the unique integer such that

$$\Delta h_t \equiv e_t \pmod{d}, \quad 0 \leq e_t \leq d-1.$$

We define Δk_t as follows:

- if $0 \leq t \leq dt_1$ we set $\Delta k_t = t + 1$;
- if $dt_1 + 1 \leq t \leq d(t_1 + 1)$ we construct Δk_t according to Lemma 3.2;
- if $t \geq d(t_1 + 1) + 1$ we set

$$\begin{aligned} \Delta k_{dt+i} &= \left\lceil \frac{\Delta h_{t+1}}{d} \right\rceil, \quad t \geq t_1 + 1, 1 \leq i \leq e_{t+1} \\ \Delta k_{dt+i} &= \left\lfloor \frac{\Delta h_{t+1}}{d} \right\rfloor, \quad t \geq t_1 + 1, e_{t+1} + 1 \leq i \leq d. \end{aligned}$$

Under our assumptions, this choice guarantees that Δk_t is the first difference function of a set of reduced points in \mathbb{P}^2 (see [KK98] Proposition 1.1). Moreover, for $t \leq t_1 - 1$, we have

$$\sum_{i=1}^d \Delta k_{dt+i} = \sum_{i=1}^d (dt + i + 1) = d^2 t + \frac{d(d+3)}{2} = \Delta h_{t+1},$$

and, for $t = t_1$ we have by construction

$$\sum_{i=1}^d \Delta k_{dt_1+1} = \Delta h_{t_1+1}$$

while for $t \geq t_1 + 1$ we have

$$\begin{aligned} \sum_{i=1}^d \Delta k_{dt+i} &= \sum_{i=1}^{e_{t+1}} \Delta k_{dt+i} + \sum_{i=e_{t+1}+1}^d \Delta k_{dt+i} = e_{t+1} \left\lceil \frac{\Delta h_{t+1}}{d} \right\rceil + (d - e_{t+1}) \left\lfloor \frac{\Delta h_{t+1}}{d} \right\rfloor = \\ &= e_{t+1} \left(\frac{\Delta h_{t+1} - e_{t+1}}{d} + 1 \right) + (d - e_{t+1}) \left(\frac{\Delta h_{t+1} - e_{t+1}}{d} \right) = \Delta h_{t+1}. \end{aligned}$$

Hence we have

$$k_{dt} = \sum_{i=0}^{dt} \Delta k_t = \Delta k_0 + \sum_{i=0}^{t-1} \left(\sum_{j=1}^d \Delta k_{di+j} \right) = \Delta h_0 + \sum_{i=0}^{t-1} \Delta h_{i+1} = \sum_{i=0}^t \Delta h_t = h_t$$

for all t . This concludes the proof. ■

We now give our effective characterization of the Hilbert functions of reduced sets of points on Veronese surfaces.

Theorem 3.4 *Let $(h_t)_{t \in \mathbb{N}}$ be the Hilbert function of a finite set of m reduced points in $\mathbb{P}^{\frac{d(d+3)}{2}}$ and set*

$$t_1 = \max \{t \mid h(t) = H_{V_{2,d}}(t)\} \quad t_2 = \min \{t \mid h(t) = m\}.$$

Then there exists $\mathbb{X} \subseteq V_{2,d} \subseteq \mathbb{P}^N$, $|\mathbb{X}| = m$ such that $H_{\mathbb{X}}(t) = h_t$ if and only if the following conditions hold

i.

$$\mu_2(d, t_1, \Delta h_{t_1+1}) \geq \left\lceil \frac{\Delta h_{t_1+2}}{d} \right\rceil;$$

ii. For all $t_1 + 2 \leq t \leq t_2 - 1$

$$\left\lfloor \frac{\Delta h_t}{d} \right\rfloor \geq \left\lfloor \frac{\Delta h_{t+1}}{d} \right\rfloor.$$

Proof It follows from Theorem 2.6 and Proposition 3.3. ■

Let us see now two explicit instances of use of Theorem 3.4 in determining whether a given Hilbert function can be realised as the Hilbert function of a subvariety of a Veronese surface.

Example 3.5 Let us consider the sequence $(h_t)_{t \in \mathbb{N}}$ defined as follows

t	0	1	2	3	4	5	6	7	8	9	10	11
h_t	1	36	120	253	435	666	946	1256	1531	1744	1956	2022

and $h_t = 2022$ for $t \geq 12$. It is easy to check, using Theorem 3.3. in [GMR83], that this is the Hilbert function of a set of 2022 reduced points in \mathbb{P}^{35} . We ask whether there exists $\mathbb{X} \subseteq V_{2,7} \subseteq \mathbb{P}^{35}$ such that $H_{\mathbb{X}}(t) = h_t$ for all $t \geq 0$. To answer we use Theorem 3.4. First we determine t_1 and t_2 . Since the Hilbert function of $V_{2,7}$ is $H_{V_{2,7}}(t) = \binom{2+7t}{2}$, we have that

t	0	1	2	3	4	5	6	7	8	9	10	11	12
$H_{V_{2,7}}$	1	36	120	253	435	666	946	1275	1653	2080	2556	3081	3655

so that $t_1 = 6$ and $t_2 = 11$. To determine $\mu_1(7, 6, \Delta h_{t_1+1})$ we compute Δh_{t_1+1} . We have that

t	0	1	2	3	4	5	6	7	8	9	10	11	12
Δh_t	1	35	84	133	182	231	280	310	275	213	212	66	0

and thus $\mu_1(7, 6, 310) = 7^2 \cdot 6 + \frac{7(7+3)}{2} - 310 = 19$. Finally, since $19 \leq \binom{7+1}{2} = 28$, we get

$$\mu_2(7, 6, 310) = \left\lfloor \frac{2 \cdot 7(6+1) + 3 - \sqrt{1 + 8 \cdot 19}}{2} \right\rfloor = 44.$$

To check conditions (i.) and (ii.) of Theorem 3.4, we compute $\lfloor \frac{\Delta h_t}{7} \rfloor$ and $\lceil \frac{\Delta h_t}{7} \rceil$ obtaining the following table

t	0	1	2	3	4	5	6	7	8	9	10	11	12
$\lceil \frac{\Delta h_t}{7} \rceil$	1	5	12	19	26	33	40	45	40	31	31	10	0
$\lfloor \frac{\Delta h_t}{7} \rfloor$	0	5	12	19	26	33	40	44	39	30	30	9	0

Since $\mu_2(7, 6, 310) = 44$ and $\lceil \frac{\Delta h_8}{7} \rceil = 40$ condition (i.) is satisfied. However condition (ii.) is not satisfied for $t = 9$ and hence such an \mathbb{X} does not exist.

Example 3.6 Now, we consider the sequence $(h_t)_{t \in \mathbb{N}}$ defined as follows

t	0	1	2	3	4	5	6	7	8	9	10	11
h_t	1	36	120	253	435	666	946	1256	1531	1744	1915	2022

and $h_t = 2022$ for $t \geq 12$; note that this function coincides with the one of the previous example, but for $t = 10$. We ask whether there exists $\mathbb{X} \subseteq V_{2,7} \subseteq \mathbb{P}^{35}$ such that $H_{\mathbb{X}}(t) = h_t$ for all $t \geq 0$. As in the previous example we have $t_1 = 6$ and $t_2 = 11$. Moreover, we get

t	0	1	2	3	4	5	6	7	8	9	10	11	12
Δh_t	1	35	84	133	182	231	280	310	275	213	201	107	0
$\lceil \frac{\Delta h_t}{7} \rceil$	1	5	12	19	26	33	40	45	40	31	29	16	0
$\lfloor \frac{\Delta h_t}{7} \rfloor$	0	5	12	19	26	33	40	44	39	30	28	15	0

and thus $\mu_1(7, 6, 310) = 19$ and $\mu_2(7, 6, 310) = 44$. Thus, condition (i.) is satisfied and condition (ii.) is satisfied for $t = 8, 9, 10$. Hence such an \mathbb{X} exists.

Remark 3.7 One could deal with both the previous examples without using Theorem 3.4, but just using Macaulay's inequalities for Hilbert functions. However, this requires a trial and error approach. In fact, since $d = 7$ one should try to fill step by step six gaps between h_i and h_{i+1} satisfying Macaulay's inequalities. Thus, for each possible choice at each step, one should compute the appropriate binomial expansion. Moreover, one should check also that the first difference function of the constructed Hilbert function is still an Hilbert function: doing this will require a very large number of computations.

4 Complete intersections

In this section we focus on the study of complete intersection varieties of \mathbb{P}^N which lie on some Veronese variety $V_{n,d}$ for $n = 1$ and $n = 2$; the case $n = 3$ and $d = 2$ is treated in Proposition 5.1. For generalities on complete intersections we refer to [Eis95]. To compare with similar existence and non-existence results for complete intersection on hypersurfaces and their applications we refer to [CCG08, ST22].

Definition 4.1 Let \mathbb{X} be a projective variety, let $\mathcal{I}(\mathbb{X}) = (f_1, f_2, \dots, f_r)$ be its defining ideal, and let $a_i = \deg f_i$. We say that \mathbb{X} is a *complete intersection of type* (a_1, \dots, a_r) if $\text{codim } \mathbb{X} = r$.

Proposition 4.2 *If $\mathbb{X} \subseteq V_{n,d} \subseteq \mathbb{P}^N$ is a reduced complete intersection of type (a_1, \dots, a_r) , with $a_1 \leq \dots \leq a_r$, then $a_1 = 1$. Moreover, either $r = N$ and $a_1 = \dots = a_N = 1$ or $a_i = 2$ for some i .*

Proof Since $\mathbb{X} \subseteq V_{n,d}$, we have that $I(\mathbb{X}) \supseteq I(V_{n,d})$. Recall that $I(V_{n,d})$ is generated by the 2×2 minors of a matrix of linear forms, see [Puc98]. Thus, either $a_1 = 1$ or $a_1 = 2$. Assume by contradiction that $a_1 = 2$, that is $a_i > 1$ for all i , and let $I(\mathbb{X}) = (f_1, \dots, f_r)$. Thus it is possible to assume that $I(V_{n,d}) = (f_1, \dots, f_q)$ for some q . Thus any syzygy of the generators of $I(V_{n,d})$ gives a syzygy of the generators of $I(\mathbb{X})$. Thus, the determinantal representation of $I(V_{n,d})$ yields that the generators of $I(\mathbb{X})$ have a syzygy of linear forms. Hence a contradiction since f_1, \dots, f_r are a regular sequence and their syzygies, given by the Koszul complex, only contain elements of degree at least $a_1 = 2$. In conclusion, $a_1 = 1$. To complete the proof, we let p be the largest index such that $a_p = 1$. Assume by contradiction that $p < r$ and $a_{p+1} > 2$. Thus, $I(V_{n,d}) \subseteq (f_1, \dots, f_p)$ and the latter is the ideal of a linear space Λ of codimension p such that $\Lambda \subseteq V_{n,d}$. Hence a contradiction since no Veronese variety contains a positive dimensional linear spaces. In conclusion, either $p = r = N$ or $p < r$ and $a_{p+1} = 2$. ■

As a straightforward application of this proposition we get the following result.

Theorem 4.3 *Let $\mathbb{X} \subseteq V_{1,d} \subseteq \mathbb{P}^d$ be a reduced complete intersection with $d \geq 3$. Then either \mathbb{X} is a point or \mathbb{X} is a set of two points.*

Proof Since $V_{1,d}$ is not a complete intersection and $\dim V_{1,d} = 1$, it must be $\dim \mathbb{X} = 0$. Since \mathbb{X} is a complete intersection, it follows from Proposition 4.2 that \mathbb{X} is contained in the intersection of a hyperplane and $V_{1,d}$. In particular we have $|\mathbb{X}| \leq d$ and using Remark 2.9 we get the Hilbert function of \mathbb{X} :

$$\begin{array}{c|cccc} t & 0 & 1 & 2 & \dots \\ \hline H_{\mathbb{X}}(t) & 1 & |\mathbb{X}| & |\mathbb{X}| & \dots \\ \Delta H_{\mathbb{X}}(t) & 1 & |\mathbb{X}| - 1 & 0 & \dots \end{array}$$

Finally, since \mathbb{X} is a complete intersection, $\Delta H_{\mathbb{X}}(t)$ is the Hilbert function of an artinian Gorenstein ideal. Thus it is symmetric and hence either $\Delta H_{\mathbb{X}}(1) = 0$ or $\Delta H_{\mathbb{X}}(1) = 1$, that is $|\mathbb{X}| = 1$ or $|\mathbb{X}| = 2$. ■

Remark 4.4 We note that the proof of Theorem 4.3 also shows that any Gorenstein reduced zero-dimensional subscheme of $V_{1,d}$ must be degenerate.

We can now finally describe reduced complete intersection subvarieties of Veronese surfaces.

Theorem 4.5 *If $\mathbb{X} \subseteq V_{2,d} \subseteq \mathbb{P}^N$ is a reduced complete intersection of type (a_1, \dots, a_r) , with $a_1 \leq \dots \leq a_r$ then one of the following holds:*

1. $(d, r, (a_1, a_2, \dots, a_r)) = (2, 4, (1, 1, 1, 2))$, that is \mathbb{X} is a conic lying on $V_{2,2}$;
2. $(d, r, (a_1, a_2, \dots, a_r)) = (2, 5, (1, 1, 1, 2, a_5))$, any $a_5 \in \mathbb{N}$, that is \mathbb{X} is a set of $2a_5$ complete intersection points of a conic lying on $V_{2,2}$ and a hypersurface of degree a_5 ;
3. $(d, r, (a_1, a_2, \dots, a_r)) = (d, N, (1, 1, \dots, 1))$ for any $d \geq 2$, that is \mathbb{X} is a reduced point;

4. $(d, r, (a_1, a_2, \dots, a_r)) = (d, N, (1, 1, \dots, 1, 2))$ for any $d \geq 2$, that is \mathbb{X} is a set of two reduced points.

Proof Let $\mathcal{I}(\mathbb{X}) = (f_1, \dots, f_r)$ be the ideal of \mathbb{X} . We can suppose without loss of generality that $a_1 \leq a_2 \leq \dots \leq a_r$. Moreover, we let p be the number of a_i equal to 1 and q be the number of a_i equal to 2, that is $a_1 = a_2 = \dots = a_p = 1$, $a_{p+1} = a_{p+2} = \dots = a_{p+q} = 2$. By Proposition 4.2 we have that $p \geq 1$. If $p = N$ we are trivially in case (3) thus from now on we suppose $p < N$ and thus, using again Proposition 4.2, we also have $q \geq 1$. Since $\mathbb{X} \subseteq V_{2,d}$, $\Delta H_{\mathbb{X}}(t)$ must satisfy conditions (i.) and (ii.) of Theorem 3.4. We now use condition (i.) to rule out some cases. We start by computing $\Delta H_{\mathbb{X}}(1)$ and $\Delta H_{\mathbb{X}}(2)$. We have that

$$H_{\mathbb{X}}(1) = \dim \mathbb{K}[y_0, \dots, y_N] - \dim(f_1, \dots, f_r)_1 = N - p + 1$$

$$\Delta H_{\mathbb{X}}(1) = H_{\mathbb{X}}(1) - H_{\mathbb{X}}(0) = N - p.$$

Now, let us consider the map

$$\begin{array}{ccc} \mathbb{K}[y_0, \dots, y_N] & \rightarrow & \mathbb{K}[y_0, \dots, y_N]/(f_1, \dots, f_p) \cong \mathbb{K}[y_0, \dots, y_{N-p}] \\ f & \mapsto & \tilde{f} \end{array}.$$

Since (f_1, f_2, \dots, f_r) is a regular sequence we have that

$$\begin{aligned} H_{\mathbb{X}}(2) &= \dim (\mathbb{K}[y_0, \dots, y_N]/(f_1, \dots, f_r))_2 = \dim (\mathbb{K}[y_0, \dots, y_{N-p}]/(\tilde{f}_{p+1}, \dots, \tilde{f}_r))_2 \\ &= \dim \mathbb{K}[y_0, \dots, y_{N-p}]_2 - q = \binom{N-p+2}{2} - q \end{aligned}$$

and hence

$$\Delta H_{\mathbb{X}}(2) = \binom{N-p+2}{2} - q - (N-p+1) = \frac{(N-p+1)(N-p)}{2} - q.$$

Note that $H_{\mathbb{X}}(1) = N - p + 1 < N + 1 = H_{V_{2,d}}(1)$, thus $t_1 = 0$. Since $t_1 = 0$ we have that

$$\mu_2(d, 0, N-p) = \left\lfloor \frac{2d+3-\sqrt{1+8p}}{2} \right\rfloor,$$

so that by (i.) we get

$$\left\lfloor \frac{2d+3-\sqrt{1+8p}}{2} \right\rfloor \geq \left\lceil \frac{(N-p+1)(N-p)}{2d} - \frac{q}{d} \right\rceil. \quad (5)$$

Since $q \leq N-p$, if (p, d, q) is a solution of (5), then (p, d) is a solution of

$$\frac{2d+3-\sqrt{1+8p}}{2} \geq \frac{(N-p)(N-p-1)}{2d}$$

and setting $\alpha = N-p$ the previous inequality yields

$$\underbrace{2d+3-\sqrt{(2d+3)^2-8(\alpha+1)}}_{f(\alpha)} \geq \underbrace{\frac{\alpha(\alpha-1)}{d}}_{g(\alpha)}. \quad (6)$$

A standard calculus argument on $f(\alpha)$ and $g(\alpha)$ shows that if (α, d) is a solution of (6) then $\alpha \leq 3$. As a consequence, if (p, d, q) is a solution of (5) then $p \geq N-3$, thus one has to look for solutions (p, d, q) only for $p = N-3, N-2, N-1$. Since $r \leq N$, one can easily check that the solutions of (5) are the following:

- $(p, d, q) = (N - 1, d, 1)$ for each $d \geq 2$.
In this case \mathbb{X} is a complete intersection of type (a_1, \dots, a_N) with $a_1 = \dots = a_{N-1} = 1$ and $a_N = 2$, that is \mathbb{X} is a set of two reduced points and we are in case 4.
- $(p, d, q) = (3, 2, 1)$.
In this case $N = 5$ and we know that $a_1 = a_2 = a_3 = 1$ and $a_4 = 2$, thus we distinguish two subcases:
 - If $r = 4$ then \mathbb{X} is a complete intersection of type $(1, 1, 1, 2)$, that is \mathbb{X} is a conic lying on $V_{2,2}$ and we are in case 1.
 - If $r = 5$ then \mathbb{X} is a complete intersection of type $(1, 1, 1, 2, a_5)$, that is \mathbb{X} is the intersection of a conic lying on $V_{2,2}$ with a hypersurface of degree a_5 and we are in case 2.
- $(p, d, q) = (3, 2, 2)$.
In this case \mathbb{X} is a complete intersection of type $(1, 1, 1, 2, 2)$, thus it this is a special case of 2.
- $(p, d, q) = (6, 3, 3)$.
In this case \mathbb{X} is a complete intersection of type $(1, 1, 1, 1, 1, 1, 2, 2, 2)$. We want to show that such \mathbb{X} can not lie on $V_{2,3}$. Let us suppose by contradiction that $\mathbb{X} \subseteq V_{2,3}$ and set $\mathbb{Y} = \nu_{2,3}^{-1}(\mathbb{X})$. From Lemma 2.3 it follows that

$$H_{\mathbb{Y}}(3) = H_{\mathbb{X}}(1) = N - p + 1 = 9 - 6 + 1 = 4.$$

Thus, the only way to complete the gaps of $H_{\mathbb{Y}}$ is

$$\begin{array}{c|cccccccccc} t & 0 & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \dots \\ \hline H_{\mathbb{Y}}(t) & 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & 8 & \dots \end{array}$$

and this shows that \mathbb{Y} lies on a line. As a consequence, \mathbb{X} lies on a rational normal curve. Hence, since \mathbb{X} is a complete intersection and $|\mathbb{X}| = 8$, this is a contradiction by Theorem 4.3.

The result is now proved. ■

5 More results and open problems

In this section, we show some more results about complete intersections on Veronese varieties. A complete characterization of the Hilbert functions of subvarieties of Veronese varieties of dimension larger than two seems, presently, to be out of reach. Nevertheless, we can deal with the case of the threefold $V_{3,2}$. This case leads us to formulate Conjecture 5.2.

Proposition 5.1 *Let $\mathbb{X} \subseteq V_{3,2} \subseteq \mathbb{P}^9$ be a reduced subvariety. Then \mathbb{X} is a complete intersection of type (a_1, \dots, a_r) , with $a_1 \leq \dots \leq a_r$ if and only if \mathbb{X} is one of the following*

- $r = 9, a_1 = \dots = a_9 = 1$, that is \mathbb{X} is a reduced point;
- $r = 9, a_1 = \dots = a_8 = 1, a_9 = 2$, that is \mathbb{X} is a set of two reduced points;
- $r = 9, a_1 = \dots = a_7 = 1, a_8 = 2, a_9 = b$, any $b \geq 2$, that is $\mathbb{X} = \mathcal{C} \cap H_b$ for $\mathcal{C} \subseteq V_{3,2}$ a conic and H_b a degree b hypersurface;
- $r = 8, a_1 = \dots = a_7 = 1, a_8 = 2$, that is \mathbb{X} is a conic.

Proof First we consider the case $\dim \mathbb{X} = 0$. Let $\mathbb{X} \subseteq V_{3,2}$ be a reduced complete intersection of type (a_1, \dots, a_9) with $a_1 \leq a_2 \leq \dots \leq a_9$. Also let $p, q \in \mathbb{N}$ such that $a_1 = \dots = a_p = 1$ and $a_{p+1} = \dots = a_{p+q+1} = 2$. By Proposition 4.2 either $p = 9$ or $p \geq 1$ and $q \geq 1$. If $p = 9$ then \mathbb{X} is just a reduced point, thus from now on we suppose $p, q \geq 1$. By the same argument used in the proof of Theorem 4.5 we get

$$H_{\mathbb{X}}(1) = 10 - p, \quad H_{\mathbb{X}}(2) = \binom{11-p}{2} - q = \frac{p^2 - 21p + 110}{2} - q.$$

Since $\mathbb{X} \subseteq V_{3,2}$, by Lemma 2.3 there exists $\mathbb{Y} = \nu_{3,2}^{-1}(\mathbb{X}) \subseteq \mathbb{P}^3$ such that $H_{\mathbb{X}}(t) = H_{\mathbb{Y}}(td)$ for all $t \geq 0$. In particular, we have that

$$H_{\mathbb{Y}}(2) = 10 - p, \quad H_{\mathbb{Y}}(4) = \frac{p^2 - 21p + 110}{2} - q.$$

Now fix $1 \leq p \leq 8$. By Macaulay's theorem for Hilbert functions (see [Sta78], Theorem 2.2) it follows that if the 3-binomial expansion of $H_{\mathbb{Y}}(2)$ is

$$H_{\mathbb{Y}}(2) = \binom{m_2}{2} + \binom{m_1}{1}$$

where $m_2 > m_1$, then

$$H_{\mathbb{Y}}(4) \leq \binom{m_2 + 2}{4} + \binom{m_1 + 2}{3} = M(p)$$

On the other hand, since $1 \leq q \leq 9 - p$, we have that

$$H_{\mathbb{Y}}(4) \geq \frac{p^2 - 21p + 110}{2} - (9 - p) = m(p).$$

Thus, if $M(p) - m(p) < 0$, then \mathbb{X} does not exist. Computing we get the following table

p	$H_{\mathbb{Y}}(2)$	$M(p)$	$m(p)$	$M(p) - m(p)$
1	9	25	37	-12
2	8	19	29	-10
3	7	16	22	-6
4	6	15	16	-1
5	5	9	11	-2
6	4	6	7	-1
7	3	5	4	1
8	2	2	2	0

Hence, \mathbb{X} is either of type $(1, 1, 1, 1, 1, 1, 1, 2, a_9)$ or of type $(1, 1, 1, 1, 1, 1, 1, 1, 2)$. In the first case \mathbb{X} is a set of $2a_9$ reduced points lying on a conic $\mathcal{C} \subseteq V_{3,2}$ and in the second case \mathbb{X} is a set of 2 reduced points. The discussion for reduced 0-dimensional complete intersection is now completed. Now let $\mathbb{X} \subseteq V_{3,2}$ be a positive dimensional reduced complete intersection of type (a_1, \dots, a_r) . For any choice of integers a_{r+1}, \dots, a_9 , we can choose suitable hypersurfaces H_{a_i} of degree a_i in such a way that

$$\mathbb{X}' = \mathbb{X} \cap H_{a_{r+1}} \cap \dots \cap H_{a_9}$$

is a complete intersection of type $(a_1, \dots, a_r, a_{r+1}, \dots, a_9)$. Moreover, we can choose the degrees a_i in a such a way that

$$a_i \leq a_{i+1}$$

for all i and $3 \leq a_{r+1}$. Thus $\mathbb{X}' \subseteq V_{3,2}$ is a zero dimensional complete intersection of type (a_1, \dots, a_9) . As a consequence, since we can freely choose the degrees a_i for $i \geq r + 1$ we have that $r = 8$ and

$$(a_1, \dots, a_8, a_9) = (1, 1, 1, 1, 1, 1, 1, 2, a_9).$$

Hence, \mathbb{X} is of type $(1, 1, 1, 1, 1, 1, 2)$, that is \mathbb{X} is a conic. ■

We can now state the following conjecture, which we already proved for $n \leq 2$ any d and for $n = 3$ and $d = 2$, see Proposition 5.1 and Theorems 4.3 and 4.5.

Conjecture 5.2 *Let $\mathbb{X} \subseteq V_{n,d} \subseteq \mathbb{P}^N$ be a reduced subvariety with $d > 1$. Then \mathbb{X} is a complete intersection of type (a_1, \dots, a_r) , with $a_1 \leq \dots \leq a_r$ if and only if*

- $r = N, a_1 = \dots = a_N = 1$, any n, d , that is \mathbb{X} is a reduced point;
- $r = N, a_1 = \dots = a_{N-1} = 1, a_N = 2$, any n, d , that is \mathbb{X} is a set of two reduced points;
- $r = N, a_1 = \dots = a_{N-2} = 1, a_{N-1} = 2, a_N = b$, any $n, d = 2$, any $a \geq 2$, that is $\mathbb{X} = \mathcal{C} \cap H_b$ for $\mathcal{C} \subseteq V_{n,2}$ a conic and H_b a degree b hypersurface;
- $r = N - 1, a_1 = \dots = a_{N-2} = 1, a_{N-1} = 2, d = 2$, any n , that is \mathbb{X} is a conic.

In the case of the Veronese threefold $V_{3,2}$, see proof of Proposition 5.1, the complete knowledge of the zero dimensional complete case allows us to complete the proof. This is true in general as shown by the following Lemma.

Lemma 5.3 *If Conjecture 5.2 holds for all reduced zero dimensional subvariety of $V_{n,d}$, then it holds for all reduced subvarieties of $V_{n,d}$.*

Proof Let $\mathbb{X} \subseteq V_{n,d} \subseteq \mathbb{P}^N$ be a reduced complete intersection of type (a_1, \dots, a_r) with $r < N$, that is \mathbb{X} is positive dimensional. Then, for any choice of integers a_{r+1}, \dots, a_N we can choose suitable hypersurfaces H_{a_i} of degree a_i in such a way that

$$\mathbb{X}' = \mathbb{X} \cap H_{a_{r+1}} \cap \dots \cap H_{a_N}$$

is a complete intersection of type $(a_1, \dots, a_r, a_{r+1}, \dots, a_N)$. Moreover, we can choose the degrees a_i in a such a way that

$$a_i \leq a_{i+1}$$

for all i and $3 \leq a_{r+1}$. Thus $\mathbb{X}' \subseteq V_{n,d}$ is a zero dimensional complete intersection of type (a_1, \dots, a_N) . Since we are assuming that the conjecture holds for such an \mathbb{X}' and since we can freely choose the degrees a_i for $i \geq r + 1$ we have that $r = N - 1$ and

$$(a_1, \dots, a_{N-1}, a_N) = (1, \dots, 1, 2, a_N)$$

and thus $d = 2$ and \mathbb{X} is a conic. Hence the conjecture holds for \mathbb{X} . ■

A Appendix

We now prove Lemma 3.2.

Proof We distinguish four cases depending on the value of p . For each of them we give a function $\tilde{h}(i)$ satisfying 1. and 2. and such that

$$\tilde{h}(d) = \begin{cases} \lfloor \frac{2d(t+1)+3-\sqrt{1+8p}}{2} \rfloor, & \text{if } 1 \leq p \leq \binom{d+1}{2} \\ dt - n, & \text{if } \binom{d+1}{2} + dn < p \leq \binom{d+1}{2} + d(n+1), 0 \leq n \leq dt \end{cases}$$

Then we show that for any function $h'(i)$ satisfying 1. and 2. it holds that $h'(d) \leq \tilde{h}(d)$. We do this in detail in case (A) and for the remaining cases we produce the function \tilde{h} .

1. $p = \binom{n}{2}$, $1 \leq n \leq d+1$

In this case we set

$$\tilde{h}(i) = \begin{cases} dt + i + 1, & \text{if } 1 \leq i \leq d - n \\ d(t+1) - n + 2, & \text{if } d - n + 1 \leq i \leq d \end{cases}$$

We have

$$\sum_{i=1}^d \tilde{h}(i) = d^2t + \frac{d(d+3)}{2} - \frac{n(n-1)}{2} = d^2t + \frac{d(d+3)}{2} - p = \sum_{i=1}^d h(i)$$

and

$$\tilde{h}(d) = d(t+1) - n + 2 = \left\lfloor \frac{2d(t+1) + 3 - \sqrt{1+8p}}{2} \right\rfloor$$

hence $\tilde{h}(i)$ is as we want. Now let us suppose that there exists $h'(i) : \{1, 2, \dots, d\} \rightarrow \mathbb{N}$ satisfying 1. and 2. and such that $h'(d) > \tilde{h}(d)$, that is $h'(d) = d(t+1) - n + 2 + a$, $a \geq 1$. Since $n \leq d+1$, we have $h'(d) \geq dt + 1 + a \geq dt + 2$. As a consequence (observe that $dt + 2$ is the maximum value of $h(1)$), by 1. and 2., it follows that $h'(i)$ is increasing at least until reaching the value $h'(d)$. In particular, if we set $i' = \min \{1 \leq i \leq d \mid h(i) = d(t+1) - n + 2 + a\}$ we have

$$dt + i' + 1 = d(t+1) - n + 2 + a$$

so that $i' = d - n + 1 + a$ and $i_0 \geq i'$. Hence, using again 1. and 2., we get

$$\begin{aligned} h'(i) &= \tilde{h}(i) = dt + i + 1 & \text{if } i \leq d - n \\ h'(i) &> \tilde{h}(i) & \text{if } i \geq d - n + 1 \end{aligned}$$

Hence:

$$\sum_{i=1}^d h'(i) = \sum_{i=1}^{d-n} \tilde{h}(i) + \sum_{i=d-n+1}^d \underbrace{h'(i)}_{> \tilde{h}(i)} > \sum_{i=1}^d \tilde{h}(i) = \sum_{i=1}^d h(i)$$

and this is a contradiction.

2. $\binom{n}{2} < p < \binom{n+1}{2}$, $1 \leq n \leq d$

Let $b \in \mathbb{Z}$ be such that $p = \binom{n}{2} + b$. In this case we set

$$\tilde{h}(i) = \begin{cases} dt + i + 1, & \text{if } 1 \leq i \leq d - n + 1 \\ d(t+1) - n + 2, & \text{if } d - n + 2 \leq i \leq d - b \\ d(t+1) - n + 1, & \text{if } d - b + 1 \leq i \leq d \end{cases}$$

3. $p = \binom{d+1}{2} + d(n+1)$, $0 \leq n \leq dt$

In this case we set

$$\tilde{h}(i) = dt - n, \quad 1 \leq i \leq n.$$

We have

$$\sum_{i=1}^d \tilde{h}(i) = d(dt - n) = d^2t - dn = d^2t + \frac{d(d+3)}{2} - p = \sum_{i=1}^d h(i)$$

and

$$\tilde{h}(d) = dt - n$$

hence $\tilde{h}(i)$ is as we want.

4. $\binom{d+1}{2} + nd < p < \binom{d+1}{2} + (n+1)d$, $1 \leq n \leq dt$

Let $b \in \mathbb{Z}$ be such that $p = \binom{d+1}{2} + nd + b$. In this case we set

$$\tilde{h}(i) = \begin{cases} dt + 1 - n, & \text{if } 1 \leq i \leq d - b \\ dt - n, & \text{if } d - b + 1 \leq i \leq d \end{cases}$$

■

References

- [ABC18] Elena Angelini, Cristiano Bocci, and Luca Chiantini. Real identifiability vs. complex identifiability. *Linear Multilinear Algebra*, 66(6):1257–1267, 2018.
- [AC20] Elena Angelini and Luca Chiantini. On the identifiability of ternary forms. *Linear Algebra Appl.*, 599:36–65, 2020.
- [AGMO18] Elena Angelini, Francesco Galuppi, Massimiliano Mella, and Giorgio Ottaviani. On the number of Waring decompositions for a generic polynomial vector. *J. Pure Appl. Algebra*, 222(4):950–965, 2018.
- [BBOV21] Edoardo Ballico, Arthur Bik, Alessandro Oneto and Emanuele Ventura. *Israel J. Math.*, Strength and slice rank of forms are generically equal, 2021. To appear.
- [BGM94] Anna Bigatti, Anthony V. Geramita, and Juan C. Migliore. Geometric consequences of extremal behavior in a theorem of Macaulay. *Trans. Amer. Math. Soc.*, 346(1):203–235, 1994.
- [BO21] Arthur Bik and Alessandro Oneto. On the strength of general polynomials. *Linear and Multilinear Algebra*, 2021.
- [CCC⁺18] Enrico Carlini, Maria Virginia Catalisano, Luca Chiantini, Anthony V. Geramita, and Youngho Woo. Symmetric tensors: rank, Strassen’s conjecture and e -computability. *Ann. Sc. Norm. Super. Pisa Cl. Sci. (5)*, 18(1):363–390, 2018.
- [CCG08] Enrico Carlini, Luca Chiantini, and Anthony V. Geramita. Complete intersections on general hypersurfaces. volume 57, pages 121–136. 2008. Special volume in honor of Melvin Hochster.

- [CCG12] Enrico Carlini, Maria Virginia Catalisano, and Anthony V. Geramita. The solution to the Waring problem for monomials and the sum of coprime monomials. *J. Algebra*, 370:5–14, 2012.
- [CMM18] Alex Casarotti, Alex Massarenti, and Massimiliano Mella. On comon’s and strassen’s conjectures. *Mathematics*, 6(11):217, 2018.
- [CN17] Gianfranco Casnati and Roberto Notari. Examples of rank two aCM bundles on smooth quartic surfaces in \mathbb{P}^3 . *Rendiconti del Circolo Matematico di Palermo Series 2*, volume 66, pages 19-41. 2017.
- [DM84] Edward D. Davis and Paolo Maroscia. Complete intersections in \mathbb{P}^2 : Cayley-Bacharach characterizations. In: Greco, S., Strano, R. (eds) *Complete Intersections:: Lectures given at the 1st 1983 Session of the Centro Internazionale Matematico Estivo (C.I.M.E.) held at Acireale (Catania), Italy, June 13–21, 1983*. Lecture Notes in Mathematics, vol 1092. Springer, Berlin, Heidelberg.
- [Eis95] David Eisenbud. *Commutative algebra*, volume 150 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995. With a view toward algebraic geometry.
- [EGH96] David Eisenbud, Mark Green and Joe Harris. *Cayley-Bacharach theorems and conjectures* *Bull. Am. Math. Soc. New Series*, volume 33, pages 295-324. 1996.
- [Eul53] Leonhard Euler. *Commentationes Geometricae. Opera omnia*, ser. 1 vol. 26, (A. Speiser, editor), Zürich: Orell Füssli, 1953.
- [GMP11] Vesselin Gasharov, Satoshi Murai, and Irena Peeva. Hilbert schemes and maximal Betti numbers over Veronese rings. *Math. Z.*, 267(1-2):155–172, 2011.
- [GMR83] A. V. Geramita, P. Maroscia, and L. G. Roberts. The Hilbert function of a reduced k -algebra. *J. London Math. Soc. (2)*, 28(3):443–452, 1983.
- [GS84] Silvio Greco and Rosario Strano. *Complete Intersections: Lectures given at the 1st 1983 Session of the Centro Internazionale Matematico Estivo (C.I.M.E.) held at Acireale (Catania), Italy, June 13–21, 1983*. Berlin: Springer-Verlag, 1984. Print. Lecture Notes in Mathematics Fondazione C.I.M.E., Firenze 1092.
- [Har70] Robin Hartshorne. Ample subvarieties of algebraic varieties. *Lecture Notes in Math.*, No. 156, 1970.
- [KK98] Bettina Kreuzer and Martin Kreuzer. Extremal zero-dimensional subschemes of \mathbf{P}^2 . *J. Pure Appl. Algebra*, 131(2):159–177, 1998.
- [KNR] Martin Kreuzer, Le Ngoc Long and Lorenzo Robbiano. Algorithms for checking zero-dimensional complete intersections. *J. Commut. Algebra*, 2019 (in press).
- [Mig07] Juan C. Migliore. The geometry of Hilbert functions. In *Syzygies and Hilbert functions*, volume 254 of *Lect. Notes Pure Appl. Math.*, pages 179–208. Chapman & Hall/CRC, Boca Raton, FL, 2007.
- [Puc98] Mario Pucci. The Veronese variety and catalecticant matrices. *J. Algebra*, 202(1):72–95, 1998.

- [Shi19] Yaroslav Shitov. Counterexamples to Strassen's direct sum conjecture. *Acta Math.*, 222(2):363–379, 2019.
- [ST22] Ronnie Sebastian and Amit Tripathi. Rank 2 Ulrich bundles on general double plane covers. *J. Pure Appl. Algebra*, 226(2):Paper No. 106823, 7, 2022.
- [Sta78] Richard P. Stanley. Hilbert functions of graded algebras. *Advances in Math.*, 28(1):57–83, 1978.