## POLITECNICO DI TORINO Repository ISTITUZIONALE

### Novel Test Integral Quadrature Scheme for the Method of Moments

Original

Novel Test Integral Quadrature Scheme for the Method of Moments / Rivero, J.; Vipiana, F.; Wilton, D. R.; Johnson, W. A. - ELETTRONICO. - (2023), pp. 1-3. (Intervento presentato al convegno 17th European Conference on Antennas and Propagation, EuCAP 2023 tenutosi a Florence, Italy nel 26-31 March 2023) [10.23919/EuCAP57121.2023.10133393].

Availability: This version is available at: 11583/2982033 since: 2023-09-12T09:01:03Z

Publisher: IEEE

Published DOI:10.23919/EuCAP57121.2023.10133393

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright IEEE postprint/Author's Accepted Manuscript

©2023 IEEE. Personal use of this material is permitted. Permission from IEEE must be obtained for all other uses, in any current or future media, including reprinting/republishing this material for advertising or promotional purposes, creating new collecting works, for resale or lists, or reuse of any copyrighted component of this work in other works.

(Article begins on next page)

# Novel Test Integral Quadrature Scheme for the Method of Moments

Javier Rivero\*, Francesca Vipiana\*, Donald R. Wilton<sup>†</sup>, William A. Johnson<sup>‡</sup>

\*Dept. of Electronics and Telecommunications, Politecnico di Torino, Torino, Italy, {javier.rivero,francesca.vipiana}@polito.it <sup>†</sup>Dept. of Electrical and Computer Engineering, University of Houston,Houston, TX, USA, wilton@uh.edu

<sup>‡</sup>Consultant, Jemez Springs, NM, USA, w.johnson24@comcast.net

Abstract—An extensive literature exists on the efficient and accurate evaluation of the double surface integrals that arise in the Method of Moments. Most papers have focused on the evaluation of the inner (source) integral with the idea that once that integral is evaluated, the test (outer) integral's integrand is sufficiently smooth that it should be much easier to integrate numerically. However, that affirmation is not always true. Here, we propose an integration scheme that improves the numerical evaluation of the test integral without affecting the treatment of the source integral. The method is numerically validated for static and dynamic kernels in the reaction integrals arising in electric field integral equations.

*Index Terms*—integral equations, moment methods, numerical analysis, singular integrals.

#### I. INTRODUCTION

A cost-effective numerical evaluation of double surface reaction integrals is required to achieve an accurate solution of direct or inverse electromagnetic problems using surface integral equation formulations. The inner integral is commonly known as *source* integral, and the outer one is the *test* integral. The most common approaches for dealing with the source integral are the singularity subtraction or singularity cancellation methods. For singularity subtraction [1]-[3], a simplified asymptotic form of the integrand is first identified and subtracted from integrand. The resulting difference integrand should be less singular than the original, and the subtracted term should be analytically integrable (or at least easily evaluated numerically); adding the analytical integral to the difference integral restores the original value of the integral. On the other hand, for singularity cancellation [3]-[7], variable transforms are chosen whose Jacobian cancels or regularizes any singularities. More recently, a paper has considered the possibility of using both methods to bring together the advantages of both previous approaches [8] handling the singular behavior that arise when evaluating the magnetic field integral equation (MFIE). As can be seen in the literature, the effort to evaluate the double integrals has been focused on the efficient evaluation of the source integral [1]-[7] and simply assuming that for the test integral any singular behavior has been smoothed by the source integral, making the test integral relatively easy to integrate numerically. In cases where

the source and test domains share points in common, this affirmation does not hold, as shown in [9].

Here, we focus on the numerical evaluation of the outer test integral that arises in the reaction integrals. We will use the vertex functions described in [10] to discern the behavior of the test integral integrand. The vertex functions are elementindependent functions describing the (possibly) singular behavior of potentials or their gradients near a given vertex or edge for constant or linear sources on a planar element. Once the behavior of the vertex functions is understood, we can design an efficient quadrature scheme for the test integral.

#### II. VERTEX FUNCTION

The evaluation of the electromagnetic interaction between a pair of triangles in the Method of Moments (MoM) leads to the evaluation of the double surface integral

$$\int_{S} \int_{S'} F(\mathbf{r}, \mathbf{r}') \, dS' \, dS, \tag{1}$$

where typically  $F(\mathbf{r}, \mathbf{r}')$  takes the form

$$F(\mathbf{r}, \mathbf{r}') = t(\mathbf{r})g(\mathbf{r}, \mathbf{r}') b(\mathbf{r}'), \qquad (2)$$

and where  $t(\mathbf{r})$  is either a scalar or a vector component of a testing function,  $b(\mathbf{r}')$  is similarly defined for a basis function, and  $g(\mathbf{r}, \mathbf{r}')$  is either a scalar or a scalar component of a vector or dyadic Green's function, with a  $\mathcal{O}(|\mathbf{r} - \mathbf{r}'|^{-1})$  or  $\mathcal{O}(\nabla |\mathbf{r} - \mathbf{r}'|^{-1})$  singularity.

We assume the inner integral in (1) may be evaluated to sufficiently high accuracy so its influence is dismissed. Its singular behavior follows from is static (k = 0) form. This, in terms, can be represented as a sum over vertex function pairs, with the superposition describing the essential behavior of the various potentials (scalar and vector) involved. For a case of a pair of triangles with a common vertex, the static vertex potential functions for the common vertices associated with a pair of adjacent source triangle edges for a constant (unit) source density are given by

$$I_{V_{1}}^{l_{3}} = R_{1} \sin \alpha \left[ \cos \beta \sinh^{-1} \left( \cot \alpha \right) - \left| \sin \beta \right| \tan^{-1} \left( \frac{\cos \alpha + \cos \beta}{\sin \alpha + \left| \sin \beta \right|} \right) \right], \quad (3)$$



Fig. 1. The vertex function of each adjacent vertex-edge pairs shows a singular behavior close to the edge that defines the function (thicker lines) and smoother behavior far from the edge (thinner lines). Dashed lines are not in the test integral domain.

where  $R_1$  is the distance from the obs. point to the vertex,  $\alpha$  is the angle between the edge  $l_3$  and  $R_1$  and  $\beta$  is the angle between the two triangles (or the extended planes in which they lie). Analyzing (3), we find that the vertex function has a linear dependence with distance  $R_1$  and exhibits a singular behavior as  $\alpha \rightarrow 0$ . The remaining vertex functions, though with more complex expressions, exhibit similar singular behavior, that is, a linear dependence with respect to the distance to the vertex and a singular behavior with respect to angle. Figure 1 shows graphically this behavior, where the vertex functions exhibit more nearly- singular along the thicker lines (with dashed lines indicating points outside the test integral domain). Considering this behavior, we can better handle the singularities if we concentrate more sample points close to the common vertex as well as closer to the source triangle. The vertex functions also reveal a logarithmic angular singularity, hence the proposed scheme uses MWR [11] quadrature with logarithmically weighted sample points for the angular integration and a parameterization of the edge opposite the common vertex to form an angular scheme. Using these points as endpoints, we now parameterize radially from the common vertex also using the logarithmic MWR scheme. This procedure is described graphically in Fig. 2. The new quadrature points  $\xi_i$  can be mathematically expressed as

$$\xi_i = V_1^T (1 - \xi_j^r) + V_2^T (\xi_j^r - \xi_j^r \xi_k^a) + V_3^T (\xi_j^r \xi_k^a)$$
(4)



Fig. 2. Procedure to obtain the proposed scheme, a) parametrize the edge oposite to the common vertex, b) define radial integration dimension, c) parametrize radial integration dimension, d) obtain quadrature points.



Fig. 3. Near-field convergence of test integrals. Inset: Orientation of a pair of triangle elements in space.

where  $V_{i=1,2,3}^T$  are the vertices of the test triangle and  $\xi_j^r$  and  $\xi_k^a$  are the 1-D MRW quadrature points used to parameterize the radial and angular directions, respectively.

#### **III. PRELIMINARY NUMERICAL RESULTS**

To demonstrate the accuracy of the proposed scheme, we analyze the convergence behavior of the test scalar potential integral in (1). We consider a pair of triangles with a common vertex as shown in the Fig. 3 inset. We compare the proposed method with the standard Gauss-triangle (GT) quadrature scheme [12]. The reference for each of the plot is evaluated with the highest number of points we have available. In the case of GT, we do not have more than 166 points so the behavior over that number of points is estimated with the plot trend. The proposed method reaches machine precision at a faster rate in the medium accuracy range than GT.

#### IV. CONCLUSION AND PERSPECTIVES

Preliminary results show better accuracy and efficiency of the proposed method with respect to the GT quadrature scheme.With small modifications, the method can also be applied to test triangles with a test triangle edge in common with the source triangle and self-term triangles for all the reaction integrals arising in the electric field integral equation and magnetic field integral equation.

#### References

- R. Graglia, "On the numerical integration of the linear shape functions times the 3-D Green's function or its gradient on a plane triangle," *IEEE Transactions on Antennas and Propagation*, vol. 41, no. 10, pp. 1448–1455, Oct. 1993. [Online]. Available: http://ieeexplore.ieee.org/document/247786/
- [2] S. Järvenpää, M. Taskinen, and P. Ylä-Oijala, "Singularity Subtraction Technique for High-Order Polynomial Vector Basis Functions on Planar Triangles," *IEEE Transactions on Antennas and Propagation*, vol. 54, no. 1, pp. 42–49, Jan. 2006.
- [3] L. Li and T. F. Eibert, "Radial-Angular Singularity Cancellation Transformations Derived by Variable Separation," *IEEE Transactions on Antennas and Propagation*, vol. 64, pp. 189–200, Jan. 2016.
  [4] M. A. Khayat, D. R. Wilton, and P. W. Fink, "An Improved Transformation".
- [4] M. A. Khayat, D. R. Wilton, and P. W. Fink, "An Improved Transformation and Optimized Sampling Scheme for the Numerical Evaluation of Singular and Near-Singular Potentials," *IEEE Antennas and Wireless Propagation Letters*, vol. 7, pp. 377–380, 2008.
  [5] F. Vipiana and D. R. Wilton, "Numerical Evaluation via Singularity
- [5] F. Vipiana and D. R. Wilton, "Numerical Evaluation via Singularity Cancellation Schemes of Near-Singular Integrals Involving the Gradient of Helmholtz-Type Potentials," *IEEE Transactions on Antennas and Propagation*, vol. 61, no. 3, pp. 1255–1265, Mar. 2013.
- [6] M. M. Botha, "A Family of Augmented Duffy Transformations for Near-Singularity Cancellation Quadrature," *IEEE Transactions on Antennas* and Propagation, vol. 61, pp. 3123–3134, Jun. 2013.
- [7] —, "Numerical Integration Scheme for the Near-Singular Green Function Gradient on General Triangles," *IEEE Transactions on Antennas and Propagation*, vol. 63, no. 10, pp. 4435–4445, Oct. 2015. [Online]. Available: http://ieeexplore.ieee.org/document/7160690/
- [8] J. Rivero, F. Vipiana, D. R. Wilton, and W. A. Johnson, "Hybrid Integration Scheme for the Evaluation of Strongly Singular and Near-Singular Integrals in Surface Integral Equations," *IEEE Transactions on Antennas and Propagation*, vol. 67, no. 10, pp. 6532–6540, Oct. 2019.
- [9] F. Vipiana, D. R. Wilton, and W. A. Johnson, "Advanced Numerical Schemes for the Accurate Evaluation of 4-D Reaction Integrals in the Method of Moments," *IEEE Transactions on Antennas and Propagation*, vol. 61, no. 11, pp. 5559–5566, Nov. 2013.
- [10] D. R. Wilton, J. Rivero, W. A. Johnson, and F. Vipiana, "Evaluation of Static Potential Integrals on Triangular Domains," *IEEE Access*, vol. 8, pp. 99806–99819, 2020.
- [11] J. Ma, V. Rokhlin, and S. Wandzura, "Generalized Gaussian quadrature rules for systems of arbitrary functions," *SIAM J. Numer. Anal.*, vol. 33, no. 3, pp. 971–996, 1996. [Online]. Available: http://www.jstor.org/stable/2158491
- [12] L. Zhang, T. Cui, and H. Liu, "A Set of Symmetric Quadrature Rules on Triangles and Tetrahedra," *Journal of Computational Mathematics*, vol. 27, no. 1, pp. 89–96, 2009.