

Implicitisation and Parameterisation in Polynomial Functors

Original

Implicitisation and Parameterisation in Polynomial Functors / Blatter, Andreas; Draisma, Jan; Ventura, Emanuele. - In: FOUNDATIONS OF COMPUTATIONAL MATHEMATICS. - ISSN 1615-3383. - (2024), pp. 1567-1593. [10.1007/s10208-023-09619-6]

Availability:

This version is available at: 11583/2982769 since: 2023-10-05T09:31:58Z

Publisher:

Springer

Published

DOI:10.1007/s10208-023-09619-6

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



Implicitisation and Parameterisation in Polynomial Functors

Andreas Blatter¹ · Jan Draisma^{2,3} · Emanuele Ventura⁴

Received: 2 June 2022 / Revised: 14 June 2023 / Accepted: 6 July 2023

© The Author(s) 2023

Abstract

In earlier work, the second author showed that a closed subset of a polynomial functor can always be defined by finitely many polynomial equations. In follow-up work on GL_∞ -varieties, Bik–Draisma–Eggermont–Snowden showed, among other things, that in characteristic zero every such closed subset is the image of a morphism whose domain is the product of a finite-dimensional affine variety and a polynomial functor. In this paper, we show that both results can be made algorithmic: there exists an algorithm **implicitise** that takes as input a morphism into a polynomial functor and outputs finitely many equations defining the closure of the image; and an algorithm **parameterise** that takes as input a finite set of equations defining a closed subset of a polynomial functor and outputs a morphism whose image is that closed subset.

Keywords Polynomial functor · Schur functors · Elimination · Parameterisation

Communicated by Teresa Krick.

Andreas Blatter was supported by Swiss National Science Foundation Grant 200021_191981. Jan Draisma was partially supported by SNSF Grant 200021_191981 and Vici Grant 639.033.514 from the Netherlands Organisation for Scientific Research. Emanuele Ventura is a member of GNSAGA of INdAM (Italy).

✉ Jan Draisma
jan.draisma@unibe.ch
Andreas Blatter
andreas.blatter@unibe.ch
Emanuele Ventura
emanuele.ventura@polito.it

- 1 Mathematical Institute, University of Bern, Alpeneggstrasse 22, 3012 Bern, Switzerland
- 2 Mathematical Institute, University of Bern, Sidlerstrasse 5, 3012 Bern, Switzerland
- 3 Department of Mathematics and Computer Science, Eindhoven University of Technology, P.O. Box 513, 5600MB Eindhoven, The Netherlands
- 4 Dipartimento di Scienze Matematiche “G.L. Lagrange”, Politecnico di Torino, Corso Duca degli Abruzzi 24, 10129 Turin, Italy

Mathematics Subject Classification 13P99 · 15A69 · 20G05 · 14M99

1 Introduction

1.1 Implicitisation

An important theme in computational algebraic geometry is *implicitisation*: given a list $(\varphi_1, \dots, \varphi_n)$ of polynomials in $K[x_1, \dots, x_m]$, which represent a polynomial map φ from the m -dimensional affine space \mathbb{A}^m to \mathbb{A}^n over the field K , the challenge is to compute equations for the Zariski closure $\overline{\text{im}(\varphi)}$ of the image of φ . This challenge is solved, at least theoretically, by *elimination* using Buchberger’s algorithm; e.g. see [7, §3.3].

1.2 Implicitisation in Families

But now consider a scenario where one is given not a single polynomial map, but rather a family $\varphi_i : \mathbb{A}^{m_i} \rightarrow \mathbb{A}^{n_i}$ of polynomial maps depending on a discrete parameter i that takes infinitely many values. If the ambient spaces \mathbb{A}^{m_i} and \mathbb{A}^{n_i} and the polynomial map φ_i vary favourably with i , it is sometimes possible to find finitely many equations that capture the image closures of all φ_i at once.

1.3 Implicitisation Over Categories

To make the phrase “vary favourably” concrete, assume that i ranges through the objects of a category C and $\mathbb{A}(i)$ is a finite-dimensional affine space varying functorially with i . This means that for each $\pi \in \text{Hom}_C(i, j)$ we have a linear map $\mathbb{A}(\pi) : \mathbb{A}(i) \rightarrow \mathbb{A}(j)$ such that $\mathbb{A}(\text{id}_i) = \text{id}_{\mathbb{A}(i)}$ and $\mathbb{A}(\sigma \circ \pi) = \mathbb{A}(\sigma) \circ \mathbb{A}(\pi)$ for $\sigma \in \text{Hom}_C(j, k)$. Suppose that $\mathbb{A}'(i)$ is another affine space depending functorially on $i \in C$, and finally that, for any $i \in C$, we have a polynomial map $\varphi_i : \mathbb{A}(i) \rightarrow \mathbb{A}'(i)$ such that the following diagram commutes for every $\pi \in \text{Hom}_C(i, j)$:

$$\begin{array}{ccc}
 \mathbb{A}(i) & \xrightarrow{\varphi_i} & \mathbb{A}'(i) \\
 \mathbb{A}(\pi) \downarrow & & \downarrow \mathbb{A}'(\pi) \\
 \mathbb{A}(j) & \xrightarrow{\varphi_j} & \mathbb{A}'(j).
 \end{array}$$

In this case, if f is a polynomial equation for the image closure $\overline{\text{im}(\varphi_j)}$ of φ_j , then $f \circ \mathbb{A}'(\pi)$ is a polynomial equation for $\overline{\text{im}(\varphi_i)}$ —and, since we assumed that $\mathbb{A}'(\pi)$ is linear, of the same degree. We then ask:

- (1) Do there exist finitely many $j \in C$ such that the equations for those $\overline{\text{im}(\varphi_j)}$, by pulling back along the linear maps $\mathbb{A}'(\pi)$ for all relevant π , define $\overline{\text{im}(\varphi_i)}$ for all $i \in C$?

(2) If so, does there exist an *algorithm* for computing these finitely many j ?

A well-known case where the answer to the two questions above is “yes” is that where C is the opposite category \mathbf{FI}^{op} of the category \mathbf{FI} of finite sets with injections, $\mathbb{A}(i) = (\mathbb{A}^n)^i$, $\mathbb{A}'(i) = (\mathbb{A}^{n'})^i$ for some fixed n and n' , and the maps $\mathbb{A}(i) \rightarrow \mathbb{A}(j)$, $\mathbb{A}'(i) \rightarrow \mathbb{A}'(j)$ corresponding to an injection $j \rightarrow i$ are the canonical projections. In that case, it is known that the kernel of the \mathbf{FI} -homomorphism φ^* dual to the \mathbf{FI}^{op} -polynomial map φ is finitely generated and can be computed using a version of Buchberger’s algorithm, and this has been applied to problems in algebraic statistics [4, 5, 15–17]. The setting discussed in the current paper is of a very different flavour in that it involves continuous symmetries rather than discrete symmetries, and it is also significantly more complicated. One cause for trouble is that in the setting below, we do not actually know whether the kernel of the relevant algebra homomorphism is finitely generated, and so we have to settle for finding set-theoretic equations for the $\overline{\text{im}(\varphi_i)}$.

1.4 Implicitisation in Polynomial Functors

In the case where C is the category \mathbf{Vec} of finite-dimensional K -vector spaces, and both \mathbb{A} and \mathbb{A}' are polynomial functors $\mathbf{Vec} \rightarrow \mathbf{Vec}$ (see Sect. 2.3 for definitions), the second author established a positive answer to the first question above [9]; see Theorem 2.4.2.

Example 1.4.1 Here are two instances of this setting:

- (1) $\mathbb{A}(V) = V^k$ for some positive integer k , $\mathbb{A}'(V) = S^3 V$, the third symmetric power of V , and

$$\varphi_V(v_1, \dots, v_k) := v_1^3 + \dots + v_k^3.$$

The image of φ_V is the set of cubic homogeneous polynomials in $\dim(V)$ variables of *Waring rank* $\leq k$, and its closure is the variety of polynomials of *border Waring rank* $\leq k$.

- (2) $\mathbb{A}(V) = (S^2 V)^k \times V^k$, $\mathbb{A}'(V) = S^3 V$, and

$$\varphi_V(q_1, \dots, q_k, v_1, \dots, v_k) := q_1 v_1 + \dots + q_k v_k.$$

In this case, the image of φ_V is closed [8] and consists of the cubics of *q-rank* (also called *slice rank* or *Schmidt rank* in the literature) $\leq k$.

In the first case, it is well known that generators of the vanishing ideal of $\overline{\text{im}(\varphi_U)}$ for U of dimension $k + 1$ pull back to generators of the ideal of $\overline{\text{im}(\varphi_V)}$ for all V ; this is called *symmetric inheritance* in [19]. Much more is known for small values of k ; in particular, for $k \leq 2$, the ideal is generated by 3×3 -minors of catalecticant matrices [22].

In the second case, we do not know whether the *ideal* is finitely generated in this sense, but [8] assures the existence of a number $\ell = \ell(k)$ such that equations for

$\overline{\text{im}(\varphi_U)}$ with U of dimension ℓ pull back to equations that define $\overline{\text{im}(\varphi_V)}$ for all V *set-theoretically*. No explicit value of $\ell(k)$ is known, even for $k = 2$. \diamond

In many respects, the q -rank example is much more difficult than the Waring rank example. The main reason for this is that the source \mathbb{A} is a polynomial functor of degree 2, whereas \mathbb{A} has degree 1 in the first case. Nevertheless, Theorem 2.4.2 implies, for general polynomial functors \mathbb{A}, \mathbb{A}' and morphisms φ , the existence of a U such that in $\mathbb{A}'(U)$ one sees enough set-theoretic equations to define $\overline{\text{im}(\varphi_V)}$ for all V .

1.5 Main Result

The main result of this paper makes Theorem 2.4.2 effective.

Theorem 1.5.1 (Main Theorem; see Theorem 4.1.1 for a more precise statement) *There exists an algorithm **implicitise** that, on input polynomial functors \mathbb{A}, \mathbb{A}' and a morphism $\varphi : \mathbb{A} \rightarrow \mathbb{A}'$, computes a $U \in \mathbf{Vec}$ such that the equations for $\overline{\text{im}(\varphi_U)}$ pull back to set-theoretic defining equations for $\overline{\text{im}(\varphi_V)}$ for all V .*

This means, in particular, that the number $\ell(k)$ for the cubics of q -rank at most any fixed k can in principle be computed.

1.6 Structure of the Algorithm

The structure of **implicitise** is as follows. For $U = K^0, K^1, K^2, \dots$, one computes equations for $\overline{\text{im}(\varphi_U)}$ using Buchberger’s algorithm. By Theorem 2.4.2, we know that at some point these equations, via pull-back, define $\overline{\text{im}(\varphi_V)}$ for all V . However, Theorem 2.4.2 does not give a criterion for when we may stop—this is different from the algorithm in the **FI**-setting: there, if one has not seen new equations arise between the finite set $\{1, \dots, n\}$ and the finite set $\{1, \dots, 2n\}$, one is guaranteed to have found a generating set of equations [4].

To obtain a stopping criterion in the polynomial functor setting, we derive an algorithm **parameterise** that, on input the polynomial equations on $\mathbb{A}'(U)$ found so far, computes a parameterisation ψ of the closed subset of \mathbb{A}' defined by those equations. If we can check that $\text{im}(\psi) \subseteq \overline{\text{im}(\varphi)}$, then we are done.

That such a parameterisation ψ exists is guaranteed by the *unirationality theorem* from [2]. The algorithm **parameterise** makes this theorem effective.

Finally, to check that $\text{im}(\psi)$ lies in the closure of $\text{im}(\varphi)$, we pass to infinite dimensions and use a result from [3] that says that this happens if and only if a suitably generic point of $\text{im}(\psi_\infty)$ can be reached as the limit of a curve in $\text{im}(\varphi_\infty)$. In the current paper, we show that this curve, which lives in infinite-dimensional space, can be represented in finite terms, and searched for, on a computer.

If such a curve does not exist, i.e. if $\text{im}(\psi)$ is not contained in $\overline{\text{im}(\varphi)}$, then this is because the current space U is too small. In this case, the search for a curve does not terminate. So for our algorithm **implicitise** to terminate, it is imperative to run the search for witness curves in parallel to the search for equations: in each step, U is increased with one dimension, new equations are computed, and a new curve search is

started. We model this behaviour by running the algorithm on countably many parallel processors. Of course, standard results in the theory of computation imply that this algorithm can then also run on an ordinary Turing machine (see, e.g. [20]).

1.7 Outlook

The algorithms **implicitise** and **parameterise** are very much theoretical algorithms, and—apart from recent research on varieties over categories—they rely on many powerful results from classical, finite-dimensional, computational algebra: Buchberger’s algorithm, of course, but also algorithms for computing the radical of an ideal and for computing the minimal primes containing a radical ideal. Also, a fair amount of representation theory of the general linear group goes into the algorithm.

At present, a general implementation of **implicitise** and **parameterise** seems entirely out of reach. However, we believe that our computational approach to implicitisation in polynomial functors will in the future serve as a guide towards finding equations in concrete settings, e.g. for the variety of cubics of q -rank ≤ 2 from Example 1.4.1.

1.8 Organisation of the Paper

In Sect. 2, we collect material that we will need in the rest of the paper: some assumptions on the ground field, computer representations of finite-dimensional affine varieties, polynomial functors and their closed subsets, and morphisms between these.

In Sect. 3, we derive the algorithm **parameterise**; see Theorem 3.1.1. In Sect. 4, we derive the algorithm **implicitise** and prove our Main Theorem; see Theorem 4.1.1 for a more precise statement.

Finally, **implicitise** depends on a procedure called **certify** that aims at certifying the existence of a curve as discussed in Sect. 1.6. This procedure requires tools from infinite-dimensional algebraic geometry, both from the paper [3] and from more classical sources. In particular, it is Greenberg’s approximation theorem (Theorem 5.5.1) from [13] and its effective version in [23] which allows us to represent this curve in finite terms.

2 Preliminaries

2.1 The Ground Field

Let K be a field of characteristic zero in which we can do computations on a computer. More precisely, we want K to be a computable field, and we further require that there exists an algorithm for factoring polynomials in $K[X]$. The class of such fields is quite large; it includes \mathbb{Q} and its finitely generated extensions [14, Appendix B].

2.2 Finite-Dimensional Affine Varieties

A finite-dimensional affine variety over K is represented by a finite list of generators of a radical ideal in some polynomial ring over K with finitely many variables. We will only consider varieties over K , and in particular the adjective irreducible will refer to irreducibility over K . A morphism between affine varieties over K is represented by a finite list of polynomials. Our algorithms will intensively use existing algorithms for dealing with affine varieties. Good general references are [6, 7]. In particular, we will need algorithms for computing the radical of an ideal; see, e.g. [18].

If R is a K -algebra and h is an element of R , then we write $R[1/h]$ for the localisation. Similarly, if B is a finite-dimensional affine variety and h an element of $K[B]$, then we write $B[1/h]$ for the basic open subset defined by h , i.e. the affine variety with coordinate ring $K[B][1/h]$.

2.3 Polynomial Functors

Let \mathbf{Vec} be the category of finite-dimensional K -vector spaces. We write $\text{Hom}(U, V)$ and $\text{End}(U)$ for the spaces of K -linear maps $U \rightarrow V$ and $U \rightarrow U$, respectively.

Definition 2.3.1 A polynomial functor of degree at most d over K is a covariant functor $P : \mathbf{Vec} \rightarrow \mathbf{Vec}$ such that for any $U, V \in \mathbf{Vec}$ the map $P : \text{Hom}(U, V) \rightarrow \text{Hom}_{P(U), P(V)}$ is polynomial of degree at most d . Polynomial functors of degree at most d form an abelian category in which a homomorphism $P \rightarrow Q$ is a natural transformation, i.e. given by a linear map $\psi_U : P(U) \rightarrow Q(U)$ for each $U \in \mathbf{Vec}$, such that for all $\varphi \in \text{Hom}(U, V)$ we have $Q(\varphi) \circ \psi_U = \psi_V \circ P(\varphi)$. \diamond

When we say polynomial functor, we will always mean a polynomial functor of degree at most some integer. A priori, a polynomial functor seems to be given by an infinite amount of data. But up to isomorphism, the following lemma due to Friedlander–Suslin [11, Lemma 3.4] implies that it is given by a finite amount of data only.

Lemma 2.3.2 [11] *The map from polynomial functors of degree at most d to GL_d -representations that assigns to P the vector space $P(K^d)$ with the algebraic group homomorphism $\text{GL}_d \rightarrow \text{GL}(P(K^d)), \varphi \mapsto P(\varphi)$ is an equivalence of abelian categories from polynomial functors of degree at most d to polynomial GL_d -representations of degree at most d .*

Here, a homomorphism $\rho : \text{GL}_d \rightarrow \text{GL}(W)$ of algebraic groups is called polynomial of degree at most d if it extends to a polynomial map $\text{End}(K^d) \rightarrow \text{End}(W)$ of degree at most d . For our algorithmic purposes, it is important that the proof of the Friedlander–Suslin lemma, and in particular the map back, is completely constructive. We sketch this map back.

Proof (Construction of the inverse map in the Friedlander–Suslin Lemma) By polynomiality, ρ extends to a polynomial map $\rho : M_d \rightarrow \text{End}(W)$, where $M_d := \text{End}(K^d)$, and this extension is a monoid homomorphism. Dually, this gives an algebra homomorphism $K[\text{End}(W)] \rightarrow K[M_d]$ that restricts to a linear map from $\text{End}(W)^*$ to

the space $K[M_d]_{\leq d}$ of polynomials of degree $\leq d$ in the standard coordinates x_{ij} on M_d . Dualising once again, we obtain a map $\tilde{\rho} : S := K[M_d]_{\leq d}^* \rightarrow \text{End}(W)$. By a similar construction, now applied to the multiplication $M_d \times M_d \rightarrow M_d$ instead of ρ , S becomes an associative algebra, and $\tilde{\rho}$ is an algebra homomorphism, so W is a left S -module. (This actually gives an equivalence of categories between polynomial GL_d -representations of degree $\leq d$ and S -modules.) Now let $V \in \mathbf{Vec}$ be arbitrary. Then, $\text{Hom}(K^d, V)$ is a right M_d -module, and by a construction like above applied to the map $\text{Hom}(K^d, V) \times M_d \rightarrow \text{Hom}(K^d, V)$, the space $K[\text{Hom}(K^d, V)]_{\leq d}^*$ becomes a right S -module. Finally, define $P(V) := K[\text{Hom}(K^d, V)]_{\leq d}^* \otimes_S W$. This is the polynomial functor corresponding to ρ . \square

The Friedlander–Suslin lemma is relevant to us for two reasons. First, since polynomial representations of GL_d are completely reducible and the irreducible ones are completely classified by combinatorial data, the same holds for polynomial functors. The consequence is that each polynomial functor of degree at most d is a direct sum of *Schur functors*: functors of the form $S_\lambda : V \mapsto \text{Hom}_{S_e}(U_\lambda, V^{\otimes e})$, where $e \leq d$, λ is a partition of e , and U_λ is the corresponding irreducible representation of the symmetric group S_e on e letters. So we may represent a polynomial functor by a finite tuple of partitions. Second, the above shows that the Friedlander–Suslin lemma is completely constructive: it can be transformed into algorithms that, using basic linear algebra, compute the following data from a polynomial representation ρ of GL_d of degree $\leq d$ corresponding to a polynomial functor P :

- (a basis for) $P(V)$ on input V ;
- (a matrix for) $P(\varphi) : P(U) \rightarrow P(V)$ on input a linear map $\varphi : U \rightarrow V$; and
- (a matrix for) $\psi_V : P(V) \rightarrow Q(V)$ on input a second polynomial GL_d -representation of degree $\leq d$ representing the polynomial functor Q , as well as homomorphism of GL_d -representations.

We will not make these algorithms explicit here.

Each polynomial functor P of degree at most d is a direct sum of *homogeneous* polynomial functors: $P = P_0 \oplus \dots \oplus P_d$, where

$$P_e(V) := \{p \in P(V) \mid P(t \cdot \text{id}_V)p = t^e p\}.$$

In particular, P_0 is a degree-0 polynomial functor, which assigns a fixed vector space, also denoted P_0 , to every V and the identity id_{P_0} to every linear map $\varphi \in \text{Hom}(U, V)$. We call P *pure* if $P_0 = 0$, and we call $P_1 \oplus \dots \oplus P_d$ the *pure part* of P .

2.4 Closed Subsets of Polynomial Functors

Definition 2.4.1 A *closed subset* X of a polynomial functor P over K is the data of a closed subvariety $X(V)$ of $\overline{K} \otimes P(V)$ defined over K such that for each $\varphi \in \text{Hom}(U, V)$ the linear map $1 \otimes P(\varphi)$ maps $X(U)$ into $X(V)$. \diamond

We take points with coordinates in \overline{K} because K might not be large enough to see all points. On the other hand, our algorithm will always work over K itself, and all

varieties will be defined over K . In fact, we shall usually drop the \overline{K} from the notation and just write *closed subvariety* $X(V)$ of $P(V)$ and $P(\varphi) : X(U) \rightarrow X(V)$ in the above setting. This is not different from the classical setting of implicitisation, where one computes the equations of the Zariski closure in \overline{K}^n of the image of a polynomial map $\overline{K}^m \rightarrow \overline{K}^n$ which is defined over K .

For a closed subset X of a polynomial functor P and a linear map $\varphi \in \text{Hom}(U, V)$, we will write $X(\varphi) : X(U) \rightarrow X(V)$ for the restriction of $P(\varphi)$ to $X(U)$. Note that, in particular, the map $\text{GL}(V) \times X(V) \rightarrow X(V)$, $(g, x) \mapsto X(g)(x)$ defines an algebraic action of $\text{GL}(V)$ on $X(V)$, for each $V \in \mathbf{Vec}$. So this paper is concerned with (typically) highly symmetric varieties, related by linear maps coming from linear maps between distinct vector spaces.

If X is a closed subset of P , then the variety $B := X(0)$ is a closed subvariety of $P(0) = P_0$. For each $V \in \mathbf{Vec}$, the zero map $0_{V,0} : V \rightarrow 0$ maps $X(V)$ to $X(0)$, and indeed *onto* $X(0)$, because for any $p_0 \in X(0) \subseteq P(0)$ we have

$$p_0 = P(0_{V,0} \circ 0_{0,V})p_0 = P(0_{V,0})(P(0_{0,V})p_0) \in P(0_{V,0})(X(V))$$

where we have used that P is a functor and X is a closed subset of P . Hence for each V , $X(V)$ is a closed subset of $B \times P'(V)$, where P' is the pure part of P , such that $X(V)$ maps surjectively to B . We will also say that X is a *closed subset of* $B \times P'$.

The following theorem with its corollary implies that a closed subset X of a polynomial functor P can be represented on a computer.

Theorem 2.4.2 [9] *Let P be a polynomial functor and let $X_1 \supseteq X_2 \supseteq \dots$ be a descending chain of closed subsets of P . Then, this chain stabilises, i.e. there exists n_0 such that $X_{n_0} = X_{n_0+1} = \dots$*

Corollary 2.4.3 *For each closed subset $X \subseteq P$, there exists a vector space $U \in \mathbf{Vec}$ with the property that for all $V \in \mathbf{Vec}$ we have*

$$X(V) = \bigcap_{\varphi \in \text{Hom}(V,U)} P(\varphi)^{-1}(X(U)).$$

Proof For every $n \in \mathbb{N}$ consider the closed subset X_n in P defined by

$$X_n(V) := \bigcap_{\varphi \in \text{Hom}(V,K^n)} P(\varphi)^{-1}(X(K^n)).$$

Using that P is a polynomial functor and X is a closed subset, we can see that for all n , $X_n(K^n) = X(K^n)$ and that $X_1 \supseteq X_2 \supseteq \dots$ is a descending chain. By Theorem 2.4.2, this chain stabilises at, say, X_{n_0} , and hence for $n \geq n_0$, $X(K^n) = X_n(K^n) = X_{n_0}(K^n)$. Using that X is a closed subset, we get the same for $n < n_0$. Hence, $X = X_{n_0}$, and the corollary holds for $U = K^{n_0}$. \square

Consequently, if $f_1, \dots, f_k \in K[P(U)]$ are defining equations for $X(U)$, then the pull-backs of the f_i along all linear maps $P(\varphi) : P(V) \rightarrow P(U)$ cut out $X(V)$. We then write $X = \mathcal{V}_P(f_1, \dots, f_k)$. The tuple consisting of P , U , and (f_1, \dots, f_k)

together form a computer representation of X . Slightly more generally, we will often represent X as follows.

Definition 2.4.4 Let B be a finite-dimensional affine variety, P a pure polynomial functor, U a finite-dimensional vector space, and $f_1, \dots, f_k \in K[B \times P(U)]$. The tuple $(B, P, U, f_1, \dots, f_k)$ is called the *implicit representation* of the closed subset X of $B \times P$ defined by

$$X(V) = \{(b, p') \mid \forall \varphi \in \text{Hom}(V, U) \forall i = 1, \dots, r : f_i(b, P(\varphi)p') = 0\}. \quad \diamond$$

Again by Corollary 2.4.3, every closed subset of $B \times P$ admits an implicit representation. Here, we allow that some of the f_i are nonzero elements of $K[B]$, so that X does not map surjectively to B but rather onto a closed subvariety of B .

2.5 Irreducibility

Definition 2.5.1 Let B be a finite-dimensional affine variety, Q a pure polynomial functor, and X a closed subset of $B \times Q$. Then, X is called *irreducible* if $X \neq \emptyset$ and whenever $X = X_1 \cup X_2$ where X_1, X_2 are closed subsets of $B \times Q$ (as always, defined over K), we have $X = X_1$ or $X = X_2$. \(\diamond\)

A straightforward check shows that X is irreducible if and only if $X(V)$ is irreducible for each $V \in \mathbf{Vec}$ (i.e. not the union of two closed proper subsets defined over K); in particular, $B \times Q$ is irreducible if and only if B is irreducible. Note that since K may not be algebraically closed, irreducibility in our sense may not imply irreducibility over \overline{K} .

2.6 Gradings and Ideals

For any polynomial functor $P = P_0 \oplus \dots \oplus P_d$, and any $V \in \mathbf{Vec}$, the coordinate ring $K[P(V)] \cong \bigotimes_{e=0}^d K[P_e(V)]$ is a graded polynomial ring in which the coordinates on $K[P_e(V)]$ are given the degree e . A polynomial f is homogeneous of degree n with respect to this grading if and only if $f(P(t \cdot \text{id}_V)p) = t^n f(p)$ for all $p \in P(V)$. If X is a closed subset of P , then $X(V)$ is preserved under $P(t \cdot \text{id}_V)$ for all t , and hence the ideal of $X(V)$ is homogeneous.

Similarly, for B a finite-dimensional affine variety and P a pure polynomial functor, $K[B \times P(V)]$ has a standard grading in which the elements of $K[B]$ have degree 0, and the ideal of any closed subset X of $B \times P$ is homogeneous.

We find that the coordinate rings $K[X(V)]$, for X a closed subset of a polynomial functor P of degree at most d (or for X a closed subset of $B \times P$ with P a pure polynomial functor of degree d), have standard gradings and are generated in degree at most d . A straightforward computation shows that for each $\varphi \in \text{Hom}(U, V)$, the pull-back $X(\varphi)^* : K[X(V)] \rightarrow K[X(U)]$ is a graded K -algebra homomorphism.

2.7 Morphisms

Definition 2.7.1 Let X, Y be closed subsets of polynomial functors. Then, a morphism $\alpha : X \rightarrow Y$ is given by a morphism $\alpha_V : X(V) \rightarrow Y(V)$ of affine varieties over K for each $V \in \mathbf{Vec}$ such that for all $\varphi \in \text{Hom}(U, V)$ we have $Y(\varphi) \circ \alpha_U = \alpha_V \circ X(\varphi)$. \diamond

By taking for φ scalar multiples of the identity, one finds that the pull-back α_V^* is a graded K -algebra homomorphism.

Lemma 2.7.3 ensures that we can represent morphisms on a computer. First, the following generalises a well-known property of finite-dimensional affine varieties (see [1, Proposition 1.3.22]).

Lemma 2.7.2 *If X is a closed subset of a polynomial functor P and Y is a closed subset of a polynomial functor Q , then any morphism $X \rightarrow Y$ extends to a morphism $P \rightarrow Q$.*

Lemma 2.7.3 *Let X, Y be closed subsets of polynomial functors of degree at most d . Then, a morphism $\alpha : X \rightarrow Y$ is uniquely determined by $\alpha_{K^d} : X(K^d) \rightarrow Y(K^d)$, and this unique determination is algorithmic in the sense that if α_{K^d} is known, then α_V can be computed for any $V \in \mathbf{Vec}$.*

Proof By Lemma 2.7.2, α extends to a morphism $\beta : P \rightarrow Q$, where X, Y are closed subsets in the degree- $\leq d$ polynomial functors P, Q . Now for each $e = 0, \dots, d$, the restriction of $\beta_V^* : Q_e(V)^* \rightarrow K[P(V)]_e$ defines, as V varies, a homomorphism from the polynomial functor $V^* \mapsto Q_e(V)^*$ to the polynomial functor $V^* \mapsto K[P(V)]_e$, both of degree e (except that $K[P(V)]_e$ is not finite-dimensional if $P_0 \neq 0$, but we may replace it by its image). Then, we apply the Friedlander–Suslin lemma to conclude that this homomorphism is uniquely determined by its evaluation at $V = K^d$. The algorithmicity follows from the algorithmicity of the Friedlander–Suslin lemma. \square

Remark 2.7.4 A special instance of the lemma is when $X = P$ and $Y = Q$, with P, Q pure polynomial functors of degree $\leq d$. In this case, the space of morphisms $\mathbf{Map}(P, Q)$ is a finite-dimensional vector space over K , namely, the direct sum for $e = 1, \dots, d$ of the space of GL_d -equivariant linear maps $Q_e(V)^* \rightarrow K[P(V)]_e$, where $V = K^d$. This space will be important to us towards the end of the paper. \diamond

Remark 2.7.5 We will often consider morphisms $\alpha : A \times P \rightarrow B \times Q$, where A, B are finite-dimensional varieties over K and P, Q are pure polynomial functors. Such a morphism decomposes into a morphism $\alpha^{(0)} : A \rightarrow B$ and a morphism $\alpha^{(1)} : A \times P \rightarrow Q$; here we use that α_V^* preserves the degree and that the coordinates on B have degree zero (see Sect. 2.6), and hence their images cannot involve the positive-degree coordinates on P . If A is irreducible, then $\alpha^{(1)}$ can be thought of as a $K(A)$ -valued point of the finite-dimensional affine space $\mathbf{Map}(P, Q)$; this will be a useful point of view later. \diamond

2.8 Shifting

Definition 2.8.1 Any fixed $U \in \mathbf{Vec}$ defines a covariant polynomial functor $\text{Sh}_U : \mathbf{Vec} \rightarrow \mathbf{Vec}$ of degree 1 by $\text{Sh}_U(V) = U \oplus V$ and, for $\varphi : V \rightarrow W, \text{Sh}_U(\varphi) = \text{id}_U \oplus \varphi$. If P is a polynomial functor of degree d , then $\text{Sh}_U P := P \circ \text{Sh}_U$ is also a polynomial functor of degree d , called the *shift over U* of P . \diamond

The top-degree parts of $\text{Sh}_U P$ and P are canonically isomorphic [9, Lemma 14]. If X is a closed subset of P , then $\text{Sh}_U X := X \circ \text{Sh}_U$ is a closed subset of $\text{Sh}_U P$. Note that $(\text{Sh}_U X)(0) = X(U)$, so shifting has the effect of making the finite-dimensional base variety larger. More precisely, if X is a closed subset of $B \times P$ with P a pure polynomial functor, then $\text{Sh}_U X$ is a closed subset of $(B \times P(U)) \times P'$ where P' is the pure part of $\text{Sh}_U P$.

2.9 An Order on Polynomial Functors

Definition 2.9.1 A polynomial functor Q is a subfunctor of P when $Q(V) \subseteq P(V)$ for all V and $Q(\varphi) = P(\varphi)|_{Q(V)}$ for all $\varphi \in \text{Hom}(V, W)$. In this case, the quotient P/Q is defined by $(P/Q)(V) := P(V)/Q(V)$. \diamond

Definition 2.9.2 We call a polynomial functor Q smaller than a polynomial functor P if the two are not isomorphic and for the largest e such that Q_e is not isomorphic to P_e , the former is a quotient of the latter. \diamond

Using the Friedlander–Suslin lemma, one can show that this is a well-founded order on polynomial functors [9, Lemma 12].

2.10 Summary

We have now indicated computer representations for all the mathematical objects that we will need below: finite-dimensional affine varieties, polynomial functors, closed subsets of the latter and morphisms between these. Furthermore, we have introduced two tools in the design and analysis of our algorithms: shifting and a well-founded order on polynomial functors.

3 Parameterisation

3.1 The Result

The goal of this section is to prove the following theorem.

Theorem 3.1.1 *There exists an algorithm **parameterise** that, on input a finite-dimensional affine variety B , a pure polynomial functor Q , a finite-dimensional vector space U over K , and elements $f_i \in K[B \times Q(U)]$ for $i = 1, \dots, k$, computes $(A; P; \beta)$ where A is a finite-dimensional affine variety, P is a pure polynomial functor, and β is a morphism $A \times P \rightarrow B \times Q$, defined over K , such that for each*

finite-dimensional K -vector space V we have

$$\beta(A(\overline{K}) \times \overline{K} \otimes P(V)) = \{(b, q) \in B(\overline{K}) \times \overline{K} \otimes Q(V) \mid \forall i \forall \varphi \in \text{Hom}_K(V, U) : f_i(b, 1 \otimes Q(\varphi)q) = 0\}.$$

Here, $Z(\overline{K})$ means the \overline{K} -points of a finite-dimensional affine variety Z defined over K , i.e. the set of K -algebra homomorphisms $K[Z] \rightarrow \overline{K}$. If Z is just a vector space over K , then this means $\overline{K} \otimes Z$. In what follows, to simplify notation, we will suppress \overline{K} and write $(b, q) \in B \times Q(V)$ when we really mean $(b, q) \in B(\overline{K}) \times Q(V)(\overline{K})$, and for a $\varphi \in \text{Hom}_K(V, W)$ we will write $(b, Q(\varphi)q) \in B \times Q(W)$ instead of $(b, (1 \otimes Q(\varphi))q) \in B(\overline{K}) \times Q(W)(\overline{K})$. This is analogous to Definition 2.4.1 in the case of closed subsets, where $X(V)$ denotes the \overline{K} -points of the given variety.

We write $X := \mathcal{V}_{B \times Q}(f_1, \dots, f_k)$, so that $X(V)$ is the set on the right-hand side above. This is the closed subset of $B \times Q$ that we want to parameterise.

Remark 3.1.2 We allow A to be reducible, even when B is irreducible. The fact that $\alpha : A \times P \rightarrow B \times Q$ as in the theorem exists is [1, Theorem 4.2.5]; see also [2, Proposition 5.6]. If B is indeed irreducible, then for some irreducible component A' of A the restriction of α to $A' \times P$ is dominant into $\mathcal{V}_{B \times Q}(f_1, \dots, f_k)$. \diamond

3.2 Smearing Out Equations

Before proving the theorem, we introduce an algorithm that computes the equations for any single instance $X(V)$.

Proposition 3.2.1 *There exists an algorithm **smear**, that, on the same input as **parameterise** plus a finite-dimensional vector space V , outputs generators of the radical ideal of $X(V) \subseteq K[B \times Q(V)]$.*

Proof The algorithm **smear**($B; Q; U; f_1, \dots, f_k; V$) proceeds as follows: choose identifications $U = K^m$ and $V = K^n$ and construct the generic matrix

$$\psi := \sum_{i=1}^{\dim U} \sum_{j=1}^{\dim V} z_{ij} E_{ij} \in K[(z_{ij})_{ij}] \otimes \text{Hom}_K(V, U),$$

where the E_{ij} form the standard basis of $\text{Hom}_K(V, U)$. Then compute $Q(\psi)y \in K[(z_{ij})_{ij}] \otimes Q(U)$, where $y = (y_1, \dots, y_{\dim Q(V)})^T$ represents a point of $Q(V) \cong K^{\dim Q(V)}$ with variables as coordinates, substitute $Q(\psi)y$ into the f_i , and expand as a polynomial in the z_{ij} with coefficients in $K[B][y_1, \dots, y_{\dim Q(V)}] \cong K[B \times Q(V)]$. Finally, return generators of the radical of the ideal generated by all these coefficients. The correctness of this algorithm follows from the fact that those coefficients span the same space as the images of the f_i under pull-back along the linear maps $Q(\varphi) : Q(V) \rightarrow Q(U)$ for all $\varphi \in \text{Hom}(V, U)$. \square

3.3 The Parameterisation Algorithm

The algorithm **parameterise** is recursive and proceeds as follows; the algorithmic part is written in normal font, text that will be used in the analysis in *italic*. The proofs of termination and correctness are below.

- (1) If $Q = 0$, then compute the variety $A \subseteq B$ defined by f_1, \dots, f_k via a radical ideal computation, return $(A; 0; A \leftrightarrow B)$, and exit.
- (2) Decompose $Q = Q' \oplus R$ where R is an irreducible subfunctor of the top-degree part of Q .

This corresponds to choosing a partition in the tuple representing Q . Let x_1, \dots, x_n be a basis of $R(U)^$. We regard elements in $K[B \times Q(U)] \cong K[B \times Q'(U)] \otimes K[R(U)] \cong K[B \times Q'(U)][x_1, \dots, x_n]$ as polynomials in x_1, \dots, x_n with coefficients in $K[B \times Q'(U)]$.*

- (3) Compute

$$(f'_1, \dots, f'_r) := \mathbf{smear}(B; Q; U; f_1, \dots, f_k; U)$$

and a Gröbner basis (g_1, \dots, g_l) of the elimination ideal in $K[B \times Q'(U)]$ obtained by eliminating all x_i from f'_1, \dots, f'_r .

The elements f'_1, \dots, f'_r generate the radical ideal of $X(U)$ in $B \times (Q'(U) \oplus R(U))$. Let $X'(V)$ be the closure of the image of $X(V)$ under projection $B \times Q(V) \rightarrow B \times Q'(V)$, so X' is a closed subset of $B \times Q'$. Then g_1, \dots, g_l generate the radical ideal of $X'(U)$ by construction; and, as we will see below in the proof of correctness, we have $X' = \mathcal{V}_{B \times Q'}(g_1, \dots, g_l)$.

- (4) In each f_i , replace each coefficient by its normal form modulo g_1, \dots, g_l .
This has the effect that all coefficients that vanish identically on $X'(U)$ are set to zero.
- (5) If all f_i are now zero, then compute

$$(A'; P'; \alpha') := \mathbf{parameterise}(B; Q'; U; g_1, \dots, g_l),$$

output $(A'; P' \oplus R; \alpha' \times \text{id}_R)$, and exit.

- (6) Pick the minimal i for which f_i is nonzero and let x_j be a variable that appears in some monomial in f_i with a nonzero coefficient.
There cannot be degree-0 elements among the f_i , because these would lie in the elimination ideal and hence have been reduced to zero.
- (7) Compute the partial derivative $h := \partial f_i / \partial x_j \in K[B \times Q'(U)][x_1, \dots, x_n]$.
By construction, this h is nonzero and its coefficients, which are a subset of the coefficients of f_i up to some positive integer scalars, do not lie in the ideal generated by g_1, \dots, g_l .
- (8) Compute

$$(A'; P'; \alpha') := \mathbf{parameterise}(B; Q; U; h, f_1, \dots, f_k).$$

- (9) Compute $\tilde{Q} := (\text{Sh}_U Q)/R$ via [12, Exercise 6.11], determine the pure part Q'' of \tilde{Q} , and compute $B'' := (B \times Q(U))[1/h]$.
 So we have $(B \times \text{Sh}_U Q)[1/h] = B'' \times (Q'' \oplus R)$.
- (10) Compute

$$(f_1'', \dots, f_s'') := \mathbf{smear}(B; Q; U; f_1, \dots, f_k; U \oplus U)$$

and replace each f_i'' by its image in $K[B'' \times (Q''(U) \oplus R(U))]$ under the map $K[B \times Q(U \oplus U)] \rightarrow K[B'' \times (Q''(U) \oplus R(U))]$ dual to the inclusion

$$\begin{aligned} B'' \times (Q''(U) \oplus R(U)) &\subseteq B \times Q(U) \times (Q''(U) \oplus R(U)) \\ &= B \times (\text{Sh}_U Q)(U) = B \times Q(U \oplus U). \end{aligned}$$

The elements f_1'', \dots, f_s'' generate the radical ideal of $(\text{Sh}_U X)(U)[1/h]$ in $B'' \times (Q''(U) \oplus R(U))$. As we will see in the proof of correctness, we have $(\text{Sh}_U X)[1/h] = \mathcal{V}_{B'' \times (Q'' \oplus R)}(f_1'', \dots, f_s'')$.

- (11) Using Buchberger’s algorithm to eliminate the coordinates on $R(U)$ from f_1'', \dots, f_s'' , compute equations g_1'', \dots, g_t'' for the projection of the finite-dimensional affine variety $(\text{Sh}_U X)(U)[1/h]$ into $B'' \times Q''(U)$.
 Recall from [9, Lemma 25] that the projection $(B'' \times (Q'' \oplus R)) \rightarrow B'' \times Q''$ restricts to a closed embedding from $(\text{Sh}_U X)[1/h]$ to a closed subset X'' of the latter space. We will see below that $X'' = \mathcal{V}_{B'' \times Q''}(g_1'', \dots, g_t'')$.
- (12) Compute the inverse $\iota : X'' \rightarrow (\text{Sh}_U X)[1/h]$.

By Lemma 2.7.3, this inverse is uniquely determined by its instance ι_V with $\dim V = \deg(Q)$.

- (13) Compute

$$(A''; P''; \alpha'') := \mathbf{parameterise}(B''; Q''; U; g_1'', \dots, g_t''),$$

output

$$(A' \sqcup A''; P' \oplus P''; \alpha' \sqcup (\pi \circ \iota \circ \alpha'')),$$

where $\pi : B \times (\text{Sh}_U Q) \rightarrow B \times Q$, evaluated at V , equals $\text{id}_B \times Q(0_{U,0} \oplus \text{id}_V)$, and exit.

Here, α' is regarded as a map $A' \times (P' \oplus P'') \rightarrow B \times Q$ that ignores the argument from P'' , and similarly α'' ignores the component in P' .

3.4 Termination of Parameterise

Proof of termination of parameterise. If, on some input, **parameterise** would not terminate, then this was due to an infinite chain of recursive calls to itself. In the recursive calls in steps 5 and 13, the polynomial functors Q' and Q'' , respectively, are smaller than Q in the order from Sect. 2.9, whereas Q remains the same in the call in step 8.

Since the order on polynomial functors is well founded, apart from a finite initial segment, the infinite chain consists entirely of consecutive calls in step 8. Now note that, after each such call, $X'(U)$ either remains constant or becomes smaller. As long as it remains constant, i.e. the list (g_1, \dots, g_l) remains constant, the degree in x_1, \dots, x_n of the first equation keeps dropping. Hence after finitely many such steps, $X'(U)$ becomes strictly smaller. It follows that $X'(U)$ becomes smaller infinitely often, but this contradicts the Noetherianity of the finite-dimensional variety $B \times Q'(U)$. \square

3.5 Correctness of Parameterise

Proof that parameterise returns the correct output. We now prove that the output of the algorithm is correct. This is immediate if the algorithm exits in step 1.

If it exits in step 5, then by Lemma 3.5.1, $X = X' \times R$, which is parameterised by $\alpha' \times \text{id}_R$ for a parameterisation α' of X' . Moreover, by the same lemma, X' is defined by g_1, \dots, g_l .

Finally, assume that the algorithm exits in step 13. We need to show that the union of the images of α' and $\pi \circ \iota \circ \alpha''$ (on \overline{K} -points) equals X .

First consider α' , which is computed in step 8. The closed subset $Y := \mathcal{V}_{B \times Q}(f_1, \dots, f_k, h)$ parameterised by α' is clearly \overline{K} -contained in X , so $\alpha' : A' \times P' \rightarrow B \times Q$ has image on \overline{K} -points contained in the \overline{K} -points of X .

Next we argue that $\mathcal{V}_{B'' \times (Q'' \oplus R)}(f''_1, \dots, f''_s)$ is precisely $(\text{Sh}_U X)[1/h]$. First, by construction, f''_1, \dots, f''_s generate the ideal of $(\text{Sh}_U X)[1/h](U)$ in $(B \times Q(U \oplus U))[1/h]$; this shows the containment \supseteq . For the opposite containment, we argue that if V is any finite-dimensional K -vector space and $(b, q) \in B \times Q(U \oplus V)$ has the property that

$$(\forall \varphi \in \text{Hom}(V, U) : (b, Q(\text{id}_U \oplus \varphi)q) \in X(U \oplus U)) \Rightarrow (b, q) \in X(U \oplus V).$$

To show that $(b, q) \in X(U \oplus V)$, since X is defined by its equations in $K[B \times Q(U)]$, it suffices to prove that for each $\psi \in \text{Hom}(U \oplus V, U)$ we have $(b, Q(\psi)q) \in X(U)$. Let $\varphi := \psi|_V \in \text{Hom}(V, U)$ be the restriction of ψ to V . Then,

$$\ker(\text{id}_U \oplus \varphi : U \oplus V \rightarrow U \oplus U) = \ker(\varphi) \subseteq \ker(\psi)$$

and hence the linear map ψ factors as $\psi' \circ (\text{id}_U \oplus \varphi)$ for some $\psi' \in \text{Hom}(U \oplus U, U)$. Hence, since Q is a functor,

$$(b, Q(\psi)q) = (b, Q(\psi')(Q(\text{id}_U \oplus \varphi)q)) \in X(U)$$

because $(b, Q(\text{id}_U \oplus \varphi)q) \in X(U \oplus U)$ and $\text{id}_B \times Q(\psi')$ maps $X(U \oplus U)$ into $X(U)$. This concludes the proof that $\mathcal{V}_{B'' \times (Q'' \oplus R)}(f''_1, \dots, f''_s) = (\text{Sh}_U X)[1/h]$.

Next, by [9, Lemma 25], the projection $B'' \times (Q'' \oplus R) \rightarrow B'' \times Q''$ restricts to an isomorphism embedding from $(\text{Sh}_U X)[1/h]$ to a closed subset of $B'' \times Q''$.

By Lemma 3.5.2, the equations of X'' can be pulled back from the equations of $X''(U)$, so we have indeed $X'' = \mathcal{V}_{B'' \times Q''}(g''_1, \dots, g''_l)$.

Finally, consider the image of $\pi \circ \iota \circ \alpha''$. A straightforward calculation shows that setting $\pi_V := \text{id}_B \times Q(0_{U,0} \oplus \text{id}_V)$ does indeed yield a morphism $B \times \text{Sh}_U Q \rightarrow B \times Q$ that maps $\text{Sh}_U X$ into X . We need to show that if $\alpha'' : A'' \times P'' \rightarrow B'' \times Q''$ is a morphism parameterising X'' , then $\pi \circ \iota \circ \alpha''$ is a morphism $A'' \times P'' \rightarrow B \times Q$ whose image contains all points in X that are not in the subset Y parameterised by α' . So assume that $(b, q) \in X(V) \subseteq B \times Q(V)$ does not lie in $Y(V)$. Then, there exists a linear map $\varphi : V \rightarrow U$ such that $h(b, Q(\varphi)q) \neq 0$. Let $\psi \in \text{Hom}(V, U \oplus V)$ be the map $v \mapsto (\varphi(v), v)$. Then, since $(\text{id}_U \oplus 0_{V,0}) \circ \psi = \varphi$, the point $q' := Q(\psi)q \in X(U \oplus V)$ satisfies

$$h(b, Q(\text{id}_U \oplus 0_{V,0})q') = h(b, Q(\varphi)q) \neq 0,$$

i.e. q' lies in $(\text{Sh}_U X)[1/h]$, which is the image of $\iota \circ \alpha''$. Moreover, since $(0_{U,0} \oplus \text{id}_V) \circ \psi = \text{id}_V$, we have $\pi_V(q') = q$. Hence, q lies in the image of $\pi \circ \iota \circ \alpha''$. This concludes the proof of correctness of **parameterise**. \square

Lemma 3.5.1 *Let Q be a pure polynomial functor and $X \subseteq B \times Q$ a closed subset. Assume that X is defined by its equations in $K[B \times Q(U)]$, let R be a subfunctor of Q , and set $Q' := Q/R$. Define X' as the closure of the image of X under the projection $\text{id}_B \times \pi : B \times Q \rightarrow B \times Q'$. Assume that $X(U) = (\text{id}_B \times \pi_U)^{-1}(X'(U))$. Then, $X(V) = (\text{id}_B \times \pi_V)^{-1}(X'(V))$ for all $V \in \mathbf{Vec}$, so that $X = X' \times R$, and moreover X' is defined by its equations in $K[B \times Q'(U)]$.*

Proof Let $(b, q) \in B \times Q(V)$ satisfy $(b, \pi_V(q)) \in X'(V)$. Then for all $\varphi \in \text{Hom}(V, U)$, we have

$$(b, \pi_U(Q(\varphi)(q))) = (b, Q'(\varphi)(\pi_V(q))) \in X'(U)$$

and hence, by assumption, $(b, Q(\varphi)(q)) \in X(U)$. But then, since X is defined by its equations in $K[B \times Q(U)]$, we have $(b, q) \in X(V)$. This proves the first statement.

For the last statement, suppose that $(b, q') \in B \times Q'(V)$ is such that $(b, Q'(\varphi)(q')) \in X'(U)$ for all $\varphi \in \text{Hom}(V, U)$. Pick any $q \in Q(V)$ with $\pi_V(q) = q'$. Then, the same computation as above shows that $(b, q) \in X(V)$, hence *a fortiori* $(b, q') \in X'(V)$. Hence, X' is defined by its equations in $K[B \times Q'(U)]$. \square

Lemma 3.5.2 *Let Q be a pure polynomial functor, R a (not necessarily irreducible) subfunctor, B an affine variety. Set $Y := B \times Q$, $Y' := B \times (Q/R)$. Let X be a closed subset of Y such that the projection $\pi : Y \rightarrow Y'$ restricts to an isomorphism from X to a closed subset X' of Y' . Let U be a vector space such that X is defined by its equations in $K[Y(U)]$. Then, X' is defined by its equations in $K[Y'(U)]$.*

Proof The idea is to find an inverse to π , i.e. a morphism $\psi : Y' \rightarrow Y$ such that

- (1) $\pi_V \circ \psi_V = \text{id}_{Y'(V)}$
- (2) $\psi|_{X'} = (\pi|_X)^{-1}$

Once we have found this ψ , we are done: Let $y' \in Y'(V)$, such that for every linear map $\varphi : V \rightarrow U$, $Y'(\varphi)(y') \in X'(U)$, and hence, by property (2), $\psi_U(Y'(\varphi)(y')) \in X(U)$. Since ψ is a morphism, we get

$$\psi_U(Y'(\varphi)(y')) = (Y(\varphi)(\psi_V(y'))) \in X(U).$$

Since X is defined by its equations in $K[Y(U)]$, we get that $\psi_V(y') \in X(V)$, and hence, with property (1) of ψ , $\pi_V(\psi_V(y')) = y' \in X'(V)$.

By the Friedlander–Suslin Lemma (Lemma 2.3.2 and proof of Lemma 2.7.3), it is enough to find a GL_d -equivariant map $\tilde{\psi} : Y'(K^d) \rightarrow Y(K^d)$, where d is the degree of Q , that fulfils properties (1) and (2) (with $V = K^d$ and $\psi_{K^d} = \tilde{\psi}$).

It is easy to find a possibly non- GL_d -equivariant map $\hat{\psi} : Y'(K^d) \rightarrow Y(K^d)$ with properties (1) and (2): Note that $X'(K^d)$ is a closed subvariety of the affine space $\mathbb{A}_B^{n'}$, where $n' = \dim(Q(K^d)/R(K^d))$ and $X(K^d)$ is a closed subvariety of the affine space \mathbb{A}_B^n , where $n = \dim(Q(K^d))$. By basic properties of affine spaces, the map $(\pi_{K^d}|_{X(K^d)})^{-1}$ extends to a morphism $\hat{\psi}$ from the ambient affine space $\mathbb{A}_B^{n'}$ of $X'(K^d)$ to the ambient affine space \mathbb{A}_B^n of $X(K^d)$. Now $\hat{\psi}$ lives in some finite-dimensional space equipped with a linear action of GL_d given by $(g, \eta) \mapsto Y(g) \circ \eta \circ Y'(g^{-1})$. Since this space is an algebraic group representation of GL_d and GL_d is linearly reductive, there exists a GL_d -equivariant linear map ρ from the space onto its subspace of GL_d -invariant elements—the so-called *Reynolds operator*. Then, $\tilde{\psi} := \rho(\hat{\psi})$ is GL_d -equivariant, and we claim that it still satisfies (1) and (2).

By the Lefschetz principle, it is sufficient to check this when $K = \mathbb{C}$. In this case, $\tilde{\psi}$ takes the more explicit form

$$\tilde{\psi} := \int_{U_d(\mathbb{C})} Y(\varphi) \circ \hat{\psi} \circ Y'(\varphi)^{-1} d\varphi$$

where $U_d(\mathbb{C}) \subseteq GL_d(\mathbb{C})$ is the unitary group, and the integral is over the Haar measure (see, e.g. [21]) with $\int_{U_d(\mathbb{C})} d\varphi = 1$ (note that the Haar integral needs to be taken over a compact group, so we have to use $U_d(\mathbb{C})$ instead of $GL_d(\mathbb{C})$).

Indeed, by construction, the morphism $\tilde{\psi}$ defined by this integral is U_d -equivariant. To see that it is, indeed, GL_d -equivariant, one uses that $U_d(\mathbb{C})$ is Zariski dense in $GL_d(\mathbb{C})$ [21, Section 8.6.1]. This is a version of Weyl’s well-known *unitary trick*.

The following chain of equations now verifies that $\tilde{\psi}$ meets property (1):

$$\begin{aligned} \pi_{\mathbb{C}^d} \circ \tilde{\psi} &= \pi_{\mathbb{C}^d} \circ \int_{U_d(\mathbb{C})} Y(\varphi) \circ \hat{\psi} \circ Y'(\varphi)^{-1} d\varphi \\ &= \int_{U_d(\mathbb{C})} \pi_{\mathbb{C}^d} \circ Y(\varphi) \circ \hat{\psi} \circ Y'(\varphi)^{-1} d\varphi \\ &= \int_{U_d(\mathbb{C})} Y'(\varphi) \circ \pi_{\mathbb{C}^d} \circ \hat{\psi} \circ Y'(\varphi)^{-1} d\varphi = \int_{U_d(\mathbb{C})} \text{id}_{Y'(\mathbb{C}^d)} d\varphi. \end{aligned}$$

For property (2), let $x \in X'(\mathbb{C}^d)$. Then,

$$\begin{aligned} \tilde{\psi}(x) &= \int_{U_d(\mathbb{C})} Y(\varphi)(\underbrace{\hat{\psi}(Y'(\varphi)^{-1}(x))}_{\in X'(\mathbb{C}^d)}) \, d\varphi \\ &= \int_{U_d(\mathbb{C})} Y(\varphi)((\pi_{\mathbb{C}^d}|_{X(\mathbb{C}^d)})^{-1}(Y'(\varphi)^{-1}(x))) \, d\varphi \\ &= \int_{U_d(\mathbb{C})} Y(\varphi)(Y(\varphi)^{-1}((\pi_{\mathbb{C}^d}|_{X(\mathbb{C}^d)})^{-1}(x))) \, d\varphi = (\pi_{\mathbb{C}^d}|_{X(\mathbb{C}^d)})^{-1}(x). \end{aligned}$$

Note that for the second to last equality we have used that $(\pi|_X)^{-1}$ is a morphism, as is easy to verify. □

4 Implicitisation

4.1 The Result

The goal of this section is to prove the following more precise version of the Main Theorem.

Theorem 4.1.1 *There exists an algorithm **implicitise** that, on input finite-dimensional affine varieties A, B , pure polynomial functors P, Q , and a morphism $\alpha : A \times P \rightarrow B \times Q$, computes a $U \in \mathbf{Vec}$ and elements $f_1, \dots, f_k \in K[B \times Q(U)]$ that define the image closure of α , i.e. such that, for all $V \in \mathbf{Vec}$, we have*

$$\begin{aligned} &\overline{\alpha(A(\overline{K}) \times \overline{K} \otimes P(V))} \\ &= \{(b, q) \in B(\overline{K}) \times \overline{K} \otimes Q(V) \mid \forall i \forall \varphi \in \text{Hom}_K(V, U) : f_i(b, 1 \otimes Q(\varphi)q) = 0\}. \end{aligned}$$

4.2 The Implicitisation Algorithm

The algorithm **implicitise**(A, P, B, Q, α) is run on countably many parallel processors: the one where the original call is handled, plus countably many labelled $0, 1, 2, 3, \dots$ where call to an auxiliary procedure **certify** is made. The latter has the following specification.

Proposition 4.2.1 *There exists a procedure **certify** that, on input finite-dimensional affine varieties A, A', B , pure polynomial functors P, P', Q , and morphisms $\alpha : A \times P \rightarrow B \times Q$ and $\alpha' : A' \times P' \rightarrow B \times Q$, has the following behaviour: for all $V \in \mathbf{Vec}$ we have*

$$\alpha'(A'(\overline{K}) \times (\overline{K} \otimes P'(V))) \subseteq \overline{\alpha(A(\overline{K}) \times (\overline{K} \otimes P(V)))},$$

then **certify**($B; Q; A; P; \alpha; A'; P'; \alpha'$) terminates and returns “true”. Otherwise, it does not terminate.

The proof of Proposition 4.2.1 is deferred to Sect. 5. We can now present the steps for **implicitise**(A, P, B, Q, α).

- (1) Set $n := 0$.
- (2) While no instance of **certify** has returned “true”, perform steps (3)–(6):
- (3) Set $U := K^n$ and, by classical elimination, compute defining equations $f_{n,1}, \dots, f_{n,k_n} \in K[B \times Q(U)]$ for the image closure of α_U . *This is a finite-dimensional affine variety.*
- (4) Compute $(A_n, P_n, \alpha_n) := \mathbf{parameterise}(B; Q; U; f_{n,1}, \dots, f_{n,k_n})$.
- (5) On the n -th processor, start **certify**($B; Q; A; P; \alpha; A_n; P_n; \alpha_n$).
- (6) Set $n := n + 1$.
- (7) if the m -th processor has returned “true”, then return $(K^m; f_{m,1}, \dots, f_{m,k_m})$.

4.3 Correctness and Termination of implicitise

Proof of Theorem 4.1.1 By Corollary 2.4.3, there exists a value of n such that the tuple $(K^n; f_{n,1}, \dots, f_{n,k_n})$ computed in iteration n is a correct output for **implicitise**. It then follows that α_n , which, by virtue of **parameterise**, parameterises the closed subset of $B \times Q$ defined by $f_{n,1}, \dots, f_{n,k_n}$, has its image contained in the closure of the image of α . Hence, by Proposition 4.2.1, the n -th call to **certify** terminates and returns “true”. This shows that **implicitise** terminates.

Next, when it terminates with output $(K^m; f_{m,1}, \dots, f_{m,k_m})$, then this is because the image of α_m , which equals the closed subset of $B \times Q$ defined by $f_{m,1}, \dots, f_{m,k_m}$, is contained in the image closure of α . Since, conversely, the image closure of α is contained in the closed subset defined by $f_{m,1}, \dots, f_{m,k_m}$, the output is correct. \square

5 Certifying Inclusion of Image Closures

5.1 An Instructive Example

Example 5.1.1 Let $\alpha : (S^1)^2 \rightarrow S^3$ be the morphism defined by $\alpha(u, v) = u^3 + v^3$. The image closure of α is the set of symmetric three-tensors of *border Waring rank* at most 2. On the other hand, let $\beta : (S^1)^2 \rightarrow S^3$ be the morphism defined by $\beta(u, v) = 6u^2v$. Then, we have

$$\begin{aligned} \beta(u, v) &= 6u^2v = \lim_{t \rightarrow 0} [(t^2v + t^{-1}u)^3 + (t^2v - t^{-1}u)^3] \\ &= \lim_{t \rightarrow 0} \alpha(t^2v + t^{-1}u, t^2v - t^{-1}u) \end{aligned}$$

and this implies that $\text{im}(\beta) \subseteq \overline{\text{im}(\alpha)}$. However, $\text{im}(\beta) \not\subseteq \text{im}(\alpha)$ —indeed, it is well known that the Waring rank of cubics of the form u^2v with u, v linearly independent vectors is equal to 3. \diamond

This example shows a *certificate* for $\text{im } \beta \subseteq \overline{\text{im } \alpha}$, namely, that β is a limit of a composition $\alpha \circ \gamma_t$ with

$$\gamma_t : (S^1)^2 \rightarrow (S^1)^2, (u, v) \mapsto (t^2v + t^{-1}u, t^2v - t^{-1}u).$$

Roughly speaking, whenever $\text{im}(\beta)$ is contained in $\overline{\text{im}(\alpha)}$, where β, α are morphisms into $B \times Q$, with B a finite-dimensional variety and Q a pure polynomial functor, there is a certificate of this inclusion such as the one above—see below for the precise statement. However, we are not aware of any a priori lower bound on the (negative) exponents of t in such a certificate. This is why, in Proposition 4.2.1, the procedure **certify** does not terminate when no certificate exists.

5.2 An Excursion to Infinite Dimensions

We collect some material on GL-varieties. The results stated here appeared in [2, 3] or can directly be derived from results there.

Given a finite-dimensional variety A and a pure polynomial functor P , we construct the inverse limit $\lim_{\leftarrow n} (A \times P(K^n)) = A \times P_\infty$, where the projections $P(K^{n+1}) \rightarrow P(K^n)$ are of the form $P(\pi)$ with π the standard projection $K^{n+1} \rightarrow K^n$. Rather than as a set of K -valued points, we will regard $A \times P_\infty$ as a reduced, affine K -scheme, namely, the spectrum of the ring $K[A] \otimes_K R$, where R is the symmetric algebra of the countable-dimensional vector space $\lim_{n \rightarrow \infty} P(K^n)^*$. The group $\text{GL} := \bigcup_{n=0}^\infty \text{GL}_n(K)$ acts on $A \times P_\infty$ by means of automorphisms and $A \times P_\infty$ is a GL-variety in the sense of [2]. More generally, if X is a closed subset of a polynomial functor, then the inverse limit X_∞ of all $X(K^n)$ is a GL-variety. The association $X \mapsto X_\infty$ is an equivalence of categories with the category of affine GL-varieties, which sends a morphism $\alpha : X \rightarrow Y$ to a GL-equivariant morphism $\alpha_\infty : X_\infty \rightarrow Y_\infty$ of affine schemes over K .

Let $\alpha : A \times P \rightarrow B \times Q$ and $\alpha' : A' \times P' \rightarrow B \times Q$ be morphisms as in Proposition 4.2.1. Then, the following two statements are equivalent:

- (1) $\overline{\text{im}(\alpha_\infty)} \supseteq \text{im}(\alpha'_\infty)$ and
- (2) $\overline{\text{im}(\alpha_V)} \supseteq \text{im}(\alpha'_V)$ for all $V \in \mathbf{Vec}$.

It is (2) which we want to certify in **certify**($B; Q; A; P; \alpha; A', P', \alpha'$). From now on, we assume that A, A' are irreducible—in the procedure **certify** we will reduce to this case.

Now the article by Bik, Eggermont, Snowden and the second author [3] contains the following useful criterion for (1). Let a' be the generic point of A' and let $p' \in P'_\infty(K)$ be a point whose GL-orbit is dense in P_∞ (such points exist, see [2]). Let $x := \alpha'_\infty(a', p')$; this is an Ω -point of $B \times Q_\infty =: X_\infty$, where $\Omega = K(A')$, and the GL-orbit of x is dense in $\text{im}(\alpha'_\infty)$. Write $Y_\infty := A \times P_\infty$.

Theorem 5.2.1 (Theorem 6.6 from [3]) *We have $\text{im}(\alpha'_\infty) \subseteq \overline{\text{im}(\alpha_\infty)}$ if and only if there exists a finite-dimensional field extension $\tilde{\Omega}$ of Ω and a bounded $\tilde{\Omega}((t))$ -point $y(t)$ of Y_∞ such that $\lim_{t \rightarrow 0} \alpha_\infty(y(t)) = x$.*

Here *bounded* means that the exponents of t in the countably many coordinates of the P_∞ -component of $y(t)$ are uniformly bounded from below, see [3, Definition 6.1]. This ensures that $\alpha_\infty(y(t))$ is a well-defined $\tilde{\Omega}((t))$ -point of X_∞ . The theorem says that we can choose $y(t)$ such that $\alpha_\infty(y(t))$ is in fact an $\tilde{\Omega}[[t]]$ -point of X_∞ and that setting t to zero yields x .

The procedure **certify** should certify the existence of $y(t)$. To this end, we will narrow down the space in which to search for $y(t)$ to an increasing chain of finite-dimensional varieties. First, we will show that $y(t)$ needs not use more of P_∞ than can be obtained by applying morphisms to p' . To do so (see Proposition 5.4.1 below), we now introduce *systems of variables* in Schur functors.

5.3 Systems of Variables in Schur Functors

Fix a field extension Ω of K . For every nonempty partition λ , $S_{\lambda,\infty}$ is an affine scheme over K whose Ω -valued points form an Ω -vector space of uncountable dimension. Let $V_\lambda \subseteq S_{\lambda,\infty}(\Omega)$ be the set of all points s for which there exist k , partitions μ_1, \dots, μ_k with $0 < |\mu_i| < |\lambda|$, an Ω -valued point α of $\mathbf{Map}(S_{\mu_1} \oplus \dots \oplus S_{\mu_k}, S_\lambda)$, and an Ω -valued point p of $S_{\mu_1,\infty} \times \dots \times S_{\mu_k,\infty}$ such that $\alpha_\infty(p) = s$. (Recall the definition of $\mathbf{Map}(P', P)$ from Remark 2.7.4.)

Example 5.3.1 If $\lambda = (2)$, then $S_{\lambda,\infty}(\Omega)$ is the space of infinite-by-infinite symmetric matrices with entries in Ω , and V_λ is the subspace of matrices of finite rank. \diamond

Now V_λ is a proper Ω -vector subspace of $S_{\lambda,\infty}(\Omega)$, and we choose any Ω -basis $(\xi_{\lambda,i})_{i \in I_\lambda}$ of a vector space complement to V_λ in $S_{\lambda,\infty}(\Omega)$, where I_λ is an uncountable index set. We call the $\xi_{\lambda,i}$ *variables*. We choose these variables for every λ and write ξ for the resulting uncountable tuple; ξ is what we call a *system of variables* (over Ω) for all Schur functors. If f is an Ω -valued point of $\mathbf{Map}(S_{\mu_1} \oplus \dots \oplus S_{\mu_k}, S_\lambda)$ and we fix indices $i_1 \in I_{\mu_1}, \dots, i_k \in I_{\mu_k}$, then we will write $f(\xi)$ for $f(\xi_{\mu_1,i_1}, \dots, \xi_{\mu_k,i_k})$. This is slight abuse of notation, since it is not apparent from the formula $f(\xi)$ which indices were chosen, but the notation is compatible with the notation $f(x)$ for a polynomial that uses finitely many of an uncountable set of variables x .

Remark 5.3.2 We need to fix the extension Ω first and then choose the system of variables. It is not true that a system of variables chosen over K is also a system of variables over field extensions of K , as $S_{\lambda,\infty}(K) \otimes_K \Omega = S_{\lambda,\infty}(\Omega)$ only when Ω is finite-dimensional over K . \diamond

The following proposition follows readily from the material in [2].

Proposition 5.3.3 *Let $S = S_{\mu_1} \oplus \dots \oplus S_{\mu_k}$ be a pure polynomial functor, and let $s = (s_1, \dots, s_k) \in S_\infty(\Omega)$. Then, the s_i can be chosen as part of a system of variables if and only if the GL-orbit of s is dense in $S_\infty(\Omega)$.*

The following theorem expresses that the variables in a system are independent and generate all vectors in all Schur functors.

Theorem 5.3.4 Fix a field extension Ω of K and a system of variables ξ over Ω for all Schur functors. Then for every nonempty partition λ and any $p \in S_{\lambda, \infty}(\Omega)$, there exist partitions $\lambda_1, \dots, \lambda_k$ and an Ω -valued point f of $\mathbf{Map}(S_{\lambda_1} \oplus \dots \oplus S_{\lambda_k}, S_\lambda)$ and variables ξ_{λ_j, i_j} for $j = 1, \dots, k$ and $i_j \in I_{\lambda_j}$ such that $p = f_\infty(\xi_{\lambda_1, i_1}, \dots, \xi_{\lambda_k, i_k})$. Moreover, if f really depends on all ξ_{λ_j, i_j} in the sense that replacing one of them by zero changes the outcome, then, up to permutations of $\{1, \dots, k\}$, the partitions λ_j , the variables ξ_{λ_j, i_j} , and f are unique.

Proof The existence of f follows by induction on $|\lambda|$: we may write p

$$p = c_1 \xi_{\lambda, i_1} + c_2 \xi_{\lambda, i_2} + \dots + c_l \xi_{\lambda, i_l} + \tilde{p}$$

with $\tilde{p} \in V_\lambda$ and (unique) $i_1, \dots, i_l \in I_\lambda$ and $c_1, \dots, c_l \in \Omega$. Now $\tilde{p} = \alpha_\infty(q)$, for a suitable α , where q is an Ω -valued point of $S_{\mu_1, \infty} \times \dots \times S_{\mu_k, \infty}$ with $0 < |\mu_i| < |\lambda|$ for all i . By the induction hypothesis, the components $q_i, i = 1, \dots, k$ of q are of the form $f_{i, \infty}(\xi)$, and then p equals

$$\alpha_\infty(f_{1, \infty}(\xi), \dots, f_{k, \infty}(\xi)) + \sum_{i=1}^l c_i \xi_{\lambda, i},$$

so that $f := \alpha(f_1, \dots, f_k) + \sum_{i=1}^l c_i \text{id}_{S_\lambda}$ does the trick for the obvious choice of variables.

For uniqueness, it suffices to show that if f is a nonzero Ω -valued point of $\mathbf{Map}(S_{\mu_1} \oplus \dots \oplus S_{\mu_k}, S_\lambda)$ and $\xi_{\mu_1, i_1}, \dots, \xi_{\mu_k, i_k}$ are distinct variables, then $f_\infty(\xi) \neq 0$. Indeed, by Proposition 5.3.3, the GL-orbit of $(\xi_{\mu_1, 1}, \dots, \xi_{\mu_k, i_k})$ is dense, and f_∞ is GL-equivariant, so that $f_\infty(\xi) = 0$ implies that $f = 0$, a contradiction. \square

Remark 5.3.5 Our notion of systems of variables is inspired by [10]. Indeed, If we restrict a system of variables in Schur functors to the symmetric powers S^d with $d = 0, 1, \dots$, then the proof of [10, Theorem 1.1] shows that these are algebraically independent (over Ω) generators of the ring $\bigoplus_d S_\infty^d(\Omega)$, and that this ring is therefore abstractly isomorphic to a polynomial ring. This fact can also be deduced from Theorem 5.3.4 above. \diamond

5.4 Narrowing Down the Search for $y(t)$

In this section, we retain the notation from Sect. 5.2. The following diagram represents the situation:

$$\begin{array}{ccc}
 & & (a', p') \in A' \times P' \\
 & & \downarrow \alpha' \\
 y(t) = (a(t), p(t)) \in (A \times P_\infty)(\tilde{\Omega}((t))) & \xrightarrow{\alpha_\infty} & x = (b, q) \in (B \times Q_\infty)(\Omega)
 \end{array}$$

where $\Omega = K(A')$, a' is the generic point of A' , p' is a K -point of P_∞ with dense GL-orbit, and we want to certify the existence of $y(t)$, defined over a finite extension $\tilde{\Omega}$ of Ω , such that $\lim_{t \rightarrow 0} \alpha_\infty(y(t)) = x$.

Proposition 5.4.1 *If a $y(t)$ as in Theorem 5.2.1 exists, then it can be chosen of the form $(a(t), \gamma(t)_\infty(p'))$ with $a(t) \in A(\tilde{\Omega}(t))$ and $\gamma(t)$ a $\tilde{\Omega}[t, t^{-1}]$ -valued point of the finite-dimensional affine space $\mathbf{Map}(P', P)$.*

Proof Write $y(t) = (a(t), p(t))$. First, terms in $p(t)$ of sufficiently high degree in t do not contribute to $\lim_{t \rightarrow 0} \alpha_\infty(a(t), p(t))$, so we may truncate $p(t)$ and assume that it is a finite sum $\sum_{d=m_1}^{m_2} t^d p_d$ where each $p_d \in P_\infty(\tilde{\Omega})$.

Now write $P' = S_{\lambda_1} \oplus \dots \oplus S_{\lambda_k}$, where the λ_i are partitions. Accordingly, decompose $p' = (p'_1, \dots, p'_k)$ with $p'_i \in S_{\lambda_i, \infty}$. Over the field extension $\tilde{\Omega}$ of K , choose a system of variables ξ in such a manner that p'_1, \dots, p'_k are among these variables; this can be done by Proposition 5.3.3 because $\text{GL} \cdot p'$ is dense in P'_∞ . Then, by Theorem 5.3.4, we have $p_d = f_{d, \infty}(\xi)$ for all d , where f_d is an (essentially unique) morphism into P with coefficients in $\tilde{\Omega}$.

Recall from Remark 2.7.5 that α splits as $\alpha^{(0)} : A \rightarrow B$ and $\alpha^{(1)} : A \times P \rightarrow Q$, and similarly for α' . The limit $\lim_{t \rightarrow 0} \alpha^{(1)}(a(t), p(t))$ equals

$$g_\infty(p_{m_1}, \dots, p_{m_2}) = g_\infty(f_{m_1, \infty}(\xi), \dots, f_{m_2, \infty}(\xi))$$

for some $\tilde{\Omega}$ -point g of $\mathbf{Map}(P^{m_2-m_1+1}, Q)$. On the other hand, by the choice of $y(t)$, it equals $(\alpha')_\infty^{(1)}(p')$. In the latter expression, only the variables p'_1, \dots, p'_k appear. By the uniqueness statement in Theorem 5.3.4, the same must apply to $g_\infty(f_{m_1, \infty}(\xi), \dots, f_{m_2, \infty}(\xi))$.

Therefore, replacing each p_d by $\tilde{p}_d := f_{d, \infty}(p', 0)$, where all variables not among the variables p'_1, \dots, p'_k are set to zero, yields a $\tilde{y}(t)$ with the same property as $y(t)$: $\lim_{t \rightarrow 0} \alpha_\infty(\tilde{y}(t)) = x$. Now $\gamma(t) := \sum_d t^d f_d(\cdot, 0)$ is the desired $\tilde{\Omega}[t, t^{-1}]$ -valued point of $\mathbf{Map}(P', P)$. □

Recall from Remark 2.7.5 that $(\alpha')^{(1)}$ can be regarded an $\mathbf{Map}(P', Q)$. Similarly, $\alpha^{(1)}(a(t), \cdot)$ can be regarded an $\tilde{\Omega}(t)$ -point of $\mathbf{Map}(P, Q)$.

Lemma 5.4.2 *A point $(a(t), \gamma(t)_\infty(p'))$ as in Proposition 5.4.1 satisfies the property $\lim_{t \rightarrow 0} \alpha_\infty(a(t), \gamma(t)_\infty(p')) = \alpha'(a', p') =: (b, q) \in (B \times Q_\infty)(\Omega)$ if and only if, first, $\lim_{t \rightarrow 0} \alpha^{(0)}(a(t)) = b$ and, second, the $\tilde{\Omega}(t)$ -point $\alpha^{(1)}(a(t), \cdot) \circ \gamma(t)$ of $\mathbf{Map}(P', Q)$ satisfies*

$$\lim_{t \rightarrow 0} \alpha^{(1)}(a(t), \cdot) \circ \gamma(t) = (\alpha')^{(1)};$$

an equality of $\tilde{\Omega}$ -points in $\mathbf{Map}(P', Q)$.

Proof The statement “if” is immediate, by substituting p' ; and the statement “only if” follows from the fact that the GL-orbit of p' is dense in P'_∞ . □

5.5 Greenberg’s Approximation Theorem

We have almost arrived at a countable chain of finite-dimensional varieties in which we can look for $y(t)$. The only problem is that the point $a(t) \in A(\tilde{\Omega}(t))$ does not yet have a finite representation. For concreteness, assume that A is given by a prime ideal $I = (f_1, \dots, f_r)$ in $K[x_1, \dots, x_m]$, B is embedded in K^n , and $\alpha^{(0)} : A \rightarrow B$ is the restriction of some polynomial map $\alpha^{(0)} : K^m \rightarrow K^n$.

Then, $a(t)$ is an m -tuple in $\tilde{\Omega}(t)^m$, and together with $\gamma(t)$ it is required to satisfy the following properties from Lemma 5.4.2:

- (i) $f_i(a(t)) = 0$ for $i = 1, \dots, r$;
- (ii) $\lim_{t \rightarrow 0} \alpha^{(0)}(a(t)) = b$;
- (iii) and $\lim_{t \rightarrow 0} \alpha^{(1)}(a(t), \cdot) \circ \gamma(t) = (\alpha')^{(1)}$.

Suppose that we fix a lower bound $-d_1$, with $d_1 \in \mathbb{Z}_{\geq 0}$, on the exponents of t appearing in $a(t)$ or in $\gamma(t)$. From the data of α and d_1 , one can compute a bound $d_2 \in \mathbb{Z}_{\geq 0}$ such that the validity of (ii) and (iii) does not depend on the terms in $a(t)$ or $\gamma(t)$ with exponents $> d_2$. However, (i) does depend on all (typically infinitely many) terms of $a(t)$. Here, Greenberg’s approximation theorem comes to the rescue. As this theorem requires formal power series rather than Laurent series, we put $\tilde{a}(t) := t^{d_1} a(t)$. Accordingly, replace each f_i by $\tilde{f}_i := t^e f_i(t^{-d_1} x_1, \dots, t^{-d_1} x_n)$ where e is large enough such that all coefficients of \tilde{f}_i for all i are in $\tilde{\Omega}[[t]]$. Note that $a(t)$ is a root of all f_i if and only if $\tilde{a}(t)$ is a root of all \tilde{f}_i .

Theorem 5.5.1 (Greenberg, [13]) *There exist numbers $N_0 \geq 1, c \geq 1, s \geq 0$ such that for all $N \geq N_0$ and $\bar{a}(t) \in \tilde{\Omega}[[t]]^n$ with $\tilde{f}_i(\bar{a}(t)) \equiv 0 \pmod{t^N}$ for all $i = 1, \dots, r$ there exists an $\tilde{a}(t) \in \tilde{\Omega}[[t]]^n$ such that $\tilde{a}(t) \equiv \bar{a}(t) \pmod{t^{\lceil \frac{N}{c} \rceil - s}}$ and moreover $\tilde{f}_i(\tilde{a}(t)) = 0$ for all i . Moreover, N_0, c, s can be computed from $\tilde{f}_1, \dots, \tilde{f}_r$.*

As a matter of fact, the computability, which is crucial to our work, is only implicit in [13]; it is made explicit in the overview paper [23].

Corollary 5.5.2 *There exist natural numbers d_2, N_1 , which can be computed from d_1 and f_1, \dots, f_r, α , such that the following statements are equivalent:*

- (1) *a pair $a(t), \gamma(t)$ with properties (i)–(iii) exists that has no exponents of t smaller than $-d_1$;*
- (2) *a pair $a(t), \gamma(t)$ exists with all exponents of t in the interval $\{-d_1, \dots, d_2\}$ that satisfies (ii) and (iii), and that satisfies (i) modulo t^{N_1} .*

Proof The implication (1) \Rightarrow (2) holds for any choice of N_1 if d_2 is chosen large enough so that the terms of $a(t), \gamma(t)$ with degree $> d_2$ in t do not affect (ii),(iii), and do not contribute to the terms of degree $< N_1$ in $f_i(a(t))$ for any i .

For the converse, first we compute \tilde{f}_i and e as above; they depend on the choice of d_1 . Then, we compute N_0, c, s as in Greenberg’s theorem. Compute $N_1 \geq N_0 - e$ such that terms in $a(t), \gamma(t)$ in which t has exponent at least $\lceil \frac{N_1+e}{c} \rceil - s - d_1$ do not affect properties (ii)–(iii), and then compute d_2 as in the first paragraph.

Given a pair $a(t), \gamma(t)$ as in (2), set $\bar{a}(t) := t^{d_1} a(t)$. Then, for each i ,

$$\tilde{f}_i(\bar{a}(t)) = t^e f_i(a(t)) \equiv 0 \pmod{t^{N_1+e}}.$$

Then, since $N_1 + e \geq N_0$, Greenberg’s theorem yields $\tilde{a}(t) \in \tilde{\Omega}[[t]]^n$ such that $\tilde{f}_i(\tilde{a}(t)) = 0$ for all i and such that

$$\tilde{a}(t) = \bar{a}(t) \pmod{t^{\lceil \frac{N_1+e}{c} \rceil - s}}.$$

Now set $a_1(t) := t^{-d_1} \tilde{a}(t)$, so that $f_i(a_1(t)) = 0$ for all i —this is property (i)—and

$$a_1(t) \equiv a(t) \pmod{t^{\lceil \frac{N_1+e}{c} \rceil - s - d_1}}.$$

Since the terms of $a(t)$ with exponent of degree at least $\lceil \frac{N_1+e}{c} \rceil - s - d_1$ do not affect (ii) and (iii), the pair $a_1(t), \gamma(t)$ also satisfy these conditions. □

5.6 The Procedure Certify

To compute **certify**($B; Q; A; P; \alpha; A'; P'; \alpha'$), we proceed as follows. For convenience, we again assume that we have sufficiently many processors working in parallel.

- (1) If A and A' are not both irreducible, decompose A into irreducible components A_i and A' into irreducible components A'_j , and assign the computation of **certify**($B; Q; A_i; P; \alpha|_{A_i \times P}; A'_j, P', \alpha'|_{A'_j \times P'}$) for all i, j to distinct processors. As soon as for each j there exists at least one i such that the computation returns “true”, return “true”. *So in what follows we may assume that A, A' are irreducible. They are given by prime ideals $I \subseteq K[x_1, \dots, x_n]$ and $J \subseteq K[y_1, \dots, y_m]$, respectively.*
- (2) Let f_1, \dots, f_r be generators of I .
- (3) Compute $b := (\alpha')^{(0)}(a')$ where a' is the generic point of A' . So a' is just the m -tuple $(y_1 + J, \dots, y_m + J) \in \Omega^m$, where Ω is the fraction field of $K[y_1, \dots, y_m]/J$.
- (4) Compute the Ω -valued point $(\alpha')^{(1)}$ of $\mathbf{Map}(P', Q)$.
- (5) Construct a K -basis $\gamma_1, \dots, \gamma_m$ of the vector space $\mathbf{Map}(P', P)$.
- (6) Set $d_1 := 0, r :=$ “false”.
- (7) While not r , perform the steps (8)–(10):
- (8) From α and d_1 , compute the natural numbers N_1, d_2 from Corollary 5.5.2 and make the Ansatz $\gamma(t) = \sum_{i=1}^m c_i(t)\gamma_m$ where $c_i(t)$ is a linear combination of t^{-d_1}, \dots, t^{d_2} with coefficients to be determined in an extension of Ω ; and the Ansatz $a(t) = (a_1(t), \dots, a_n(t))$, where a_i is also a linear combination of t^{-d_1}, \dots, t^{d_2} with coefficients to be determined.
- (9) The desired properties of $(a(t), \gamma(t))$ from the second item of Corollary 5.5.2 translate into a system of polynomial equations for the $(m + n) \cdot (d_2 + d_1 + 1)$ coefficients of the $c_i(t)$ and the $a_i(t)$. By a Gröbner basis computation, test whether a solution exists over an algebraic closure of Ω . If so, set $r :=$ “true”.
- (10) Set $d_1 := d_1 + 1$.
- (11) Return “true”.

Proof of Proposition 4.2.1 The first step is justified by the observation that the image closure of α contains the image of α' if and only if for each j , the image of $\alpha'|_{A'_j \times P'}$ is contained in the image closure of some $\alpha|_{A_i \times P}$.

If the image closure of α contains the image of α' , then by Theorem 5.2.1, Proposition 5.4.1, Lemma 5.4.2, and Corollary 5.5.2, the procedure **certify** terminates and returns “true”. Otherwise, by the same results, the system of equations in step (9) does not have a solution, and the procedure does not terminate. \square

6 Conclusion

The Noetherianity theorem [9] and the unirationality theorem [2] imply that closed subsets of polynomial functors admit two different descriptions: an implicit description by finitely many equations, and a description as the image (closure) of some morphism. In this paper, we have described highly nontrivial algorithms **parameterise** and **implicitise** that translate between these two descriptions. Interestingly, **implicitise** requires calls to **parameterise** as well as Greenberg’s approximation theorem for a stopping criterion. We do not expect these two algorithms to be implemented in full generality anytime soon, but we do believe that their structure may guide us in finding equations for closed subsets of polynomial functors such as “cubics of q -rank ≤ 2 ”.

Acknowledgements The second author thanks Arthur Bik, Rob Eggermont, and Andrew Snowden for numerous discussions on polynomial functors, especially on the topic of systems of variables from Sect. 5.3 and on formal curves approximating a point in a closure, Theorem 5.2.1. We also thank the referees for many valuable remarks about the original version of our paper.

Funding Open access funding provided by University of Bern.

Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article’s Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article’s Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

References

1. Arthur Bik. Strength and noetherianity for infinite tensors. PhD thesis, Universität Bern, 2020.
2. Arthur Bik, Jan Draisma, Rob H. Eggermont, and Andrew Snowden. The geometry of polynomial representations. *Int. Math. Res. Not.*, 2021. To appear, [arXiv:2105.12621](https://arxiv.org/abs/2105.12621).
3. Arthur Bik, Jan Draisma, Rob H. Eggermont, and Andrew Snowden. Uniformity for limits of tensors. 2023. Preprint, [arXiv:2305.19866](https://arxiv.org/abs/2305.19866).
4. Andries E. Brouwer and Jan Draisma. Equivariant Gröbner bases and the two-factor model. *Math. Comput.*, 80:1123–1133, 2011.
5. Daniel E. Cohen. Closure relations, Buchberger’s algorithm, and polynomials in infinitely many variables. In *Computation theory and logic*, volume 270 of *Lect. Notes Comput. Sci.*, pages 78–87, 1987.
6. David A. Cox, John Little, and Donal O’Shea. *Using algebraic geometry*, volume 185 of *Graduate Texts in Mathematics*. Springer, New York, second edition, 2005.

7. David A. Cox, John Little, and Donal O'Shea. *Ideals, varieties, and algorithms*. Undergraduate Texts in Mathematics. Springer, Cham, fourth edition, 2015. An introduction to computational algebraic geometry and commutative algebra. <https://doi.org/10.1007/978-3-319-16721-3>.
8. Harm Derksen, Rob H. Eggermont, and Andrew Snowden. Topological noetherianity for cubic polynomials. *Algebra Number Theory*, 11(9):2197–2212, 2017. <https://doi.org/10.2140/ant.2017.11.2197>.
9. Jan Draisma. Topological Noetherianity of polynomial functors. *J. Am. Math. Soc.*, 32:691–707, 2019.
10. Daniel Erman, Steven V. Sam, and Andrew Snowden. Big polynomial rings and Stillman's conjecture. *Invent. Math.*, 218(2):413–439, 2019. <https://doi.org/10.1007/s00222-019-00889-y>.
11. Eric M. Friedlander and Andrei Suslin. Cohomology of finite group schemes over a field. *Invent. Math.*, 127(2):209–270, 1997.
12. William Fulton and Joe Harris. *Representation Theory. A First Course*. Number 129 in Graduate Texts in Mathematics. Springer-Verlag, New York, 1991.
13. M. J. Greenberg. Rational points in Henselian discrete valuation rings. *Publ. Math., Inst. Hautes Étud. Sci.*, 31:563–568, 1966. <https://doi.org/10.1007/BF02684802>.
14. Gert-Martin Greuel and Gerhard Pfister. *A Singular introduction to commutative algebra*. Springer, Berlin, extended edition, 2008. With contributions by Olaf Bachmann, Christoph Lossen and Hans Schönemann.
15. Christopher J. Hillar, Robert Krone, and Anton Leykin. EquivariantGB Macaulay2 package. Webpage: <http://people.math.gatech.edu/~rkrone3/EquivariantGB.html>, 2013.
16. Christopher J. Hillar, Robert Krone, and Anton Leykin. Equivariant Gröbner bases. In The 50th anniversary of Gröbner bases. Proceedings of the 8th Mathematical Society of Japan-Seasonal Institute (MSJ-SI 2015), Osaka, Japan, July 1–10, 2015, pages 129–154. Tokyo: Mathematical Society of Japan (MSJ), 2018.
17. Christopher J. Hillar and Seth Sullivant. Finite Gröbner bases in infinite dimensional polynomial rings and applications. *Adv. Math.*, 221:1–25, 2012.
18. Teresa Krick and Alessandro Logar. An algorithm for the computation of the radical of an ideal in the ring of polynomials. In *Applied algebra, algebraic algorithms and error-correcting codes* (New Orleans, LA, 1991), volume 539 of *Lecture Notes in Comput. Sci.*, pages 195–205. Springer, Berlin, 1991. https://doi.org/10.1007/3-540-54522-0_108.
19. J.M. Landsberg and Giorgio Ottaviani. Equations for secant varieties of Veronese and other varieties. *Ann. Mat. Pura Appl. (4)*, 192(4):569–606, 2013.
20. Toby Ord. How to simulate everything (all at once). URL: <http://www.amirrorclear.net/academic/ideas/simulation/index.html>.
21. Claudio Procesi. *Lie Groups*. New York: Springer, 2007.
22. Claudiu Raicu. 3×3 minors of catalecticants. *Math. Res. Lett.*, 20(4):745–756, 2013. <https://doi.org/10.4310/MRL.2013.v20.n4.a10>.
23. Guillaume Rond. Artin approximation. *J. Singul.*, 17:108–192, 2018. <https://doi.org/10.5427/jsing.2018.17g>.

Publisher's Note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.