

Acute effects of isotonic eccentric exercise on the neuromuscular function of knee extensors vary according to the motor task: impact on muscle strength profiles,

Original

Acute effects of isotonic eccentric exercise on the neuromuscular function of knee extensors vary according to the motor task: impact on muscle strength profiles, proprioception and balance / Vila-Chã, Carolina; Bovolini, Antonio; Francisco, Cristiana; Costa-Brito, Ana R; Vaz, Cláudia; Rua-Alonso, María; de Paz, José Antonio; Vieira, Taian; Mendonca, Goncalo V. - In: FRONTIERS IN SPORTS AND ACTIVE LIVING. - ISSN 2624-9367. - ELETTRONICO. - 5:(2023). [10.3389/fspor.2023.1273152]

Availability:

This version is available at: 11583/2984776 since: 2023-12-29T17:47:25Z

Publisher:

Frontiers Media SA

Published

DOI:10.3389/fspor.2023.1273152

Terms of use:

This article is made available under terms and conditions as specified in the corresponding bibliographic description in the repository

Publisher copyright

(Article begins on next page)



OPEN ACCESS

EDITED BY

Yaodong Gu,
Ningbo University, China

REVIEWED BY

Xuanzhen Cen,
Óbuda University, Hungary
Yuqi He,
University of Pannonia, Hungary

*CORRESPONDENCE

Carolina Vila-Chã
✉ cvilacha@ipg.pt

RECEIVED 05 August 2023

ACCEPTED 18 October 2023

PUBLISHED 03 November 2023

CITATION

Vila-Chã C, Bovolini A, Francisco C,
Costa-Brito AR, Vaz C, Rua-Alonso M,
de Paz JA, Vieira T and Mendonça GV (2023)

Acute effects of isotonic eccentric exercise on the neuromuscular function of knee extensors vary according to the motor task: impact on muscle strength profiles, proprioception and balance.

Front. Sports Act. Living 5:1273152.

doi: 10.3389/fspor.2023.1273152

COPYRIGHT

© 2023 Vila-Chã, Bovolini, Francisco, Costa-Brito, Vaz, Rua-Alonso, de Paz, Vieira and Mendonça. This is an open-access article distributed under the terms of the [Creative Commons Attribution License \(CC BY\)](https://creativecommons.org/licenses/by/4.0/). The use, distribution or reproduction in other forums is permitted, provided the original author(s) and the copyright owner(s) are credited and that the original publication in this journal is cited, in accordance with accepted academic practice. No use, distribution or reproduction is permitted which does not comply with these terms.

Acute effects of isotonic eccentric exercise on the neuromuscular function of knee extensors vary according to the motor task: impact on muscle strength profiles, proprioception and balance

Carolina Vila-Chã^{1,2*}, Antonio Bovolini^{1,2}, Cristiana Francisco¹, Ana R. Costa-Brito¹, Cláudia Vaz¹, María Rua-Alonso^{1,2,3}, José Antonio de Paz⁴, Taian Vieira^{5,6} and Gonçalo V. Mendonça^{7,8}

¹Laboratory for the Assessment of Sports Performance, Physical Exercise and Health (Labmov), Polytechnic of Guarda, Guarda, Portugal, ²Research Center in Sports Sciences, Health Sciences, and Human Development, Vila Real, Portugal, ³Performance and Health Group, Department of Physical Education and Sport, Faculty of Sports Sciences and Physical Education, University of A Coruña, A Coruña, Spain, ⁴Institute of Biomedicine, University of León, León, Spain, ⁵Laboratorio di Ingegneria del Sistema Neuromuscolare (LISIN), Dipartimento di Elettronica e Telecomunicazioni, Politecnico di Torino, Turin, Italy, ⁶PoliToBIOMed Lab, Politecnico di Torino, Turin, Italy, ⁷Neuromuscular Research Laboratory, Faculdade de Motricidade Humana, Universidade de Lisboa, Lisbon, Portugal, ⁸Interdisciplinary Center for the Study of Human Performance (CIPER), Faculdade de Motricidade Humana, Universidade de Lisboa, Lisbon, Portugal

Introduction: Eccentric exercise has often been reported to result in muscle damage, limiting the muscle potential to produce force. However, understanding whether these adverse consequences extend to a broader, functional level is of apparently less concern. In this study, we address this issue by investigating the acute and delayed effects of supramaximal isotonic eccentric exercise on neuromuscular function and motor performance of knee extensors during tasks involving a range of strength profiles, proprioception, and balance.

Methods: Fifteen healthy volunteers (23.2 ± 2.9 years old) performed a unilateral isotonic eccentric exercise of the knee extensors of their dominant lower limb (4×10 reps at 120% of one Repetition Maximum (1RM)). The maximum voluntary isometric contraction (MVC), rate of force development (RFD), force steadiness of the knee extensors, as well as knee joint position sense and mediolateral (MLI) and anteroposterior stability (API) of the dominant lower limb, were measured pre-, immediately, and 24 h after the eccentric exercise. The EMG amplitude of the vastus medialis (VM) and biceps femoris (BF) were concomitantly evaluated.

Results: MVC decreased by 17.9% immediately after exercise ($P < 0.001$) and remained reduced by 13.6% 24 h following exercise ($P < 0.001$). Maximum RFD decreased by 20.4% immediately after exercise ($P < 0.001$) and remained reduced by 15.5% at 24 h ($P < 0.001$). During the MVC, EMG amplitude of the VM increased immediately after exercise while decreasing during the RFD task. Both values returned to baseline 24 h after exercise. Compared to baseline, force steadiness during submaximal isometric tasks reduced immediately after exercise, and it was accompanied by an increase in the EMG amplitude of the VM. MLI and knee joint position sense were impaired immediately after isotonic eccentric exercise ($P < 0.05$). While MLI returned to baseline values 24 h later, the absolute error in the knee repositioning task did not.

Discussion: Impairments in force production tasks, particularly during fast contractions and in the knee joint position sense, persisted 24 h after maximal isotonic eccentric training, revealing that neuromuscular functional outputs were affected by muscle fatigue and muscle damage. Conversely, force fluctuation and stability during the balance tasks were only affected by muscle fatigue since fully recovered was observed 24 h following isotonic eccentric exercise.

KEYWORDS

eccentric exercise, isotonic load, rate of force development, force steadiness, joint position sense, postural control

1. Introduction

In the last decades, eccentric exercise has gained a growing interest in sports and rehabilitation fields, leading to the development of a variety of training protocols through the manipulation of different training variables, including mechanical loads (1, 2). Eccentric exercise can be performed at constant angular velocity movement (isokinetic) or against a constant external load (isotonic), providing distinct mechanical loads as a stimulus, which might lead to specific neuromuscular adaptations. Most studies have used isokinetic mechanical loads to investigate the chronic and acute effects of eccentric training on different parameters of health and sports performance (3–5). However, isotonic loads are more commonly applied in the training field (2, 6). Isotonic eccentric training with external loads exceeding the load that subjects are able to lift for one repetition (1 RM load) has become popular among coaches, athletes, and fitness practitioners (6, 7). Several studies have shown that this training mode is more effective than isokinetic eccentric training in improving muscle strength, muscle architecture, and activation (5, 6, 8). The greater effectiveness of this type of training has been justified by the mechanical overload and the greater limb acceleration at the beginning of the isotonic contractions (6, 8). On the other hand, such mechanical stress can favor muscle damage and related delayed onset muscle soreness (DOMS) (3, 9, 10). Several studies have revealed that eccentric contractions are strongly associated with skeletal muscle impairments, consisting of structural disruption of sarcomeres, disturbed excitation–contraction coupling, and calcium signaling, leading to an inflammatory response (10–13). Moreover, these alterations are frequently accompanied by altered neural drive to the muscles (4, 14).

The neurophysiological changes induced by eccentric exercise can negatively impact neuromuscular function (3, 15, 16). Evidence has shown that isokinetic eccentric contractions of the knee extensors induce an acute and delayed decrease of different muscle force profiles, including maximal muscle strength (10, 13), rate of force development (4, 17), and force steadiness (18, 19). However, the magnitude of changes might vary depending on the muscle strength profile measured. Besides alterations in contractile properties, eccentric contractions can also disturb proprioception (20, 21). Significant disturbances in the joint position sense were found after maximal and sub-maximal eccentric exercise (22–24). Such impairments may contribute to an increased risk of injury and impaired postural control. Nevertheless, despite extensive

research on the effects of eccentric exercise on proprioception, no study has investigated its impact on postural balance, especially when isotonic eccentric exercise with supramaximal loads is applied.

The extent to which eccentric exercise and related DOMS affects neuromuscular integrity and motor performance may vary depending on the mechanical characteristics of the eccentric load (5, 15). Moreover, the performance of different motor tasks seems to rely on distinct neuromuscular pathways and neural drive delivered to specific muscle groups. As a result, the impact of eccentric exercise and DOMS on motor performance can vary depending on the task specificity. Certain motor tasks may be more sensitive to the effects of DOMS, leading to a longer recovery period before optimal performance is restored. On the other hand, some motor tasks may exhibit resilience to DOMS or may recover more rapidly. Therefore, it is essential to consider the specific demands of different motor tasks when designing eccentric exercise programs and determining appropriate recovery strategies. Thus, this study aimed to investigate the acute effects of supramaximal isotonic eccentric exercise on different force profiles of the knee extensors, specifically maximum strength, rate of force development and force steadiness. Additionally, we aimed to evaluate the impact of this eccentric exercise mode on knee proprioception and dynamic balance. Changes in agonist and antagonist activity during force and balance tasks were also examined. By comprehensively investigating these factors, our study seeks to enhance the understanding of the implications of isotonic eccentric exercise and DOMS on neuromuscular function and motor performance. It was postulated that the magnitude of changes in the motor performance induced by a supramaximal isotonic eccentric exercise would depend on the motor task specificity. Moreover, we also hypothesized that changes in motor performance would be accompanied by alterations in the activation of the agonist and antagonist muscles.

2. Methods

2.1. Participants

Fifteen volunteers participated in the study (nine men and six women, age 23.2 ± 2.8 years; height $169.3 \text{ cm} \pm 6.5 \text{ cm}$; weight $63.8 \pm 8.4 \text{ kg}$). All participants were practitioners of exercise, familiarized with strength training, and without any pathology in the dominant lower limb. The study was approved by the Polytechnic of Guarda Committee on Research Ethics and

performed according to the Declaration of Helsinki. All subjects gave written informed consent before undertaking testing and training.

2.2. Experimental design

The participants visited the laboratory 3 times: 48 h before the experimental sessions to get familiarized with the experimental setup and to estimate the 1RM of the knee extensor of the dominant lower limb (session 1); to perform the isotonic eccentric exercise session and assess the neuromuscular function before and immediately after exercise session (session 2) and 24 h after eccentric exercise for neuromuscular function reassessment and DOMS evaluation (session 3). All experimental measurements were assessed by following the same order: (i) dynamic balance; (ii) knee proprioception; (iii) maximum voluntary isometric contraction; (iv) explosive isometric contraction; and (v) force steadiness. The flow chart in **Figure 1** summarizes the experimental protocol.


2.3. Isotonic eccentric exercise protocol

In order to determine the exercise intensity of the isotonic eccentric exercise, the 1RM of the knee extensors of the dominant lower limb was estimated for all participants using a multiple repetition test protocol and a regular leg extension machine (1MTH081, Panatta SLR, Italy). After a 5-min warm-up in a cycle ergometer, familiarization trials were performed to ensure proper execution of the test. The assessment protocol followed the guidelines outlined by National Strength and Conditioning Association Guidelines (25). The 1RM was estimated based on the Lombardi linear regression equation (26). The estimated 1RM was then used to define the exercise intensity

of the isotonic eccentric protocol. After at least 48 h, the participants visited the laboratory to perform a single exercise session consisting of 4 sets of 10 repetitions at 120% of 1RM, with a recovery interval of 1 min and 30 s between sets. To guarantee proper technical execution and physical integrity of the participants, the following criteria were established: (i) participants were well seated in the machine, with their back well supported on the machine seat (support cushion adjusted individually); (ii) the participants were instructed to exert maximal force only during the descending phase of the movement (eccentric action) for 2 s; (iii) joint range of motion was established between 180° (complete extension) and 90° of knee flexion; (iv) the non-dominant lower limb remained stationary, in a neutral and supported position; and (v) the concentric phase of the movement was executed by two assistants.

2.4. Dynamic balance test

Unilateral dynamic balance of the dominant lower limb was measured with the Biodex Balance System (BBS) (model 945-300, United States of America). Participants were instructed to stand in their dominant lower limb on the BBS locked foot platform, keeping their body straight and the upper limbs lateral to the body and with the unsupported leg placed in a comfortable knee-flexed position without touching any surface. Then, the foot platform was released, and participants were asked to maintain an upright standing position for 20 s without contact with any other surfaces. During the test, the platform stability decreased progressively from level 8 to level 2. Each participant performed five to six trials with an interval in-between of 40 s. Deviations in the anteroposterior (x-axis) and mediolateral (y-axis) relative to the platform center were sampled at frequency of 20 Hz. Then, anteroposterior stability index (API), mediolateral stability index



(MLI) and global stability index (SI) were computed with the BBS software, as described by Biodex Medical Systems, Inc. (27). API and MLI represent the variance of foot platform displacement in degrees for motion in the sagittal and frontal plans, respectively (27). SI represents the variance of foot platform displacement in degrees, in all motions during a test. These indices were calculated using the degree of oscillation of the platform, in which low values indicated that the individual had good stability (28). For each participant the indexes corresponded to the average of the values obtained in the five to six trials. Electromyography (EMG) data of the vastus medialis (VM), and biceps femoris (BF) of the dominant leg were recorded while the participants were performing the test.

2.5. Knee proprioception test

The participants were seated in an elevated chair and blindfolded, with the knees approximately at 90° and the legs freely moving, without contact with any surface. Markers were placed on the external condyles of the femur (approximately 20 cm above the knee) and tibia (approximately 20 cm below the knee) for further knee angle position identification. The protocol consisted of positioning the non-dominant leg at an angle of 130° (manually measured with a goniometer), and subsequently participant was requested to move their dominant leg to match the position non-dominant leg as accurately as possible. Participants were allowed to practice the tasks and then repeated it three times. The trials were recorded with a video camera (Nikon, D3200, 1,920 × 1,080 pixels, 30 fps), positioned perpendicular to the subject, at two meters from the chair. The joint angle was then later assessed using the Kinovea program (version 0.8.15). For each trial the joint position error was assessed by computing: (i) absolute (shows the magnitude of the error); (ii) constant (indicates the direction of the error); and (iii) variable (indicates the response variability).

2.6. Muscle strength profiles

Maximal, explosive and submaximal isometric contractions of the knee extensors of the dominant lower limb were performed to assess the impact of the isotonic eccentric exercise on the muscle strength and motor control profiles. MVC was measured with the participants seated in an elevated chair (customized chair made by INEGI-UP), with their arms crossed in front of their chest and with the trunk and hips firmly strapped to the chair. The dominant leg was positioned at 90° of flexion and adjustable padded strap was placed around the tibiotarsal region, attached to a load cell (model load cell 614, SENSOR, United Kingdom), and fixed to the chair. The unassessed lower limb was positioned atop a box, ensuring it remained flexed, while exercising utmost caution to avoid applying any force. The participants were then instructed to progressively increase force against the padded strap and encouraged to exert their maximum within the first 5 s. Each participant performed 3 repetitions with 2 min of rest in between. The MVC value corresponded to the maximal force exerted in the three trials. Ten

minutes after the MVC test, and in the same previous position, the participants performed three explosive isometric contractions, and they were encouraged to exert their maximum force as fast as possible. A rest of 2 min was given between trials.

After 15 min of rest, the participants performed submaximal isometric contractions at 10%, 20%, 30%, 40%, and 60% MVC, keeping it as stable as possible for 10 s. The participants completed two trials for each submaximal isometric contraction at 10%, 20%, 30%, and 40% MVC, as well as one trial for the submaximal isometric contraction at 60% MVC. A two-minute recovery interval was given between each trial. Participants were provided with visual feedback of the force exerted, that was displaced in computer screen of 22 inches in from them. It was required to the participants to maintain the target force level within two error bars of 2% MVC, centered around the target force level. EMG signals from the VM and BF and force were concomitantly record.

2.7. Assessment of muscle pain

To confirm the presence of DOMS 24 h post-eccentric exercise, participants verbally rated their perceived pain on a scale from 0 (“no soreness”) to 10 (“worst soreness imaginable”). The subjects were asked to rate the average pain intensity in the quadriceps during their regular activities of daily living (e.g., descending stairs) since their last visit to the laboratory (over the past 24 h).

2.8. EMG and force recordings

Surface EMG signals were acquired from the VM and BF muscles during the dynamic balance and isometric tasks used to measure the different muscle strength profiles. Signals from the selected muscles were recorded with three pairs Ag-AgCl electrodes (Ambu Neuroline, Denmark; conductive area 28 mm²), placed as recommended by Hermens et al. (29). Before placement of the electrodes, the skin was shaved, lightly abraded, and cleansed with water. A ground electrode was placed around the ankle of the dominant lower limb. Surface EMG signals were amplified as bipolar derivations (EMG amplifier, OT Bioelettronica, Turin, Italy), band-pass filtered (−3 dB bandwidth, 10–500 Hz), sampled at 2048 samples/s, and converted to digital data by a 12-bit A/D converter board. The EMG electrodes remained in place during the exercise session and were used for all EMG measurements on the same day (before and immediately after training). At the end of the experiment, the electrodes were removed, and the electrode positions were marked on the skin to ensure that the electrodes were placed in the same location for the EMG recordings obtained 24 h after training. The force signals were measured with a load cell (load sensitivity = 0.0048 V/N; SENSOR, load cell 614, United Kingdom) and collected through an auxiliary channel of the EMG amplifier (OT Bioelettronica, Italy), sampled at 2048 samples/s, and converted to digital data by a 12-bit A/D converter board and recorded in the OT Biolab software (OT Bioelettronica, Italy), thus allowing the simultaneous collection of force and EMG signals.

2.9. EMG and force signal analysis

For the reference MVC, the average rectified value (ARV) was computed from a time interval of 250 ms centered at instant of the maximal force. During the explosive contractions, the ARV was computed from two-time intervals: (i) 70 ms prior to the onset of the contraction and (ii) 50 ms centered at the time instant of the maximal slope (19). The ARV values for each muscle and task were normalized by the respective muscle's ARV obtained during the MVC [(task ARV/ARV MVC) × 100]. During dynamic balance tests the percent of activation time of the VM and BF was determined by using the algorithm described by Vincent and Soile (30). The analog signals from the load cell were converted into force (N, Newton) and the maximum rate of force development (RFD) was computed as the maximum slope of the force-time curve. The onset of the contraction was defined as the time instant that force exceeded 3 times the standard deviation value observed at baseline. For each submaximal contraction, the coefficient of variation (CoV) of force was calculated over the entire duration of the contraction. The CoV of force was computed by dividing the standard deviation (SD) of the force signal by the mean force multiplied by 100 [(SD/mean force) × 100]. This measure expresses the absolute force variability as a fraction of the mean force exerted (31).

2.10. Statistical analysis

Statistical analyses were performed using the Statistical Package for Social Sciences (SPSS Version 24, IBM Corporation, Armonk, New York, USA) software. The normality of the dependent variables was confirmed using the Shapiro-Wilk test. The acute effects of the isotonic eccentric training on muscle force profile (MVC and maximum RFD), dynamic balance (GBI, API and MLI),

proprioception (absolute, constant and variable errors) and muscle activity were assessed with one-way repeated-measures ANOVA, with factor time (Pre-, Post-, and 24 h after isotonic eccentric exercise). During submaximal isometric contractions, changes in the CoV of the force and ARV of the VM and BF were assessed with a two-way repeated measures ANOVA with factor time and target force levels (10%, 20%, 30%, 40% and 60% of MVC). Pairwise comparisons were performed with the Bonferroni *post hoc* test when ANOVA was significant. Partial eta-squared (η^2) was used to calculate the effect size of the statistical results, which were classified as weak ($\eta^2 < 0.01$), medium ($\eta^2 0.01 < 0.06$) or high ($\eta^2 > 0.14$) (32). The significance level was set to $P < 0.05$. Results are reported as means ± SD.

3. Results

3.1. Pain perception

Twenty-four hours after isotonic eccentric exercise, the participants rated their perceived pain intensity as 3.6 ± 2.6 on a scale from 0 to 10.

3.2. Maximum voluntary contraction

The MVC of the knee extensors decreased immediately after (-17.9% ; $P < 0.001$) and 24 h after training session (-13.6% ; $P < 0.001$) compared to pre-exercise values (Figure 2A; main effect: $P < 0.001$, $\eta^2 = 0.671$). These alterations were accompanied by a decline in the ARV of the vastus medialis ($P < 0.040$, $\eta^2 = 0.238$; Table 1), while no changes were observed in the ARV of the biceps femoris ($P < 0.207$, $\eta^2 = 0.106$;




TABLE 1 Average rectified value (ARV) for the vastus medialis (VM) and biceps femoris (BF) obtained during the maximal voluntary contraction (MVC), the isometric explosive contractions at onset of the contraction and at time instant that maximum rate of force development (RFD) was reached and during submaximal isometric contractions at target force levels of 10%, 20%, 30%, 40% and 60% MVC obtained at baseline (pre), immediately post, and 24 h after isotonic eccentric exercise.

Variable	Vastus medialis			Biceps femoris		
	Pre	Post	24 h	Pre	Post	24 h
MVC [ARV (μV)]	367.72 ± 122.65	312.66 ± 110.25 [§]	347.92 ± 130.38	113.04 ± 63.50	120.61 ± 45.01	97.28 ± 25.21
RFD onset (% ARV MVC)	59.63 ± 23.70	79.80 ± 20.27 [§]	62.99 ± 24.92	71.99 ± 25.74	86.23 ± 34.92	76.70 ± 36.34
RFD max (% ARV MVC)	69.89 ± 15.33	92.50 ± 33.49	72.58 ± 33.49	69.78 ± 13.05	82.82 ± 24.88	80.01 ± 26.23
10% MVC (% ARV MVC)	12.53 ± 3.64	16.22 ± 5.29 ^{§,†}	11.93 ± 3.97	14.42 ± 4.52	17.13 ± 8.31	15.97 ± 5.25
20% MVC (% ARV MVC)	21.62 ± 6.24	29.59 ± 8.86 ^{§,‡}	20.52 ± 6.37	22.51 ± 4.95	28.27 ± 10.31	25.04 ± 7.84
30% MVC (% ARV MVC)	33.06 ± 6.93	41.50 ± 9.72 ^{§,†}	32.89 ± 10.59	35.30 ± 7.03	39.01 ± 12.43	36.92 ± 13.48
40% MVC (% ARV MVC)	46.03 ± 11.20	56.39 ± 10.68 ^{§,}	45.98 ± 12.05	50.39 ± 16.05	39.32 ± 8.11	41.91 ± 10.39
60% MVC (% ARV MVC)	48.97 ± 8.84	53.83 ± 7.65	49.62 ± 11.97	39.78 ± 8.11	41.91 ± 10.39	39.78 ± 11.27
Dynamic balance (% time activation)	46.62 ± 15.53	57.17 ± 13.86	49.82 ± 23.10	38.83 ± 16.13	45.67 ± 15.88	39.73 ± 21.80

[†]Post significantly different from 24 h ($P < 0.01$).

[‡]Post significantly different from pre-condition ($P < 0.01$).

[§]Post significantly different from pre-condition ($P < 0.05$).

^{||}Post significantly different from pre and after 24 h ($P < 0.05$).

Table 1). Following exercise session, the vastus medialis ARV significantly decreased ($P = 0.017$) returning to baseline values 24 h after ($P = 1.000$).

3.3. Rate of force development

As for MVC, the maximum RFD was substantially reduced immediately after the isotonic eccentric exercise (-20.3% ; $P < 0.001$), remaining diminished 24 h following exercise (-15.5% ; $P < 0.001$) when compared to pre-exercise values (**Figure 2B**; time effect: $P < 0.001$, $\eta^2 = 0.608$). The

time to reach maximum RFD did not differ between pre (179 ± 3.6 ms), post (183 ± 4.0 ms) and 24 h after eccentric exercise (174 ± 2.9 ms) (time effect: $P = 0.615$.) The EMG amplitude of the VM, at onset of the explosive isometric contraction (time effect: $P = 0.025$, $\eta^2 = 0.435$) and at the time instant that maximum RFD was reached (time effect: $P = 0.014$, $\eta^2 = 0.483$) increased immediately post ($P < 0.05$, for both time intervals) and returned to baseline 24 h after exercise session (**Table 1**). No alterations were observed in the ARV of the BF for both time intervals of the explosive contractions (**Table 1**, time effect: $P > 0.180$, $\eta^2 > 0.122$, in both time intervals).




FIGURE 3 Mean and SD of the coefficient of variation (CoV) of force during submaximal isometric contractions at target force levels of 10%, 20%, 30%, 40% and 60% MVC obtained at baseline (pre), immediately post, and 24 h after eccentric exercise. *Indicates that CoV post eccentric exercise is significantly different from the CoV measured before (pre) and 24 h after exercise ($P < 0.05$).

3.4. Force steadiness

Figure 3 shows the CoV of force at the different target force levels measured pre, immediately post and 24 h after isotonic eccentric exercise. Data analysis revealed a significant effect for both factors load ($P=0.004$, $\eta^2=0.289$) and time ($P<0.001$, $\eta^2=0.491$). CoV at 10% MVC was higher than the CoV observed at 20% and 30% of MVC ($P=0.028$ and $P=0.042$, respectively). At 60% MVC, there was a significantly higher CoV compared to the CoV at 20% and 30% ($P=0.020$ and $P=0.013$, respectively).

The pooled data showed that CoV significantly increased post exercise (+14.7%; $P=0.025$), returning to baseline values in the 24 h after (**Figure 3**). No interaction time \times load was observed ($P=0.343$, $\eta^2=0.075$). When stratified by force levels, the results revealed that only at 10% and 20% MVC, the CoV was significantly higher immediately after exercise ($P=0.014$), returning to baseline values 24 h after exercise. Changes in the force steadiness were accompanied by significant alterations in the ARV of the VM (time \times load effect: $P=0.041$, $\eta^2=0.166$; **Table 1**). The ARV of the VM increased immediately after exercise for at all force levels ($P<0.030$), except for the 60%MVC ($P=0.331$). Twenty-four hours later the ARV values returned to baseline (**Table 1**). No significant changes were observed in the EMG amplitude of the biceps femoris either immediately post or 24 h after the eccentric exercise (time effect: $P=0.441$, $\eta^2=0.057$; **Table 1**).

3.5. Proprioception

Figure 4 illustrates the absolute, constant, and variable errors during the knee repositioning task, before, immediately post, and 24 h after the isotonic eccentric session. The results showed that participants significantly increased the magnitude of the repositioning error following eccentric exercise (time effect: $P=0.002$, $\eta^2=0.358$; **Figure 4A**). The absolute error increased immediately after exercise ($+2.4 \pm 2.1^\circ$; $P=0.021$) and remained higher 24 h after

when compared to baseline ($+2.7 \pm 2.7^\circ$; $P=0.020$) (**Figure 4A**). After training, participants moved into a more extended knee position relative to the reference leg, as indicated by the constant error (time effect: $P=0.012$, $\eta^2=0.270$; **Figure 4B**). Immediately after the training session, the constant error increased by $+5.5 \pm 3.1^\circ$ when compared to baseline ($P=0.015$). Twenty-four hours later the constant error still increased, but no statistical differences were observed between pre and 24 h after training conditions ($P=0.133$; **Figure 4B**). The position consistency, as indicated by the variable error, although declined was not significantly affected by the present exercise protocol (time effect: $P=0.435$, $\eta^2=0.057$; **Figure 4C**).

3.6. Dynamic balance

Figure 5 shows the results on the global balance, anteroposterior and mediolateral stability indexes before, immediately post and 24 h after eccentric isotonic exercise. Global balance and anteroposterior indexes were not affected by the exercise protocol (time effect: $P=0.483$, $\eta^2=0.051$ and $P=0.685$, $\eta^2=0.025$, respectively; **Figures 5A,B**). Nonetheless, significant alterations were observed in the mediolateral direction (time effect: $P=0.002$, $\eta^2=0.399$; **Figure 5C**). The instability in the mediolateral direction significantly increased immediately post exercise ($P<0.001$). Following 24 h, the mediolateral index remained higher, but no statistical differences were observed when compared to baseline values ($P=0.092$) (**Figure 5C**).

Although there was an increase in the relative time of VM and BF activation during balance tests immediately after the eccentric training, the results were not statistically significant (time effect: $P>0.434$; $\eta^2<0.080$, for both muscles; **Table 1**).

4. Discussion

The present study showed that eccentric exercise with a supramaximal isotonic load caused DOMS and impaired motor output across different motor tasks. Force output assessed either





FIGURE 4

Mean and SD of the absolute joint position error (A), constant error (B) and variable error (C) during the knee repositioning task recorded at baseline (pre), immediately post and 24 h post eccentric exercise. *Indicates that absolute joint position error pre-eccentric exercise significantly lower than immediate post and 24 h after eccentric exercise ($P<0.05$). †Denotes a significant difference between post and pre constant error values ($P<0.05$).



by MVC, RFD or force steadiness declined immediately after eccentric training. Twenty-four hours later, MVC and RFD remained impaired, while isometric force steadiness, at different target levels, was recovered. These alterations were accompanied by adjustments in agonist muscle activity that followed a different time course and pattern depending on the motor task characteristics. Postural control and proprioception were also impaired by the isotonic eccentric exercise as well. The stability in the mediolateral direction decreased, and the error magnitude of knee positioning increased immediately after exercise and in the presence of DOMS. Collectively, these findings demonstrate that the magnitude and the time course of the impairments induced by eccentric exercise with a supramaximal isotonic vary depending on the motor task specificity.

4.1. Muscle soreness and muscle strength performance

The participants reported soreness in the quadriceps 24 h after training, confirming the presence of DOMS. The average soreness level was 3.6 out of 10, which is in agreement with studies on acute responses to isokinetic eccentric exercise (3, 19, 33). Both MVC and RFD of the knee extensors decreased immediately after isotonic eccentric exercise and remained impaired 24 h later. Nonetheless, the magnitude of decline in the RFD post and 24 h after exercise was greater than in the MVC (20.3% and 15.5% vs. 17.9% and 13.6%, respectively). Several studies have reported similar results, suggesting that RFD is more sensitive to muscle fatigue (34, 35) and muscle damage after eccentric exercise (15, 17, 19, 36). The observed decrease in force-generating capacity is a well-known consequence of performing unaccustomed and/or vigorous eccentric exercise. This type of exercise can lead to disruptions in cytoskeletal structures and microlesions in muscle fibers, thereby affecting the excitation-contraction coupling mechanisms (16, 37). Consequently, a long-lasting deficit in force production (> 24 h) is commonly observed (38, 39). The magnitude of the force production deficit and time

course of the recovery seems to be partially determined by the content and size of the muscle fibers type II (4). Several studies have shown that type II fibers are particularly susceptible to damage from intense eccentric exercise, mainly due to their higher tension generating capacity and shorter optimum length for tension (10, 40). Although we did not directly measure muscle fiber damage, it is very likely that the greater changes observed in the RFD are a consequence of subcellular perturbances within type II fibers. Fiber type is often considered a major factor influencing RFD, in which type II fibers play an important role in producing faster rates of tension development (9). Muscle activation is also a critical determinant of RFD, particularly in the earlier phases of the rising force during rapid contractions (41). Given the potential for muscle damage caused by eccentric exercise, neural adaptations may occur as an effort to overcome muscle impairments and generate the required muscle tension. In the current study, the EMG amplitude of the vastus medialis exhibited a significant increase at the onset of rapid contraction and around the time of maximum RFD immediately after eccentric exercise, while no changes were observed in the antagonist activity. Still, EMG amplitude of the vastus medialis returned to baseline levels within 24 h, even though RFD impairment persisted.

The findings indicate that changes in EMG patterns are strongly associated with neuromuscular fatigue, rather than the neuromuscular mechanisms that cause maximum RFD impairments from isotonic eccentric exercise. While several factors could potentially account for the immediate post-eccentric exercise increment in the EMG amplitude, it is highly plausible that these changes are a consequence of an increment of the recruitment and/or firing rate of the motor units (18, 42). Because at the contraction onset and at the instant of maximum RFD, force is submaximal, the neural drive can be increased to compensate the force impairments induced by muscle fatigue. In fact, increased EMG amplitude of the agonist muscle was also observed during the sustained submaximal contractions immediately post, but not in the 24 h after eccentric exercise. The presence of muscle fatigue is also indirectly confirmed by

the decline in EMG activity during the MVC. Contrary to submaximal efforts, during maximal contractions, the EMG amplitude typically decreases in the presence of muscle fatigue, indicating the limitations of the central nervous system to continually increase synaptic input to the motoneurons (43).

Peñailillo et al. (17) did not observe any change in the EMG amplitude of the vastus lateralis during explosive contractions following eccentric cycling protocols at 60% of maximal concentric contraction. Conversely, Vila-Chã et al. (19) reported a decline of the EMG amplitude of the vasti muscles at the contraction onset and at the time of maximum RFD, which persisted in the 24 h after a maximal isokinetic eccentric exercise protocol. The conflicting results might be explained by differences arising from variations in the modes of eccentric exercise employed across each study, that might have induced different magnitudes of fatigue and muscle damage. It is well known that both phenomena are dependent of modulation based on task-specific factors, which in turn govern the neurophysiological changes (44, 45). In the current study, the immediate decline in the maximum RFD and MVC after isotonic eccentric exercise is very likely explained by a combination of fatigue and initial muscle damage, leading to changes in muscle activation. However, the sustained alteration in RFD and MVC values observed 24 h after isotonic eccentric exercise appears to be predominantly stem from peripheral adaptations prompted by muscle damage, with negligible impact on the neural drive.

Another important parameter about muscle strength is the force steadiness measured during submaximal contractions. It has been shown that force steadiness during submaximal isometric contractions is moderately associated with performance on different motor tasks, such as standing balance and walking performance (31). Therefore, gaining insight into the influence of isotonic eccentric exercise on force fluctuation can yield valuable functional and physiological information regarding complementary indicators of function recovery after eccentric exercise. Such significance is accentuated by the fact that this parameter is predominantly influenced by neural factors that are not directly associated with MVC or RFD. The present study showed that force steadiness during different submaximal isometric levels significantly decreased immediately after eccentric exercise, but returned to baseline values in the 24 h after exercise. The EMG amplitude of the VM followed a similar pattern and no changes were observed in the activity of the antagonist muscle. The results are in line with previous studies that applied isokinetic eccentric protocols (19, 23, 46). In these studies, the force steadiness of the elbow flexors (23, 46) and knee extensors (19) decreased immediately after eccentric exercise for all submaximal levels (between 2.5% and 50% MVC). Twenty-four hours later the force steadiness remained impaired only during the submaximal contractions at very low intensities ($\leq 10\%$ MVC). The observed alterations were accompanied by an increased EMG amplitude of the agonist muscles, indicating the presence of low-frequency fatigue following eccentric exercise (46, 47). In the current study, the lowest intensity used was 10% MVC. Even for this very low intensity, the force steadiness of the

knee extensors was recovered 24 h after the eccentric exercise. The findings suggest that the CoV of force is a better indicator of muscle fatigue than potential muscle damage resulting from the isotonic eccentric protocol applied in this study.

4.2. Proprioception and balance

Immediately after isotonic eccentric exercise, an increase of the position errors was observed. The magnitude of the error (measured as absolute error) remained higher after 24 h after exercise, while direction of the error (constant error) and position consistency (variable error) returned to pre-exercise values. The results indicate that the participants matched the predetermined knee position by adopting a more extended position. Similar results have been obtained previously (20–22, 48–50). It is well known that a period of intense exercise increases errors in limb position sense (51). The disturbance in joint position sense seems to arise from the altered sense of effort due to the presence of fatigue, either after concentric or eccentric exercise (24, 52). Moreover, exercise effects on position sense are directly dependent on the level of muscle fatigue produced by the exercise (51). It has been shown that the accumulation of metabolites, stimulation of small muscle afferents in the group III and IV due to muscle fatigue might be involved in the sense of effort, perturbing the sense of position (51, 53). However, following eccentric exercise it was observed that position matching errors might persist in the subsequent days, where chemical products of fatigue were long gone (21, 24, 50). This suggests that factors beyond fatigue affected the position sense. In the past it has been hypothesized that eccentric exercise would cause also damage to muscle spindles, leading to perturbations in proprioception. Nonetheless, animal studies have shown that after intense eccentric exercise the responsiveness of muscle spindles is not disturbed despite extensive muscle damage (51, 54).

Tsay et al. (50) observed an association between decrease in force and increase position errors, suggesting that exercise-induced decline in force (of 20%–30%) can disrupt limb position sense. It has been suggested that the effects of exercise on proprioception resides on the operation of an internal feed-forward model (53, 55). This hypothesis indicates that the expected sensory feedback for a particular limb position is based on past memories and compares it with the actual sensory feedback from the fatigued limb. Based on previous experience, the sensory feedback from the fatigued muscle might be greater than anticipated from the motor command, altering the neural drive to the fatigued limb (23, 50). Conversely Da Silva et al. (48) observed a better association between position matching errors and voluntary activation deficits than with MVC decline. However, in their study, and despite a long-lasting MVC decrease and the presence of DOMS up to 48 h postexercise, position errors returned to baseline values within in the 24 h after eccentric exercise. In line with Tsay et al. (50) results, in the present study, the magnitude of errors increased immediately after eccentric

exercise and remained higher 24 h later, which accompanied the observed changes in MVC and RFD.

Notwithstanding the observed changes in position sense, during the balance task only alterations in the mediolateral stability index were observed. This parameter was increased immediately after isotonic eccentric exercise but returned to pre-exercise values 24 h after. The global and the anteroposterior stability index was not affected by the exercise protocol. Dabbs & Chander (56) investigated the effects of eccentric exercise on balance performance and despite the significant reductions in knee extensor torque up to at least 48 h, no changes were observed during the balance task. The lower sensitivity of the balance task to fatigue and muscle damage induced by the eccentric exercise might be explained by the fact that postural control is not only dependent of the proprioceptive and somatosensory system (57). The maintenance of upright balance is a complex task that involves afferent visual and vestibular systems that are not primarily affected by muscle fatigue or muscle damage (58).

Although this study provides insights into the complex effects of supramaximal isotonic eccentric exercise on neuromuscular function, motor performance, and proprioception, it is essential to acknowledge its specific limitations. Firstly, muscle damage induced by the isokinetic eccentric exercise was not directly measured, enabling further investigation of the relationship between muscle damage characteristics and the degree of motor performance impairments across a variety of motor tasks. The study focused on the 24 hours after exercise recovery period. However, extending the follow-up duration could shed light on the longer-term effects and the persistence of neuromuscular alterations induced by eccentric exercise. The evaluation of postural control during a balance task revealed a degree of insensitivity to fatigue and muscle damage in this study. To provide a more comprehensive understanding on the impact of the eccentric exercise on balance, a thorough analysis of additional balance-related parameters would be necessary. Lastly, given the highly specific nature of the eccentric exercise protocol used in this study, it is important to acknowledge that the results are inherently constrained by the confines of the defined experimental protocol. Consequently, the applicability of these findings to broader eccentric exercise regimens or different populations may be limited.

5. Conclusion

The supramaximal isotonic eccentric exercise affected differently the performed motor tasks. When compared to baseline, both, MVC and RFD were impaired immediately after exercise and in the subsequent 24 h. However, the magnitude of changes was more pronounced in the RFD than in the MVC knee extensors profile. Proprioception of knee position was also affected by the isotonic eccentric exercise and remained altered 24 hours after the exercise protocol. The results revealed that both RFD, MVC and position sense are affected by muscle fatigue and muscle damage. Conversely, force steadiness during

submaximal contractions and mediolateral stability index during the balance task were only impaired immediately after eccentric exercise, returning to baseline values within 24 h. The current study showed that isotonic eccentric exercise induces specific alterations on different neuromuscular functional outputs. The findings underscore the importance of understanding the specific alterations and recovery timelines associated with eccentric exercise for effective exercise and rehabilitation planning. However, future research needs to be performed to enhance understanding of these phenomena.

Data availability statement

The raw data supporting the conclusions of this article will be made available by the authors, without undue reservation, under request.

Ethics statement

The studies involving humans were approved by Polytechnic of Guarda Committee on Research Ethics. The studies were conducted in accordance with the local legislation and institutional requirements. The participants provided their written informed consent to participate in this study.

Author contributions

CV-C: Conceptualization, Formal Analysis, Methodology, Supervision, Writing – original draft, Writing – review & editing. AB: Conceptualization, Formal Analysis, Methodology, Writing – original draft, Writing – review & editing. CF: Conceptualization, Formal Analysis, Methodology, Writing – original draft. AC-B: Conceptualization, Formal Analysis, Project administration, Writing – review & editing. CV: Formal Analysis, Writing – original draft, Writing – review & editing. MR-A: Conceptualization, Formal Analysis, Methodology, Writing – original draft, Writing – review & editing. JP: Methodology, Writing – original draft, Writing – review & editing. TV: Conceptualization, Formal Analysis, Methodology, Writing – review & editing. GM: Conceptualization, Formal Analysis, Writing – original draft, Writing – review & editing.

Funding

The author(s) declare financial support was received for the research, authorship, and/or publication of this article.

This study was funded by the Portuguese Foundation for Science and Technology, IP, under project UID04045/2020. The Polytechnic of Guarda partly supported the research reported in this publication.

Acknowledgments

ARB and AB are part of the Highly Qualified Human Resources Project, reference number CENTRO-04-3559-FSE-000162. MR-A acknowledges the financial support received from the Spanish Ministry of Universities through the Grants for the Requalification of the Spanish University System under the Postdoctoral Margarita Salas Programme (RSUC.UDC.MS09), funded by the European Union – Next Generation.

Conflict of interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

References

- Lates AD, Greer BK, Wagle JP, Taber CB. Accentuated eccentric loading and cluster set configurations in the bench press. *J Strength Cond Res.* (2022) 36(6):1485–9. doi: 10.1519/JSC.0000000000003664
- Wagle JP, Taber CB, Cunanan AJ, Bingham GE, Carroll KM, DeWeese BH, et al. Accentuated eccentric loading for training and performance: a review. *Sports Med.* (2017) 47(12):2473–95. doi: 10.1007/s40279-017-0755-6
- Douglas J, Pearson S, Ross A, McGuigan M. Eccentric exercise: physiological characteristics and acute responses. *Sports Med.* (2017b) 47(4):663–75. doi: 10.1007/s40279-016-0624-8
- Farup J, Rahbek SK, Bjerre J, de Paoli F, Vissing K. Associated decrements in rate of force development and neural drive after maximal eccentric exercise. *Scand J Med Sci Sports.* (2016) 26(5):498–506. doi: 10.1111/sms.12481
- Guilhem G, Cornu C, Guével A. Neuromuscular and muscle-tendon system adaptations to isotonic and isokinetic eccentric exercise. *Ann Phys Rehabil Med.* (2010) 53(5):319–41. doi: 10.1016/j.rehab.2010.04.003
- Guilhem G, Cornu C, Maffiuletti NA, Guével A. Neuromuscular adaptations to isoload versus isokinetic eccentric resistance training. *Med Sci Sports Exerc.* (2013) 45(2):326–35. doi: 10.1249/MSS.0b013e31826e7066
- Douglas J, Pearson S, Ross A, McGuigan M. Chronic adaptations to eccentric training: a systematic review. *Sports Med.* (2017a) 47(5):917–41. doi: 10.1007/s40279-016-0628-4
- Vogt M, Hoppeler HH. Eccentric exercise: mechanisms and effects when used as training regime or training adjunct. *J Appl Physiol.* (2014) 116(11):1446–54. doi: 10.1152/jappphysiol.00146.2013
- Harridge SDR, Bottinelli R, Canepari M, Pellegrino MA, Reggiani C, Esbjörnsson M, et al. Whole-muscle and single-fibre contractile properties and myosin heavy chain isoforms in humans. *Pflug Arch Eur J Phy.* (1996) 432(5):913–20. doi: 10.1007/s004240050215
- Jones DA, Newham DJ, Round JM, Tolfree SE. Experimental human muscle damage: morphological changes in relation to other indices of damage. *J Physiol.* (1986) 375(1):435–48. doi: 10.1113/jphysiol.1986.sp016126
- Allen DG, Lamb GD, Westerblad H. Skeletal muscle fatigue: cellular mechanisms. *Physiol Rev.* (2008) 88(1):287–332. doi: 10.1152/physrev.00015.2007
- Baumert P, Lake MJ, Stewart CE, Drust B, Erskine RM. Genetic variation and exercise-induced muscle damage: implications for athletic performance, injury and ageing. *Eur J Appl Physiol.* (2016) 116(9):1595–625. doi: 10.1007/S00421-016-3411-1
- Friden J, Sjöström M, Ekblom B. Myofibrillar damage following intense eccentric exercise in man. *Int J Sports Med.* (1983) 4(3):170–6. doi: 10.1055/s-2008-1026030
- Goodall S, Thomas K, Barwood M, Keane K, Gonzalez JT, St Clair Gibson A, et al. Neuromuscular changes and the rapid adaptation following a bout of damaging eccentric exercise. *Acta Physiol.* (2017) 220(4):486–500. doi: 10.1111/apha.12844
- Aleman JA, Delgado-Díaz DC, Mathews H, Davis JM, Kostek MC. Comparison of acute responses to isotonic or isokinetic eccentric muscle action: differential outcomes in skeletal muscle damage and implications for rehabilitation. *Int J Sports Med.* (2014) 35(1):1–7. doi: 10.1055/s-0032-1327652
- Proske U, Allen TJ. Damage to skeletal muscle from eccentric exercise. *Exerc Sport Sci Rev.* (2005) 33(2):98–104. doi: 10.1097/00003677-200504000-00007
- Peñailillo L, Blazeovich A, Numazawa H, Nosaka K. Rate of force development as a measure of muscle damage. *Scand J Med Sci Sports.* (2015) 25(3):417–27. doi: 10.1111/SMS.12241
- Dartnall TJ, Rogasch NC, Nordstrom MA, Semmler JG. Eccentric muscle damage has variable effects on motor unit recruitment thresholds and discharge patterns in elbow flexor muscles. *J Neurophysiol.* (2009) 102(1):413–23. doi: 10.1152/jn.91285.2008
- Vila-Chã C, Hassanlouei H, Farina D, Falla D. Eccentric exercise and delayed onset muscle soreness of the quadriceps induce adjustments in agonist-antagonist activity, which are dependent on the motor task. *Exp Brain Res.* (2012) 216(3):385–95. doi: 10.1007/s00221-011-2942-2
- Torres R, Vasques J, Duarte JA, Cabri JMH. Knee proprioception after exercise-induced muscle damage. *Int J Sports Med.* (2010) 31(06):410–5. doi: 10.1055/s-0030-1248285
- Vila-Chã C, Riis S, Lund D, Møller A, Farina D, Falla D. Effect of unaccustomed eccentric exercise on proprioception of the knee in weight and non-weight bearing tasks. *J Electromyogr Kinesiol.* (2011) 21(1):141–7. doi: 10.1016/j.jelekin.2010.10.001
- Givoni NJ, Pham T, Allen TJ, Proske U. The effect of quadriceps muscle fatigue on position matching at the knee. *J Physiol.* (2007) 584(1):111–9. doi: 10.1113/jphysiol.2007.134411
- Semmler JG, Tucker KJ, Allen TJ, Proske U. Eccentric exercise increases EMG amplitude and force fluctuations during submaximal contractions of elbow flexor muscles. *J Appl Physiol.* (2007) 103(3):979–89. doi: 10.1152/jappphysiol.01310.2006
- Walsh LD, Allen TJ, Gandevia SC, Proske U. Effect of eccentric exercise on position sense at the human forearm in different postures. *J Appl Physiol.* (2006) 100(4):1109–16. doi: 10.1152/jappphysiol.01303.2005
- Haff GG, Triplett NT. *Essentials of strength training and conditioning 4th edition.* Champaign, IL: Human kinetics (2015).
- Lombardi VP. *Beginning weight training: The safe and effective way.* Dubuque: Wm. C. Brown Company Publishers (1989).
- Biodex Medical Systems, I. Balance System SD Operation/Service Manual (Shirley, N, Vol. 6339). www.biodex.com (2011).
- Cachupe WJC, Shifflett B, Kahanov L, Wughalter EH. Reliability of biodex balance system measures. *Meas Phys Educ Exerc Sci.* (2001) 5(2):97–108. doi: 10.1207/S15327841MPEE0502_3
- Hermens H, Freriks B, Merletti R, Stegeman DF, Blok J, Rau G, et al. *European recommendations for surface electromyography. Results of the SENIAM project. Roessingh Research and Development.* Enschede: Roessingh Research and Development (1999).
- Vincent L, Soille P. Watersheds in digital spaces: an efficient algorithm based on immersion simulations. *IEEE Trans Pattern Anal Mach Intell.* (1991) 13(06):583–98. doi: 10.1109/34.87344
- Enoka RM, Farina D. Force steadiness: from motor units to voluntary actions. *Physiology.* (2021) 36(2):114–30. doi: 10.1152/physiol.00027.2020
- Cohen J. Statistical power analysis for the behavioral sciences. (1988):18–74. doi: 10.4324/9780203771587

The author(s) declared that they were an editorial board member of Frontiers, at the time of submission. This had no impact on the peer review process and the final decision.

Publisher's note

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article, or claim that may be made by its manufacturer, is not guaranteed or endorsed by the publisher.

33. Hedayatpour N, Hassanlouei H, Arendt-Nielsen L, Kersting UG, Falla D. Delayed-onset muscle soreness alters the response to postural perturbations. *Med Sci Sports Exerc.* (2011) 43(6):1010–6. doi: 10.1249/MSS.0b013e3182058628
34. Andersen LL, Andersen JL, Zebis MK, Aagaard P. Early and late rate of force development: differential adaptive responses to resistance training? *Scand J Med Sci Sports.* (2010) 20(1):e162–9. doi: 10.1111/j.1600-0838.2009.00933.x
35. D'Emanuele S, Maffiuletti NA, Tarperi C, Rainoldi A, Schena F, Boccia G. Rate of force development as an indicator of neuromuscular fatigue: a scoping review. *Front Hum Neurosci.* (2021) 15:701916. doi: 10.3389/fnhum.2021.701916
36. Cramer RM, Aagaard P, Qvortrup K, Langberg H, Olesen J, Kjaer M. Myofibre damage in human skeletal muscle: effects of electrical stimulation versus voluntary contraction. *J Physiol.* (2007) 583(1):365–80. doi: 10.1113/jphysiol.2007.128827
37. Prasartwuth O, Taylor JL, Gandevia SC. Maximal force, voluntary activation and muscle soreness after eccentric damage to human elbow flexor muscles. *J Physiol.* (2005) 567(1):337–48. doi: 10.1113/jphysiol.2005.087767
38. Clarkson PM, Nosaka K, Braun B. Muscle function after exercise-induced muscle damage and rapid adaptation. *Med Sci Sports Exerc.* (1992) 24(5):512–20.
39. Butterfield TA. Eccentric exercise in vivo: strain-induced muscle damage and adaptation in a stable system. *Exerc Sport Sci Rev.* (2010) 38(2):51–60. doi: 10.1097/JES.0b013e3181d496eb
40. Vijayan K, Thompson JL, Norenberg KM, Fitts RH, Riley DA. Fiber-type susceptibility to eccentric contraction-induced damage of hindlimb-unloaded rat AL muscles. *J Appl Physiol.* (2001) 90(3):770–6. doi: 10.1152/JAPPL.2001.90.3.770
41. Folland JP, Buckthorpe MW, Hannah R. Human capacity for explosive force production: neural and contractile determinants. *Scand J Med Sci Sports.* (2014) 24(6):894–906. doi: 10.1111/sms.12131
42. Proske U, Morgan DL. Muscle damage from eccentric exercise: mechanism, mechanical signs, adaptations and clinical applications. *J Physiol.* (2001) 537(2):333–45. doi: 10.1111/j.1469-7793.2001.00333.x
43. Gandevia SC. Spinal and supraspinal factors in human muscle fatigue. *Physiol Rev.* (2001) 81(4):1725–89. doi: 10.1152/physrev.2001.81.4.1725
44. Enoka RM, Stuart DG. Neurobiology of muscle fatigue. *J Appl Physiol.* (1992) 72(5):1631–48. doi: 10.1152/jap.1992.72.5.1631
45. Prasartwuth O, Allen TJ, Butler JE, Gandevia SC, Taylor JL, Prasartwuth O. Length-dependent changes in voluntary activation, maximum voluntary torque and twitch responses after eccentric damage in humans. *J Physiol.* (2006) 571(Pt 1):243–52. doi: 10.1113/jphysiol.2005.101600
46. Dartnall TJ, Nordstrom MA, Semmler JG. Motor unit synchronization is increased in Biceps brachii after exercise-induced damage to elbow flexor muscles. *J Neurophysiol.* (2008) 99(2):1008–19. doi: 10.1152/jn.00686.2007
47. Dundon JM, Cirillo J, Semmler JG. Low-frequency fatigue and neuromuscular performance after exercise-induced damage to elbow flexor muscles. *J Appl Physiol.* (2008) 105(4):1146–55. doi: 10.1152/jappphysiol.01339.2007
48. Da Silva F, Monjo F, Zghal F, Chorin F, Guérin O, Colson SS. Altered position sense after submaximal eccentric exercise-inducing central fatigue. *Med Sci Sports Exerc.* (2021) 53(1):218–27. doi: 10.1249/MSS.0000000000002444
49. Paschalis V, Nikolaidis MG, Giakas G, Jamurtas AZ, Pappas A, Koutedakis Y. The effect of eccentric exercise on position sense and joint reaction angle of the lower limbs. *Muscle Nerve.* (2007) 35(4):496–503. doi: 10.1002/mus.20723
50. Tsay A, Allen TJ, Leung M, Proske U. The fall in force after exercise disturbs position sense at the human forearm. *Exp Brain Res.* (2012) 222(4):415–25. doi: 10.1007/s00221-012-3228-z
51. Proske U. Exercise, fatigue and proprioception: a retrospective. *Exp Brain Res.* (2019) 237(10):2447–59. doi: 10.1007/s00221-019-05634-8
52. Brockett C, Warren N, Gregory JE, Morgan DL, Proske U. A comparison of the effects of concentric versus eccentric exercise on force and position sense at the human elbow joint. *Brain Res.* (1997) 771(2):251–8. doi: 10.1016/S0006-8993(97)00808-1
53. Allen TJ, Ansems GE, Proske U. Effects of muscle conditioning on position sense at the human forearm during loading or fatigue of elbow flexors and the role of the sense of effort. *J Physiol.* (2007) 580(2):423–34. doi: 10.1113/jphysiol.2006.125161
54. Gregory JE, Morgan DL, Proske U. Response of muscle spindles following a series of contractions. *Exp Brain Res.* (2004) 157(2):234–40. doi: 10.1007/S00221-004-1838-9
55. Bays PM, Wolpert DM. Computational principles of sensorimotor control that minimize uncertainty and variability. *J Physiol.* (2007) 578(2):387–96. doi: 10.1113/jphysiol.2006.120121
56. Dabbs NC, Chander H. The effects of exercise induced muscle damage on knee joint torque and balance performance. *Sports (Basel).* (2018) 6(3):101. doi: 10.3390/sports6030101
57. Deliagina TG, Zelenin PV, Orlovsky GN. Physiological and circuit mechanisms of postural control. *Curr Opin Neurobiol.* (2012) 22(4):646–52. doi: 10.1016/j.conb.2012.03.002
58. Caron O. Effects of local fatigue of the lower limbs on postural control and postural stability in standing posture. *Neurosci Lett.* (2003) 340(2):83–6. doi: 10.1016/S0304-3940(02)01455-6